

## Week 11: Linear mixed models & autocorrelation

ANTH 674: Research Design & Analysis in Anthropology

Professor Andrew Du

[Andrew.Du2@colostate.edu](mailto:Andrew.Du2@colostate.edu)

Office Hours: Thursdays, 9:00am–12:00pm

In person: GSB 312

Virtual: <https://tinyurl.com/F22ANTH674>

1

## Lecture outline

1. Linear mixed models
  1. What are they? What are they used for?
  2. Variance partitioning w/ crossed or nested designs
2. Measuring & dealing with autocorrelation
  1. Correlograms
  2. Association between two variables
    1. Cross-correlograms
    2. Generalized least squares

2

## Linear mixed models

What are they? What are they used for?

3

## What is a linear mixed model?


- A linear model with **fixed** AND **random** factors/effects
- I've always found the distinction between the two factors confusing, so let's go through them carefully



4

## Fixed factors


- Measured levels are the only ones of interest
- E.g., want to know how intercept of growth rate ~ temperature changes w/ **different levels of nitrogen** (i.e., ANCOVA)
- Thus far, we have only looked at fixed factors



5

## Random factors


- Measured levels are a random, representative sample of some population of levels
- E.g., want to infer something about population of intercepts (i.e., SD of intercepts across levels)



6

## Ultimately, depends on research question!

- **Fixed factor:** estimate how intercept and/or slope changes ~ different N levels (i.e., ANCOVA)
- **Random factor:** estimate SD of population of intercepts and/or slopes across N levels




7

## For example...




- Study measured reaction times of 18 subjects each day after full night's sleep & then 9 days of sleep deprivation (180 total observations)



8

## Fixed factor approach

- Estimates intercept & slope of reaction ~ days for each subject (factor level)
- ANCOVA**: `lm(reaction ~ days * subject)`



9

## Random factor approach

- Estimates SD of intercepts & slopes across levels **ALONG WITH** intercept & slope fixed effects (via maximum likelihood)

Random effects:

|             | StdDev    | Corr   |
|-------------|-----------|--------|
| (Intercept) | 24.740241 | (Intr) |
| Days        | 5.922103  | 0.066  |
| Residual    | 25.591843 |        |

Fixed effects: Reaction ~ Days


|             | Value     | Std.Error | DF  | t-value  | p-value |
|-------------|-----------|-----------|-----|----------|---------|
| (Intercept) | 251.40510 | 6.824516  | 161 | 36.83853 | 0       |
| Days        | 10.46729  | 1.545783  | 161 | 6.77151  | 0       |

Can only calculate P-values in simple mixed models (e.g., balanced design, no crossed random effects)

10

## Random factor approach


- Can get intercept & slope for each subject (**NOT** estimated coefficients!)



11


## Random factor approach

- Can get intercept & slope for each subject (**NOT** estimated coefficients!)




12

## Compared to ANCOVA



13



14

## Shrinkage or regularization

- In ANCOVA, each line is fit to data for one factor level **only**
- In LMM, levels are drawn from a population, so they are expected to share characteristics (w/ a distribution mean = fixed effect)
  - This is how you can predict for new factor levels w/ LMM (can't do this w/ ANCOVA)
- If a level has few observations, it borrows more information from other levels and is pulled more towards the average (fixed effect)

15

## Fixed vs. random factors

### • Fixed factors

- Interested in how intercept and slope changes **independently for each subject**
- Potentially need to estimate **A LOT** of coefficients

### • Random factors

- Interested in the SD of intercepts and slopes **across all possible subjects**
- Estimates fixed effects AND SDs
- Shrinks intercepts & slopes towards means

<https://dynamicecology.wordpress.com/2015/11/04/is-it-a-fixed-or-random-effect/>

16


## Questions?



17

## Reasons for doing LMM

1. Factor levels are random sample from larger population → interested in estimating SD of population of intercepts/slopes
2. Ratio of data points to # coefficients in ANCOVA too small → overfitting (only need to estimate SDs in LMM)



36 estimated  
coefficients for  
180 data points!

18

## Reasons for doing LMM


1. Factor levels are random sample from larger population → interested in estimating SD of population of intercepts/slopes
2. Ratio of data points to # coefficients in ANCOVA too small → overfitting (only need to estimate SDs in LMM)
3. Shrink intercepts and slopes towards mean
4. Want to account for non-independence in data using a blocking factor (very popular reason)

19

## E.g., sleep study




- Each subject is treated as a block, each w/ treatments of different days of sleep deprivation (AKA **repeated measures design**)



20

## Analyzing block design

- Can do ANCOVA, but problematic if lots of levels & not enough data
- Also, many times blocks are not of interest (i.e., don't care how intercept/slope change w/ block)
- **Can use LMM & include block as a random factor to account for it**



21

## Random factor rules

- Random factor **MUST** be categorical (i.e., it must be a factor w/ levels)
  - A continuous IV can be entered into model as covariate (only one extra estimated coefficient)
- Need at least five levels (would you calculate SD of a sample w/ < 5 data points?)
  - Otherwise, just make factor a fixed effect

22

## Generalized linear mixed models (GLMMs)

- Mixed model w/ non-normally distributed errors (cf. generalized linear models)
- E.g., presence/absence of lions ~ wildebeest population density w/ plot as blocking factor




23

## Questions?



24



25

## Variance partitioning


- Can partition DV variance **not accounted for by fixed effects** among multiple random factors
- Or, can fit an intercept-only model to partition DV variance among random factors



26


## Multiple random factors

1. **Crossed random factors:** each level from a factor is w/ each level from the other
- E.g., how penicillin inhibiting growth of *Bacillus* varies across plates & penicillin samples



27

## Crossed random factors



- Fully crossed! Every combination of sample and plate is represented
- Unreplicated and completely balanced (only one observation per sample/plate combo)

28




## Crossed random factors

- Intercept-only LMM w/ plate & sample as crossed random effects

- Random effects:

| Groups   | Name        | Variance |
|----------|-------------|----------|
| plate    | (Intercept) | 0.7169   |
| sample   | (Intercept) | 3.7311   |
| Residual |             | 0.3024   |




29

## Multiple random factors


2. **Nested random factors:** factors are nested within each other

- E.g., strength of chemical paste, where two samples were taken from each cask from each batch



30

## Nested random factors



- 10 batches (capital letters)
- 3 casks per batch (lower case)
- Two samples per cask


31

## Nested random factors

- Intercept-only LMM w/ cask nested within batch

- Random effects:

| Groups   | Name        | Variance |
|----------|-------------|----------|
| cask     | (Intercept) | 8.434    |
| batch    | (Intercept) | 1.657    |
| Residual |             | 0.678    |



32



## Questions?



33


Average Potential ET during March-August (mm)\_Quantile

## Measuring & dealing w/ autocorrelation

34

## Non-independent data

- Data across space, time, or a phylogeny will almost always be non-independent
- Data closer in space/time/relatedness will be more similar




35

## Three components

- Patterns in space & time (& phylogeny) can be decomposed into three components
- $z(d) = \mu(d) + \eta(d) + \varepsilon$ , where  $(d)$  means component is a function of distance
  - i.e., pattern = trend + autocorrelated (red) noise + independent (white) noise
- Temporal data can also include a cyclical component too (e.g., monthly, seasonality, orbital cycles)


36

## Simulated example for time



37

## Simulated example for space



38

## Autocorrelated noise: nuisance or interesting?


- **Nuisance:** non-independence violates one of the assumptions of linear models (increases Type I error rate)
  - Must correct for this (e.g., w/ blocking)!
- **Interesting:** How does strength of autocorrelation vary as a function of distance in space/time/phylogeny?
  - May tell us something about the underlying process generating autocorrelation!

39

## Questions?



40




41

## Detrending

- If temporal or spatial data are strongly trended, need to first remove trend (i.e., **detrending**) to focus on autocorrelated noise
- 1. Can fit a trend to data and take the residuals
- 2. Calculate first differences:  $x'_i = x_{i+1} - x_i$

42


## Example w/ temporal data



43

## Autocorrelation function


- Quantifies how time series is correlated with itself at different lags
- E.g., `c(1, 2, 3, 4, 5)`
  - time lag = 1, `c(1, 2, 3, 4)` vs. `c(2, 3, 4, 5)`
  - time lag = 2, `c(1, 2, 3)` vs. `c(3, 4, 5)`



44

## A real example


- Quarterly approval rating of US president from 1945 to 1974
- Not strongly trended, so no need to detrend
- Can begin to hypothesize what's generating AC!



45

## Spatial correlogram

- Quantifies correlation between pairs of points as you increase the distance (lag) between them



46


## Questions?



47

## Measures of association

Cross-correlograms & generalized least squares



48


## Comparing two variables

1. **Cross-correlograms**: looks at how two variables are correlated at different lags
  - Perhaps one variable responds to another only after a certain amount of time/distance
  - As w/ correlation, symmetry between variables
2. **Generalized least squares**: models a DV as a function of one or more IVs, while explicitly modeling errors as non-independent
  - Good if variables are obviously DV or IV, or if you want to do prediction (as w/ linear models)

49

## Cross-correlation function


- E.g., climate variability vs. mammal extinction rates in Plio-Pleistocene eastern Africa



50

## Spatial cross-correlogram

- E.g., soil organic matter & cadmium level in floodplain of Meuse River, Netherlands



51


## Generalized least squares

- Relaxes GLM assumption of independent errors
- GLS estimates extra parameters to model non-independent errors
- Parameters estimated using maximum likelihood

52


## Temporal data

- AR1 autocorrelation error structure is popular
- Correlation between errors one time step apart estimated as  $\rho$  (usually positive), two steps apart is  $\rho^2$ , three steps apart is  $\rho^3$ , etc.



53

## E.g., Hawaiian gallinule





54

## `sqrt(gallinule)~Rainfall+Year`

Coefficients are very similar!


| lm coefficients: |            |          | GLS coefficients: |           |         |
|------------------|------------|----------|-------------------|-----------|---------|
|                  | Estimate   | Pr(> t ) |                   | Value     | p-value |
| (Intercept)      | -4.777e+02 | 1.26e-10 | (Intercept)       | -436.4326 | 0.0030  |
| Rainfall         | 8.604e-04  | 0.986    | Rainfall          | -0.0098   | 0.7649  |
| Year             | 2.450e-01  | 8.15e-11 | Year              | 0.2241    | 0.0026  |



55

## Spatial data

- Non-independence between errors modeled as a function of distance between them




- Five common models
- Different parameterizations will slightly change shape of each model

56



## One last note

- If an IV captures autocorrelation in DV, then there will be no autocorrelation remaining in residuals (GLM assumptions not violated!)



57

## Questions?



58

## Summary

- Use LMMs (random effects) if:
  1. Factor levels are random, representative samples from some larger population and are interested in estimating SD of population of intercepts/slopes
  2. Have too few data (prevent overfitting w/ ANCOVA)
  3. Want to account for non-independence w/ blocking (and have too few data)
  4. Interested in shrinkage of estimates
  5. You want to partition variance in DV
- Correlograms quantify autocorrelation structure, which is interesting in itself!
- Cross-correlograms & GLS quantify association btw variables in time/space/phylogeny (but remember, OLS estimates are unbiased!)

59

60



## Statistics vignette

Let's play a game...


### Rules

1. Pick an outcome of three coin flips, e.g., HHH
2. I pick my own outcome
3. If your outcome appears first in  $n$  flips, you win

61

## What's the secret?

- This is a game called "Penney's game"



62

## Why does it work? (an example)

You: HHH

Me: THH

You win only if HHH comes up in first three flips:  
HHH...

Otherwise, T *must* precede any HHH, and I win:  
...THHH...

<https://www.youtube.com/watch?v=Sa9jLWkrX0c>

63