

Week 9: Modern regression models

ANTH 674: Research Design & Analysis in Anthropology

Professor Andrew Du

Andrew.Du2@colostate.edu

Office Hours: Thursdays, 9:00am–12:00pm

In person: GSB 312

Virtual: <https://tinyurl.com/F22ANTH674>

1

Lecture outline

1. What are modern regression models?
2. The models we'll go over:
 - Model II regression
 - Segmented regression
 - Non-linear least squares
 - Locally estimated scatterplot smoothing

2


What are modern regression models?

*Just a “wastebasket” phrase to describe newer regression methods

3

Quick review: general linear models (GLM)

- Models continuous DV as a linear/additive function of one or more IVs
- $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 \dots + \beta_n X_n + \varepsilon$
- IVs can be continuous or categorical



4

Review: GLM assumptions

1. Relationship between DV and IVs is linear
2. IVs measured without error
3. Error terms have mean = 0 and are normally distributed
4. Error terms drawn from population with the same variance (homoskedasticity)
5. Error terms are independent

5

Relaxing GLM assumptions

1. Relationship between DV and IVs is linear
 - Segmented regression
 - Non-linear least squares
 - Locally estimated scatterplot smoothing
2. IVs measured without error
 - Model II regression

6

Questions?




7



8

GLM: IV measured w/o error

- Makes most sense in experiments where IV is controlled/manipulated



El Omari et al., 2016

9

GLM: IV measured w/o error

- Makes most sense in experiments where IV is controlled/manipulated
- What if this is not the case & IV varies due to unmeasured factors (i.e., “has error”)?
- E.g., Measuring human weight ~ height
- Or unclear which variable is IV or DV
- E.g., Petal length ~ width
- Need Model II regression methods


10

<https://cran.r-project.org/web/packages/lmodel2/vignettes/mod2user.pdf>

Model II regression


- Minimizes errors in BOTH variables to estimate regression intercept & slope
- Can hypothesis test (i.e., 95% CIs & P-values on coefficients), but cannot predict

Model I regression
(least squares)




Model II regression

Major axis regression (MA)



Standard major axis regression (SMA)




11

Standard major axis regression

- MA assumes X & Y in same units
- SMA is MA done on standardized X & Y
- SMA called reduced major axis regression (RMA) in bio. anth. (and other fields)
- Slope = $\pm SD_y / SD_x$ (sign determined by corr., r)
- Intercept = $\bar{y} - \bar{x} \times \text{slope}$ (as in OLS)


- RMA slope also = OLS slope / $|r|$
- So, RMA slope magnitude \geq OLS slope magnitude



12

Be careful!

- Even if no relationship between two variables, RMA will plot a positive or negative fitted line (b/c slope = $\pm SD_y/SD_x$)



"If r is not significantly different from zero, [RMA slope] should not be computed."

-Legendre & Legendre 2012:550

13

Final Model II thoughts

- I find RMA slope difficult to interpret: what does SD_y/SD_x mean?
- Perhaps Model II best for comparing coefficient estimate(s) to value(s) expected from theory
- If error in DV is much greater than that in IV, okay to use OLS (Model I)
- I never use Model II regressions

14


Questions?



15

Segmented regression


AKA piecewise regression
AKA breakpoint regression
AKA change point regression



16

When it's used


- When relationship between DV & IV can be described w/ ≥ 2 linear segments
 - E.g., threshold effect as IV increases
 - E.g., change in relationship w/ time
- Regression segments separated by breakpoints



17

How it works


- Need to specify number of breakpoints (n)
 - Changing intercepts only $\rightarrow 2n + 1$ coefficients
 - Changing intercepts & slopes $\rightarrow 2n + 2$ coeff.
- Coefficients estimated algorithmically using maximum likelihood estimation
- Can hypothesis test all estimated coefficients, including breakpoints (95% CI & P-values)



18

Recommendations

- Don't use to approximate non-linear functions
- Best to use when breakpoint is of main interest




19

Questions?




20



21

GLM: coefficients linearly enter model


- $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 \dots + \beta_n X_n + \varepsilon$
- May want to fit function where this isn't the case
 - E.g., power function: $Y = \beta_0 X^{\beta_1}$



22

Non-linear least squares

- Coefficients estimated algorithmically → computer tries different values to minimize residuals (i.e., least squares criterion)
- Need prespecified function (e.g., based on theory)
- Can do hypothesis testing & prediction




23

Questions?




24



25

When it's used


- No *a priori* function to explore or test (e.g., linear, segmented, exponential)
- Simply want to draw audience's eye to average trend, using a smoothed curve



26

How it works in principle



- Data are smoothed by iteratively moving a window & fitting a linear model each time



27

Adjusting the smoothing

- Window size set as proportion of total data pts.
 - Smoother as window size increases
- Window linear model can be simple linear regression or quadratic regression
 - Smoother if simple linear regression



28

Questions?

