

# Genus Bounds for Some Dynatomic Modular Curves

UWO Final Thesis Exam Public Lecture

---

Andrew W. Herring

30 August, 2021

1. Introduction
2. Function Fields and Curves
3. Flavors of Maximal Subgroups and Genus Bounds

# Introduction

---

- ⊙ Let  $\mathbb{F}$  be any perfect field with a fixed algebraic closure  $\bar{\mathbb{F}}$ ;

# Polynomial Dynamical Systems

- ⊙ Let  $\mathbb{F}$  be any perfect field with a fixed algebraic closure  $\bar{\mathbb{F}}$ ;
- ⊙ For a non-constant polynomial  $g(x) \in \mathbb{F}[x]$  and a positive integer  $k$ , let  $g^k(x)$  denote the  $k$ -th iterate of  $g$ ;

# Polynomial Dynamical Systems

- ⊙ Let  $\mathbb{F}$  be any perfect field with a fixed algebraic closure  $\bar{\mathbb{F}}$ ;
- ⊙ For a non-constant polynomial  $g(x) \in \mathbb{F}[x]$  and a positive integer  $k$ , let  $g^k(x)$  denote the  **$k$ -th iterate of  $g$** ;
- ⊙ Define  $\text{Per}(g) := \{\alpha \in \bar{\mathbb{F}} : g^k(\alpha) = \alpha, \text{ some } k \geq 1\}$ ;

# Polynomial Dynamical Systems

- ⊙ Let  $\mathbb{F}$  be any perfect field with a fixed algebraic closure  $\bar{\mathbb{F}}$ ;
- ⊙ For a non-constant polynomial  $g(x) \in \mathbb{F}[x]$  and a positive integer  $k$ , let  $g^k(x)$  denote the  **$k$ -th iterate of  $g$** ;
- ⊙ Define  $\text{Per}(g) := \{\alpha \in \bar{\mathbb{F}} : g^k(\alpha) = \alpha, \text{ some } k \geq 1\}$ ;
- ⊙ Elements of  $\text{Per}(g)$  are **periodic points of  $g$** ;

# Polynomial Dynamical Systems

- ⊙ Let  $\mathbb{F}$  be any perfect field with a fixed algebraic closure  $\bar{\mathbb{F}}$ ;
- ⊙ For a non-constant polynomial  $g(x) \in \mathbb{F}[x]$  and a positive integer  $k$ , let  $g^k(x)$  denote the  **$k$ -th iterate of  $g$** ;
- ⊙ Define  $\text{Per}(g) := \{\alpha \in \bar{\mathbb{F}} : g^k(\alpha) = \alpha, \text{ some } k \geq 1\}$ ;
- ⊙ Elements of  $\text{Per}(g)$  are **periodic points of  $g$** ;
- ⊙ The **period** of  $\alpha \in \text{Per}(g)$  is  $n := \min\{k \geq 1 : g^k(\alpha) = \alpha\}$ ;



# Polynomial Dynamical Systems

- ⊙ Let  $\mathbb{F}$  be any perfect field with a fixed algebraic closure  $\bar{\mathbb{F}}$ ;
- ⊙ For a non-constant polynomial  $g(x) \in \mathbb{F}[x]$  and a positive integer  $k$ , let  $g^k(x)$  denote the  **$k$ -th iterate of  $g$** ;
- ⊙ Define  $\text{Per}(g) := \{\alpha \in \bar{\mathbb{F}} : g^k(\alpha) = \alpha, \text{ some } k \geq 1\}$ ;
- ⊙ Elements of  $\text{Per}(g)$  are **periodic points of  $g$** ;
- ⊙ The **period** of  $\alpha \in \text{Per}(g)$  is  $n := \min\{k \geq 1 : g^k(\alpha) = \alpha\}$ ;
- ⊙ Define  $\text{Per}_n(g) := \{\alpha \in \bar{\mathbb{F}} : \alpha \text{ has period } n\}$ ;

# Polynomial Dynamical Systems

- ⊙ Let  $\mathbb{F}$  be any perfect field with a fixed algebraic closure  $\bar{\mathbb{F}}$ ;
- ⊙ For a non-constant polynomial  $g(x) \in \mathbb{F}[x]$  and a positive integer  $k$ , let  $g^k(x)$  denote the  **$k$ -th iterate of  $g$** ;
- ⊙ Define  $\text{Per}(g) := \{\alpha \in \bar{\mathbb{F}} : g^k(\alpha) = \alpha, \text{ some } k \geq 1\}$ ;
- ⊙ Elements of  $\text{Per}(g)$  are **periodic points of  $g$** ;
- ⊙ The **period** of  $\alpha \in \text{Per}(g)$  is  $n := \min\{k \geq 1 : g^k(\alpha) = \alpha\}$ ;
- ⊙ Define  $\text{Per}_n(g) := \{\alpha \in \bar{\mathbb{F}} : \alpha \text{ has period } n\}$ ;
- ⊙ For any subset  $S \subseteq \bar{\mathbb{F}}$ , define  $\text{Per}_n(g; S) := \text{Per}_n(g) \cap S$

# Motivating Problem

For  $c \in \mathbb{Q}$ , let  $f_c(x) := x^2 + c \in \mathbb{Q}[x]$

# Motivating Problem

For  $c \in \mathbb{Q}$ , let  $f_c(x) := x^2 + c \in \mathbb{Q}[x]$

**Conjecture:** Flynn, Poonen, and Schaefer–1997; [3]

If  $n \geq 4$ , then  $\text{Per}_n(f_c; \mathbb{Q})$  is empty for every  $c \in \mathbb{Q}$ .

# Motivating Problem

For  $c \in \mathbb{Q}$ , let  $f_c(x) := x^2 + c \in \mathbb{Q}[x]$

**Conjecture: Flynn, Poonen, and Schaefer–1997; [3]**

If  $n \geq 4$ , then  $\text{Per}_n(f_c; \mathbb{Q})$  is empty for every  $c \in \mathbb{Q}$ .

The conjecture is known to hold for:

# Motivating Problem

For  $c \in \mathbb{Q}$ , let  $f_c(x) := x^2 + c \in \mathbb{Q}[x]$

**Conjecture:** Flynn, Poonen, and Schaefer–1997; [3]

If  $n \geq 4$ , then  $\text{Per}_n(f_c; \mathbb{Q})$  is empty for every  $c \in \mathbb{Q}$ .

The conjecture is known to hold for:

- ⊙  $n = 4$ ; [6]

# Motivating Problem

For  $c \in \mathbb{Q}$ , let  $f_c(x) := x^2 + c \in \mathbb{Q}[x]$

**Conjecture: Flynn, Poonen, and Schaefer–1997; [3]**

If  $n \geq 4$ , then  $\text{Per}_n(f_c; \mathbb{Q})$  is empty for every  $c \in \mathbb{Q}$ .

The conjecture is known to hold for:

- ⊙  $n = 4$ ; [6]
- ⊙  $n = 5$ ; [3]

# Motivating Problem

For  $c \in \mathbb{Q}$ , let  $f_c(x) := x^2 + c \in \mathbb{Q}[x]$

**Conjecture: Flynn, Poonen, and Schaefer–1997; [3]**

If  $n \geq 4$ , then  $\text{Per}_n(f_c; \mathbb{Q})$  is empty for every  $c \in \mathbb{Q}$ .

The conjecture is known to hold for:

- ⊙  $n = 4$ ; [6]
- ⊙  $n = 5$ ; [3]
- ⊙  $n = 6^*$ ; [9]

---

\*Conditional on Birch and Swinnerton-Dyer Conjecture



## Example

Let  $c = -1 \in \mathbb{Q}$  and consider  $f_c(x) = x^2 - 1 \in \mathbb{Q}[x]$ .

## Example

Let  $c = -1 \in \mathbb{Q}$  and consider  $f_c(x) = x^2 - 1 \in \mathbb{Q}[x]$ .

- ⊙ Suppose  $f_c(\alpha) = \alpha$ . Then  $\alpha$  is a root of  $f_c(x) - x = x^2 - x - 1$ .

## Example

Let  $c = -1 \in \mathbb{Q}$  and consider  $f_c(x) = x^2 - 1 \in \mathbb{Q}[x]$ .

- ⊙ Suppose  $f_c(\alpha) = \alpha$ . Then  $\alpha$  is a root of  $f_c(x) - x = x^2 - x - 1$ .
- ⊙ But the discriminant  $\Delta(x^2 - x - 1) = 5$  is not a square in  $\mathbb{Q}$ .

# Example

Let  $c = -1 \in \mathbb{Q}$  and consider  $f_c(x) = x^2 - 1 \in \mathbb{Q}[x]$ .

- ⊙ Suppose  $f_c(\alpha) = \alpha$ . Then  $\alpha$  is a root of  $f_c(x) - x = x^2 - x - 1$ .
- ⊙ But the discriminant  $\Delta(x^2 - x - 1) = 5$  is not a square in  $\mathbb{Q}$ .
- ⊙ Thus  $\alpha \notin \mathbb{Q}$  and  $\text{Per}_1(f_c; \mathbb{Q})$  is empty.

# Example

Let  $c = -1 \in \mathbb{Q}$  and consider  $f_c(x) = x^2 - 1 \in \mathbb{Q}[x]$ .

- ⊙ Suppose  $f_c(\alpha) = \alpha$ . Then  $\alpha$  is a root of  $f_c(x) - x = x^2 - x - 1$ .
- ⊙ But the discriminant  $\Delta(x^2 - x - 1) = 5$  is not a square in  $\mathbb{Q}$ .
- ⊙ Thus  $\alpha \notin \mathbb{Q}$  and  $\text{Per}_1(f_c; \mathbb{Q})$  is empty.
- ⊙ Meanwhile, we have

$$f_c(0) = (0)^2 - 1 = -1; \quad f_c(-1) = (-1)^2 - 1 = 0; \quad \text{and}$$

$$f_c^2(x) - x = x^4 - 2x^2 - x = x(x+1)(x^2 - x - 1).$$

# Example

Let  $c = -1 \in \mathbb{Q}$  and consider  $f_c(x) = x^2 - 1 \in \mathbb{Q}[x]$ .

- ⊙ Suppose  $f_c(\alpha) = \alpha$ . Then  $\alpha$  is a root of  $f_c(x) - x = x^2 - x - 1$ .
- ⊙ But the discriminant  $\Delta(x^2 - x - 1) = 5$  is not a square in  $\mathbb{Q}$ .
- ⊙ Thus  $\alpha \notin \mathbb{Q}$  and  $\text{Per}_1(f_c; \mathbb{Q})$  is empty.
- ⊙ Meanwhile, we have

$$f_c(0) = (0)^2 - 1 = -1; \quad f_c(-1) = (-1)^2 - 1 = 0; \text{ and}$$

$$f_c^2(x) - x = x^4 - 2x^2 - x = x(x+1)(x^2 - x - 1).$$

- ⊙ Hence,  $\text{Per}_2(f_c; \mathbb{Q}) = \{0, -1\}$ .

# Motivating Problem—Finitely Many Exceptions

For  $n \geq 1$  consider the statement

F.E.( $n$ ) = “there are at most finitely many  $c \in \mathbb{Q}$  such that  $\text{Per}_n(f_c; \mathbb{Q})$  is non-empty”

# Motivating Problem—Finitely Many Exceptions

For  $n \geq 1$  consider the statement

F.E.( $n$ ) = “there are at most finitely many  $c \in \mathbb{Q}$  such that  $\text{Per}_n(f_c; \mathbb{Q})$  is non-empty”

**Theorem: Krumm–2019; [5]**

F.E.( $n$ ) holds for  $n \in \{5, 6, 7, 9\}$ .



# Motivating Problem—Finitely Many Exceptions

For  $n \geq 1$  consider the statement

$\text{F.E.}(n)$  = “there are at most finitely many  $c \in \mathbb{Q}$  such that  $\text{Per}_n(f_c; \mathbb{Q})$  is non-empty”

**Theorem: Krumm–2019; [5]**

$\text{F.E.}(n)$  holds for  $n \in \{5, 6, 7, 9\}$ .

**Theorem: H.–2021**

$\text{F.E.}(n)$  holds for every  $n \geq 10$ .

# Motivating Problem—Finitely Many Exceptions

For  $n \geq 1$  consider the statement

F.E.( $n$ ) = “there are at most finitely many  $c \in \mathbb{Q}$  such that  $\text{Per}_n(f_c; \mathbb{Q})$  is non-empty”

**Theorem: Krumm–2019; [5]**

F.E.( $n$ ) holds for  $n \in \{5, 6, 7, 9\}$ .

**Theorem: H.–2021**

F.E.( $n$ ) holds for every  $n \geq 10$ .

Thus F.E.( $n$ ) holds for every  $n \geq 5$  except possibly for  $n = 8$

# Dynatomic Polynomials

- ⊙ Let  $t$  be transcendental over  $\mathbb{C}$  and consider  $K_0 := \mathbb{Q}(t)$ ,  $A_0 := \mathbb{Q}[t]$ , and  $f(x) := x^2 + t \in A_0[x]$ .

# Dynatomic Polynomials

- ⊙ Let  $t$  be transcendental over  $\mathbb{C}$  and consider  $K_0 := \mathbb{Q}(t)$ ,  $A_0 := \mathbb{Q}[t]$ , and  $f(x) := x^2 + t \in A_0[x]$ .
- ⊙ For  $n \geq 1$ , the  $n$ -th dynatomic polynomial is

$$\Phi_n(x) := \prod_{d|n} \left( f^d(x) - x \right)^{\mu(n/d)}$$

where  $\mu$  is the Mobius function.

# Dynatomic Polynomials

- Let  $t$  be transcendental over  $\mathbb{C}$  and consider  $K_0 := \mathbb{Q}(t)$ ,  $A_0 := \mathbb{Q}[t]$ , and  $f(x) := x^2 + t \in A_0[x]$ .
- For  $n \geq 1$ , the  $n$ -th dynatomic polynomial is

$$\Phi_n(x) := \prod_{d|n} \left( f^d(x) - x \right)^{\mu(n/d)}$$

where  $\mu$  is the Mobius function.

## Example:

$$\Phi_1(x) = x^2 - x + t; \quad \Phi_2(x) = x^2 + x + t + 1$$

$$\begin{aligned} \Phi_3(x) = & x^6 + x^5 + (3t + 1)x^4 + (2t + 1)x^3 + (3t^2 + 3t + 1)x^2 \\ & + (t^2 + 2t + 1)x + t^3 + 2t^2 + t + 1 \end{aligned}$$

# Dynatomic Polynomials

Fix  $n \geq 1$ . Let  $L_0$  denote the **splitting field of  $\Phi_n(x)$  over  $K_0$**  and let  $Z := \{\alpha \in L_0 : \Phi_n(\alpha) = 0\}$  denote the **zero set of  $\Phi_n$** .

# Dynatomic Polynomials

Fix  $n \geq 1$ . Let  $L_0$  denote the **splitting field** of  $\Phi_n(x)$  over  $K_0$  and let  $Z := \{\alpha \in L_0 : \Phi_n(\alpha) = 0\}$  denote the **zero set** of  $\Phi_n$ .

## Proposition

The following hold:

# Dynatomic Polynomials

Fix  $n \geq 1$ . Let  $L_0$  denote the **splitting field** of  $\Phi_n(x)$  over  $K_0$  and let  $Z := \{\alpha \in L_0 : \Phi_n(\alpha) = 0\}$  denote the **zero set** of  $\Phi_n$ .

## Proposition

The following hold:

1.  $\Phi_n(x) \in A_0[x]$  and  $\Phi_n(x)$  is monic in  $x$ ; [7]



# Dynatomic Polynomials

Fix  $n \geq 1$ . Let  $L_0$  denote the **splitting field** of  $\Phi_n(x)$  over  $K_0$  and let  $Z := \{\alpha \in L_0 : \Phi_n(\alpha) = 0\}$  denote the **zero set** of  $\Phi_n$ .

## Proposition

The following hold:

1.  $\Phi_n(x) \in A_0[x]$  and  $\Phi_n(x)$  is monic in  $x$ ; [7]
2.  $\Phi_n(x)$  is irreducible over  $\mathbb{C}(t)$ ; [1]

# Dynatomic Polynomials

Fix  $n \geq 1$ . Let  $L_0$  denote the **splitting field** of  $\Phi_n(x)$  over  $K_0$  and let  $Z := \{\alpha \in L_0 : \Phi_n(\alpha) = 0\}$  denote the **zero set** of  $\Phi_n$ .

## Proposition

The following hold:

1.  $\Phi_n(x) \in A_0[x]$  and  $\Phi_n(x)$  is monic in  $x$ ; [7]
2.  $\Phi_n(x)$  is irreducible over  $\mathbb{C}(t)$ ; [1]
3.  $Z = \text{Per}_n(f; \bar{\mathbb{Q}})$ ;

# Dynatomic Polynomials

Fix  $n \geq 1$ . Let  $L_0$  denote the **splitting field** of  $\Phi_n(x)$  over  $K_0$  and let  $Z := \{\alpha \in L_0 : \Phi_n(\alpha) = 0\}$  denote the **zero set** of  $\Phi_n$ .

## Proposition

The following hold:

1.  $\Phi_n(x) \in A_0[x]$  and  $\Phi_n(x)$  is monic in  $x$ ; [7]
2.  $\Phi_n(x)$  is irreducible over  $\mathbb{C}(t)$ ; [1]
3.  $Z = \text{Per}_n(f; \bar{\mathbb{Q}})$ ;
4.  $\Phi_n(x)$  divides  $\Phi_n(f(x))$ . [7]

## Corollary

Let  $r := \frac{\deg \Phi_n(x)}{n}$ . Then

$$Z = \sqcup_{i=1}^r A_i$$

where  $A_i = \{\alpha_i, f(\alpha_i), \dots, f^{n-1}(\alpha_i)\}$  for some  $\alpha_i \in L_0$  for every  $i = 1, \dots, r$ .

## Corollary

Let  $r := \frac{\deg \Phi_n(x)}{n}$ . Then

$$Z = \sqcup_{i=1}^r A_i$$

where  $A_i = \{\alpha_i, f(\alpha_i), \dots, f^{n-1}(\alpha_i)\}$  for some  $\alpha_i \in L_0$  for every  $i = 1, \dots, r$ .

Each set  $A_i$  is an  $f$ -orbit.

# Specializations

Let  $c \in \mathbb{Q}$

# Specializations

Let  $c \in \mathbb{Q}$

- ⊙  $f_c(x) = x^2 + c = (x^2 + t) |_{t=c} = f(x) |_{t=c}$ , i.e.,  $f_c(x)$  is a **specialization** of  $f(x)$ .

# Specializations

Let  $c \in \mathbb{Q}$

- ⊙  $f_c(x) = x^2 + c = (x^2 + t) |_{t=c} = f(x) |_{t=c}$ , i.e.,  $f_c(x)$  is a **specialization** of  $f(x)$ .
- ⊙ Define

$$\Phi_{n,c}(x) := \Phi_n(x) |_{t=c} = \prod_{d|n} \left( f_c^d(x) - x \right)^{\mu(n/d)} \in \mathbb{Q}[x]$$

the  **$n$ -th dynatomic polynomial at  $c$** .



# Specializations

Let  $c \in \mathbb{Q}$

- ⊙  $f_c(x) = x^2 + c = (x^2 + t) |_{t=c} = f(x) |_{t=c}$ , i.e.,  $f_c(x)$  is a **specialization** of  $f(x)$ .
- ⊙ Define

$$\Phi_{n,c}(x) := \Phi_n(x) |_{t=c} = \prod_{d|n} \left( f_c^d(x) - x \right)^{\mu(n/d)} \in \mathbb{Q}[x]$$

the  **$n$ -th dynatomic polynomial at  $c$** .

- ⊙ Let  $L_{0,c}$  denote the **splitting field of  $\Phi_{n,c}(x)$  over  $\mathbb{Q}$** , and let  $Z_c := \{\alpha \in L_{0,c} : \Phi_{n,c}(\alpha) = 0\}$  denote the **zero set of  $\Phi_{n,c}$** .

# Specializations

Let  $c \in \mathbb{Q}$

- ⊙  $f_c(x) = x^2 + c = (x^2 + t) |_{t=c} = f(x) |_{t=c}$ , i.e.,  $f_c(x)$  is a **specialization** of  $f(x)$ .
- ⊙ Define

$$\Phi_{n,c}(x) := \Phi_n(x) |_{t=c} = \prod_{d|n} \left( f_c^d(x) - x \right)^{\mu(n/d)} \in \mathbb{Q}[x]$$

the  **$n$ -th dynatomic polynomial at  $c$** .

- ⊙ Let  $L_{0,c}$  denote the **splitting field of  $\Phi_{n,c}(x)$  over  $\mathbb{Q}$** , and let  $Z_c := \{\alpha \in L_{0,c} : \Phi_{n,c}(\alpha) = 0\}$  denote the **zero set of  $\Phi_{n,c}$** .
- ⊙ We have  $\text{Per}_n(f_c) \subseteq Z_c$ , but equality need not hold.

# Galois Groups

- ⊙  $L_0/K_0$  is a finite Galois extension; let  $G := \text{Gal}(L_0/K_0)$  and call it the  *$n$ -th dynatomic Galois group*.

# Galois Groups

- ⊙  $L_0/K_0$  is a finite Galois extension; let  $G := \text{Gal}(L_0/K_0)$  and call it the  *$n$ -th dynatomic Galois group*.
- ⊙ For every  $c \in \mathbb{Q}$ , also have that  $L_{0,c}/\mathbb{Q}$  is a finite Galois extension; let  $G_c := \text{Gal}(L_{0,c}/\mathbb{Q})$  and call it the  *$n$ -th dynatomic Galois group at  $c$* .

# Galois Groups

- ⊙  $L_0/K_0$  is a finite Galois extension; let  $G := \text{Gal}(L_0/K_0)$  and call it the  **$n$ -th dynatomic Galois group**.
- ⊙ For every  $c \in \mathbb{Q}$ , also have that  $L_{0,c}/\mathbb{Q}$  is a finite Galois extension; let  $G_c := \text{Gal}(L_{0,c}/\mathbb{Q})$  and call it the  **$n$ -th dynatomic Galois group at  $c$** .
- ⊙ Define

$$D_n := \{c \in \mathbb{Q} : \Phi_{n,c} \text{ is inseparable}\}.$$

# Galois Groups

- ⊙  $L_0/K_0$  is a finite Galois extension; let  $G := \text{Gal}(L_0/K_0)$  and call it the  **$n$ -th dynatomic Galois group**.
- ⊙ For every  $c \in \mathbb{Q}$ , also have that  $L_{0,c}/\mathbb{Q}$  is a finite Galois extension; let  $G_c := \text{Gal}(L_{0,c}/\mathbb{Q})$  and call it the  **$n$ -th dynatomic Galois group at  $c$** .
- ⊙ Define

$$D_n := \{c \in \mathbb{Q} : \Phi_{n,c} \text{ is inseparable}\}.$$

## Proposition

If  $c \in \mathbb{Q} - D_n$ , then  $G_c \leq G$ .

# Galois Groups

- ⊙  $L_0/K_0$  is a finite Galois extension; let  $G := \text{Gal}(L_0/K_0)$  and call it the  **$n$ -th dynatomic Galois group**.
- ⊙ For every  $c \in \mathbb{Q}$ , also have that  $L_{0,c}/\mathbb{Q}$  is a finite Galois extension; let  $G_c := \text{Gal}(L_{0,c}/\mathbb{Q})$  and call it the  **$n$ -th dynatomic Galois group at  $c$** .
- ⊙ Define

$$D_n := \{c \in \mathbb{Q} : \Phi_{n,c} \text{ is inseparable}\}.$$

## Proposition

If  $c \in \mathbb{Q} - D_n$ , then  $G_c \leq G$ .

$D_n$  is the zero set of the discriminant of  $\Phi_n$ , and is therefore finite.

# Exceptional Values

Define

$$E_n := \{c \in \mathbb{Q} - D_n : G_c \leq G\}$$

and call it the *n*-th exceptional set. Elements of  $E_n$  are exceptional values



# Exceptional Values

Define

$$E_n := \{c \in \mathbb{Q} - D_n : G_c \preceq G\}$$

and call it the  *$n$ -th exceptional set*. Elements of  $E_n$  are *exceptional values*

## Proposition

If  $c \in \mathbb{Q} - D_n$  and  $\text{Per}_n(f_c; \mathbb{Q})$  is non-empty, then  $c \in E_n$ .

# Exceptional Values

Define

$$E_n := \{c \in \mathbb{Q} - D_n : G_c \leq G\}$$

and call it the  **$n$ -th exceptional set**. Elements of  $E_n$  are **exceptional values**

## Proposition

If  $c \in \mathbb{Q} - D_n$  and  $\text{Per}_n(f_c; \mathbb{Q})$  is non-empty, then  $c \in E_n$ .

Thus to prove F.E.( $n$ ), it suffices to show that  $E_n$  is finite.

# Exceptional Values

Define

$$E_n := \{c \in \mathbb{Q} - D_n : G_c \leq G\}$$

and call it the *n*-th exceptional set. Elements of  $E_n$  are exceptional values

## Proposition

If  $c \in \mathbb{Q} - D_n$  and  $\text{Per}_n(f_c; \mathbb{Q})$  is non-empty, then  $c \in E_n$ .

Thus to prove F.E.( $n$ ), it suffices to show that  $E_n$  is finite.

## Remark

By Hilbert's Irreducibility Theorem,  $E_n$  is known to be "thin." Thin sets are small (in a suitable sense), but can still be infinite.

# Function Fields and Curves

---

# Equivalent Diagrams

- ⊙ For a subgroup  $H \leq G$ , let  $L_0^H$  denote the fixed field of  $H$  in  $L_0$ .

# Equivalent Diagrams

- ⊙ For a subgroup  $H \leq G$ , let  $L_0^H$  denote the **fixed field of  $H$  in  $L_0$** .
- ⊙ Each of the fields  $K_0$ ,  $L_0$ , and  $L_0^H$  is a **function field over  $\mathbb{Q}$** . That is, for every  $\mathbb{F} \in \{K_0, L_0, L_0^H\}$ ,  $\mathbb{F}$  is a finitely generated extension of  $\mathbb{Q}$  with transcendence degree 1 over  $\mathbb{Q}$  and  $\mathbb{Q}$  is algebraically closed in  $\mathbb{F}$ .

# Equivalent Diagrams

- ⊙ For a subgroup  $H \leq G$ , let  $L_0^H$  denote the **fixed field of  $H$  in  $L_0$** .
- ⊙ Each of the fields  $K_0$ ,  $L_0$ , and  $L_0^H$  is a **function field over  $\mathbb{Q}$** . That is, for every  $\mathbb{F} \in \{K_0, L_0, L_0^H\}$ ,  $\mathbb{F}$  is a finitely generated extension of  $\mathbb{Q}$  with transcendence degree 1 over  $\mathbb{Q}$  and  $\mathbb{Q}$  is algebraically closed in  $\mathbb{F}$ .
- ⊙ By a well-known equivalence of categories, there are curves  $X, Y, W$  defined over  $\mathbb{Q}$  corresponding to  $L_0, L_0^H$ , and  $K_0$ , respectively.

# Equivalent Diagrams

- ⊙ For a subgroup  $H \leq G$ , let  $L_0^H$  denote the **fixed field of  $H$  in  $L_0$** .
- ⊙ Each of the fields  $K_0$ ,  $L_0$ , and  $L_0^H$  is a **function field over  $\mathbb{Q}$** . That is, for every  $\mathbb{F} \in \{K_0, L_0, L_0^H\}$ ,  $\mathbb{F}$  is a finitely generated extension of  $\mathbb{Q}$  with transcendence degree 1 over  $\mathbb{Q}$  and  $\mathbb{Q}$  is algebraically closed in  $\mathbb{F}$ .
- ⊙ By a well-known equivalence of categories, there are curves  $X, Y, W$  defined over  $\mathbb{Q}$  corresponding to  $L_0, L_0^H$ , and  $K_0$ , respectively.
- ⊙ The curve  $W$  corresponding to  $K_0 = \mathbb{Q}(t)$  is  $\mathbb{P}_{\mathbb{Q}}^1$ , the **projective line defined over  $\mathbb{Q}$** .




# Equivalent Diagrams

- ⊙ The curve  $Y$  corresponding to  $L_0^H$  is the **quotient curve**  
 $X_H \equiv X/H$  whose points are  $H$ -orbits under the action of  
 $H$  on  $X$ .

# Equivalent Diagrams

- ⊙ The curve  $Y$  corresponding to  $L_0^H$  is the **quotient curve**  $X_H \equiv X/H$  whose points are  $H$ -orbits under the action of  $H$  on  $X$ .

We therefore have equivalent diagrams




of function fields and curves over  $\mathbb{Q}$ .

# Equivalent Diagrams

- ⊙ The curve  $Y$  corresponding to  $L_0^H$  is the **quotient curve**  $X_H \equiv X/H$  whose points are  $H$ -orbits under the action of  $H$  on  $X$ .

We therefore have equivalent diagrams



of function fields and curves over  $\mathbb{Q}$ .

- ⊙ Any curve of the form  $X_H$  is called a **dynamotic modular curve**.

# Genus of a curve

Let  $C$  be any curve defined over  $\mathbb{Q}$ . We will use  $g(C)$  to denote the **genus of  $C$**  which is one measure of the geometric complexity of  $C$ .


# Genus of a curve

Let  $C$  be any curve defined over  $\mathbb{Q}$ . We will use  $g(C)$  to denote the **genus of  $C$**  which is one measure of the geometric complexity of  $C$ .


By considering  $C$  over  $\mathbb{C}$ , we obtain a compact Riemann surface  $X$ ; then the genus  $g(C)$  counts the number of “holes/handles” of  $X$ .

# Genus of a curve


$$\mathbb{P}^1 = \mathbb{C} \cup \{\infty\}$$



# Genus of a curve




# Genus of a curve



"Stereographic Projection"




# Genus of a curve




"Stereographic Projection"

# Genus of a curve




"Stereographic Projection"

# Genus of a curve




"Stereographic Projection"

# Genus of a curve




"Stereographic Projection"

# Genus of a curve



"Stereographic Projection"

# Genus of a curve




$$\text{Thus, } g(\mathbb{P}^1) = g(\text{Sphere}) = 0$$

# Genus of a curve

"elliptic curves"

$$y^2 = x^3 + ax + b$$




# Genus of a curve

"elliptic curves"

$$y^2 = x^3 + ax + b$$



"lattice"







# Genus of a curve

"elliptic curves"


$$y^2 = x^3 + ax + b$$



# Genus of a curve



# Genus of a curve



thus,  $g(\mathcal{C}) = g(\text{torus}) = 1$

# Genus of a curve

$$g(\text{circle}) = 0$$

$$g(\text{torus}) = 1$$

$$g(\text{double torus}) = 2$$

⋮

$$g(\text{surface with } g_0 \text{ handles}) = g_0$$

# Rational Points and Exceptions

Let  $C$  be any curve defined over  $\mathbb{Q}$ . Let  $g(C)$  denote the genus of  $C$ , and let  $C(\mathbb{Q})$  denote the set of  $\mathbb{Q}$ -rational points of  $C$ .

# Rational Points and Exceptions

Let  $C$  be any curve defined over  $\mathbb{Q}$ . Let  $g(C)$  denote the **genus of  $C$** , and let  $C(\mathbb{Q})$  denote the **set of  $\mathbb{Q}$ -rational points of  $C$** .

**Theorem: Faltings–1983; [2]**

If  $g(C) \geq 2$ , then  $C(\mathbb{Q})$  is finite.

# Rational Points and Exceptions

Let  $C$  be any curve defined over  $\mathbb{Q}$ . Let  $g(C)$  denote the **genus of  $C$** , and let  $C(\mathbb{Q})$  denote the **set of  $\mathbb{Q}$ -rational points of  $C$** .

**Theorem: Faltings–1983; [2]**

If  $g(C) \geq 2$ , then  $C(\mathbb{Q})$  is finite.

**Theorem: (see Propositions 3.3.1 and 3.3.5 of [8])**

Let  $c \in \mathbb{Q} - D_n$ . Then  $c \in E_n$  if and only if  $c \in \pi_H(X_H(\mathbb{Q}))$  for some proper subgroup  $H \subsetneq G$ .

# Rational Points and Exceptions

## Proposition

If  $g(X_M) \geq 2$  for every maximal subgroup  $M$  of  $G$ , then  $E_n$  is finite.



# Rational Points and Exceptions

## Proposition

If  $g(X_M) \geq 2$  for every maximal subgroup  $M$  of  $G$ , then  $E_n$  is finite.

## Proof sketch:

# Rational Points and Exceptions

## Proposition

If  $g(X_M) \geq 2$  for every maximal subgroup  $M$  of  $G$ , then  $E_n$  is finite.

## Proof sketch:

1. Suffices that  $g(X_H) \geq 2$  for every proper subgroup  $H \subsetneq G$ .

# Rational Points and Exceptions

## Proposition

If  $g(X_M) \geq 2$  for every maximal subgroup  $M$  of  $G$ , then  $E_n$  is finite.

## Proof sketch:

1. Suffices that  $g(X_H) \geq 2$  for every proper subgroup  $H \subsetneq G$ .
  - If this holds, then  $X_H(\mathbb{Q})$  is finite for every  $H \subsetneq G$  by Faltings.

# Rational Points and Exceptions

## Proposition

If  $g(X_M) \geq 2$  for every maximal subgroup  $M$  of  $G$ , then  $E_n$  is finite.

## Proof sketch:

1. Suffices that  $g(X_H) \geq 2$  for every proper subgroup  $H \subsetneq G$ .
  - If this holds, then  $X_H(\mathbb{Q})$  is finite for every  $H \subsetneq G$  by Faltings.
  - Thus  $\pi_H(X_H(\mathbb{Q})) \subset \mathbb{P}_{\mathbb{Q}}^1$  is finite.

# Rational Points and Exceptions

## Proposition

If  $g(X_M) \geq 2$  for every maximal subgroup  $M$  of  $G$ , then  $E_n$  is finite.

## Proof sketch:

1. Suffices that  $g(X_H) \geq 2$  for every proper subgroup  $H \subsetneq G$ .
  - If this holds, then  $X_H(\mathbb{Q})$  is finite for every  $H \subsetneq G$  by Faltings.
  - Thus  $\pi_H(X_H(\mathbb{Q})) \subset \mathbb{P}_{\mathbb{Q}}^1$  is finite.
  - Thus  $E_n$  is finite by the previous Theorem.

# Rational Points and Exceptions

## Proposition

If  $g(X_M) \geq 2$  for every maximal subgroup  $M$  of  $G$ , then  $E_n$  is finite.

## Proof sketch:

1. Suffices that  $g(X_H) \geq 2$  for every proper subgroup  $H \subsetneq G$ .
  - If this holds, then  $X_H(\mathbb{Q})$  is finite for every  $H \subsetneq G$  by Faltings.
  - Thus  $\pi_H(X_H(\mathbb{Q})) \subset \mathbb{P}_{\mathbb{Q}}^1$  is finite.
  - Thus  $E_n$  is finite by the previous Theorem.
2. For every  $H \subsetneq G$ , there is some maximal subgroup  $M$  of  $G$  containing  $H$ .

# Rational Points and Exceptions

## Proposition

If  $g(X_M) \geq 2$  for every maximal subgroup  $M$  of  $G$ , then  $E_n$  is finite.

## Proof sketch:

1. Suffices that  $g(X_H) \geq 2$  for every proper subgroup  $H \subsetneq G$ .
  - If this holds, then  $X_H(\mathbb{Q})$  is finite for every  $H \subsetneq G$  by Faltings.
  - Thus  $\pi_H(X_H(\mathbb{Q})) \subset \mathbb{P}_{\mathbb{Q}}^1$  is finite.
  - Thus  $E_n$  is finite by the previous Theorem.
2. For every  $H \subsetneq G$ , there is some maximal subgroup  $M$  of  $G$  containing  $H$ .
3.  $M \supseteq H \iff L^M \subseteq L^H \iff X_H \rightarrow X_M$ , and genus is non-decreasing along curve morphisms:  $g(X_H) \geq g(X_M)$ .

# Flavors of Maximal Subgroups and Genus Bounds

---



# Semi-Direct Product Structure

Recall our earlier

## Corollary

$Z = \sqcup_{i=1}^r A_i$ , where  $A_i = \{\alpha_i, f(\alpha_i), \dots, f^{n-1}(\alpha_i)\}$  is the  $i$ -th  $f$ -orbit for every  $i \in [r] := \{1, \dots, r\}$ .

# Semi-Direct Product Structure

Recall our earlier

## Corollary

$Z = \sqcup_{i=1}^r A_i$ , where  $A_i = \{\alpha_i, f(\alpha_i), \dots, f^{n-1}(\alpha_i)\}$  is the  $i$ -th  $f$ -orbit for every  $i \in [r] := \{1, \dots, r\}$ .

- ⊙ Let  $\Omega := \{A_1, \dots, A_r\}$  the **set of all  $f$ -orbits**, and let  $\Gamma := \text{Sym}(\Omega) \cong S_r$ .

# Semi-Direct Product Structure

Recall our earlier

## Corollary

$Z = \sqcup_{i=1}^r A_i$ , where  $A_i = \{\alpha_i, f(\alpha_i), \dots, f^{n-1}(\alpha_i)\}$  is the  $i$ -th  $f$ -orbit for every  $i \in [r] := \{1, \dots, r\}$ .

- ⊙ Let  $\Omega := \{A_1, \dots, A_r\}$  the **set of all  $f$ -orbits**, and let  $\Gamma := \text{Sym}(\Omega) \cong S_r$ .
- ⊙ For every  $\sigma \in G = \text{Gal}(L_0/K_0)$  we have that  $\sigma \circ f^j = f^j \circ \sigma$  for every  $j$ .

# Semi-Direct Product Structure

Recall our earlier

## Corollary

$Z = \sqcup_{i=1}^r A_i$ , where  $A_i = \{\alpha_i, f(\alpha_i), \dots, f^{n-1}(\alpha_i)\}$  is the  $i$ -th  $f$ -orbit for every  $i \in [r] := \{1, \dots, r\}$ .

- ⊙ Let  $\Omega := \{A_1, \dots, A_r\}$  the **set of all  $f$ -orbits**, and let  $\Gamma := \text{Sym}(\Omega) \cong S_r$ .
- ⊙ For every  $\sigma \in G = \text{Gal}(L_0/K_0)$  we have that  $\sigma \circ f^j = f^j \circ \sigma$  for every  $j$ .
- ⊙ Thus  $\sigma(A_i) \in \Omega$  for every  $i \in [r]$  and there's a permutation  $\sigma$  on  $\Omega \equiv [r]$  given by  $A_i \mapsto \sigma(A_i) \equiv A_{\sigma(i)}$

# Semi-Direct Product Structure

Recall our earlier

## Corollary

$Z = \sqcup_{i=1}^r A_i$ , where  $A_i = \{\alpha_i, f(\alpha_i), \dots, f^{n-1}(\alpha_i)\}$  is the  $i$ -th  $f$ -orbit for every  $i \in [r] := \{1, \dots, r\}$ .

- ⊙ Let  $\Omega := \{A_1, \dots, A_r\}$  the **set of all  $f$ -orbits**, and let  $\Gamma := \text{Sym}(\Omega) \cong S_r$ .
- ⊙ For every  $\sigma \in G = \text{Gal}(L_0/K_0)$  we have that  $\sigma \circ f^j = f^j \circ \sigma$  for every  $j$ .
- ⊙ Thus  $\sigma(A_i) \in \Omega$  for every  $i \in [r]$  and there's a permutation  $\sigma$  on  $\Omega \equiv [r]$  given by  $A_i \mapsto \sigma(A_i) \equiv A_{\sigma(i)}$
- ⊙ Get a permutation representation  $\phi : G \rightarrow \Gamma$ ; let  $N := \ker(\phi)$ .

## Claim

The following hold:

## Claim

The following hold:

1.  $N \cong (\mathbb{Z}/n)^r$ ;

## Claim

The following hold:

1.  $N \cong (\mathbb{Z}/n)^r$ ;
2.  $\phi$  is surjective. Thus  $\phi$  is precisely the quotient homomorphism  $\pi : G \twoheadrightarrow G/N \cong \Gamma$ ;



## Claim

The following hold:

1.  $N \cong (\mathbb{Z}/n)^r$ ;
2.  $\phi$  is surjective. Thus  $\phi$  is precisely the quotient homomorphism  $\pi : G \twoheadrightarrow G/N \cong \Gamma$ ;
3.  $\Gamma = S_r$  acts by automorphisms on  $N = (\mathbb{Z}/n)^r$  via  ${}^\tau(a_1, \dots, a_r) := (a_{\tau(1)}, \dots, a_{\tau(r)})$ ;

## Claim

The following hold:

1.  $N \cong (\mathbb{Z}/n)^r$ ;
2.  $\phi$  is surjective. Thus  $\phi$  is precisely the quotient homomorphism  $\pi : G \twoheadrightarrow G/N \cong \Gamma$ ;
3.  $\Gamma = S_r$  acts by automorphisms on  $N = (\mathbb{Z}/n)^r$  via  ${}^\tau(a_1, \dots, a_r) := (a_{\tau(1)}, \dots, a_{\tau(r)})$ ;
4. The semi-direct product  $N \rtimes \Gamma = (\mathbb{Z}/n)^r \rtimes S_r$  is defined, and in fact  $G \cong N \rtimes \Gamma [1]$ .

# Flavors of Maximal Subgroups

We partition the maximal subgroups  $M$  of  $G$  according to whether  $\pi(M) = \Gamma$ , or  $\pi(M) \leq \Gamma$ .

# Flavors of Maximal Subgroups

We partition the maximal subgroups  $M$  of  $G$  according to whether  $\pi(M) = \Gamma$ , or  $\pi(M) \leq \Gamma$ .

## Lemma: Vanilla Maximal Subgroups

If  $\pi(M) = \Gamma$ , then  $N \cap M$  is maximal amongst all  $\Gamma$ -invariant subgroups of  $N$ .

# Flavors of Maximal Subgroups

We partition the maximal subgroups  $M$  of  $G$  according to whether  $\pi(M) = \Gamma$ , or  $\pi(M) \leq \Gamma$ .

## Lemma: Vanilla Maximal Subgroups

If  $\pi(M) = \Gamma$ , then  $N \cap M$  is maximal amongst all  $\Gamma$ -invariant subgroups of  $N$ .

We call any maximal  $M$  of  $G$  with  $\pi(M) = \Gamma$  a **vanilla maximal subgroup** of  $G$ .

# Flavors of Maximal Subgroups

We partition the maximal subgroups  $M$  of  $G$  according to whether  $\pi(M) = \Gamma$ , or  $\pi(M) \leq \Gamma$ .

## Lemma: Vanilla Maximal Subgroups

If  $\pi(M) = \Gamma$ , then  $N \cap M$  is maximal amongst all  $\Gamma$ -invariant subgroups of  $N$ .

We call any maximal  $M$  of  $G$  with  $\pi(M) = \Gamma$  a **vanilla maximal subgroup** of  $G$ .

## Lemma: Chocolate Maximal Subgroups

If  $\pi(M) \leq \Gamma$ , then  $N \subseteq M$  and  $\pi(M)$  is a maximal subgroup of  $\Gamma$ .

# Flavors of Maximal Subgroups

We partition the maximal subgroups  $M$  of  $G$  according to whether  $\pi(M) = \Gamma$ , or  $\pi(M) \leq \Gamma$ .

## Lemma: Vanilla Maximal Subgroups

If  $\pi(M) = \Gamma$ , then  $N \cap M$  is maximal amongst all  $\Gamma$ -invariant subgroups of  $N$ .

We call any maximal  $M$  of  $G$  with  $\pi(M) = \Gamma$  a **vanilla maximal subgroup** of  $G$ .

## Lemma: Chocolate Maximal Subgroups

If  $\pi(M) \leq \Gamma$ , then  $N \subseteq M$  and  $\pi(M)$  is a maximal subgroup of  $\Gamma$ .

We call any maximal  $M$  of  $G$  with  $\pi(M) \leq \Gamma$  a **chocolate maximal subgroup** of  $G$ .

# Genus Bounds for Chocolate Maximals

## Theorem

For every  $n \geq 5$  and every chocolate maximal subgroup  $M$  of  $G$  we have  $g(X_M) \geq 2$ .



# Genus Bounds for Chocolate Maximals

## Theorem

For every  $n \geq 5$  and every chocolate maximal subgroup  $M$  of  $G$  we have  $g(X_M) \geq 2$ .

## Ideas:

# Genus Bounds for Chocolate Maximals

## Theorem

For every  $n \geq 5$  and every chocolate maximal subgroup  $M$  of  $G$  we have  $g(X_M) \geq 2$ .

## Ideas:

1. Krumm ([5]) showed that for  $n \in \{5, 6, 7, 9\}$ , we have  $g(X_M) \geq 2$  for *every* maximal subgroup  $M$  of  $G$ ; in particular this holds for the chocolate maximals.

# Genus Bounds for Chocolate Maximals

## Theorem

For every  $n \geq 5$  and every chocolate maximal subgroup  $M$  of  $G$  we have  $g(X_M) \geq 2$ .

## Ideas:

1. Krumm ([5]) showed that for  $n \in \{5, 6, 7, 9\}$ , we have  $g(X_M) \geq 2$  for *every* maximal subgroup  $M$  of  $G$ ; in particular this holds for the chocolate maximals.
2. For  $n \geq 8$ , we can use a powerful theorem of Guralnick and Shareshian ([4]) to show that  $g(X_M) \geq 2$  for every chocolate maximal  $M$ .

# Genus Bounds for Vanilla Maximals

## Theorem

For every  $n \geq 10$ , we have that  $g(X_M) \geq 2$  for every vanilla maximal subgroup  $M$  of  $G$ .

# Genus Bounds for Vanilla Maximals

## Theorem

For every  $n \geq 10$ , we have that  $g(X_M) \geq 2$  for every vanilla maximal subgroup  $M$  of  $G$ .

## Ideas:

# Genus Bounds for Vanilla Maximals

## Theorem

For every  $n \geq 10$ , we have that  $g(X_M) \geq 2$  for every vanilla maximal subgroup  $M$  of  $G$ .

## Ideas:

1. For every vanilla maximal  $M$ , there is a prime  $\ell$  dividing  $n$  and a positive integer  $e$  such that  $[G : M] = \ell^e$ .

# Genus Bounds for Vanilla Maximals

## Theorem

For every  $n \geq 10$ , we have that  $g(X_M) \geq 2$  for every vanilla maximal subgroup  $M$  of  $G$ .

## Ideas:

1. For every vanilla maximal  $M$ , there is a prime  $\ell$  dividing  $n$  and a positive integer  $e$  such that  $[G : M] = \ell^e$ .
2. Further, for one type of ramified point of  $X_M \rightarrow \mathbb{P}^1$  we will always have ramification index  $\ell$ , and there will be  $\varphi(n)$  many such ramified points.

# Genus Bounds for Vanilla Maximals

## Theorem

For every  $n \geq 10$ , we have that  $g(X_M) \geq 2$  for every vanilla maximal subgroup  $M$  of  $G$ .

## Ideas:

1. For every vanilla maximal  $M$ , there is a prime  $\ell$  dividing  $n$  and a positive integer  $e$  such that  $[G : M] = \ell^e$ .
2. Further, for one type of ramified point of  $X_M \rightarrow \mathbb{P}^1$  we will always have ramification index  $\ell$ , and there will be  $\varphi(n)$  many such ramified points.
3. Plug this data into the Riemann-Hurwitz genus formula to prove the Theorem in the cases  $n = 11$  and  $n \geq 13$ .



# Genus Bounds for Vanilla Maximals

## Ideas:

1. For every vanilla maximal  $M$ , there is a prime  $\ell$  dividing  $n$  and a positive integer  $e$  such that  $[G : M] = \ell^e$ .
2. Further, for one type of ramified point of  $X_M \rightarrow \mathbb{P}^1$  we will always have ramification index  $\ell$ , and there will be  $\varphi(n)$  many such ramified points.
3. Plug this data into the Riemann-Hurwitz genus formula to prove the Theorem for  $n = 11$  and  $n \geq 13$ .
4. For  $n = 10$  and  $n = 12$ , we have  $\varphi(n) = 4$  and  $\ell = 2$  is a possibility—these values do not contribute enough in Riemann-Hurwitz to deduce  $g(X_M) \geq 2$

# Genus Bounds for Vanilla Maximals

## Ideas:

1. For every vanilla maximal  $M$ , there is a prime  $\ell$  dividing  $n$  and a positive integer  $e$  such that  $[G : M] = \ell^e$ .
2. Further, for one type of ramified point of  $X_M \rightarrow \mathbb{P}^1$  we will always have ramification index  $\ell$ , and there will be  $\varphi(n)$  many such ramified points.
3. Plug this data into the Riemann-Hurwitz genus formula to prove the Theorem for  $n = 11$  and  $n \geq 13$ .
4. For  $n = 10$  and  $n = 12$ , we have  $\varphi(n) = 4$  and  $\ell = 2$  is a possibility—these values do not contribute enough in Riemann-Hurwitz to deduce  $g(X_M) \geq 2$
5. In these cases we consider a second type of ramified point corresponding to prime divisors  $p$  of  $n$  other than 2, i.e.,  $p = 5$  for  $n = 10$  and  $p = 3$  for  $n = 12$ .

- [1] Thierry Bousch. “Sur quelques problemes de dynamique holomorphe”. PhD thesis. Paris 11, 1992.
- [2] G. Faltings. “Endlichkeitssätze für abelsche Varietäten über Zahlkörpern”. In: *Invent. Math.* 73.3 (1983), pp. 349–366.
- [3] E. V. Flynn, Bjorn Poonen, and Edward F. Schaefer. “Cycles of quadratic polynomials and rational points on a genus-2 curve”. In: *Duke Math. J.* 90.3 (1997), pp. 435–463.

- [4] Robert M. Guralnick and John Shareshian. “Symmetric and alternating groups as monodromy groups of Riemann surfaces. I. Generic covers and covers with many branch points”. In: *Mem. Amer. Math. Soc.* 189.886 (2007), pp. vi+128.
- [5] David Krumm. “A finiteness theorem for specializations of dynatomic polynomials”. In: *Algebra & Number Theory* 13.4 (2019), pp. 963–993.
- [6] Patrick Morton. “Arithmetic properties of periodic points of quadratic maps, II”. In: *Acta Arithmetica* 87.2 (1998), pp. 89–102.

- [7] Patrick Morton and Pratiksha Patel. “The Galois theory of periodic points of polynomial maps”. In: *Proc. London Math. Soc.* (3) 68.2 (1994), pp. 225–263. ISSN: 0024-6115.
- [8] Jean-Pierre Serre. *Topics in Galois theory*. Second. Vol. 1. Research Notes in Mathematics. A K Peters, Ltd., Wellesley, MA, 2008.
- [9] Michael Stoll. “Rational 6-cycles under iteration of quadratic polynomials”. In: *LMS Journal of Computation and Mathematics* 11 (2008), pp. 367–380.

THE  
END