Unreal Is Not Open:
prediction, horizons and the future in relativistic becoming

Andrew Impellitteri

25 October 2026 (draft 3)

Abstract

Local theories of relativistic becoming make what is real from an event its causal past. Does the unreality of the future from an event make the future open from it? We separate the two questions. Call an event guaranteed from b if every past-inextendible causal curve from it meets b's chronological past; then every deterministic law with the domain-of-dependence property fixes it from what is real from b. The notion is essentially Geroch's domain of prediction. We characterize it by a condition on the slices of any Cauchy time function, show that it is conformally invariant, and compute it for the standard cosmological models. In Minkowski spacetime, spatially flat and open big-bang universes, the closed radiation universe and de Sitter spacetime, nothing outside an event's past is guaranteed. In Minkowski spacetime and the flat universes no local law need fix more than the causal past: a conformally coupled scalar field is fixed from an event exactly there. In the closed dust model everything is guaranteed from every event after maximum expansion, the known moment at which the particle horizon disappears. Before it, the same scalar field is already fixed on three diamonds: the event's past, an antipodal echo of it, and a region of the event's own timelike future just before the big crunch, while the intervening future stays open. For this field, which travels on light rays alone, what is fixed is decided by null geodesics rather than by all causal curves. Other local laws fix more: one with a preferred flow fixes everything even in Minkowski spacetime. So unreality does not entail openness. How much of the unreal future is fixed from an event depends on the spacetime's causal structure, whose guarantee can change with cosmic epoch, and on the fields, which that structure does not decide. A closing remark records what the corresponding standpoint logic does and does not show about cosmological horizons.

1 Introduction

Stein's theorem says that in Minkowski spacetime a transitive, symmetry-invariant relation of becoming that makes some past event definite is the causal order or trivial [6, 18, 19]. A companion paper [20] extends this to general relativity under a locality principle: what has become for an event b always includes its chronological past and never goes beyond its causal past. On views that read "has become for b" as "is real from b" [7, 13], the future of b is not real from b.

It is tempting to infer that the future is *open* from b. The inference fails, and the way it fails is instructive. Openness in the sense relevant here is a matter of what the real fixes, and that depends on the laws and on the global causal structure. The point is old in outline: a growing block [4] combined with deterministic laws makes the future unreal yet fixed, and Rakíc's sea-fight case turns on an event outside an observer's past that a decision inside it appears to settle [16]. What is missing is a precise account, for relativistic spacetimes, of when what is real from an event fixes anything beyond itself. Geroch's domain of prediction supplies the tool [8], and Manchak's work on prediction its limits [14]. We use them to answer three questions:

1. Which events does the causal structure alone guarantee to be fixed, under any deterministic local law, by what is real from b (Theorem 3)?

2. In the standard cosmological models, how much is that, and how much more can a particular field fix (Section 3)?

3. What follows for the relation between unreality and openness (Section 4)?

Conventions. Spacetimes are smooth, connected, time-oriented and globally hyperbolic. $I^-_{\mathrm{s}}(p)$ is the strict chronological past (via timelike curves), $J^-(p)$ the causal past; $x \le y$ means $x \in J^-(y)$, and $x \ll y$ means $x \in I^-_{\mathrm{s}}(y)$. We use that $\le$ is closed, that $J^-(x) \cap S$ is compact for every Cauchy surface S, push-up ($y \le z \ll b$ implies $y \ll b$), and that $M = I^-_{\mathrm{s}}(S) \cup S \cup I^+_{\mathrm{s}}(S)$ for a Cauchy surface S [15]. A *Cauchy time function* is a continuous surjection $t : M \to (\alpha, \beta)$, $-\infty \le \alpha < \beta \le \infty$, strictly increasing along future-directed causal curves, whose level sets $S_c = t^{-1}(c)$ are Cauchy surfaces [5].

2 Guaranteed from an event

Definition 1. An event x is *guaranteed from* b if every past-inextendible causal curve from x meets $I^-_{\mathrm{s}}(b)$. Write $\mathrm{Det}(b)$ for the set of such events. Clearly $I^-_{\mathrm{s}}(b) \subseteq \mathrm{Det}(b)$.

$\mathrm{Det}(b)$ is a property of the spacetime and the event. To connect it with what the laws fix, let $\mathcal{L}$ be the set of solutions of some field equations on M. Say that x is *fixed from* b *under* $\mathcal{L}$, and write $x \in F_{\mathcal{L}}(b)$, if any two solutions in $\mathcal{L}$ that agree on $I^-_{\mathrm{s}}(b)$ agree at x. Say that $\mathcal{L}$ has the *domain-of-dependence property* if, for every open $U \subseteq M$, any two solutions that agree on U agree at every event all of whose past-inextendible causal curves meet U. This is a locality assumption on the dynamics: nothing propagates outside the light cones.

Remark 2 (What is guaranteed and what is fixed)**.** If $\mathcal{L}$ has the domain-of-dependence property and consists of continuous fields, then

$$\mathrm{Det}(b) \cup J^-(b) \subseteq F_{\mathcal{L}}(b),$$

since $J^-(b)$ is the closure of $I^-_{\mathrm{s}}(b)$ in a globally hyperbolic spacetime [15]. So $\mathrm{Det}(b)$ is fixed under every such law: it is what the causal structure guarantees, whatever the fields. The inclusion is in general strict, and nothing about $F_{\mathcal{L}}(b)$ beyond it follows from $\mathrm{Det}(b)$ being small. For example, on Minkowski spacetime with the unit timelike field $u = \partial_t$, the local equations $u \cdot \nabla Q = P$, $u \cdot \nabla P = 0$ have the solutions $Q = A(\vec{X}) + tB(\vec{X})$, $P = B(\vec{X})$ for arbitrary smooth A, B. The values of (Q, P) at one point of an integral curve fix them along it, and the past integral curve from x is a past-inextendible causal curve, so the law has the domain-of-dependence property; and every integral curve of u enters $I^-_{\mathrm{s}}(b)$, so $F_{\mathcal{L}}(b) = M$ for every b. A law of this kind uses background structure beyond the metric, but nothing in the domain-of-dependence property excludes it. Neither notion is Belnap's settledness, on which a proposition is settled at a point event if it holds in every possible history through it [2]; under indeterministic laws they come apart from it, and we say nothing about such laws. We use $I^-_{\mathrm{s}}(b)$, which every local becoming theory counts as real from b, rather than $J^-(b)$; the characterization below uses that $I^-_{\mathrm{s}}(b)$ is open.

Relation to Geroch. Geroch's domain of prediction $P(q)$ is the set of x such that every past-directed timelike curve from x without past endpoint enters $I^-(q)$, but $I^-(x) \not\subseteq I^-(q)$ [8, sec. 3]. $\mathrm{Det}(b)$ is the causal-curve analogue of the union of $P(b)$ with the events whose pasts lie in that of b; we expect the two to differ at most on null boundaries, but have not proved it. Geroch gives an equivalent form by achronal surfaces in $I^-(q)$ [8, sec. 3]; Theorem 3 is a slicing form for globally hyperbolic spacetimes.

Theorem 3. *Let $t : M \to (\alpha, \beta)$ be a Cauchy time function. Then $x \in \mathrm{Det}(b)$ if and only if there is $c \in (\alpha, t(x))$ with $J^-(x) \cap S_c \subseteq I^-_{\mathrm{s}}(b)$.*

Proof. ($\Leftarrow$) A past-inextendible causal curve from x crosses the Cauchy surface S_c, at a point of $J^-(x) \cap S_c \subseteq I^-_{\mathrm{s}}(b)$.

($\Rightarrow$) Suppose that for every $c \in (\alpha, t(x))$ the set $K_c = (J^-(x) \cap S_c) \setminus I^-_{\mathrm{s}}(b)$ is nonempty; it is compact, since $I^-_{\mathrm{s}}(b)$ is open. Fix $t(x) > c_1 > c_2 > \dots$ with $c_n \to \alpha$, and let

$$P_n = \{(z_1, \dots, z_n) \in K_{c_1} \times \cdots \times K_{c_n} : x \geq z_1 \geq z_2 \geq \cdots \geq z_n\}.$$

P_n is compact, since $\leq$ is closed. It is nonempty: pick $y \in K_{c_n}$ and a causal curve from x to y; it crosses each S_{c_i}, $i < n$, at some $z_i \leq x$, and $z_i \notin I^-_{\mathrm{s}}(b)$, for otherwise $y \leq z_i \ll b$ would give $y \ll b$. The maps $P_{n+1} \to P_n$ forgetting the last coordinate are continuous, so the inverse limit of the nonempty compact spaces P_n is nonempty, giving $x \geq z_1 \geq z_2 \geq \cdots$ with $z_n \in K_{c_n}$. Join consecutive terms by causal curves and concatenate. Each segment avoids $I^-_{\mathrm{s}}(b)$ by push-up, as before. Along the concatenation $t \to \alpha$, so it has no past endpoint in M: it is a past-inextendible causal curve from x missing $I^-_{\mathrm{s}}(b)$. □

Corollary 4. *If $I^-_{\mathrm{s}}(b)$ contains a Cauchy surface S, then $\mathrm{Det}(b) = M$.*

Proof. If $x \in I^-_{\mathrm{s}}(S) \cup S$, then $x \leq s \ll b$ for some $s \in S$, so $x \in I^-_{\mathrm{s}}(b)$. If $x \in I^+_{\mathrm{s}}(S)$, extend a past-inextendible causal curve from x to an inextendible one; it meets S, and not to the future of x, since then $s' \ll x \leq s$ for some $s, s' \in S$, contradicting achronality. So the curve from x meets $S \subseteq I^-_{\mathrm{s}}(b)$. □

Since $\mathrm{Det}(b)$ is defined by causal curves alone, it depends only on the conformal class of the metric; conformal diagrams decide it.

3 The standard models

Lemma 5. *Let (Σ, d) be $\mathbb{R}^3$, hyperbolic space H^3, or the unit sphere S^3 with $r_2 \leq \pi$. Then the closed ball $\bar{B}(X, r_1)$ lies in the open ball $B(Y, r_2)$ if and only if $d(X, Y) + r_1 < r_2$.*

Proof. ($\Leftarrow$) is the triangle inequality. ($\Rightarrow$) Let $r_1 \geq 0$ and suppose $d(X, Y) + r_1 \geq r_2$. On $\mathbb{R}^3$ and H^3, where geodesics extend indefinitely and minimize, or on S^3 when $d(X, Y) + r_1 \leq \pi$, follow a minimizing geodesic from Y through X (any geodesic from X if $X = Y$) a further distance r_1; the endpoint lies in $\bar{B}(X, r_1)$ at distance $d(X, Y) + r_1 \geq r_2$ from Y. If $d(X, Y) + r_1 > \pi$ on S^3, the antipode of Y is at distance $\pi - d(X, Y) < r_1$ from X and $\pi \geq r_2$ from Y. □

Proposition 6. *Let M be conformal to $(\eta_-, \eta_+) \times \Sigma$ with metric $-d\eta^2 + d\Sigma^2$, where Σ is $\mathbb{R}^3$, H^3, or the unit S^3 with $\eta_+ - \eta_- \leq \pi$. Then $\mathrm{Det}(b) = I^-_{\mathrm{s}}(b)$ for every b.*

Proof. Use $t = \eta$. For $x = (\eta_x, X)$, $b = (\eta_b, Y)$: $x \ll b$ iff $d(X, Y) < \eta_b - \eta_x$ (on S^3, $\eta_b - \eta_x < \pi$). For $\eta_- < c < \eta_x$, $J^-(x) \cap S_c = \bar{B}(X, \eta_x - c)$ and $I^-_{\mathrm{s}}(b) \cap S_c = B(Y, \eta_b - c)$, with $\eta_b - c < \pi$ on S^3. By Lemma 5 the inclusion holds iff $d(X, Y) < \eta_b - \eta_x$, i.e. iff $x \ll b$. Apply Theorem 3. □

This covers Minkowski spacetime, spatially flat and open FLRW universes with a big bang (conformal to $\eta > 0$), the closed radiation universe ($a \propto \sin\eta$, $\eta \in (0, \pi)$) and global de Sitter spacetime ($\Sigma = S^3$, $\eta \in (-\pi/2, \pi/2)$) [12]. In none of them is anything outside an event's chronological past guaranteed from it. In Minkowski spacetime and the spatially flat universes this bound is attained by a natural field.

Proposition 7. *Let $M = (\eta_-, \eta_+) \times \mathbb{R}^3$ with $g = a(\eta)^2(-d\eta^2 + d\vec{X}^2)$, $a > 0$ smooth, and let $\mathcal{L}$ be the smooth solutions of the conformally coupled scalar equation $\Box_g\phi - \frac{1}{6}R\phi = 0$. Then $F_{\mathcal{L}}(b) = J^-(b)$ for every b.*

Proof. $J^-(b) \subseteq F_{\mathcal{L}}(b)$ by continuity, as in Remark 2. Conversely, ϕ is a solution iff $\psi = a\phi$ solves the flat wave equation $\partial_\eta^2\psi = \Delta\psi$ on M [21, app. D]. Let $b = (\eta_b, \vec{Y})$ and $x = (\eta_x, \vec{X}) \notin J^-(b)$, so $\eta_x - \eta_b + |\vec{X} - \vec{Y}| > 0$, since the causal curves of M are those of the conformal Minkowski metric and the slab is causally convex in Minkowski spacetime. Choose a unit vector $\hat{k}$ with $\hat{k}\cdot(\vec{X} - \vec{Y}) = -|\vec{X} - \vec{Y}|$, a smooth f vanishing on $(-\infty, 0]$ and positive on $(0, \infty)$, and put $\psi = f(u)$ with $u = \eta - \eta_b - \hat{k}\cdot(\vec{Z} - \vec{Y})$ at $(\eta, \vec{Z})$. This solves the wave equation. On $I_{\mathrm{s}}^-(b)$, $\eta - \eta_b < -|\vec{Z} - \vec{Y}|$, so $u < 0$ and $\psi = 0$; at x, $u > 0$ and $\psi \neq 0$. So ψ/a and 0 are solutions that agree on $I_{\mathrm{s}}^-(b)$ but not at x. □

Proposition 8. *Let M be conformal to $(\eta_-, \eta_+) \times S^3$ with $\eta_+ - \eta_- > \pi$. Then $\mathrm{Det}(b) = I_{\mathrm{s}}^-(b)$ if $\eta_b \leq \eta_- + \pi$, and $\mathrm{Det}(b) = M$ if $\eta_b > \eta_- + \pi$. In the Einstein static universe ($\eta_\pm = \pm\infty$), $\mathrm{Det}(b) = M$ for every b.*

Proof. If $\eta_b > \eta_- + \pi$, take $c \in (\eta_-, \eta_b - \pi)$: $I_{\mathrm{s}}^-(b) \cap S_c$ is an open ball of radius greater than π, hence all of S_c, and Corollary 4 applies. If $\eta_b \leq \eta_- + \pi$, every $c > \eta_-$ has $\eta_b - c < \pi$, and the proof of Proposition 6 applies. □

For the closed dust FLRW model, $a(\eta) = \frac{1}{2}a_m(1 - \cos\eta)$ with $\eta \in (0, 2\pi)$, the threshold is $\eta = \pi$, the moment of maximum expansion. That the particle horizon of the closed dust model disappears at maximum expansion, so that an observer can then in principle see the whole universe, is known [3, 10]; the Einstein static case is Geroch's [8, Fig. 5]. What Proposition 8 adds is the consequence for determination: with Corollary 4, once the particle horizon has vanished ($\eta_b > \pi$), everything is guaranteed from b, so that every law with the domain-of-dependence property fixes everything from what is real from b. This fits Manchak's theorem that if some event of b's timelike future lies in Geroch's domain of prediction of b, the spacetime admits a compact spacelike slice [14, p. 319]. That theorem concerns the geometric notion. The flow law of Remark 2 shows that it has no analogue for what a particular law fixes: in Minkowski spacetime, which has no compact slice, that law fixes all of b's timelike future. Manchak also records the view that "Maxwell's equations constrain some spacetimes in such a way as to allow for prediction" [14, p. 321 and n. 15], so the law-relative notion has antecedents. We have not been able to consult the two works he cites there, Ellis and Sciama (1972, pp. 41–42) and Earman (1995, pp. 125–130), and Theorem 10 below may overlap with them.

Before maximum expansion a particular field can fix more than is guaranteed. Light leaving a point of the closed model refocuses at its antipode after conformal time π, and the rescaled conformal scalar $\psi = a\phi$ returns there with its sign reversed.

Proposition 9. *Let M be the closed dust model, $g = a(\eta)^2(-d\eta^2 + d\Omega_3^2)$ with $a = \frac{1}{2}a_m(1 - \cos\eta)$, $\eta \in (0, 2\pi)$, and $\mathcal{L}$ as in Proposition 7. Let $A(\eta, X) = (\eta + \pi, -X)$, where $-X$ is the antipode of X. Then $J^-(b) \cup \big(A(J^-(b)) \cap M\big) \subseteq F_{\mathcal{L}}(b)$. The set $A(J^-(b)) \cap M$ is disjoint from $I_{\mathrm{s}}^+(b)$, and if $\eta_b \leq \pi$ it is nonempty and disjoint from $\mathrm{Det}(b)$.*

Proof. Here $\psi = a\phi$ solves the conformally invariant equation of the Einstein static universe, $\partial_\eta^2\psi = \Delta\psi - \psi$ on the unit S^3 [21, app. D], and by global well-posedness extends uniquely to $\mathbb{R}\times S^3$. Expanding in spherical harmonics of degree n, which have frequency $n+1$ and parity $(-1)^n$ under the antipodal map, each mode, and so every smooth solution, satisfies $\psi(\eta+\pi,-X) = -\psi(\eta,X)$. Two solutions that agree on $I_{\mathrm{s}}^-(b)$ agree, by continuity, on $J^-(b)$, and hence on $A(J^-(b))\cap M$. If $(\eta,X)\in J^-(b)$ and $\eta+\pi<2\pi$, then $d(X,Y)\le\eta_b-\eta$, so $d(-X,Y)=\pi-d(X,Y)\ge(\eta+\pi)-\eta_b$; thus $A(\eta,X)\notin I_{\mathrm{s}}^+(b)$. If $\eta_b\le\pi$, then $\mathrm{Det}(b) = I_{\mathrm{s}}^-(b)$ by Proposition 8, which lies in $\{\eta<\eta_b\}$, while $A(J^-(b))$ lies in $\{\eta>\pi\}$ and meets M, for instance at $A(\eta_b/2,Y)$ when $b=(\eta_b,Y)$. □

For $\eta_b\le\pi$, the echo events are unreal from b and fixed from b, but they lie beside b or on its future light cone, not in its timelike future. The field fixes more than the echo, including part of b's timelike future. Let $\Diamond_b = \{(\eta,X)\in\mathbb{R}\times S^3 : d(X,Y)+|\eta|<\eta_b\}$, the double cone in the Einstein static universe between b and its mirror image $(-\eta_b,Y)$, and $U_b = \Diamond_b\cup A\Diamond_b\cup A^2\Diamond_b$.

Theorem 10. *In the setting of Proposition 9, let $0<\eta_b<\pi$. Then $F_{\mathcal{L}}(b) = M\cap\overline{U_b}$. In particular $F_{\mathcal{L}}(b)$ contains the nonempty set $\{(\eta,X)\in M : d(X,Y)\le\eta+\eta_b-2\pi\}$, which lies in $I_{\mathrm{s}}^+(b)$.*

Proof. For data (f,g) at $\eta=0$, the Einstein static solution is $\psi(\eta,X) = \partial_\eta[\sin\eta\, M_f(X,\eta)] + \sin\eta\, M_g(X,\eta)$, where $M_h(X,r)$ is the mean of h over the geodesic sphere of radius r about X; one checks this on each mode, using that the zonal function of degree n on S^3 is $\sin((n+1)r)/((n+1)\sin r)$. So $\psi(x)$ depends on the data only near the sphere $\Sigma_x = S(X,\eta_x)$, which is the point $-X$ when $\eta_x=\pi$.

Sufficiency. Solutions agreeing on $I_{\mathrm{s}}^-(b)$ agree on the ball $B(Y,\eta_b-\epsilon)$ in the slice $\eta=\epsilon$, hence on its domain of dependence in the Einstein static universe; the union over $\epsilon>0$ is $\Diamond_b$. The identities $\psi\circ A = -\psi$ and $\psi\circ A^2 = \psi$ carry the agreement to $A\Diamond_b$ and $A^2\Diamond_b$. Moreover, $F_{\mathcal{L}}(b)$ is relatively closed in M: it is the intersection of the zero sets of all smooth difference solutions vanishing on $I_{\mathrm{s}}^-(b)$. Since M is open in the Einstein cylinder, taking the relative closure gives $M\cap\overline{U_b}\subseteq F_{\mathcal{L}}(b)$.

Necessity. Let $x\in M\setminus\overline{U_b}$. Replacing x by $A^{-1}x$ if $\eta_x>\pi$, assume $\eta_x\le\pi$. Let $C = \bar{B}(Y,\eta_b)$. Data supported off C give a solution vanishing on $I_{\mathrm{s}}^-(b)$: if $z\in I_{\mathrm{s}}^-(b)$ lay in its support, then $w\le z\ll b$ for some w in the support of the data, so $d(w,Y)<\eta_b$. If $\eta_x=\pi$, then $d(-X,Y)>\eta_b$, and data $(f,0)$ with f a bump at $-X$ off C give $\psi(x) = -f(-X)\neq 0$. If $\eta_x<\pi$, then $\Sigma_x\not\subseteq C$: otherwise the connected set $S^3\setminus C$ lies in one of the two open balls bounded by Σ_x, so the other closed ball lies in C, which by the argument of Lemma 5 means $d(X,Y)+\eta_x\le\eta_b$ or $d(-X,Y)+\pi-\eta_x\le\eta_b$, that is, $x\in\overline{\Diamond_b}\cup\overline{A\Diamond_b}$. Take $p\in\Sigma_x\setminus C$ and data $(0,g)$ with $g\ge 0$ supported near p off C and $g(p)>0$; then $\psi(x) = \sin\eta_x\, M_g(X,\eta_x) > 0$.

Finally, if $d(X,Y)\le\eta+\eta_b-2\pi$, then $d(X,Y)<\eta-\eta_b$ because $\eta_b<\pi$, so $(\eta,X)\in I_{\mathrm{s}}^+(b)$; the set contains (η,Y) for $\eta\ge 2\pi-\eta_b$. □

On b's comoving geodesic $X=Y$, the field is fixed from b exactly for $\eta\le\eta_b$ and for $\eta\ge 2\pi-\eta_b$, and open in between: the end is fixed before the middle. The reason is that the past light cone of an event just before the crunch wraps around the universe and refocuses near Y at the big bang, inside what is real from b. The same sphere argument shows that, up to boundary, $M\cap U_b$ is the set of events all of whose past-inextendible null geodesics meet $I_{\mathrm{s}}^-(b)$, so for this field, which propagates on light rays alone, the null geodesics decide what is fixed. The same argument applies to global de Sitter spacetime, conformal to the slab $(-\pi/2,\pi/2)$, where the field is fixed from b on the antipodal comoving geodesic for all $\eta>-\eta_b$, although no signal from that geodesic ever reaches the comoving geodesic of b. So the geometric threshold at maximum expansion marks when everything is guaranteed, not when this field first fixes anything beyond the past.

For a law whose fixed set itself changes at the threshold, take the comoving flow $u = a^{-1}\partial_\eta$ of the closed dust model with the equations of Remark 2; here u is the dust's own four-velocity, so the law adds no structure the model lacks. The comoving line through X meets $I_{\mathrm{s}}^-(b)$ iff $d(X, Y) < \eta_b$, so, by continuity in X and by choosing data supported off the closed ball, $F_{\mathcal{L}}(b) = (0, 2\pi) \times \bar{B}(Y, \eta_b)$. This is a proper subset of M, containing events in b's timelike future, if $\eta_b < \pi$, and all of M if $\eta_b \geq \pi$. Under this law the answer to "is the future fixed from here?" differs between events before and after maximum expansion.

Whether in the closed models some local law fixes no more than $\mathrm{Det}(b) \cup J^-(b)$ we leave open.

4 Unreality and openness

Put together with the companion theorem, these results separate three properties of an event x relative to a standpoint b:

- *real from* b: x is in b's fragment, which a local becoming theory fixes to lie between $I_{\mathrm{s}}^-(b)$ and $J^-(b)$;
- *guaranteed from* b: $x \in \mathrm{Det}(b)$, so that every law with the domain-of-dependence property fixes x from what is real from b;
- *fixed from* b *under given laws*: $x \in F_{\mathcal{L}}(b)$.

Every real event is fixed under any continuous law with the domain-of-dependence property, and every guaranteed event is fixed under each such law. The standard models exhibit two geometric regimes: in the flat, open, radiation and de Sitter models nothing beyond the fragment is guaranteed; in the closed dust model after maximum expansion, and in the Einstein static universe, everything is. So:

1. *Unreality does not entail openness.* An event can be unreal from b and guaranteed from b: in the closed dust model, every future event is so from every event after maximum expansion. Under particular laws, events can be unreal and fixed without being guaranteed: the conformal scalar in the closed dust model before maximum expansion fixes the echo and part of b's timelike future near the crunch (Theorem 10), and the flow law of Remark 2 fixes all of b's timelike future even in Minkowski spacetime. This is the relativistic form of the growing block combined with determinism, and on our reading it is also where Rakíc's intuition belongs [16]: her sea-fight is, at most, fixed from the observer, not real from her.

2. *Determinism does not entail fixedness from each event.* Under deterministic laws the whole spacetime is fixed by data on any Cauchy surface, but no single event's past contains a Cauchy surface in the flat, open, radiation or de Sitter models. For the conformal scalar in Minkowski spacetime and the flat universes, what is fixed from b is exactly $J^-(b)$ (Proposition 7): from each standpoint the future is unfixed as well as unreal. In the open, closed radiation and de Sitter models we have shown only that nothing beyond the past is guaranteed.

3. *How much of the unreal future is fixed depends on two things.* The causal structure of the spacetime decides what is guaranteed, and the guarantee can change with cosmic epoch: in the closed dust model it is everything after maximum expansion and nothing beyond the past before it. The fields decide what is fixed beyond the guarantee; under the comoving flow law above, the answer to "is the future fixed from here?" differs between two events of the same spacetime. The causal structure of the given spacetime settles the first; it does not settle the second.

Three limits. First, this openness is relational and informational: an event outside $\mathrm{Det}(b)$ is still fixed by its own past, and a determinist eternalist can accept everything in this section. What the section shows is that the standpoint theorist's claim that the future is unreal from b neither secures nor excludes its being fixed from b; an ontically open future needs indeterministic laws, about which $\mathrm{Det}(b)$ and $F_{\mathcal{L}}(b)$ are silent. Second, the closed dust result concerns an idealized model with deterministic, classical dynamics; it is not a claim about our universe, which on current evidence is not of that kind. Third, Manchak shows that *genuine* prediction, required to hold in every inextendible spacetime into which the observer's causal past embeds isometrically, reaches no further than the boundary of that past, even under any local conditions on matter [14, pp. 320–321]; $\mathrm{Det}(b)$ and $F_{\mathcal{L}}(b)$ are nomological notions given M, not epistemic ones.

5 A remark on standpoint logic

Read $\Box A$ at b as "at every event $x \leq b$, A" on the frame (M, R) with bRx iff $x \leq b$. If A is a tenseless claim about what occurs at x, this is the reflexive past modality of Diodorean tense logic, and the standpoint reading of Fine and Lipman [7, 13] adds no formal content. R is reflexive and transitive, so S4 is valid, and in Minkowski spacetime the valid formulas are exactly those of S4.2 [9]. The axiom .2, $\Diamond\Box A \to \Box\Diamond A$, is valid at a world for all valuations iff any two of its successors have a common successor [1]. So .2 is valid at b iff the subframe generated by b is directed, that is, iff any two events in $J^-(b)$ have a common causal past.

In a model conformal to $(\eta_-, \eta_+)\times\mathbb{R}^3$, two events have a common past iff $|X-Y| < \eta_x+\eta_y-2\eta_-$, so if $\eta_- > -\infty$ the directedness fails in the past of every event and .2 is valid at no event. This is the particle-horizon structure of Rindler [17] in modal form, and it holds equally with or without an inflationary era of finite conformal duration. The horizon problem of cosmology [11] is therefore not the failure of .2, which occurs with and without inflation, but the failure of particular instances: pairs of points on our last-scattering surface, at comoving separation up to $2(\eta_0 - \eta_{\mathrm{ls}})$, have no common past unless $\eta_0 - \eta_{\mathrm{ls}} < \eta_{\mathrm{ls}} - \eta_-$, which fails without inflation and holds with enough of it. Two cautions. Under the view's persistence principle, on which what holds from an event holds from every later one, valuations are constrained along the future direction, and the frame property that matters may concern common futures, which these models have when $\eta_+ = \infty$; the logic of persistent valuations is a separate question. And Lewis and van der Schaaf study causal modal logics of general spacetimes [22]; we have not checked how their results bear on the past-directed frames used here. Whether S4 is exactly the logic of all past-directed globally hyperbolic frames we leave open.

6 Conclusion

The unreality of the future from an event, which local theories of relativistic becoming deliver, neither implies nor excludes its being fixed from that event. What the causal structure guarantees, that is, what every local deterministic law fixes by the domain-of-dependence property alone, is characterized by a slicing condition on Geroch's domain of prediction (Theorem 3) and computable from conformal diagrams; it is a lower bound on what any particular law fixes. In most standard models nothing beyond an event's past is guaranteed, and in Minkowski spacetime and the flat universes a conformal scalar field shows that nothing more need be fixed; in the closed dust model everything is guaranteed from every event after maximum expansion. Particular fields can fix more than is guaranteed: in the closed dust model before maximum expansion, a conformal scalar is fixed on part of an event's far future while its nearer future stays open, and with a preferred flow

a field is fixed everywhere even in Minkowski spacetime. The question whether the future is open from a standpoint thus divides into a question about the causal structure of the spacetime, which settles the guarantee, and a question about the fields, which that structure does not settle.

References

[1] P. Blackburn, M. de Rijke and Y. Venema, *Modal Logic* (Cambridge University Press, 2001).

[2] N. Belnap, Branching space-time, *Synthese* 92 (1992) 385–434, doi:10.1007/BF00414289.

[3] Yu. L. Bolotin and I. V. Tanatarov, A thousand problems in cosmology: horizons, arXiv:1310.6329 (2013).

[4] C. D. Broad, *Scientific Thought* (Kegan Paul, 1923).

[5] A. N. Bernal and M. Sánchez, Smoothness of time functions and the metric splitting of globally hyperbolic spacetimes, *Commun. Math. Phys.* 257 (2005) 43–50.

[6] R. Clifton and M. Hogarth, The definability of objective becoming in Minkowski spacetime, *Synthese* 103 (1995) 355–387.

[7] K. Fine, Tense and reality, in *Modality and Tense* (Oxford University Press, 2005) 261–320.

[8] R. Geroch, Prediction in general relativity, in J. Earman, C. Glymour and J. Stachel (eds.), *Foundations of Space-Time Theories*, Minnesota Studies in the Philosophy of Science 8 (University of Minnesota Press, 1977) 81–93.

[9] R. Goldblatt, Diodorean modality in Minkowski spacetime, *Studia Logica* 39 (1980) 219–236, doi:10.1007/BF00370321.

[10] Ø. Grøn and S. Johannesen, FRW universe models in conformally flat spacetime coordinates. III: universe models with positive spatial curvature, *Eur. Phys. J. Plus* 126 (2011) 30, arXiv:1103.4753.

[11] A. H. Guth, Inflationary universe: a possible solution to the horizon and flatness problems, *Phys. Rev. D* 23 (1981) 347–356.

[12] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, 1973).

[13] M. A. Lipman, Standpoints: a study of a metaphysical picture, *J. Phil.* 120 (2023) 117–138.

[14] J. B. Manchak, Is prediction possible in general relativity?, *Found. Phys.* 38 (2008) 317–321, doi:10.1007/s10701-008-9204-6.

[15] E. Minguzzi and M. Sánchez, The causal hierarchy of spacetimes, in *Recent Developments in Pseudo-Riemannian Geometry* (EMS, 2008) 299–358, arXiv:gr-qc/0609119.

[16] N. Rakić, Past, present, future, and special relativity, *Brit. J. Phil. Sci.* 48 (1997) 257–280.

[17] W. Rindler, Visual horizons in world-models, *Mon. Not. R. Astron. Soc.* 116 (1956) 662–677.

[18] H. Stein, On Einstein–Minkowski space–time, *J. Phil.* 65 (1968) 5–23.

[19] H. Stein, On relativity theory and openness of the future, *Phil. Sci.* 58 (1991) 147–167.

[20] A. Impellitteri, Stein's theorem in general relativity: locality, not symmetry, fixes what has become, preprint (2026).

[21] R. M. Wald, *General Relativity* (University of Chicago Press, 1984).

[22] M. Lewis and N. van der Schaaf, Some results on causal modalities in general spacetimes, arXiv:2601.14029v2 (2026).