

ELLIOTT

903

Volume 2: PROGRAMMING INFORMATION

Part 2: PROGRAM DESCRIPTIONS

Section 11: QEXP (B. 2)

Contents

	Page
Chapter 1: DESCRIPTION	
1. 1 Introduction	1
1. 1. 1 Purpose	1
1. 1. 2 Forms of Distribution	1
1. 1. 3 Method of Use	1
1. 1. 4 Accuracy	1
1. 2 Functions	1
1. 2. 1 Number Type	1
1. 2. 2 Exit and Entry	1
1. 2. 3 Identifiers	2
1. 3 Error Indications	2
1. 4 Method Used	2
1. 5 Store Used	2
1. 6 Time Taken	2

Chapter 1: DESCRIPTION

1. 1 INTRODUCTION

1. 1. 1 Purpose.

To calculate $\exp(2^p x)$

where

$$-1 \leq x < 0$$

$p \geq 0$, and p is integral.

1. 1. 2 Forms of Distribution.

The program is distributed as a machine code tape for input by T2 or SIR.

1. 1. 3 Method of Use.

The routine is assembled as a block of the user's program and used as a sub-routine. It may be run at any program level and in any store-module.

1. 1. 4 Accuracy.

The maximum error is 2^{-16} ($\approx .000015$).

1. 2 FUNCTIONS

1. 2. 1 Number Type.

x is treated as a pure fraction;
 p is treated as an integer.

The result is a pure fraction.

1. 2. 2 Exit and Entry.

On entry

x must be placed in the accumulator

and p must be placed in QEXP + 53 (53;N)
 p is not preserved by QEXP

On exit

the result is in the accumulator

and in QEXP + 54 (54;N)

x is in QEXP + 52 (52;N)

Entry is made by

(for assembly by SIR.) (for translation by T2,
where N is the number of the
block).

11 QEXP
8 QEXP + 1

11 0;N
8 1;N

1. 2. 3 Identifiers.

QEXP must be declared as a global identifier
in all blocks of the user's program which refer to it.

On the library tape, a mnemonic label and
identifier list are separated from the coding by several inches of blank
tape: the mnemonics must not be loaded into the tape reader if the tape
is to be translated by T2.

1. 3 ERROR INDICATIONS.

If $x \geq 0$ or $p < 0$ tape is output continuously.

1. 4 METHOD USED.

The program first calculates $\exp(x)$ by a Chebyshev
series.

$$\exp(x) = \sum_{n=0}^6 a_n x^n$$

The final result is calculated by repeated squaring of
 e^x .

1. 5 STORE USED.

55 locations and the appropriate B register.

1. 6 TIME TAKEN.

(3.7 + 0.26 p) milliseconds approx.