

Debiasing Word Embeddings

Andrew Maurer

July 26, 2018

Debiasing Word Embeddings

Andrew Maurer

July 26, 2018

Cohomology Rings and Geometry

Andrew Maurer

July 26, 2018

Plan

- ① Groups and Modules for Groups
- ② Cohomology
- ③ Realizability and Consequences

Finite groups

Finite groups

Recall that a finite group is a finite set G with a multiplication rule satisfying some axioms that make it behave like symmetry.

Finite groups

Recall that a finite group is a finite set G with a multiplication rule satisfying some axioms that make it behave like symmetry.

- $GL(V) =$ all linear isomorphisms of a vector space V/\mathbb{F}_q .

Finite groups

Recall that a finite group is a finite set G with a multiplication rule satisfying some axioms that make it behave like symmetry.

- $GL(V)$ = all linear isomorphisms of a vector space V/\mathbb{F}_q .
- S_n = all bijections $\{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$.

Finite groups

Recall that a finite group is a finite set G with a multiplication rule satisfying some axioms that make it behave like symmetry.

- $GL(V)$ = all linear isomorphisms of a vector space V/\mathbb{F}_q .
- S_n = all bijections $\{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$.
- $(\mathbb{Z}/p)^r$ = elementary Abelian group of rank r .

Finite groups

Recall that a finite group is a finite set G with a multiplication rule satisfying some axioms that make it behave like symmetry.

- $GL(V)$ = all linear isomorphisms of a vector space V/\mathbb{F}_q .
- S_n = all bijections $\{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$.
- $(\mathbb{Z}/p)^r$ = elementary Abelian group of rank r .
- Others

Modules

Modules

Groups are modeled on symmetries. A G -**module** is just a vector space where G acts as symmetries, i.e., linear automorphisms.

Modules

Groups are modeled on symmetries. A **G -module** is just a vector space where G acts as symmetries, i.e., linear automorphisms.

Namely, there's a map

$$\varphi : G \rightarrow \mathrm{GL}(V), g \mapsto \varphi_g$$

Modules

Groups are modeled on symmetries. A **G -module** is just a vector space where G acts as symmetries, i.e., linear automorphisms.

Namely, there's a map

$$\varphi : G \rightarrow \mathrm{GL}(V), g \mapsto \varphi_g$$

Examples:

Modules

Groups are modeled on symmetries. A **G -module** is just a vector space where G acts as symmetries, i.e., linear automorphisms.

Namely, there's a map

$$\varphi : G \rightarrow \text{GL}(V), g \mapsto \varphi_g$$

Examples:

- *Trivial Representation*: k is a G -module via the action $\varphi_g(v) = v$.

Modules

Groups are modeled on symmetries. A **G -module** is just a vector space where G acts as symmetries, i.e., linear automorphisms.

Namely, there's a map

$$\varphi : G \rightarrow \text{GL}(V), g \mapsto \varphi_g$$

Examples:

- *Trivial Representation*: k is a G -module via the action $\varphi_g(v) = v$.
- *Natural Representation*: V is a $\text{GL}(V)$ module.

Modules

Groups are modeled on symmetries. A **G -module** is just a vector space where G acts as symmetries, i.e., linear automorphisms.

Namely, there's a map

$$\varphi : G \rightarrow \text{GL}(V), g \mapsto \varphi_g$$

Examples:

- *Trivial Representation*: k is a G -module via the action $\varphi_g(v) = v$.
- *Natural Representation*: V is a $\text{GL}(V)$ module.
- *Permutation Representation*: The vector space with basis $\{e_1, \dots, e_n\}$ is a S_n -module by $\varphi_\sigma(e_i) = e_{\sigma(i)}$, and extend by linearity.

Modules

Groups are modeled on symmetries. A **G -module** is just a vector space where G acts as symmetries, i.e., linear automorphisms.

Namely, there's a map

$$\varphi : G \rightarrow \text{GL}(V), g \mapsto \varphi_g$$

Examples:

- *Trivial Representation*: k is a G -module via the action $\varphi_g(v) = v$.
- *Natural Representation*: V is a $\text{GL}(V)$ module.
- *Permutation Representation*: The vector space with basis $\{e_1, \dots, e_n\}$ is a S_n -module by $\varphi_\sigma(e_i) = e_{\sigma(i)}$, and extend by linearity.

G -modules as defined above are really just modules for a certain ring denoted kG .

Cohomology groups are Ext groups

Cohomology groups are Ext groups

The cohomology group $H^n(G; k)$ can be defined in many ways.

Cohomology groups are Ext groups

The cohomology group $H^n(G; k)$ can be defined in many ways.

Most concrete is as $H^n(G; k) \cong \text{Ext}_{kG}^n(k, M)$.

$$\text{Ext}_{kG}^n(k, M) = \{0 \rightarrow M \rightarrow E_1 \rightarrow \dots \rightarrow E_n \rightarrow k \rightarrow 0\} / \sim$$

i.e., as all n -fold extensions of k by M , modulo a certain equivalence relation.

Cohomology groups are Ext groups

The cohomology group $H^n(G; k)$ can be defined in many ways.

Most concrete is as $H^n(G; k) \cong \text{Ext}_{kG}^n(k, M)$.

$$\text{Ext}_{kG}^n(k, M) = \{0 \rightarrow M \rightarrow E_1 \rightarrow \dots \rightarrow E_n \rightarrow k \rightarrow 0\} / \sim$$

i.e., as all n -fold extensions of k by M , modulo a certain equivalence relation.

Addition in the cohomology group corresponds to Baer sum of extensions.

Cohomology groups are Ext groups

The cohomology group $H^n(G; k)$ can be defined in many ways.

Most concrete is as $H^n(G; k) \cong \text{Ext}_{kG}^n(k, M)$.

$$\text{Ext}_{kG}^n(k, M) = \{0 \rightarrow M \rightarrow E_1 \rightarrow \dots \rightarrow E_n \rightarrow k \rightarrow 0\} / \sim$$

i.e., as all n -fold extensions of k by M , modulo a certain equivalence relation.

Addition in the cohomology group corresponds to Baer sum of extensions. This is complicated.

Yoneda splicing

Yoneda splicing

Realizing cohomology groups $H^n(G; k)$ in this way means we can multiply two sequences together.

$$\mathrm{Ext}_{kG}^n(k, k) \times \mathrm{Ext}_{kG}^m(k, k) \rightarrow \mathrm{Ext}_{kG}^{m+n}(k, k)$$

Yoneda splicing

Realizing cohomology groups $H^n(G; k)$ in this way means we can multiply two sequences together.

$$\mathrm{Ext}_{kG}^n(k, k) \times \mathrm{Ext}_{kG}^m(k, k) \rightarrow \mathrm{Ext}_{kG}^{m+n}(k, k)$$

$$\begin{array}{ccccccc} \dots & \longrightarrow & E_n & \longrightarrow & k & \longrightarrow & 0 \\ & & & & \parallel & & \\ & & & & & \searrow & \\ & 0 & \longrightarrow & k & \longrightarrow & F_1 & \longrightarrow \dots \end{array}$$

Yoneda splicing

Realizing cohomology groups $H^n(G; k)$ in this way means we can multiply two sequences together.

$$\mathrm{Ext}_{kG}^n(k, k) \times \mathrm{Ext}_{kG}^m(k, k) \rightarrow \mathrm{Ext}_{kG}^{m+n}(k, k)$$

$$\begin{array}{ccccccc} \dots & \longrightarrow & E_n & \longrightarrow & k & \longrightarrow & 0 \\ & & & & \parallel & & \\ & & & & & \searrow & \\ & 0 & \longrightarrow & k & \longrightarrow & F_1 & \longrightarrow \dots \end{array}$$

And the new sequence has the form

$$0 \rightarrow k \rightarrow \underbrace{E_1 \rightarrow \dots \rightarrow E_n \rightarrow F_1 \rightarrow \dots \rightarrow F_m}_{m+n} \rightarrow k \rightarrow 0$$

$H^\bullet(G; k) = \text{Ext}_{kG}^\bullet(k, k)$ is a ring

$H^\bullet(G; k) = \text{Ext}_{kG}^\bullet(k, k)$ is a ring

Under this $\text{Ext}_{kG}^n(k, k) \times \text{Ext}_{kG}^m(k, k) \rightarrow \text{Ext}_{kG}^{n+m}(k, k)$ association,

$$\text{Ext}_{kG}^\bullet(k, k) = \bigoplus \text{Ext}_{kG}^n(k, k)$$

becomes a graded ring.

$H^\bullet(G; k) = \text{Ext}_{kG}^\bullet(k, k)$ is a ring

Under this $\text{Ext}_{kG}^n(k, k) \times \text{Ext}_{kG}^m(k, k) \rightarrow \text{Ext}_{kG}^{n+m}(k, k)$ association,

$$\text{Ext}_{kG}^\bullet(k, k) = \bigoplus \text{Ext}_{kG}^n(k, k)$$

becomes a graded ring. What can we say about it?

$H^\bullet(G; k) = \text{Ext}_{kG}^\bullet(k, k)$ is a ring

Under this $\text{Ext}_{kG}^n(k, k) \times \text{Ext}_{kG}^m(k, k) \rightarrow \text{Ext}_{kG}^{n+m}(k, k)$ association,

$$\text{Ext}_{kG}^\bullet(k, k) = \bigoplus \text{Ext}_{kG}^n(k, k)$$

becomes a graded ring. What can we say about it?

- It is *graded-commutative*, i.e., $\alpha \cdot \beta = (-1)^{\tilde{\alpha} \cdot \tilde{\beta}} \beta \cdot \alpha$.

$H^\bullet(G; k) = \text{Ext}_{kG}^\bullet(k, k)$ is a ring

Under this $\text{Ext}_{kG}^n(k, k) \times \text{Ext}_{kG}^m(k, k) \rightarrow \text{Ext}_{kG}^{n+m}(k, k)$ association,

$$\text{Ext}_{kG}^\bullet(k, k) = \bigoplus \text{Ext}_{kG}^n(k, k)$$

becomes a graded ring. What can we say about it?

- It is *graded-commutative*, i.e., $\alpha \cdot \beta = (-1)^{\tilde{\alpha} \cdot \tilde{\beta}} \beta \cdot \alpha$.
- (Evens 1961) It is finitely-generated over $\text{Ext}_{kG}^0(k, k) \cong k$.

$H^\bullet(G; k) = \text{Ext}_{kG}^\bullet(k, k)$ is a ring

Under this $\text{Ext}_{kG}^n(k, k) \times \text{Ext}_{kG}^m(k, k) \rightarrow \text{Ext}_{kG}^{n+m}(k, k)$ association,

$$\text{Ext}_{kG}^\bullet(k, k) = \bigoplus \text{Ext}_{kG}^n(k, k)$$

becomes a graded ring. What can we say about it?

- It is *graded-commutative*, i.e., $\alpha \cdot \beta = (-1)^{\bar{\alpha} \cdot \bar{\beta}} \beta \cdot \alpha$.
- (Evens 1961) It is finitely-generated over $\text{Ext}_{kG}^0(k, k) \cong k$.

Define a commutative, finitely generated ring:

$$H^c(G; k) = \begin{cases} H^\bullet(G; k) & \text{if } p = 2 \\ \bigoplus H^{2n}(G; k) & \text{if } p > 2 \end{cases}$$

One more thing about rings

One more thing about rings

$\text{Ext}_{kG}^\bullet(M, M)$ is a ring in the exact same way. It is far less nice, and usually not finitely generated.

One more thing about rings

$\text{Ext}_{kG}^\bullet(M, M)$ is a ring in the exact same way. It is far less nice, and usually not finitely generated.

$\text{Ext}_{kG}^\bullet(M, M)$ is a graded module for $H^\bullet(G; k)$ (or $H^c(G; k)$) using the tensor product.

One more thing about rings

$\text{Ext}_{kG}^\bullet(M, M)$ is a ring in the exact same way. It is far less nice, and usually not finitely generated.

$\text{Ext}_{kG}^\bullet(M, M)$ is a graded module for $H^\bullet(G; k)$ (or $H^c(G; k)$) using the tensor product.

So $\text{Ext}_{kG}^\bullet(M, M)$ has an annihilator $I_M \trianglelefteq H^\bullet(G; k)$ which is a homogeneous ideal.

One more thing about rings

$\text{Ext}_{kG}^\bullet(M, M)$ is a ring in the exact same way. It is far less nice, and usually not finitely generated.

$\text{Ext}_{kG}^\bullet(M, M)$ is a graded module for $H^\bullet(G; k)$ (or $H^c(G; k)$) using the tensor product.

So $\text{Ext}_{kG}^\bullet(M, M)$ has an annihilator $I_M \trianglelefteq H^\bullet(G; k)$ which is a homogeneous ideal.

$$I_M = \{x \in H^\bullet(G; k) \mid x \cdot \text{Ext}_{kG}^\bullet(M, M) = 0\}.$$

Cohomology varieties and support varieties

Cohomology varieties and support varieties

To every finitely generated ring R , there is a geometric space $\operatorname{Spec} R$ whose functions are R .

Cohomology varieties and support varieties

To every finitely generated ring R , there is a geometric space $\operatorname{Spec} R$ whose functions are R .

The main example is if $R = k[x_1, \dots, x_n]$, then $\operatorname{Spec} R \cong k^n$, with some “special” points added.

Cohomology varieties and support varieties

To every finitely generated ring R , there is a geometric space $\operatorname{Spec} R$ whose functions are R .

The main example is if $R = k[x_1, \dots, x_n]$, then $\operatorname{Spec} R \cong k^n$, with some “special” points added.

Define $\mathcal{V}_G(k) = \operatorname{Spec}(H^c(G; k))$. This is the *cohomology variety* of G/k .

Cohomology varieties and support varieties

To every finitely generated ring R , there is a geometric space $\operatorname{Spec} R$ whose functions are R .

The main example is if $R = k[x_1, \dots, x_n]$, then $\operatorname{Spec} R \cong k^n$, with some “special” points added.

Define $\mathcal{V}_G(k) = \operatorname{Spec}(H^c(G; k))$. This is the *cohomology variety* of G/k .

For a G -module M ,

$$\mathcal{V}_G(M) = \mathcal{Z}(I_M) \subseteq \mathcal{V}_G(k)$$

is the *support variety* of M .

Geometry via Representation Theory

Geometry via Representation Theory

Goal

Describe the representation theory of G in terms of the geometry of $\mathcal{V}_G(k)$

Geometry via Representation Theory

Goal

Describe the representation theory of G in terms of the geometry of $\mathcal{V}_G(k)$

Theorem

Let M_1 and M_2 be G -modules.

Geometry via Representation Theory

Goal

Describe the representation theory of G in terms of the geometry of $\mathcal{V}_G(k)$

Theorem

Let M_1 and M_2 be G -modules.

- $\mathcal{V}_G(M_1) = \{0\}$ if M_1 is projective.

Geometry via Representation Theory

Goal

Describe the representation theory of G in terms of the geometry of $\mathcal{V}_G(k)$

Theorem

Let M_1 and M_2 be G -modules.

- $\mathcal{V}_G(M_1) = \{0\}$ if M_1 is projective.
- $\mathcal{V}_G(M_1 \oplus M_2) = \mathcal{V}_G(M_1) \cup \mathcal{V}_G(M_2)$.

Geometry via Representation Theory

Goal

Describe the representation theory of G in terms of the geometry of $\mathcal{V}_G(k)$

Theorem

Let M_1 and M_2 be G -modules.

- $\mathcal{V}_G(M_1) = \{0\}$ if M_1 is projective.
- $\mathcal{V}_G(M_1 \oplus M_2) = \mathcal{V}_G(M_1) \cup \mathcal{V}_G(M_2)$.
- $\mathcal{V}_G(M_1 \otimes M_2) = \mathcal{V}_G(M_1) \cap \mathcal{V}_G(M_2)$.

Geometry via Representation Theory

Goal

Describe the representation theory of G in terms of the geometry of $\mathcal{V}_G(k)$

Theorem

Let M_1 and M_2 be G -modules.

- $\mathcal{V}_G(M_1) = \{0\}$ if M_1 is projective.
- $\mathcal{V}_G(M_1 \oplus M_2) = \mathcal{V}_G(M_1) \cup \mathcal{V}_G(M_2)$.
- $\mathcal{V}_G(M_1 \otimes M_2) = \mathcal{V}_G(M_1) \cap \mathcal{V}_G(M_2)$.

But these are statements about the geometry of $\mathcal{V}_G(k)$. Can we figure out the representation theory of G by studying varieties $\mathcal{V}_G(M)$?

Carlson Modules

Each nonzero element $\zeta \in H^\bullet(G; k)$ of the cohomology ring determines a surjective homomorphism $\Omega^n(k) \rightarrow k$

Each nonzero element $\zeta \in H^\bullet(G; k)$ of the cohomology ring determines a surjective homomorphism $\Omega^n(k) \rightarrow k$, which can be fit into a short exact sequence

$$0 \rightarrow L_\zeta \rightarrow \Omega^n(k) \rightarrow k \rightarrow 0.$$

Each nonzero element $\zeta \in H^\bullet(G; k)$ of the cohomology ring determines a surjective homomorphism $\Omega^n(k) \rightarrow k$, which can be fit into a short exact sequence

$$0 \rightarrow L_\zeta \rightarrow \Omega^n(k) \rightarrow k \rightarrow 0.$$

Theorem

$$\mathcal{V}_G(L_\zeta) = \mathcal{Z}(\zeta)$$

Rehash previous slide

Rehash previous slide

On the one

hand:

Rehash previous slide

On the one

Each element $\zeta \in H^\bullet(G; k)$ determines a subvariety of
 $\mathrm{Spec} H^\bullet(G; k) = \mathcal{V}_G(k).$

hand:

Rehash previous slide

On the one

Each element $\zeta \in H^\bullet(G; k)$ determines a subvariety of
 $\mathrm{Spec} H^\bullet(G; k) = \mathcal{V}_G(k).$

hand:

On the other

hand

Rehash previous slide

On the one

Each element $\zeta \in H^\bullet(G; k)$ determines a subvariety of $\text{Spec } H^\bullet(G; k) = \mathcal{V}_G(k)$.

hand:

On the other

Each element $\zeta \in H^\bullet(G; k)$ defines a morphism $\Omega^n(k) \rightarrow k$, which has a kernel L_ζ , which determines a subvariety $\mathcal{V}_G(L_\zeta) \subseteq \mathcal{V}_G(k)$.

hand

Rehash previous slide

On the one

Each element $\zeta \in H^\bullet(G; k)$ determines a subvariety of $\text{Spec } H^\bullet(G; k) = \mathcal{V}_G(k)$.

hand:

On the other

Each element $\zeta \in H^\bullet(G; k)$ defines a morphism $\Omega^n(k) \rightarrow k$, which has a kernel L_ζ , which determines a subvariety $\mathcal{V}_G(L_\zeta) \subseteq \mathcal{V}_G(k)$.

hand

And these two processes yield the same subvariety.

Realization

Theorem

Every variety that could arise as the support variety of a module, does arise as the support variety of a module.

Theorem

Every variety that could arise as the support variety of a module, does arise as the support variety of a module.

- Closed

Theorem

Every variety that could arise as the support variety of a module, does arise as the support variety of a module.

- Closed
- Conical

Theorem

Every variety that could arise as the support variety of a module, does arise as the support variety of a module.

- Closed
- Conical

Take $X \subseteq \mathcal{V}_G(k)$ to be a closed conical subvariety:

Theorem

Every variety that could arise as the support variety of a module, does arise as the support variety of a module.

- Closed
- Conical

Take $X \subseteq \mathcal{V}_G(k)$ to be a closed conical subvariety:

X

Theorem

Every variety that could arise as the support variety of a module, does arise as the support variety of a module.

- Closed
- Conical

Take $X \subseteq \mathcal{V}_G(k)$ to be a closed conical subvariety:

$$X = \bigcap_{i=1}^n \mathcal{Z}(\zeta_i)$$

Theorem

Every variety that could arise as the support variety of a module, does arise as the support variety of a module.

- Closed
- Conical

Take $X \subseteq \mathcal{V}_G(k)$ to be a closed conical subvariety:

$$X = \bigcap_{i=1}^n \mathcal{Z}(\zeta_i) = \bigcap_{i=1}^n \mathcal{V}_G(L_{\zeta_i})$$

Theorem

Every variety that could arise as the support variety of a module, does arise as the support variety of a module.

- Closed
- Conical

Take $X \subseteq \mathcal{V}_G(k)$ to be a closed conical subvariety:

$$X = \bigcap_{i=1}^n \mathcal{Z}(\zeta_i) = \bigcap_{i=1}^n \mathcal{V}_G(L_{\zeta_i}) = \mathcal{V}_G \left(\bigotimes_{i=1}^n L_{\zeta_i} \right).$$

Partial converse to direct sum theorem

Theorem

If M is a G -module, and $\mathcal{V}_G(M) = V_1 \cup V_2$, with $V_1 \cap V_2 = \{0\}$, then there are modules with $V_1 = \mathcal{V}_G(M_1)$ and $V_2 = \mathcal{V}_G(M_2)$ such that $M = M_1 \oplus M_2$ and .

Thank You