



Simple Linear Regression II


Identify relationships For each of the six plots, identify the strength of the relationship (e.g. weak, moderate, or strong) in the data and whether fitting a linear model would be reasonable.




(1)




(2)




(3)



(4)



(5)




(6)

Finding the best line


Calculating the least-squares estimates

Estimating β_1




We use s_x , s_y , and R to calculate b_1 .

Estimating β_1



We use s_x , s_y , and R to calculate b_1 .

Estimating β_0



If the line of best fit *must* pass through (\bar{x}, \bar{y}) , what is b_0 ?

Estimating β_0 , cont.

Since (11.35, 86.01) is on the line, the following relationship holds.

$$86.01 = b_0 - 0.9(11.35)$$

Then just solve for b_0 .

$$b_0 = 86.01 + 0.9(11.35) = 96.22$$

More generally:

$$b_0 = \bar{y} - b_1\bar{x}$$

Estimation in R

```
m1 <- lm(Graduates ~ Poverty, data = poverty)
summary(m1)
```

```
##
## Call:
## lm(formula = Graduates ~ Poverty, data = poverty)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -5.954  -1.820   0.544   1.515   6.199
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   96.202     1.343    71.65  < 2e-16 ***
## Poverty      -0.898     0.114   -7.86  3.1e-10 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 2.5 on 49 degrees of freedom
```

The `lm` object

```
attributes(m1)
```

```
## $names
##  [1] "coefficients"  "residuals"      "effects"        "
##  [7] "qr"            "df.residual"    "xlevels"        "
##
## $class
##  [1] "lm"
```

```
m1$coef
```

```
## (Intercept)      Poverty
##          96.202      -0.898
```

```
m1$fit
```

Interpretation of b_1

The **slope** describes the estimated difference in the y variable if the explanatory variable x for a case happened to be one unit larger.

```
m1$coef[2]
```

```
## Poverty  
## -0.898
```

For each additional percentage point of people living below the poverty level, we expect a state to have a proportion of high school graduates that is 0.898 lower.

Be Cautious: if it is observational data, you do not have evidence of a *causal link*, but of an association, which still can be used for prediction.

Interpretation of b_0

The **intercept** is the estimated y value that will be taken by a case with an x value of zero.

```
m1$coef[1]
```

```
## (Intercept)  
##          96.2
```


While necessary for prediction, the intercept often has no meaningful interpretation.

Visualizing the residuals

Residual plot

```
m1 <- lm(runs ~ at_bats, data = mlb11)
ggplot(m1, aes(x = .fitted, y = .resid)) +
  geom_point() +
  geom_hline(yintercept = 0,
             linetype = "dashed") +
  xlab("Fitted values") +
  ylab("Residuals")
```


Residual plot



Distribution of the residuals

```
ggplot(m1, aes(x = .resid)) +  
  geom_histogram(binwidth = 25) +  
  xlab("Residuals")  
  
ggplot(m1, aes(sample = .resid)) +  
  geom_point(stat = "qq")
```


Distribution of the residuals



QQ plot

