NeuroScience Survey ID #1634768030

Requested for:
Name: **Dominic Hebert**
Sex: **male**
Age: **24**
State: **California**

Fatigue Evalution

Your Score 55

Your overall fatigue level is **55**, which indicates your fatigue level is **poor**.

Fatigue area	Description	Rating
General Fatigue	General fatigue measures your overall fatigue and decreased ability to function[21]	★★★
Physical Fatigue	Physical fatigue measures physical sensations related to your fatigue[21]	★★★
Activity Fatigue	Activity measures the influence of physical and psychological factors on your level of activity[21]	★
Motivation Fatigue	Motivation measures your lack of motivation for starting activity[21]	★★
Mental Fatigue	Mental fatigue measures your cognitive functioning and difficulty concentrating or focusing[21]	★

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Fatigue Product Recommendations

Alpha GABA PM

Researched blend of botanicals and 400 mg of L-theanine shown to induce calming brain waves and naturally improve sleep during times of stress*

- ✓ Researched blend of botanicals and 400 mg of L-theanine shown to induce calming brain waves and naturally improve sleep during times of stress*
- ✓ Sleep quality has been shown to be inversely correlated with fatigue25
- ✓ Alpha GABA PM was recommended because you indicated you struggle with general fatigue

Suggested use:
Take 1-2 capsules at bedtime or as directed by your healthcare provider. Do not exceed suggested use.
If you are pregnant or nursing, consult your healthcare provider before use.

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

ExcitaPlus

Contains non-glandular ingredients important for adrenal health and reducing stress-induced fatigue.* Provides over three times the L-tyrosine and six times the Rhodiola per capsule compared to AdreCor.

- ✓ Contains ingredients important for adrenal health and reducing stress-induced fatigue*
- ✓ Contains ingredients to support the body's production of dopamine, a neurotransmitter implicated in motivation and Rhodiola, a botanical shown to reduce mental fatigue22,23
- ✓ ExcitaPlus was recommended because you indicated your struggle with general fatigue, mental fatigue, and motivation

Suggested use:
Take 1-2 capsules one to two times daily or as directed by your healthcare provider. Do not exceed suggested use.
If you are pregnant or nursing, consult your healthcare provider before use.

NAD

- NAD is central to the body's metabolism, mitochondrial health, and cellular energy production
- NAD was recommended because you indicated you struggle with physical fatigue

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Foundational Evaluation

★

Fruits & Vegetables

You report consuming 0 to 2 servings per day

★

Essential Fatty Acids

You report consuming 0 to 1 servings per week

★★★

Fiber

You report consuming 2 to 3 servings per day

★★★

Protein

You report consuming 2 to 3 servings per day

★

Calcium

You report consuming 0 to 2 servings per day

★★★★

Exercise

You reported that you are very active and/or workout almost daily

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Foundational Recommendation

Multivitamin

USDA recommends consuming 1-2 cups of whole fruits and 1-3 cups of vegetables per day[15]

Increase your daily intake of whole fruits and vegetables and consider adding a multivitamin to your daily regimen

Fish Oil

The American Heart Association recommends eating 2 servings of omega-3 rich foods weekly[15]

Increase your weekly intake of omega-3 rich food and consider adding a fish oil supplement to your daily regimen

Fiber

The American Heart Association recommends eating 25 grams of fiber per day[17]

Consider increasing your daily intake of fiber-containing foods

Protein

The American Heart Association recommends eating 8-9 servings of protein per week[18]

Consider increasing your daily intake of high protein foods

Calcium

The USDA recommends consuming 2-3 cups of calcium-rich foods daily for adequate intake[15]

Increase your daily intake of calcium rich foods and consider adding a calcium supplement to your daily regimen

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Lifestyle Suggestions

Fatigue Hygiene

- **Physical Activity:** Physical activity has been shown to help improve feelings of general fatigue[26]
- **Hydration**: Adequate hydration has been found to reduce feelings of fatigue[27]
- **Sleep**: Go to bed at regular times and get 7 - 8.5 hours of sleep per night[28]
- **Balanced Diet:** A balanced diet with whole grains high in fiber, polyphenol-rich vegetables, and omega-3 fatty acids may improve fatigue symptoms[29]
- **Positive Outlook**: A positive outlook has been correlated with better stress management and reduced fatigue[30]

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

References

1. Data on file, Kavinace OS study. 2019. NeuroScience, Osceola, WI 54020.

2. Data on file, Alpha GABA PM study. 2019. NeuroScience, Osceola, WI 54020.

3. Langade D, et al. Cureus. 2019;11(9):e5797.

4. Shaw K, et al. Cochrane Database Syst Rev. 2002;1:CD003198.

5. Cespuglio R. Sleep Med. 2018;49:31-39.

6. Dakshinamurti K, et al. Ann N Y Acad Sci. 1990;585:128-44.

7. Kakuda T, et al. Biosci Biotechnol Biochem. 2002;66(12):2683-86.

8. Cuciureanu M, et al. Magnesium in the Central Nervous System. University of Adelaide Press. 2011.

9. Data on file, Calm CP study. 2012. NeuroScience, Inc., Osceola, WI 54020.

10. Wang C, et al. J Psychiatr Res. 2019;117:62-67.

11. Wuttke W, et al. J Steroid Biochem Mol Biol. 2014;138:302-10.

12. Jarry H, et al. Maturitas. 2003;44 Suppl 1:S31-8.

13. Jiang K, et al. Climacteric. 2015;18(4):559-67.

14. VanDolah H, et al. Mayo Clin Proc. 2019;94(9):1840-51.

15. Dietary Guidelines for Americans 2015-2020. Eighth Edition. United States Department of Agriculture. https://health.gov/sites/default/files/2019-09/2015-2020_Dietary_Guidelines.pdf

16. Fish and Omega-3 Fatty Acids. American Heart Association. 2017. https://www.heart.org/en/healthy-living/healthy-eating/eat-smart/fats/fish-and-omega-3-fatty-acids.

17. Whole Grains, Refined Grains, and Dietary Fiber. American Heart Association. 2016. https://www.heart.org/en/healthy-living/healthy-eating/eat-smart/nutrition-basics/whole-grains-refined-grains-and-dietary-fiber.

18. Suggested Servings from each Food Group. American Heart Association. 2017. https://www.heart.org/en/healthy-living/healthy-eating/eat-smart/nutrition-basics/suggested-servings-from-each-food-group.

19. Shechter A, et al. J Psychiatr Res. 2018;96:196-202.

20. Irish A, et al. Sleep Med Rev. 2015;22:23-36.

21. Lin J, et al. Popul Health Metr. 2009:7:18.

22. Pardridge W. NeuroRx. 2005;2(1):3-14.

23. Spasov A, et al. Phytomedicine. 2000;7(2):85-9.

24. Cools R, et al. Handb Clin Neurol. 2019;163:123-43.

25. Christie A, et al. Exp Gerontol. 2016;77:7-11.

26. Strahler J, et al. Stress. 2016;19(5):535-41.

27. Pross N, et al. PLoS One. 2014;9(4):e94754.

28. Spaeth A, et al. Nutr Rev. 2014;72(0 1):34-47.

29. Hab U, et al. Nutrients. 2019;11(10):2315.

30. Hirsch J and Sirois F. J Health Psychol. 2016;21(4):451-6.

31. Cohen S, et al. J Health Soc Behav. 1983;24(4):385-96.

32. Duke A, et al. Pyschol Bull. 2013;139(5):1148-72.

33. Herman J, et al. Prog Brain Res. 2008;170:353-64.

34. Auddy B, et al. JANA. 2008;11(1):50-6.

35. Elder G, et al. Sleep Med Rev. 2014;18(3):215-24.

36. Kakuda T, et al. Biosci Biotechnol Biochem. 2002;66(12):2683-86.

37. Meldrum B. J Nutr. 2000;130(4S Suppl):1007S-15S.

38. Carlson L, et al. Cancr J. 2019;25(5):329-336.

39. Nasir M, et al. Front Psychiatry. 2020;11:548505.

40. Ng J. et al, Nat Rev Neurol. 2015;11:567-84.

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.