NeuroScience Survey ID #1636928433

Requested for:
Name: **Suzanne Burton**
Sex: **female**
Age: **76**
State: **California**

Sleep Evaluation

Sleep area	Description	Rating
Self-Rated Sleep Quality	This is how you rate your sleep quality	★★
Ability to Fall Asleep	This is your body's ability to fall asleep	★
Ability to Stay Asleep	This is your body's ability to stay asleep through the night	★
Temperature Regulation	This is your body's ability to regulate temperature at night	★★★★
Nighttime Discomfort	This is how you rate your nighttime discomfort	★★★
Daytime Fatigue	This is how you rate your sleepiness during the day	★★★
Daytime Motivation	This is how you rate your motivation and energy to accomplish tasks during the day	★★

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Sleep Product Recommendations

Kavinace OS

Target the cellular consequences of sleeplessness with a formulation proven to significantly improve both sleep onset and efficiency after the first dose*

- Blend of ingredients proven to significantly improve both falling asleep and effectiveness after the first dose1*
- Recommended because you indicated your body's ability to fall asleep is very poor1

Suggested use:
Take 2 capsules at bedtime or as directed by your healthcare provider. Do not exceed suggested use.
If you are pregnant or nursing, consult your healthcare provider before use.



Calm CP

Decreases cortisol levels and provides ingredients important for calm, sleep, and management of blood sugar*

- Decreases cortisol levels and provides ingredients important for calm, sleep, and management of blood sugar9*
- Elevated cortisol has been associated with difficulty sleeping10
- Calm CP is a proven blend of ingredients that has been shown to decrease cortisol levels in as little as 5 days9

Suggested use:
Take 2 capsules one to two times daily or as directed by your healthcare provider. Do not exceed suggested use.
If you are pregnant or nursing, consult your healthcare provider before use.

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Kavinace OS

Target the cellular consequences of sleeplessness with a formulation proven to significantly improve both sleep onset and efficiency after the first dose*

- Blend of ingredients proven to significantly improve both falling asleep and effectiveness after the first dose1*
- Recommended because you indicated your body's ability to fall asleep is very poor1

Suggested use:
Take 2 capsules at bedtime or as directed by your healthcare provider. Do not exceed suggested use.
If you are pregnant or nursing, consult your healthcare provider before use.



Calm CP

Decreases cortisol levels and provides ingredients important for calm, sleep, and management of blood sugar*

- Decreases cortisol levels and provides ingredients important for calm, sleep, and management of blood sugar9*
- Elevated cortisol has been associated with difficulty sleeping10
- Calm CP is a proven blend of ingredients that has been shown to decrease cortisol levels in as little as 5 days9

Suggested use:
Take 2 capsules one to two times daily or as directed by your healthcare provider. Do not exceed suggested use.
If you are pregnant or nursing, consult your healthcare provider before use.

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Kavinace OS

Target the cellular consequences of sleeplessness with a formulation proven to significantly improve both sleep onset and efficiency after the first dose*

- Blend of ingredients proven to significantly improve both falling asleep and effectiveness after the first dose1*
- Recommended because you indicated your body's ability to fall asleep is very poor1

Suggested use:
Take 2 capsules at bedtime or as directed by your healthcare provider. Do not exceed suggested use.
If you are pregnant or nursing, consult your healthcare provider before use.



Calm CP

Decreases cortisol levels and provides ingredients important for calm, sleep, and management of blood sugar*

- Decreases cortisol levels and provides ingredients important for calm, sleep, and management of blood sugar9*
- Elevated cortisol has been associated with difficulty sleeping10
- Calm CP is a proven blend of ingredients that has been shown to decrease cortisol levels in as little as 5 days9

Suggested use:
Take 2 capsules one to two times daily or as directed by your healthcare provider. Do not exceed suggested use.
If you are pregnant or nursing, consult your healthcare provider before use.

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Kavinace OS

Target the cellular consequences of sleeplessness with a formulation proven to significantly improve both sleep onset and efficiency after the first dose*

- Blend of ingredients proven to significantly improve both falling asleep and effectiveness after the first dose1*
- Recommended because you indicated your body's ability to fall asleep is very poor1

Suggested use:
Take 2 capsules at bedtime or as directed by your healthcare provider. Do not exceed suggested use.
If you are pregnant or nursing, consult your healthcare provider before use.



Calm CP

Decreases cortisol levels and provides ingredients important for calm, sleep, and management of blood sugar*

- Decreases cortisol levels and provides ingredients important for calm, sleep, and management of blood sugar9*
- Elevated cortisol has been associated with difficulty sleeping10
- Calm CP is a proven blend of ingredients that has been shown to decrease cortisol levels in as little as 5 days9

Suggested use:
Take 2 capsules one to two times daily or as directed by your healthcare provider. Do not exceed suggested use.
If you are pregnant or nursing, consult your healthcare provider before use.

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Foundational Evaluation

★★★

Fruits & Vegetables

You report consuming 3 to 5 servings per day

★★★

Essential Fatty Acids

You report consuming 2 to 3 servings per week

★★★

Fiber

You report consuming 2 to 3 servings per day

★★★

Protein

You report consuming 2 to 3 servings per day

★

Calcium

You report consuming 0 to 2 servings per day

★

Exercise

You reported that you are not very active and/or do not workout

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Foundational Recommendation

Multivitamin

USDA recommends consuming 1-2 cups of whole fruits and 1-3 cups of vegetables per day[15]

Consider increasing your daily intake of whole fruits and vegetables

Fish Oil

The American Heart Association recommends eating 2 servings of omega-3 rich foods weekly[15]

Consider increasing your weekly intake of omega-3 rich food

Fiber

The American Heart Association recommends eating 25 grams of fiber per day[17]

Consider increasing your daily intake of fiber-containing foods

Protein

The American Heart Association recommends eating 8-9 servings of protein per week[18]

Consider increasing your daily intake of high protein foods

Calcium

The USDA recommends consuming 2-3 cups of calcium-rich foods daily for adequate intake[15]

Increase your daily intake of calcium rich foods and consider adding a calcium supplement to your daily regimen

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Lifestyle Suggestions

Sleep Hygiene

- **Blue Light**: Reducing blue light exposure 2 hours before bed has been shown to improve sleep quality and total sleep time[19]
- **Exercise**: Regular exercise may improve sleep and is best completed several hours before bedtime[20]
- **Caffeine**: Limit caffeine intake several hours before bedtime as it can be disruptive on sleep[20]
- **Nicotine**: Nicotine is associated with sleep issues, contact your healthcare provider for further guidance on the use of nicotine products[20]
- **Alcohol**: While alcohol has been shown to help individuals fall asleep, it may prevent deep sleep and increase awakenings in the middle of the night[20]
- **Manage Stress**: Practice calming and relaxing techniques before bed as stress may contribute to difficulty sleeping[20]
- **Regular Sleep Times**: Practice going to bed at regular times to promote stable sleep and wakefulness[20]

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

References

1. Data on file, Kavinace OS study. 2019. NeuroScience, Osceola, WI 54020.

2. Data on file, Alpha GABA PM study. 2019. NeuroScience, Osceola, WI 54020.

3. Langade D, et al. Cureus. 2019;11(9):e5797.

4. Shaw K, et al. Cochrane Database Syst Rev. 2002;1:CD003198.

5. Cespuglio R. Sleep Med. 2018;49:31-39.

6. Dakshinamurti K, et al. Ann N Y Acad Sci. 1990;585:128-44.

7. Kakuda T, et al. Biosci Biotechnol Biochem. 2002;66(12):2683-86.

8. Cuciureanu M, et al. Magnesium in the Central Nervous System. University of Adelaide Press. 2011.

9. Data on file, Calm CP study. 2012. NeuroScience, Inc., Osceola, WI 54020.

10. Wang C, et al. J Psychiatr Res. 2019;117:62-67.

11. Wuttke W, et al. J Steroid Biochem Mol Biol. 2014;138:302-10.

12. Jarry H, et al. Maturitas. 2003;44 Suppl 1:S31-8.

13. Jiang K, et al. Climacteric. 2015;18(4):559-67.

14. VanDolah H, et al. Mayo Clin Proc. 2019;94(9):1840-51.

15. Dietary Guidelines for Americans 2015-2020. Eighth Edition. United States Department of Agriculture. https://health.gov/sites/default/files/2019-09/2015-2020_Dietary_Guidelines.pdf

16. Fish and Omega-3 Fatty Acids. American Heart Association. 2017. https://www.heart.org/en/healthy-living/healthy-eating/eat-smart/fats/fish-and-omega-3-fatty-acids.

17. Whole Grains, Refined Grains, and Dietary Fiber. American Heart Association. 2016. https://www.heart.org/en/healthy-living/healthy-eating/eat-smart/nutrition-basics/whole-grains-refined-grains-and-dietary-fiber.

18. Suggested Servings from each Food Group. American Heart Association. 2017. https://www.heart.org/en/healthy-living/healthy-eating/eat-smart/nutrition-basics/suggested-servings-from-each-food-group.

19. Shechter A, et al. J Psychiatr Res. 2018;96:196-202.

20. Irish A, et al. Sleep Med Rev. 2015;22:23-36.

21. Lin J, et al. Popul Health Metr. 2009:7:18.

22. Pardridge W. NeuroRx. 2005;2(1):3-14.

23. Spasov A, et al. Phytomedicine. 2000;7(2):85-9.

24. Cools R, et al. Handb Clin Neurol. 2019;163:123-43.

25. Christie A, et al. Exp Gerontol. 2016;77:7-11.

26. Strahler J, et al. Stress. 2016;19(5):535-41.

27. Pross N, et al. PLoS One. 2014;9(4):e94754.

28. Spaeth A, et al. Nutr Rev. 2014;72(0 1):34-47.

29. Hab U, et al. Nutrients. 2019;11(10):2315.

30. Hirsch J and Sirois F. J Health Psychol. 2016;21(4):451-6.

31. Cohen S, et al. J Health Soc Behav. 1983;24(4):385-96.

32. Duke A, et al. Pyschol Bull. 2013;139(5):1148-72.

33. Herman J, et al. Prog Brain Res. 2008;170:353-64.

34. Auddy B, et al. JANA. 2008;11(1):50-6.

35. Elder G, et al. Sleep Med Rev. 2014;18(3):215-24.

36. Kakuda T, et al. Biosci Biotechnol Biochem. 2002;66(12):2683-86.

37. Meldrum B. J Nutr. 2000;130(4S Suppl):1007S-15S.

38. Carlson L, et al. Cancr J. 2019;25(5):329-336.

39. Nasir M, et al. Front Psychiatry. 2020;11:548505.

40. Ng J. et al, Nat Rev Neurol. 2015;11:567-84.

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.