NeuroScience Survey ID #1681946444

Requested for:

Name: **INNA DUBROV**
Sex: **female**
Age: **51**
State: **Virginia**

Your Score33

Your overall stress level is **33**, which indicates your stress level is **high**.

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

TravaCor

Contains a blend of ingredients shown to improve mood and reduce stress*

- Contains a blend of ingredients shown to improve mood and reduce stress*
- TravaCor was recommended beause you indicated you are perceiving elevated levels of stress and reported feelings of upset or anger
- Low levels of serotonin have been correlated with feelings of upset or anger. TravaCor contains 5-HTP and B-vitamins important for the synthesis of serotonin in the body4,32

Suggested use:
Take 1-3 capsules one to two times daily or as directed by your healthcare provider.

If you are pregnant or nursing, consult your healthcare provider before use.

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.



ExcitaPlus

Contains non-glandular ingredients important for adrenal health and reducing stress-induced fatigue.* Provides over three times the L-tyrosine and six times the Rhodiola per capsule compared to AdreCor.

- Contains non-glandular ingredients important for adrenal health and reducing stress-induced fatigue*
- ExcitaPlus was recommended because you indicated you may experience cognitive issues and difficulty handling personal problems, coping with things you have to do, keeping on top of things
- Contains ingredients to support the synthesis of dopamine, a neurotransmitter implicated in motivation and Rhodiola, a botanical shown to reduce mental fatigue23,24

Suggested use:
Take 1-2 capsules one to two times daily or as directed by your healthcare provider. Do not exceed suggested use.

If you are pregnant or nursing, consult your healthcare provider before use.

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.



Calm CP

Decreases cortisol levels and provides ingredients important for calm, sleep, and management of blood sugar*

- Decreases cortisol levels and provides ingredients important for calm, sleep, and management of blood sugar*
- Calm CP was recommended because you indicated you are perceiving elevated levels of stress
- Cortisol secretion follows a marked circadian rhythm and increases in response to stress through activation of the HPA axis35

Suggested use:
Take 2 capsules one to two times daily or as directed by your healthcare provider. Do not exceed suggested use.

If you are pregnant or nursing, consult your healthcare provider before use.

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Fruits & Vegetables

You report consuming 0 to 2 servings per day

Essential Fatty Acids

You report consuming 2 to 3 servings per week

Fiber

You report consuming 2 to 3 servings per day

Protein

You report consuming 2 to 3 servings per day

Calcium

You report consuming 0 to 2 servings per day

Exercise

You reported that you are not very active and/or do not workout

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Multivitamin

USDA recommends consuming 1-2 cups of whole fruits and 1-3 cups of vegetables per day[15]

Increase your daily intake of whole fruits and vegetables and consider adding a multivitamin to your daily regimen

Fish Oil

The American Heart Association recommends eating 2 servings of omega-3 rich foods weekly[15]

Consider increasing your weekly intake of omega-3 rich food

Fiber

The American Heart Association recommends eating 25 grams of fiber per day[17]

Consider increasing your daily intake of fiber-containing foods

Protein

The American Heart Association recommends eating 8-9 servings of protein per week[18]

Consider increasing your daily intake of high protein foods

Calcium

The USDA recommends consuming 2-3 cups of calcium-rich foods daily for adequate intake[15]

Increase your daily intake of calcium rich foods and consider adding a calcium supplement to your daily regimen

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

Stress Hygiene

- **Cognitive-Behavioral Management**: Cognitive-behavioral stress management techniques such as relaxation techniques, cognitive restructuring, psychoeducation, and coping skills have been shown to be very effective at reducing stress[38]
- **Yoga:** Yoga has been shown to greatly reduce short-term anxiousness, perceived stress, and psychological distress[38]
- **Mindfulness**: Mindfulness as defined as nonjudgmental present-moment awareness that has been shown to significantly reduce stress when practiced daily[38]
- **Music Therapy**: Guided music therapy while listening to music meaningful to oneself has been shown to significantly reduce feelings of anxiousness[38]

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.

References

1. Data on file, Kavinace OS study. 2019. NeuroScience, Osceola, WI 54020.

2. Data on file, Alpha GABA PM study. 2019. NeuroScience, Osceola, WI 54020.

3. Langade D, et al. Cureus. 2019;11(9):e5797.

4. Shaw K, et al. Cochrane Database Syst Rev. 2002;1:CD003198.

5. Cespuglio R. Sleep Med. 2018;49:31-39.

6. Dakshinamurti K, et al. Ann N Y Acad Sci. 1990;585:128-44.

7. Kakuda T, et al. Biosci Biotechnol Biochem. 2002;66(12):2683-86.

8. Cuciureanu M, et al. Magnesium in the Central Nervous System. University of Adelaide Press. 2011.

9. Data on file, Calm CP study. 2012. NeuroScience, Inc., Osceola, WI 54020.

10. Wang C, et al. J Psychiatr Res. 2019;117:62-67.

11. Wuttke W, et al. J Steroid Biochem Mol Biol. 2014;138:302-10.

12. Jarry H, et al. Maturitas. 2003;44 Suppl 1:S31-8.

13. Jiang K, et al. Climacteric. 2015;18(4):559-67.

14. VanDolah H, et al. Mayo Clin Proc. 2019;94(9):1840-51.

15. Dietary Guidelines for Americans 2015-2020. Eighth Edition. United States Department of Agriculture. https://health.gov/sites/default/files/2019-09/2015-2020_Dietary_Guidelines.pdf

16. Fish and Omega-3 Fatty Acids. American Heart Association. 2017. https://www.heart.org/en/healthy-living/healthy-eating/eat-smart/fats/fish-and-omega-3-fatty-acids.

17. Whole Grains, Refined Grains, and Dietary Fiber. American Heart Association. 2016. https://www.heart.org/en/healthy-living/healthy-eating/eat-smart/nutrition-basics/whole-grains-refined-grains-and-dietary-fiber.

18. Suggested Servings from each Food Group. American Heart Association. 2017. https://www.heart.org/en/healthy-living/healthy-eating/eat-smart/nutrition-basics/suggested-servings-from-each-food-group.

19. Shechter A, et al. J Psychiatr Res. 2018;96:196-202.

20. Irish A, et al. Sleep Med Rev. 2015;22:23-36.

21. Lin J, et al. Popul Health Metr. 2009:7:18.

22. Pardridge W. NeuroRx. 2005;2(1):3-14.

23. Spasov A, et al. Phytomedicine. 2000;7(2):85-9.

24. Cools R, et al. Handb Clin Neurol. 2019;163:123-43.

25. Christie A, et al. Exp Gerontol. 2016;77:7-11.

26. Strahler J, et al. Stress. 2016;19(5):535-41.

27. Pross N, et al. PLoS One. 2014;9(4):e94754.

28. Spaeth A, et al. Nutr Rev. 2014;72(0 1):34-47.

29. Hab U, et al. Nutrients. 2019;11(10):2315.

30. Hirsch J and Sirois F. J Health Psychol. 2016;21(4):451-6.

31. Cohen S, et al. J Health Soc Behav. 1983;24(4):385-96.

32. Duke A, et al. Pyschol Bull. 2013;139(5):1148-72.

33. Herman J, et al. Prog Brain Res. 2008;170:353-64.

34. Auddy B, et al. JANA. 2008;11(1):50-6.

35. Elder G, et al. Sleep Med Rev. 2014;18(3):215-24.

36. Kakuda T, et al. Biosci Biotechnol Biochem. 2002;66(12):2683-86.

37. Meldrum B. J Nutr. 2000;130(4S Suppl):1007S-15S.

38. Carlson L, et al. Cancr J. 2019;25(5):329-336.

39. Nasir M, et al. Front Psychiatry. 2020;11:548505.

40. Ng J. et al, Nat Rev Neurol. 2015;11:567-84.

***These statements have not been evaluated by the Food and Drug Administration.**

Recommendations and product information provided was requested by the user and is not intended to diagnose, treat, cure or prevent any disease.