

A CRITERION FOR A HUREWICZ COFIBRATION TO BE A QUILLEN COFIBRATION

ANDREW RONAN

In this paper, we prove that h -cofibrations between q -cofibrant spaces are q -cofibrations and discuss a number of applications. In fact, since our non-equivariant proofs generalise readily to the equivariant context, we work throughout with the $\mathcal{C}$ -model structure on G -spaces. Here $\mathcal{C}$ is a set of subgroups of G which we take to be closed under conjugacy, and the $\mathcal{C}$ -model structure is that of [Sch18, Proposition B.7]. Recall that the $\mathcal{C}$ -model structure is a topological model structure cofibrantly generated by:

$$\begin{aligned} I &= \{G/H \times S^{n-1} \rightarrow G/H \times D^n \mid n \geq 0, H \in \mathcal{C}\}, \\ J &= \{G/H \times D^n \rightarrow G/H \times D^n \times I \mid n \geq 0, H \in \mathcal{C}\} \end{aligned}$$

Moreover, the $\mathcal{C}$ -fibrations (resp. $\mathcal{C}$ -equivalences) are the maps p such that p^H is a q -fibration (resp. q -equivalence) for every $H \in \mathcal{C}$. Note that our assumption that $\mathcal{C}$ is closed under conjugacy results in no loss of generality, and is convenient for some of the proofs below. For example, if H is a subgroup of G and we define $\mathcal{C} \cap H = \{K \cap H \mid K \in \mathcal{C}\}$, we then have:

Lemma 1: *If X is $\mathcal{C}$ -cofibrant as a G -space, then X is $(\mathcal{C} \cap H)$ -cofibrant as an H -space.*

Hurewicz cofibrations between q -cofibrant spaces are q -cofibrations

We now move onto the proof of our main theorem. The key point is the following generalisation of [MP12, Proposition 17.1.4], which admits the same proof:

Lemma 2: *Let $i : A \rightarrow B$ be an h -cofibration and $p : X \rightarrow Y$ be a map which has the RLP wrt $B \rightarrow B \times I$. Suppose that we are given a commutative square:*

$$\begin{array}{ccc} A & \xrightarrow{v} & X \\ i \downarrow & & \downarrow p \\ B & \xrightarrow{w} & Y \end{array}$$

and a map $\sigma : B \rightarrow X$ such that $p\sigma i = pv$, $\sigma i \simeq v$ over Y and $p\sigma \simeq w$ rel A . Then σ is homotopic to some τ with $\tau i = v$ and $p\tau = w$.

Proof. Since i is an h -cofibration, σ is homotopic to some σ' with $\sigma' i = v$ in such a way that the composite homotopy $p\sigma' \simeq w$ is still relative to A . So we have reduced to the case where $\sigma i = v$. Let (H, λ) represent (B, A) as an NDR -pair. Let $K : B \times I \rightarrow Y$ be a homotopy between $p\sigma$ and w , relative to A and normalised so that $K(b, t) = K(b, 1)$ for $t \geq \lambda(b)$ (see Lemma 26). Since p has the RLP wrt $B \rightarrow B \times I$, we can lift this homotopy to $\tilde{K} : B \times I \rightarrow X$ starting at σ . If we define $\tilde{\sigma}(b) = \tilde{K}(b, \lambda(b))$ we solve the lifting problem as desired. $\square$

Lemma 3: *An h -cofibration between $\mathcal{C}$ -cofibrant spaces is a $\mathcal{C}$ -cofibration.*

Proof. Let $i : A \rightarrow B$ be an h -cofibration between $\mathcal{C}$ -cofibrant spaces. We can factor i as a $\mathcal{C}$ -cofibration followed by a $\mathcal{C}$ -acyclic $\mathcal{C}$ -fibration between $\mathcal{C}$ -cofibrant objects, so by the retract argument it suffices to show that i has the LLP wrt $\mathcal{C}$ -acyclic $\mathcal{C}$ -fibrations between $\mathcal{C}$ -cofibrant objects.

If $p : X \rightarrow Y$ is a $\mathcal{C}$ -acyclic $\mathcal{C}$ -fibration between $\mathcal{C}$ -cofibrant objects, then there exists $s : Y \rightarrow X$ such that $ps = 1$ and $sp \simeq 1$ over Y . Since B is $\mathcal{C}$ -cofibrant p has the RLP wrt $B \rightarrow B \times I$. Therefore, we can solve a lifting problem as in Lemma 2, with the initial up to homotopy lift σ as sw . $\square$

Example 4: Suppose that $H \in \mathcal{C}$. If U is an equivariant open subspace of $G/H \times D^n$, then $i : U \cap (G/H \times \partial D^n) \rightarrow U \cap (G/H \times D^n)$ is the inclusion of the boundary of a smooth G -manifold and so is an h -cofibration between $\mathcal{C}$ -cofibrant spaces by the equivariant collar neighbourhood theorem, [Kan07, Theorem 3.5]. Therefore, Lemma 3 tells us that i is a $\mathcal{C}$ -cofibration.

Lemma 5: If $i : A \rightarrow B$ is a $\mathcal{C}$ -cofibration and $U \subset B$ is an equivariant open subspace, then $i : A \cap U \rightarrow U$ is a $\mathcal{C}$ -cofibration.

Proof. Firstly, i is a retract of a relative $\mathcal{C}$ -cell complex, $j : A \rightarrow C$, as shown below:

$$\begin{array}{ccccc} A & \longrightarrow & A & \longrightarrow & A \\ \downarrow i & & \downarrow j & & \downarrow i \\ B & \xrightarrow{i} & C & \xrightarrow{r} & B \end{array}$$

If we let $V = r^{-1}(U)$, then $i : A \cap U \rightarrow U$ is a retract of $j : A \cap V \rightarrow V$. So we have reduced to the case where i is a relative $\mathcal{C}$ -cell complex, which we now assume.

We can express i as a transfinite composite of maps $A_\lambda \rightarrow A_{\lambda+1}$, each of which is a pushout of a map of the form $G/H \times S^{n-1} \rightarrow G/H \times D^n$, with $H \in \mathcal{C}$. By [Ron23, Lemma 6.2.3] and Lemma 24, the map $i : A \cap U \rightarrow U$ can be expressed as a transfinite composite of the maps $A_\lambda \cap U \rightarrow A_{\lambda+1} \cap U$, each of which is a pushout of a map of the form $(G/H \times S^{n-1}) \cap \alpha^{-1}(U) \rightarrow (G/H \times D^n) \cap \alpha^{-1}(U)$, where α is the inclusion of the cell into B . By Example 4, $A_\lambda \cap U \rightarrow A_{\lambda+1} \cap U$ is therefore a $\mathcal{C}$ -cofibration, and, hence, so is the transfinite composite $i : A \cap U \rightarrow U$. $\square$

Corollary 6: An equivariant open subspace of a $\mathcal{C}$ -cofibrant G -space is $\mathcal{C}$ -cofibrant.

Combining Lemma 3 and Corollary 6, we can rephrase Lemma 3 as follows:

Theorem 7: If $i : A \rightarrow B$ is an h -cofibration and B is $\mathcal{C}$ -cofibrant, then A is $\mathcal{C}$ -cofibrant and i is a $\mathcal{C}$ -cofibration.

Proof. If (H, λ) represents (B, A) as a G -NDR-pair, then A is a retract of $\lambda^{-1}([0, 1])$ which is $\mathcal{C}$ -cofibrant by Corollary 6. $\square$

A pushout-product property of symmetrizable cofibrations

Let $\mathcal{C}$ be a closed symmetric monoidal, tensored topological model category with monoidal product $\boxtimes$. If $f : A \rightarrow B$ and $g : C \rightarrow D$ are maps in $\mathcal{C}$, let $f \hat{\boxtimes} g$ denote the pushout-product of f and g , $f \hat{\boxtimes} g : A \boxtimes D \cup_{A \boxtimes C} B \boxtimes C \rightarrow B \boxtimes D$. In general, a hat above an operation denotes either the corresponding pushout-product or the dual notion involving pullbacks. The n -fold composite $f \hat{\boxtimes} \dots \hat{\boxtimes} f$ is denoted by $f^{\hat{\boxtimes} n}$. The notion of a symmetrizable cofibration is then defined as follows:

Definition 8: A map $i : A \rightarrow B$ in $\mathcal{C}$ is said to be a symmetrizable (acyclic) cofibration if for every $n \geq 1$, $h^{\hat{\boxtimes} n} / \Sigma_n$ is an (acyclic) cofibration.

Example 9: Let $G = \Sigma_n$ and let $\mathcal{C}$ be the set of subgroups of G , each isomorphic to a product of symmetric groups $\Sigma_{n_1} \times \dots \times \Sigma_{n_l}$, corresponding to partitions of $\{1, \dots, n\}$ into sets of size $n_1, \dots, n_l$. Let $i : S^{k-1} \rightarrow D^k$ be the boundary inclusion. If we consider the iterated pushout product map $j : S^{k-1} \times (D^k)^{n-1} \cup \dots \cup (D^k)^{n-1} \times S^{k-1} := Q_n(i) \rightarrow (D^k)^n$, then j is an h -cofibration by [May72, Lemma A.4], and the domain and codomain are $\mathcal{C}$ -cofibrant since they are G -manifolds, hence G -CW complexes, [Ill83, Corollary 7.2], with the correct isotropy groups. Therefore, Lemma 3 tells us that j is a $\mathcal{C}$ -cofibration.

We then have the following application to the theory of symmetrizable cofibrations, which is a generalisation of [Sch18, Proposition 2.1.12]:

Lemma 10: *Let $j : A \rightarrow B$ be a symmetrizable cofibration in a tensored and cotensored closed symmetric monoidal topological model category and let $i : C \rightarrow D$ be a q -cofibration of spaces. Then the pushout product map:*

$$j \widehat{\times} i : A \times D \cup_{A \times C} B \times C \rightarrow B \times D$$

is a symmetrizable cofibration, which is a symmetrizable acyclic cofibration if either i is acyclic or j is a symmetrizable acyclic cofibration.

Proof. The assumption that $\mathcal{C}$ is tensored and cotensored implies that the bifunctor $-\times-$ preserves colimits in both variables. It follows that we can reduce to the case where i is a generating q -cofibration via [RV22, Proposition C.2.9], and using the closure properties of symmetrizable cofibrations described in [GG16, Theorem 7]. Now, for any $n \geq 1$, we want to show that $(j \widehat{\times} i)^{\widehat{\boxtimes} n} / \Sigma_n$ is an (acyclic) cofibration. Equivalently, we want to show that $(j \widehat{\times} i)^{\widehat{\boxtimes} n}$ has the Σ_n -equivariant LLP with respect to (acyclic) fibrations $p : X \rightarrow Y$, where Σ_n acts trivially on X and Y . Using the tensor adjunction, it suffices to show that $i^{\widehat{\times} n}$ has the equivariant LLP wrt $\widehat{Map}(j^{\widehat{\boxtimes} n}, p)$. In Example 9, we saw that $i^{\widehat{\times} n}$ is a $\mathcal{C}$ -cofibration, with respect to the specified set of subgroups $\mathcal{C}$. So it suffices to show that $\widehat{Map}(j^{\widehat{\boxtimes} n}, p)$ is a $\mathcal{C}$ -acyclic $\mathcal{C}$ -fibration. We need to show that for every $H \in \mathcal{C}$, $\widehat{Map}(j^{\widehat{\boxtimes} n} / H, p)$ is a q -acyclic q -fibration. Since $H \in \mathcal{C}$ it is isomorphic to a group of the form $\Sigma_{n_1} \times \dots \times \Sigma_{n_l}$, so $j^{\widehat{\boxtimes} n} / H$ is isomorphic to $(j^{\widehat{\boxtimes} n_1} / \Sigma_{n_1}) \widehat{\boxtimes} \dots \widehat{\boxtimes} (j^{\widehat{\boxtimes} n_l} / \Sigma_{n_l})$. Therefore, the result follows from the fact that the model structure is topological and monoidal, j is a symmetrizable cofibration and either p is acyclic or j is a symmetrizable acyclic cofibration. $\square$

Local gluing results for fibrations and cofibrations

In this subsection, we explain how Theorem 7 allows us to prove gluing results for $\mathcal{C}$ -fibrations and $\mathcal{C}$ -cofibrations analogous to Dold's gluing results for h -fibrations and h -cofibrations, [Dol63, Theorem 4.8] and [Dol68, Satz 2]. First recall that:

Definition 11: *A G -numerable open cover $\{U_j\}_{j \in J}$ of a G -space, B , is a locally finite open cover of B such that for every $j \in J$ there exists an equivariant map $\lambda_j : B \rightarrow I$ with $U_j = \lambda_j^{-1}((0, 1])$.*

The next lemma is central to the proofs of all four gluing theorems:

Lemma 12: *Consider a lifting problem of G -spaces as below:*

$$(1) \quad \begin{array}{ccc} A & \xrightarrow{f} & X \\ i \downarrow & & \downarrow p \\ B & \xrightarrow{g} & Y \end{array}$$

Suppose that there exists a G -numerable open cover $\{U_i\}_{i \in I}$ of B such that if $i \in I$ and $J \subset I$, then there exists a solution to the induced lifting problem:

$$\begin{array}{ccc} i^{-1}(U_i \cap (\cup_{j \in J} U_j)) & \xrightarrow{f} & X \\ i \downarrow & & \downarrow p \\ U_i \cap (\cup_{j \in J} U_j) & \xrightarrow{g} & Y \end{array}$$

and given two solutions ψ_1, ψ_2 of the induced lifting problem there exists a G -homotopy from ψ_1 to ψ_2 which is relative to both p and i .

Then there is a solution to the original lifting problem 1.

Proof. Let:

$$S = \{(A, \psi) \mid A \subset I, \psi : \cup_{\alpha \in A} U_\alpha \rightarrow X, p\psi = g, \psi i = f\}.$$

We define a partial order on S by:

$$(A, \psi_1) \leq (B, \psi_2)$$

iff $A \subset B$ and $\psi_1 = \psi_2$ on $\cup_{\alpha \in A} U_\alpha \setminus \cup_{\beta \in B \setminus A} U_\beta$.

We may as well assume that B is non-empty, in which case taking $J = \{i\}$ in the condition of the lemma shows that S is non-empty. Since $\{U_i\}$ is a locally finite cover, every chain in S has an upper bound. Hence, by Zorn's lemma, S contains a maximal element, say (A, ψ_1) . If $A \neq I$, there exists some $\beta \in I \setminus A$. By the condition in the lemma, there exists some $(\{\beta\}, \psi_2) \in S$. By the uniqueness up to relative homotopy condition, there exists a G -homotopy $K : (U_\beta \cap (\cup_{\alpha \in A} U_\alpha)) \times I \rightarrow X$ between ψ_1 and ψ_2 , relative to p and i . Pick G -maps $\lambda_1, \lambda_2 : U_\beta \cup (\cup_{\alpha \in A} U_\alpha) \rightarrow I$ such that $\cup_{\alpha \in A} U_\alpha = \lambda_1^{-1}((0, 1])$, $U_\beta = \lambda_2^{-1}((0, 1])$ and $\lambda_1 + \lambda_2 = 1$. Define:

$$\psi(u) = \begin{cases} \psi_1(u) & \text{if } \lambda_1 > \frac{3}{4} \\ \psi_2(u) & \text{if } \lambda_1 < \frac{1}{4} \\ K(u, \frac{3}{2} - 2\lambda_1) & \text{otherwise} \end{cases}$$

Then $(A \cup \{\beta\}, \psi) \in S$ with $(A, \psi_1) \leq (A \cup \{\beta\}, \psi)$, a contradiction. So $A = I$ and ψ_1 is a solution to the original lifting problem. $\square$

As an immediate consequence, we have the following gluing result for $\mathcal{C}$ -cofibrations:

Theorem 13: *If $i : A \rightarrow B$ is a map of G -spaces and there is a G -numerable open cover $\mathcal{U}$ of B such that for every $U \in \mathcal{U}$, $i|_{A \cap U} : A \cap U \rightarrow U$ is a $\mathcal{C}$ -cofibration, then i is a $\mathcal{C}$ -cofibration.*

Proof. We consider a lifting problem for i with respect to a $\mathcal{C}$ -acyclic $\mathcal{C}$ -fibration, $p : X \rightarrow Y$. The G -numerable open cover $\mathcal{U}$ ensures the existence of local lifts as in Lemma 12. Uniqueness follows from Lemma 5 and the fact that the $\mathcal{C}$ -model structure is topological, [Sch18, Proposition B.5], so solutions to such lifting problems are unique up to relative homotopy. Therefore, the result follows from Lemma 12. $\square$

The corresponding gluing theorem for h -cofibrations is [Dol68, Satz 2], whose proof uses the fact that the restriction of an h -cofibration to a numerable open subspace is an h -cofibration, [Dol68, Satz 1], in place of Lemma 5:

Theorem 14: *Suppose that $\{U_j\}_{j \in J}$ is a G -numerable open cover of B . If $f : A \rightarrow B$ is a map such that $f : f^{-1}(U_j) \rightarrow U_j$ is an hG -cofibration for all $j \in J$, then $f : A \rightarrow B$ is an hG -cofibration.*

As for h -fibrations, [Dol63, Theorem 4.8], with a little more work we can use Lemma 12 to prove a gluing result for $\mathcal{C}$ -fibrations:

Theorem 15: *If $p : X \rightarrow Y$ is a map and there exists a G -numerable open cover $\{U_i\}_{i \in I}$ of Y such that $p : p^{-1}(U_i) \rightarrow U_i$ is a $\mathcal{C}$ -fibration for all $i \in I$, then p is a $\mathcal{C}$ -fibration.*

Proof. Consider a lifting problem:

$$\begin{array}{ccc} A & \xrightarrow{f} & X \\ i \downarrow & & \downarrow p \\ A \times I & \xrightarrow{H} & Y \end{array}$$

where A is $\mathcal{C}$ -cofibrant and $i : A \rightarrow A \times I$ is the inclusion of $A \times \{0\}$.

Choose functions, $\lambda_i : Y \rightarrow I$, for every $i \in I$, such that $U_i = \lambda_i^{-1}((0, 1])$ and $\sum_{i \in I} \lambda_i = 1$. Suppose that $S = (i_0, \dots, i_{n-1})$ is a finite sequence of elements of I , so, in particular, the members of the sequence need not be distinct. Define a function $\mu_S : A \rightarrow I$ by:

$$(2) \quad \mu_S(a) = \max \left\{ \min_{j \in \{0, \dots, n-1\}} \left(\inf_{t \in [\frac{j}{n}, \frac{j+1}{n}]} \lambda_{i_j}(H(a, t)) \right) - n \sum_{T \mid |T| < n} \mu_T, 0 \right\}$$

Then $\{V_S := \mu_S^{-1}((0, 1])\}$ is a G -numerable open cover of A , indexed over finite sequences of elements of I . Therefore, $\{V_S \times I\}$ is a G -numerable open cover of $A \times I$. Moreover, by inspection of equation (2), we see that $H(V_S \times [\frac{j}{n}, \frac{j+1}{n}]) \subset U_{i_j}$ for every j . It follows that the existence condition of Lemma 12 is satisfied,

since V_S is $\mathcal{C}$ -cofibrant by Corollary 6 and so we can construct a lift inductively on these intervals. Similarly, we can construct relative homotopies between any two lifts inductively, and so the uniqueness condition of Lemma 12 is also satisfied, which gives the result. $\square$

Other Examples

The next result is the equivariant analogue of [DE72, Corollary III.2].

Lemma 16: *Suppose that $\mathcal{C}$ is closed under conjugacy and intersections. If X is $\mathcal{C}$ -cofibrant, then $\Delta : X \rightarrow X \times X$ is a $\mathcal{C}$ -cofibration.*

Proof. The condition that $\mathcal{C}$ is closed under conjugacy and intersections ensures that $X \times X$ is also $\mathcal{C}$ -cofibrant. Since any $\mathcal{C}$ -cofibrant G -space is a retract of a $\mathcal{C}$ -cell complex, by Lemma 3 it suffices to show that if X is a G -cell complex then $\Delta : X \rightarrow X \times X$ is an hG -cofibration. To this end, assume that $\Delta : Y \rightarrow Y \times Y$ is an hG -cofibration and X is obtained from Y by attaching a single cell along $\alpha : G/H \times S^{n-1} \rightarrow Y$. We will show that if (H, λ) represents $(Y \times Y, Y)$ a G -NDR-pair, then there is a G -NDR pair (K, μ) for $(X \times X, X)$ such that $K|_{Y \times Y \times I} = H$ and $\mu|_{Y \times Y} = \lambda$. Firstly, the composite $Y \rightarrow Y \times Y \rightarrow Y \times X \cup_{Y \times Y} X \times Y$ is an hG -cofibration, which can be represented by a G -NDR pair that extends (H, λ) by Lemma 27, where note $Y \times Y \rightarrow Y \times X \cup_{Y \times Y} X \times Y$ is an hG -cofibration by [Str72, Lemma 5]. We then consider the map between pushouts:

$$\begin{array}{ccccc} G/H \times D^n & \longleftarrow & G/H \times S^{n-1} & \longrightarrow & Y \\ \downarrow \Delta & & \downarrow \Delta & & \downarrow \alpha \\ G/H \times G/H \times D^n \times D^n & \longleftarrow & G/H \times G/H \times \partial D^{2n} & \longrightarrow & Y \times X \cup_{Y \times Y} X \times Y \end{array}$$

We've seen that α is an hG -cofibration. The map between pushouts is $\Delta : X \rightarrow X \times X$ and to show this is an hG -cofibration it suffices to show the pushout-product map, ϕ , below is an hG -cofibration:

$$\phi : G/H \times D^n \cup_{G/H \times S^{n-1}} G/H \times G/H \times \partial D^{2n} \rightarrow G/H \times G/H \times D^n \times D^n$$

Let $U \cong G/H \times \partial D^n \times [0, 1)$ be a numerable boundary collar in $G/H \times D^n$. Then consider the restriction of ϕ to $U \times U$, which can be identified with:

$$G/H \times \partial D^n \times [0, 1) \cup G/H \times G/H \times \partial D^n \times \partial D^n \times \partial([0, 1) \times [0, 1)) \rightarrow G/H \times G/H \times \partial D^n \times \partial D^n \times [0, 1) \times [0, 1)$$

Let $L \subset [0, 1) \times [0, 1)$ be the union of the diagonal and the boundaries, so L consists of three straight lines. The map above is the composite of the pushout product of $\Delta : G/H \times \partial D^n \rightarrow G/H \times G/H \times \partial D^n \times \partial D^n$ and $\partial([0, 1) \times [0, 1)) \rightarrow L$ with the map:

$$G/H \times G/H \times \partial D^n \times \partial D^n \times L \rightarrow G/H \times G/H \times \partial D^n \times \partial D^n \times [0, 1) \times [0, 1)$$

and is therefore an hG -cofibration as desired. If V is the complement in $G/H \times D^n$ of $G/H \times \partial D^n \times [0, 1/2] \subset U$, then V is also numerable and the restriction of ϕ to each of $U \times U$, $V \times G/H \times D^n$ and $G/H \times D^n \times V$ is an hG -cofibration. Therefore, ϕ is an hG -cofibration by [Dol68, Satz 2], and, therefore, $\Delta : X \rightarrow X \times X$ is an hG -cofibration, since it is the composite of a pushout of α with a pushout of ϕ . Moreover, $X \rightarrow X \times X$ can be represented by a G -NDR-pair that extends (H, λ) by Lemmas 27 and 28. The lemma now follows by passage to colimits. $\square$

As in [DE72], we make the following definition:

Definition 17: *We call a G -space, X , G -equivariantly locally equiconnected (or G -LEC) if $\Delta : X \rightarrow X \times X$ is an hG -cofibration.*

The next lemma is the equivariant analogue of [DE72, Theorem II.3]:

Lemma 18: *Let X be G -LEC and suppose that $x \in X$ has isotropy group H . Then there exists an H -equivariant open neighbourhood U of x such that $\pi_1, \pi_2 : U \times U \rightarrow X$ are H -homotopic. Here, π_i denotes the projection onto the i th factor followed by the inclusion into X .*

Proof. Let (H, λ) represent the inclusion of the diagonal $(X \times X, \Delta)$ as a G -NDR pair. Identify X with $\{x\} \times X$ and let $U = \lambda^{-1}([0, 1)) \cap (\{x\} \times X)$. Consider $H : \{x\} \times U \times I \rightarrow X \times X$. Then $\pi_2 H$ defines an H -homotopy from the inclusion $i : U \rightarrow X$ to some map f . Moreover, $\pi_1 H$ defines an H -homotopy from f to the constant map to x . If we now consider $\pi_1, \pi_2 : U \times U \rightarrow X$, we see that π_1 is the composite $U \times U \rightarrow X \times U \xrightarrow{\pi_1} X$, which is H -homotopic to the constant map to x . Similarly, π_2 is H -homotopic to the constant map to x , so π_1 and π_2 are H -homotopic. $\square$

Lemma 19: *Let X be $\mathcal{C}$ -cofibrant and suppose that $x \in X$ has isotropy group H . Then there exists an H -equivariant open neighbourhood U of x such that $\pi_1, \pi_2 : U \times U \rightarrow X$ are H -homotopic via an H -homotopy which is constant on Δ .*

Proof. By Lemma 16 and Lemma 18, we know that π_1 and π_2 are H -homotopic, say via an H -homotopy $L : U \times U \times I \rightarrow X$. Define $\Gamma : \Delta \times I \rightarrow X$ to be the restriction of L to $\Delta \times I$. Let Γ^{-1} denote the inverse homotopy from π_2 to π_1 . Define K to be composite (via concatenation) homotopy of L and $\Gamma^{-1}\pi_2$:

$$U \times U \times I \xrightarrow{\pi_2} U \times I \cong \Delta \times I \xrightarrow{\Gamma^{-1}} X$$

Then the restriction of K to $\Delta \times I$ is the composite of Γ and Γ^{-1} , and K is an H -homotopy from π_1 to π_2 . The composite of Γ and Γ^{-1} is itself H -homotopic, say via $\alpha : \Delta \times I \times I \rightarrow X$, to the constant homotopy on $\pi_1|_{\Delta}$, and α can be made relative to $\Delta \times \{0, 1\} \times I$. We can then define a lifting problem:

$$\begin{array}{ccc} U \times U \times I \times \{0\} \cup \Delta \times I \times I \cup U \times U \times \{0, 1\} \times I & \xrightarrow{K \cup \alpha \cup \text{const}(\{\pi_1, \pi_2\})} & X \\ \downarrow & & \downarrow \\ U \times U \times I \times I & \xrightarrow{\hspace{10em}} & * \end{array}$$

A lift $\tilde{K} : U \times U \times I \times I \rightarrow X$ exists since U is $(\mathcal{C} \cap H)$ -cofibrant, by Corollary 6 and Lemma 1, and, therefore, H -LEC by Lemma 16. This implies the left hand vertical map is an hH -acyclic hH -cofibration. Restricting $\tilde{K}$ to $U \times U \times I \times \{1\}$ gives the desired H -homotopy. $\square$

We will use the following consequence of [Las82, Lemma 2.8], which was noted in [BJ15, Lemma 6.1]:

Lemma 20: *If $p : E \rightarrow B$ is an hH -fibration, then $1 \times_H p : G \times_H E \rightarrow G \times_H B$ is an hG -fibration.*

Proof. Consider a lifting problem of G -spaces as in the right hand square below:

$$\begin{array}{ccccc} G \times_H V_0 & \xrightarrow{\cong} & X & \xrightarrow{f} & G \times_H E \\ \downarrow & & \downarrow & & \downarrow \\ G \times_H (V_0 \times I) & \xrightarrow{\cong} & X \times I & \xrightarrow{\alpha} & G \times_H B \end{array}$$

Then $X \times I \cong G \times_H \alpha^{-1}(B)$ and so [Las82, Lemma 2.8] tells us that there is an H -homeomorphism $\psi : V_0 \times I \rightarrow V$, where $V = \alpha^{-1}(B)$, $V_0 = V \cap (X \times \{0\})$ and $\psi|_{V_0 \times \{0\}}$ is the inclusion of V_0 into V . Note that $G \times_H V_0 \cong X$. The H -homeomorphism, ψ , induces a G -homeomorphism $G \times_H (V_0 \times I) \rightarrow X \times I$, as in the diagram above, which restricts at $0 \in I$ to a homeomorphism onto $X \times \{0\}$. So to solve the lifting problem of the right hand square, it suffices to solve the adjoint lifting problem of H -spaces:

$$\begin{array}{ccc} V_0 & \xrightarrow{f} & G \times_H E \\ \downarrow & & \downarrow \\ V_0 \times I & \xrightarrow{\alpha\psi} & G \times_H B \end{array}$$

Since $V = \alpha^{-1}(B)$, $\alpha\psi : V_0 \times I \rightarrow G \times_H B$ factors through B , and $f : V_0 \rightarrow G \times_H E$ factors through E . Since p is an hH -fibration, it follows that the lifting problem can be solved. $\square$

A version of the following theorem was first proved in an equivariant context by Gevorgyan and Jimenez in [GJ19, Theorem 6], using the qG -model structure and under the assumption that both E and B are G -CW complexes. The proof below corrects a minor gap in that proof, since it is not necessarily true that a G -CW complex, X , can be covered by G -equivariant open subspaces which are non-equivariantly contractible in X .

Theorem 21: *Suppose that $\mathcal{C}$ is closed under conjugacy and intersections. If $p : E \rightarrow B$ is a $\mathcal{C}$ -fibration between $\mathcal{C}$ -cofibrant spaces, then p is an hG -fibration.*

Proof. Let (H, λ) represent $(B \times B, \Delta_B)$ as a G -NDR pair. As in Lemma 19, if $b \in B$ has isotropy group H there is an H -equivariant open neighbourhood U of b and an H -homotopy $K : U \times U \times I \rightarrow B$ between π_1 and π_2 such that $K(u, v, t) = K(u, v, 1)$ for all $t \geq \lambda(u, v)$. Note that $H \in \mathcal{C}$ since there exists an inclusion of B into a $\mathcal{C}$ -cell complex. Since $p : E \rightarrow B$ is a $(\mathcal{C} \cap H)$ -fibration, $\mathcal{C}$ is closed under intersections and both U and $p^{-1}(U)$ are $(\mathcal{C} \cap H)$ -cofibrant, we can solve the lifting problem:

$$\begin{array}{ccc} U \times p^{-1}(U) & \xrightarrow{\pi_2} & E \\ \downarrow & & \downarrow p \\ U \times p^{-1}(U) \times I & \xrightarrow{K \circ (1 \times p \times 1)} & B \end{array}$$

to define a slicing function $\tilde{K} : U \times p^{-1}(U) \times I \rightarrow E$. We now show that $p : p^{-1}(U) \rightarrow U$ is an hH -fibration. Indeed, given a lifting problem of H -spaces:

$$\begin{array}{ccc} X & \xrightarrow{f} & p^{-1}(U) \\ \downarrow & & \downarrow p \\ X \times I & \xrightarrow{L} & U \end{array}$$

a lift can be defined via $\tilde{L}(x, t) = \tilde{K}(L(x, t), f(x), \lambda(L(x, t), pf(x)))$.

Since B is completely regular, b also has a G -equivariant open neighbourhood G -homeomorphic to $G \times_H S$, where S is an H -equivariant subspace of B containing b , by [Bre72, Theorem 5.4]. Replacing S by $S \cap U$ if necessary, we may as well assume $S \subset U$. In this case, it follows from the above that $p : p^{-1}(S) \rightarrow S$ is an hH -fibration and so $p : G \times_H p^{-1}(S) \rightarrow G \times_H S$ is an hG -fibration by Lemma 20. It follows that for every $b \in B$ there is a G -equivariant neighbourhood, V , of b for which $p : p^{-1}(V) \rightarrow V$ is an hG -fibration. Since B is paracompact, it follows that p is an hG -fibration. $\square$

An alternative proof of Waner's Theorem

Using the results of the previous sections, we can give an alternative proof of Waner's theorem on G -spaces of the G -homotopy type of a G -CW complex, [Wan80, Corollary 4.14]:

Theorem 22: *Let $p : E \rightarrow B$ be an hG -fibration and $* \in B$ a G -fixed basepoint. If E and B have the G -homotopy type of a G -CW complex, then $F := p^{-1}(*)$ has the G -homotopy type of a G -CW complex.*

Proof. The q -model structure induces a model structure on the category $(G - Sp) \downarrow B$, [Hir03, Theorem 7.6.5]. Let $\tilde{p}$ be a fibrant cofibrant approximation to p in this model category, [Hir03, Definition 8.1.22]. We can write $\tilde{p} = qi$, where i is a trivial cofibration and q is a fibration in $(G - Sp) \downarrow B$. Since i has fibrant domain, since p is also a qG -fibration, there is a weak equivalence r such that $ri = 1$. Since we are working in $(G - Sp) \downarrow B$, this construction yields a commutative square:

$$\begin{array}{ccc} \tilde{E} & \xrightarrow{q} & \tilde{B} \\ fr \downarrow \simeq & & \simeq \downarrow g \\ E & \xrightarrow{p} & B \end{array}$$

Since $\tilde{E}$ and $\tilde{B}$ are qG -cofibrant, q is an hG -fibration by Theorem 21. We know that p is an hG -fibration by assumption. The vertical maps are hG -equivalences since E and B are mG -cofibrant. Since g is an

acyclic qG -fibration, we can choose a G -fixed basepoint for $\tilde{B}$ which maps to $*$ under g . We also denote this basepoint by $*$. Since the hG -model structure is right proper, the induced map $q^{-1}(*) \rightarrow F$ is a G -homotopy equivalence. Since $* \rightarrow \tilde{B}$ is an hG -cofibration and q is an hG -fibration, $q^{-1}(*) \rightarrow \tilde{E}$ is an hG -cofibration by [MP12, Lemma 1.3.1]. Since $\tilde{E}$ is qG -cofibrant, it follows that $q^{-1}(*)$ is qG -cofibrant by Theorem 7. Therefore, F is mG -cofibrant as desired. $\square$

Combining the proof above with the proof we gave in [Ron23, Theorem 6.4.6], we have a new proof of the following result:

Theorem 23: *Let $p : E \rightarrow B$ be an hG -fibration, and suppose that B has the G -homotopy type of a G -CW complex. Then E has the G -homotopy type of a G -CW complex iff for every $b \in B$, $p^{-1}(b)$ has the H_b -homotopy type of an H_b -CW complex, where H_b is the isotropy group of b .*

Proof. The $\implies$ direction follows from Theorem 22, the fact that (for any subgroup H of G) an hG -fibration is also an hH -fibration, and the fact that an mG -cofibrant G -space is also mH -cofibrant, [Ron23, Lemma 6.4.5]. The $\impliedby$ direction is [Ron23, Theorem 6.4.6]. $\square$

Appendix of point set topology

Lemma 24: *Let I be a filtered category and let $D : I \rightarrow \mathcal{U}$ be a functor which takes maps in I to closed inclusions. Let $A := \operatorname{colim} D$ and $f : X \rightarrow A$ be a map. Define $Y(i)$ to be the pullback of $D(i) \rightarrow A$ along f . Then the canonical map $\operatorname{colim} Y \rightarrow X$ is a homeomorphism.*

Proof. We can express filtered colimits of closed inclusions, such as A , as quotients of the disjoint union of the colimiting subspaces, $\sqcup_{i \in I} D(i) \rightarrow A$, [Str09, Lemma 3.3]. We have a pullback:

$$\begin{array}{ccc} \sqcup_{i \in I} Y_i & \longrightarrow & X \\ \downarrow & & \downarrow f \\ \sqcup_{i \in I} D_i & \longrightarrow & A \end{array}$$

Since the pullback of a quotient map is a quotient map, [Str09, Propostion 2.36], we obtain the result. $\square$

Lemma 25: *The following properties of a space X are closed under retracts: being i) Hausdorff, ii) paracompact, iii) completely regular.*

Proof. Suppose that we have maps $i : A \rightarrow X, r : X \rightarrow A$ with $ri = 1$. If X is Hausdorff, let a and b be distinct points in A . Then there exists disjoint open sets U and V in X separating $i(a)$ and $i(b)$. So $i^{-1}(U)$ and $i^{-1}(V)$ are disjoint and separate a and b . So A is Hausdorff.

If X is paracompact and $\{U_i\}_{i \in I}$ is an open cover of A , then $\{r^{-1}(U_i)\}$ is an open cover of X which admits a locally finite open refinement $\{V_j\}_{j \in J}$. Then, for each j $i^{-1}(V_j)$ is contained within $i^{-1}r^{-1}(U_i) = U_i$ for some i , so $\{i^{-1}(V_j)\}$ is an open refinement of $\{U_i\}$. Moreover, for every $a \in A$ there is an open neighbourhood, W , of $i(a)$ which intersects only finitely many of the V_j . Then $i^{-1}(W)$ is an open neighbourhood of a , intersecting only finitely many of the $i^{-1}(V_j)$. So A is paracompact.

Next suppose that X is completely regular. Let $a \in A$ and C be a closed subspace of A not containing a . Then $r^{-1}(C)$ is a closed subspace of X not containing $i(a)$. So there exists a continuous function $\lambda : X \rightarrow [0, 1]$ such that $\lambda i(a) = 0$ and $\lambda(r^{-1}(C)) = 1 \implies \lambda i(C) = 1$. So A is completely regular. $\square$

Lemma 26: *Let C be a closed subset of X and define an equivalence relation on $X \times I$ by $(x, t) \sim (y, s)$ iff $x = y$ and either $s = t$ or $x \in C$. Let $f : A \rightarrow X$ and $g : f^{-1}(X/C) \rightarrow I$ be continuous maps. Define a function $F : A \rightarrow (X \times I)/\sim$ by:*

$$F(a) = \begin{cases} (f(a), g(a)) & \text{if } a \in f^{-1}(X/C) \\ (f(a), 1) & \text{otherwise} \end{cases}$$

Then F is continuous.

Proof. Let $V = X/C$ and $\pi : (X \times I)/\sim \rightarrow X$ be the projection. Since $\sim$ is a closed equivalence relation, $(X \times I)/\sim$ is CGWH with the usual quotient topology, [May99, pg. 40]. Since I is compact Hausdorff, π is a closed map. Let $E \subset (X \times I)/\sim$ be closed. Then $F^{-1}(E) \cap f^{-1}(V)$ is closed in $f^{-1}(V)$ by the continuity of f and g . Moreover, $F^{-1}(E) \cap f^{-1}(V) \subset f^{-1}(\pi(E))$, which is closed since π is a closed map and f is continuous. Therefore, there exists a closed subset B of $f^{-1}(\pi(E))$ such that $B \cap f^{-1}(V) = F^{-1}(E) \cap f^{-1}(V)$. Now $F^{-1}(E) \cap f^{-1}(C) = f^{-1}(C \cap \pi(E))$ is closed, since C is closed. So:

$$F^{-1}(E) = B \cup f^{-1}(C \cap \pi(E))$$

is closed, so F is continuous. $\square$

Lemma 27: *Let $i : A \rightarrow B$ be an hG -cofibration represented by an G -NDR-pair (H, λ) and suppose that $j : B \rightarrow C$ is an hG -cofibration. Then there exists a G -NDR-pair, (L, τ) , representing the composite $k := ji : A \rightarrow C$ as an hG -cofibration such that $L|_{B \times I} = H$ and $\tau|_B = \lambda$.*

Proof. Let (K, μ) be an G -NDR-pair representing j such that $K(c, t) = K(c, 0)$ for all $t \leq 1 - \mu(c)$, which is possible by Lemma 26. Let D be the closure of $\mu^{-1}([0, 1))$ and define a map $E : C \times I \cup D \times [0, 2] \rightarrow C$ by the concatenation of K and H . Define:

$$\tilde{\mu}(c) = \begin{cases} 0 & \text{if } \mu(c) \leq \frac{1}{2} \\ 2(\mu(c) - \frac{1}{2}) & \text{if } \mu(c) \geq \frac{1}{2} \end{cases}$$

Now define $L(c, t) = E(c, (1 - t)(1 - \mu(c)) + t(2 - \tilde{\mu}(c)))$ and $\tau(c) = \min\{1, \max\{2\mu(c), \lambda(K(c, 1))\}\}$, if $\mu(c) < 1$, and $\tau(c) = 1$ otherwise. $\square$

Lemma 28: *Given a pushout square:*

$$\begin{array}{ccc} A & \xrightarrow{f} & C \\ i \downarrow & & \downarrow j \\ B & \xrightarrow{g} & D \end{array}$$

where i is an hG -cofibration represented by the G -NDR-pair (H, λ) , then there exists a G -NDR-pair (K, μ) representing j as an hG -cofibration such that $K \circ (g \times 1) = H$ and $\mu g = \lambda$.

Proof. We can define K using the relations $K \circ (g \times 1) = H$ and $K \circ (j \times 1) = \pi_C$ and the universal property of pushouts. Similarly for μ . $\square$

REFERENCES

- [BJ15] Alexander Bykov and Raúl Juárez Flores. G -fibrations and twisted products. *Topology and its Applications*, 196:379–397, 2015. Proceedings of the International Conference on Topology and Geometry 2013, joint with the Sixth Japan-Mexico Topology Symposium (September 2-6, 2013, Matsue, Japan).
- [Bre72] G. Bredon. *Introduction to Compact Transformation Groups*, volume 46 of *Pure and Applied Mathematics*. Elsevier, 1972.
- [DE72] Eldon Dyer and S. Eilenberg. An adjunction theorem for locally equiconnected spaces. *Pacific Journal of Mathematics*, 41(3):669 – 685, 1972.
- [Dol63] Albrecht Dold. Partitions of unity in the theory of fibrations. *Annals of Mathematics*, 78(2):223–255, 1963.
- [Dol68] A. Dold. Die homotopieerweiterungseigenschaft (=hep) ist eine lokale eigenschaft. *Inventiones mathematicae*, 6(3):185–189, 1968.
- [GG16] S. Gorchinskiy and V. Guletskiĭ. Symmetric powers in abstract homotopy categories. *Advances in Mathematics*, 292:707–754, 2016.
- [GJ19] P. S. Gevorgyan and R. Jimenez. On equivariant fibrations of g -cw-complexes. *Sbornik: Mathematics*, 210(10):1428, oct 2019.
- [Hir03] P. Hirschhorn. *Model Categories and Their Localizations*, volume 99 of *Mathematical Surveys and Monographs*. AMS, 2003.
- [Ill83] S. Illman. The equivariant triangulation theorem for actions of compact lie groups. *Mathematische Annalen*, 262:487–501, 1983.
- [Kan07] Marja Kankaanrinta. Equivariant collaring, tubular neighbourhood and gluing theorems for proper Lie group actions. *Algebr. Geom. Topol.*, 7:1–27, 2007.
- [Las82] R. K. Lashof. Equivariant bundles. *Illinois Journal of Mathematics*, 26(2):257 – 271, 1982.
- [May72] J.P. May. *The Geometry of Iterated Loop Spaces*. Lecture Notes in Mathematics. Springer Berlin, Heidelberg, 1972.
- [May99] J.P. May. *A Concise Course in Algebraic Topology*. Chicago Lectures in Mathematics. The University of Chicago Press, 1999.

- [MP12] J. P. May and K. Ponto. *More Concise Algebraic Topology: localization, completion, and model categories*. Chicago lectures in mathematics. University of Chicago Press, 2012.
- [Ron23] A. Ronan. *Nilpotent Groups, Spaces and G-Spaces*. PhD thesis, University of Warwick, 2023. https://wrap.warwick.ac.uk/id/eprint/187006/1/WRAP_Theses_Ronan_2023.pdf.
- [RV22] Emily Riehl and Dominic Verity. *Elements of infinity-category Theory*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2022.
- [Sch18] Stefan Schwede. *Global Homotopy Theory*. New Mathematical Monographs. Cambridge University Press, 2018.
- [Str72] A. Strøm. The homotopy category is a homotopy category. *Arch. Math.*, 23:435–441, 1972.
- [Str09] Neil P. Strickland. The category of CGWH spaces. 2009. <https://ncatlab.org/nlab/files/StricklandCGHWSpaces.pdf>.
- [Wan80] Stefan Waner. Equivariant homotopy theory and Milnor’s theorem. *Transactions of the American Mathematical Society*, 258(2):351–368, 1980.