

Galois Theory

Andrew Tran

1 Polynomial Rings

1.1 Adjunctions and their Properties

Definition 1. Let \mathcal{C} and \mathcal{D} be categories. Then an adjunction from \mathcal{C} to \mathcal{D} is a 4-tuple $(\eta, \epsilon) : F \vdash G$, where $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ are functors, $\eta : 1_{\mathcal{C}} \Rightarrow GF$ and $\epsilon : FG \Rightarrow 1_{\mathcal{D}}$ are natural transformations and the following conditions hold:

1. For every $X \in \text{Ob}\mathcal{C}$, $\epsilon_{FX} \circ F\eta_X = 1_{FX}$
2. For every $Y \in \text{Ob}\mathcal{D}$, $G\epsilon_Y \circ \eta_{GY} = 1_{GY}$

1.2 Polynomial Ring Adjoints

Definition 2. Let $CRing$ be the category of commutative rings and ring homomorphisms and $CRing^*$ be the category of pointed commutative rings and pointed ring homomorphisms. Then the functor

$$U : CRing^* \rightarrow CRing : [\phi : (R, a) \rightarrow (S, b)] \mapsto [\phi : R \rightarrow S]$$

is called the forgetful functor.

Definition 3. A polynomial ring adjoint is a triple (Π, η, ϵ) such that

$$(\eta, \epsilon) : \Pi \vdash U$$

is an adjunction from $CRing$ to $CRing^*$.

For the remainder of these notes, let (Π, η, ϵ) be a fixed polynomial ring adjoint. We define $\pi = U\Pi$.

Definition 4. Let R be a commutative ring and let x be the distinguished point of the pointed ring ΠR . Then ΠR is referred to as the polynomial ring over R with indeterminate element x . The elements $p \in \Pi R$ are referred to as (formal, univariate) polynomials over R .

Definition 5. Let (R, a) be a pointed commutative ring. Then we define $eval_{(R,a)} = U\epsilon_{(R,a)} : \pi R \rightarrow R$ and refer to $eval_{(R,a)}$ as the evaluation homomorphism on R at a .

Definition 6. Let R be a commutative ring. Then we refer to $\eta_R : R \rightarrow \pi R$ as the R embedding homomorphism.

Theorem 1. For any pointed commutative ring (R, a) , $\eta_R : R \rightarrow \pi R$ is injective and $eval_{(R,a)} : \pi R \rightarrow R$ is surjective.

Proof. Let (R, a) be a pointed commutative ring. By definition of adjunctions,

$$eval_{(R,a)} \circ \eta_R = U\epsilon_{(R,a)} \circ \eta_{U(R,a)} = 1_{U(R,a)}$$

In other words, $eval_{(R,a)}$ is a right invertible function and η_R is a left invertible function. This implies that they are surjective and injective, respectively. \square

Definition 7. Let R be a commutative ring. We define the subring of constant polynomials over R , $\overline{R} \subseteq \pi R$, to be the subring of πR given by the image of R under the embedding homomorphism η_R .

Lemma 1. For any commutative ring R , $R \cong \overline{R}$.

Proof. It has already been proven that $\eta_R : R \rightarrow \overline{R}$ is injective and is, by construction, surjective on \overline{R} . \square

Theorem 2. Let R be a commutative ring. Suppose $x \in \pi R$ is the indeterminate element. Then the following are equivalent:

1. x is a unit in πR
2. $x \in \overline{R}$
3. R is a zero ring
4. πR is a zero ring

Theorem 3. Let R be a commutative ring. Then the indeterminate element $x \in \pi R$ is not a zero divisor in πR .

Theorem 4. Let R be a non-trivial commutative ring and $x \in \pi R$ be the indeterminate element. Then for any polynomial $p \in \pi R$, $xp \neq 1_{\pi R}$.