

A NOTE ON THE L^2 -NORM OF THE SECOND FUNDAMENTAL FORM OF ALGEBRAIC MANIFOLDS

ANDREA LOI, MICHELA ZEDDA

ABSTRACT. Let $M \xrightarrow{f} \mathbb{CP}^n$ be an algebraic manifold of complex dimension d and let σ_f be its second fundamental form. In this paper we address the following conjecture: *if $\|\sigma_f\|_{L^2}^2 < 2d \operatorname{vol}(\mathbb{CP}^d)$ then M is totally geodesic and equality holds iff f is congruent to the standard embedding of the complex quadric Q_d into \mathbb{CP}^n .* We prove the conjecture in the following three cases: (i) $d = 1$; (ii) M is a complete intersection; (iii) the scalar curvature of M is constant.

1. INTRODUCTION AND STATEMENT OF MAIN RESULT

In [5] M. Gromov conjectures that every *smooth* immersion $f : M \rightarrow \mathbb{CH}^n/G$ of a compact manifold M of dimension d into a compact quotient of the complex hyperbolic space \mathbb{CH}^n/G , whose second fundamental form σ_f is “small”, is homotopic to a totally geodesic submanifold.

In [2] G. Besson, G. Courtois and S. Gallot give an answer to this problem in terms of the L^2 and L^{2d} norms of the second fundamental form σ_f , when the immersion is a *holomorphic* map:

Theorem 1. *Let $f : M \rightarrow \mathbb{CH}^n/G$ be a holomorphic immersion of a compact Kähler manifold M of complex dimension d . If $\|\sigma_f\|_{L^2}^2$ and $\|\sigma_f\|_{L^{2d}}^2$ are smaller than a constant depending only on n , then M is totally geodesic.*

Date: February 21, 2010.

2000 Mathematics Subject Classification. 53C42; 53C55.

Key words and phrases. Kähler metrics; holomorphic maps into projective space; algebraic manifolds; degree .

Research partially supported by GNSAGA (INdAM) and MIUR of Italy.

It is natural to ask what happens if the ambient space is replaced by its compact dual, namely the complex projective space \mathbb{CP}^n endowed with the Fubini–Study metric g_{FS} of holomorphic sectional curvature 1. So, let $M \xrightarrow{f} \mathbb{CP}^n$ be a complex d -dimensional algebraic manifold (f is a holomorphic injective immersion) and denote by σ_f the second fundamental form of f , by $\|\sigma_f\|^2$ its length and by

$$\|\sigma_f\|_{L^2}^2 = \int_M \|\sigma_f\|^2 \frac{\omega^d}{d!}$$

its L^2 -norm, where ω is the Kähler form associated to the induced metric $g = f^*g_{FS}$. Observe that

$$\|\sigma_f\|^2 = \sum_{j,k=1}^{2d} g_{FS}(\sigma_f(e_j, e_k), \sigma_f(e_j, e_k)),$$

where $\{e_1, \dots, e_d, Je_1, \dots, Je_d\}$ is an orthonormal basis for $T_x M$ (here J denotes the complex structure on M). If $\{e_1, \dots, e_d, Je_1, \dots, Je_d\}$ is a basis which diagonalizes the quadratic form

$$\tilde{\sigma}_f(X, Y) = \sum_{j=1}^{2d} g_{FS}(\sigma_f(e_j, X), \sigma_f(e_j, Y)), \quad X, Y \in T_x M,$$

and $\eta_1^2, \dots, \eta_{2d}^2$ are its eigenvalues, then we can write

$$\|\sigma_f\|^2 = \sum_{j=1}^{2d} \eta_j^2.$$

Observe that by $\sigma_f(X, JY) = \sigma_f(JX, Y) = J\sigma_f(X, Y)$ for all $X, Y \in T_x M$ it follows that $\eta_j^2 = \eta_{j+d}^2$ for $j = 1, \dots, d$.

In this paper we address the problem of finding the optimal constant $c(d)$ (depending only on d) such that if $\|\sigma_f\|_{L^2}^2 < c(d)$ then M is totally geodesic. Similar questions for $\|\sigma_f\|^2$ have been addressed and studied by several mathematicians (cfr. [3], [4], [7], [8], [9]). In particular, in the next section we recall the result by J. Cheng [3] which proves a long standing conjecture posed by K. Ogiue [7].

We believe in the validity of the following:

Conjecture. *Let $M \xrightarrow{f} \mathbb{CP}^n$ be as above. If $\|\sigma_f\|_{L^2}^2 < 2d \operatorname{vol}(\mathbb{CP}^d)$ then M is totally geodesic and equality holds iff f is congruent to the standard embedding of the complex quadric*

$$Q_d = \{[Z_0, \dots, Z_{d+1}], Z_0^2 + \dots + Z_{d+1}^2 = 0\} \subset \mathbb{CP}^{d+1} \xrightarrow{i} \mathbb{CP}^n,$$

where i is the natural inclusion.

Remark 2. Recall that $M \xrightarrow{f} \mathbb{CP}^n$ is totally geodesic, i.e. $\sigma_f \equiv 0$, if and only if M is biholomorphic to \mathbb{CP}^d and $f = A \circ i$, where $A \in \operatorname{Aut}(\mathbb{CP}^n)$ and $i: \mathbb{CP}^d \hookrightarrow \mathbb{CP}^n$ is the natural inclusion, i.e. $i([Z_0, \dots, Z_d]) = [Z_0, \dots, Z_d, 0, \dots, 0]$. Furthermore, observe that for $d = 1$, $Q_1 = (\mathbb{CP}^1, 2g_{FS})$ and f is (congruent to) the Veronese embedding

$$[Z_0, Z_1] \mapsto [Z_0^2, Z_0Z_1, Z_1^2, 0, \dots, 0].$$

Here is the main result of the present paper:

Theorem 3. *Let $M \xrightarrow{f} \mathbb{CP}^n$ be an algebraic manifold of complex dimension d which satisfies one of the following conditions:*

- (i) $d = 1$;
- (ii) M is a complete intersection;
- (iii) the scalar curvature ρ of M is constant.

If

$$\|\sigma_f\|_{L^2}^2 < 2d \operatorname{vol}(\mathbb{CP}^d)$$

then M is totally geodesic and, if equality holds, i.e. $\|\sigma_f\|_{L^2}^2 = 2d \operatorname{vol}(\mathbb{CP}^d)$, then f is congruent to the standard embedding of the complex quadric Q_d .

The paper contains two other sections. In the next one we summarize the background material, while the last one is entirely dedicated to the proof of Theorem 3.

Acknowledgements. We wish to thank Prof. Sylvestre Gallot for interesting and stimulating discussions.

2. PRELIMINARIES

Let $\{e_1, \dots, e_d, Je_1, \dots, Je_d\}$ be an orthonormal basis of $T_x M$ as in the previous section and let us denote $Je_j = e_{d+j}$, $j = 1, \dots, d$. From the Gauss–Codazzi formula (see e.g. [6, Prop. 9.5, Ch. IX])

$$\text{Ric}_g(X, X) = \frac{1}{2}(d+1)g(X, X) - \sum_{j=1}^{2d} g_{FS}(\sigma_f(e_j, X), \sigma_f(e_j, X)), \quad (1)$$

we obtain (cfr. [2])

$$\text{Ric}_g = \frac{1}{2} \sum_{j=1}^d (d+1 - 2\eta_j^2) (e_j^* \otimes e_j^* + (Je_j)^* \otimes (Je_j)^*). \quad (2)$$

If ρ is the scalar curvature for M , namely the smooth function on M defined by

$$\rho = \sum_{j=1}^{2d} \text{Ric}_g(e_j, e_j),$$

then by (2) we get

$$\rho = d(d+1) - \|\sigma_f\|^2. \quad (3)$$

This formula together with the inequality

$$\int_M (\rho - d^2) (\rho - d(d+1)) \frac{\omega^d}{d!} \geq 0,$$

which is obtained by using algebro-geometric machinery, are the key ingredients for the proof of the following result needed in the proof of Theorem 3:

Lemma 4 (J. Cheng [3]). *Let $M \xrightarrow{f} \mathbb{CP}^n$ be as above. If $\|\sigma_f\|^2 < d$ then M is totally geodesic and equality holds iff f is congruent to the standard embedding of the complex quadric Q_d .*

The proof of Theorem 3 relies on the concept of degree $\deg(f)$ of $M \xrightarrow{f} \mathbb{CP}^n$. Given a holomorphic immersion $f: M \rightarrow \mathbb{CP}^n$, if $\dim(M) = d < n$ by Sard's Theorem there exists a point $q \notin f(M)$. Up to unitary transformation of \mathbb{CP}^n we can suppose q to be the point of coordinates $[1, 0, \dots, 0]$. Consider the projection $p_n: \mathbb{CP}^n \setminus \{q\} \rightarrow \mathbb{CP}^{n-1}$, $p_n([Z_0, \dots, Z_n]) = [Z_1, \dots, Z_n]$ and

define the map $F: M \rightarrow \mathbb{CP}^d$ by $F = \tilde{p} \circ f$, where $\tilde{p} = p_{d+1} \circ \cdots \circ p_n$. The degree $\deg(f)$ of f is by definition the degree $\deg(F)$ of the map F , which is the integer number such that

$$\langle F^* \alpha, [M] \rangle = \deg F \langle \alpha, [\mathbb{CP}^d] \rangle, \quad (4)$$

where $[\alpha] \in H^{2d}(\mathbb{CP}^d, \mathbb{R})$ and

$$\langle \alpha, [\mathbb{CP}^d] \rangle = \int_{\mathbb{CP}^d} \alpha, \quad \langle F^* \alpha, [M] \rangle = \int_M F^* \alpha.$$

What we need about $\deg(f)$ is summarized in the following:

Lemma 5 (W. Wirtinger [10], M. Barros, A. Ros, [1]). *The degree $\deg(f)$ is a positive integer such that*

$$\text{vol}(M) = \deg(f) \text{vol}(\mathbb{CP}^d), \quad (5)$$

where $\text{vol}(M) = \int_M \frac{\omega^d}{d!}$ and $\text{vol}(\mathbb{CP}^d) = (4\pi)^d/d!$. Moreover, $\deg(f) = 1$ iff M is totally geodesic and $\deg(f) = 2$ iff f is congruent to the standard embedding of Q_d .

Observe that (5) follows easily by the definition of $\deg(f)$ above. In fact, if we denote by $\omega_{FS}(n)$ (resp. $\omega_{FS}(d)$) the Fubini–Study metric on \mathbb{CP}^n (resp. \mathbb{CP}^d), we have

$$\langle f^* \omega_{FS}^d(n), [M] \rangle = \int_M \omega^d = d! \text{vol}(M).$$

Since the map $\Psi: \mathbb{CP}^n \times [0, 1] \rightarrow \mathbb{CP}^n$,

$$\Psi([Z_0, \dots, Z_n], t) = [tZ_0, \dots, tZ_{n-d-1}, Z_{n-d}, \dots, Z_n]$$

is a homotopy between the identity map of \mathbb{CP}^d , and $i \circ \tilde{p}$, where $i: \mathbb{CP}^d \rightarrow \mathbb{CP}^n$ is the canonical inclusion (cfr. Remark 2), we get

$$\begin{aligned} d! \text{vol}(M) &= \langle f^* \omega_{FS}^d(n), [M] \rangle = \langle (i \circ F)^* \omega_{FS}^d(n), [M] \rangle = \langle F^* (i^* \omega_{FS}^d(n)), [M] \rangle \\ &= \langle F^* (\omega_{FS}^d(d)), [M] \rangle = \deg(F) \langle \omega_{FS}^d(d), [\mathbb{CP}^d] \rangle \\ &= \deg(f) d! \text{vol}(\mathbb{CP}^d). \end{aligned}$$

3. PROOF OF THEOREM 3

Assume (i) holds. Then $\rho = 2K$, where K is the Gaussian curvature of M . Hence Gauss–Bonnet theorem yields

$$\int_M \rho \frac{\omega^d}{d!} = 4\pi \chi(M),$$

where $\chi(M) = 2 - 2\gamma$ denotes the Euler characteristic of M .

By (3) we have

$$\int_M \rho \frac{\omega^d}{d!} = \int_M (2 - \|\sigma_f\|^2) \frac{\omega^d}{d!} = 2\text{vol}(M) - \|\sigma_f\|_{L^2}^2,$$

thus

$$\|\sigma_f\|_{L^2}^2 = 2\text{vol}(M) - 4\pi \chi(M).$$

If $\|\sigma_f\|_{L^2}^2 < 8\pi$, then $2\text{vol}(M) - 4\pi \chi(M) < 8\pi$. By (5) one gets

$$\deg(f) < 1 + \frac{\chi(M)}{2} = 2 - \gamma.$$

It follows by Lemma 5 that $\deg(f) = 1$ and so $\gamma = 0$ and M is totally geodesic.

If $\|\sigma_f\|_{L^2}^2 = 8\pi$ then $\deg(f) = 2$, $\gamma = 0$ and again by Lemma 5 f is congruent to the Veronese embedding (cfr. Remark 2).

Assume (ii) holds. Let a_1, \dots, a_p , $p = n - d$, be the degrees of the hypersurfaces defining M . Then, by [7, Th. 7.1], we have

$$\int_M \rho \frac{\omega^d}{d!} = d \left(d + p + 1 - \sum_{j=1}^p a_j \right) \left(\prod_{j=1}^p a_j \right) \text{vol}(\mathbb{CP}^d),$$

and, since $\deg(f) = \prod_{j=1}^p a_j$, by (3) and (5) we gets

$$\|\sigma_f\|_{L^2}^2 = d \left(\sum_{j=1}^p a_j - p \right) \left(\prod_{j=1}^p a_j \right) \text{vol}(\mathbb{CP}^d).$$

If $\|\sigma_f\|_{L^2}^2 < 2d \text{vol}(\mathbb{CP}^d)$, we have

$$\left(\sum_{j=1}^p a_j - p \right) \left(\prod_{j=1}^p a_j \right) < 2,$$

and since each a_j 's is an integer greater than or equals to 1, we get $a_j = 1$ for all $j = 1, \dots, p$. So $\deg(f) = 1$ and by Lemma 5 M is totally geodesic.

If $\|\sigma_f\|_{L^2}^2 = 2d \operatorname{vol}(\mathbb{CP}^d)$ we get

$$\left(\sum_{j=1}^p a_j - p \right) \left(\prod_{j=1}^p a_j \right) = 2.$$

Thus $\deg(f) = \prod_{j=1}^p a_j = 2$ and the conclusion follows once again by the last part of Lemma 5.

Finally, assume (iii) holds which, by (3), implies $\|\sigma_f\|^2$ is constant. If $\|\sigma_f\|^2 < d$ (resp. $\|\sigma_f\|^2 = d$) then f is totally geodesic (resp. congruent to the quadric) by Lemma 4. If $\|\sigma_f\|^2 > d$ then

$$d \operatorname{vol}(M) < \|\sigma_f\|_{L^2}^2 < 2d \operatorname{vol}(\mathbb{CP}^d)$$

which, by (5), implies $\deg(f) = 1$, i.e. M is totally geodesic. □

REFERENCES

- [1] M. Barros, A. Ros, *Spectral geometry of submanifolds*, Note di Matematica IV (1984), 1–56.
- [2] G. Besson, G. Courtois, S. Gallot, *Lemme de Schwarz réel et applications géométriques*, Acta Math. 183 (1999), 145–169.
- [3] J. Cheng, *An integral formula on the scalar curvature of algebraic manifolds*, Proc. of the Amer. Math. Soc. 81 (1981), no. 3, 451–454.
- [4] S. Choi, J. Kim, Y. Pyo, *On complete complex submanifolds with constant scalar curvature in a complex projective space*, Math. Sci. Res. Hot-Line 4 (2000), no. 9, 47–62.
- [5] M. Gromov, *Asymptotic invariants of infinite groups*, Geometry Groups Theory 2 (1991), 1–295, London Math. Soc. Lecture Note Ser., 182 Cambridge Univ. Press (1993).
- [6] S. Kobayashi, K. Nomizu, *Foundations of differential geometry*, vol. II, Interscience Publishers
- [7] K. Ogiue, *Differential Geometry of Kähler Submanifolds*, Advances in Mathematics 13 (1974), 73–114.

- [8] A. Ros, *A characterization of seven compact Kaehler submanifolds by holomorphic pinching*, Annals of Math. 121 (1985), 377–382.
- [9] A. Ros, *Positively curved Kaehler submanifolds*, Proc. of the Amer. Math. Soc. 93 (1985), no. 2, 329–331.
- [10] W. Wirtinger, *Eine Determinantenidentität und ihre Anwendung auf analytische Gebilde in Euclidischer und Hermitischer Massbestimmung* Monatsh. Math. Phys. 44 (1936), 343–365.

DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI CAGLIARI, VIA OSPEDALE 72, 09124 CAGLIARI, ITALY

E-mail address: loi@unica.it; michela.zedda@gmail.com