# ASSESSING THE COMMUNICATION GAP BETWEEN AI MODELS AND HEALTHCARE PROFESSIONALS: EXPLAINABILITY, UTILITY AND TRUST IN AI-DRIVEN CLINICAL DECISION-MAKING

Oskar Wysocki\*<sup>1,2</sup>, Jessica Katharine Davies<sup>3</sup>, Markel Vigo<sup>1</sup>, Anne Caroline Armstrong<sup>3</sup>, Dónal Landers<sup>2</sup>, Rebecca Lee<sup>3</sup>, and André Freitas<sup>1,2,4</sup>

<sup>1</sup>Department of Computer Science, The University of Manchester
 <sup>2</sup>digital Experimental Cancer Medicine Team, Cancer Biomarker Centre,
 CRUK Manchester Institute, University of Manchester
 <sup>3</sup>Faculty of Biology Medicine and Health, The University of Manchester
 <sup>4</sup>Idiap Research Institute

#### **ABSTRACT**

This paper contributes with a pragmatic evaluation framework for explainable Machine Learning (ML) models for clinical decision support. The study revealed a more nuanced role for ML explanation models, when these are pragmatically embedded in the clinical context. Despite the general positive attitude of healthcare professionals (HCPs) towards explanations as a safety and trust mechanism, for a significant set of participants there were negative effects associated with confirmation bias, accentuating model over-reliance and increased effort to interact with the model. Also, contradicting one of its main intended functions, standard explanatory models showed limited ability to support a critical understanding of the limitations of the model. However, we found new significant positive effects which repositions the role of explanations within a clinical context: these include reduction of automation bias, addressing ambiguous clinical cases (cases where HCPs were not certain about their decision) and support of less experienced HCPs in the acquisition of new domain knowledge.

# 1 Introduction

Clinical predictive models based on machine learning (ML) bring the promise of integrating real world evidence into clinical decision-making, balancing individual clinical experience with data-driven evidence. The dialogue between evidence, which is systematically collected and analysed, and in clinical practice, allows for a continuous evolution of the understanding of disease and treatment response. This is particularly important in the context of new diseases (e.g., COVID-19), new treatments or in understanding personalised responses (e.g. multi-morbidity or diverse populations).

Effective communication between healthcare professionals (HCPs) and ML models depends on the ability of the former to have a faithful mental representation of the latter and a critical ability to assess the strengths and limitations of ML models. ML explainability [1, 2, 3, 4, 5, 6] emerged as a research area aiming to address interpretability bottlenecks in ML models resulting from the fact that many of them can be effective in predicting an outcome, but not in explaining their underlying reasoning, which limits their practical application in critical areas. Whilst recent explainable ML methods [7, 8] are instrumenting these models to be more transparent, the perceived utility and suitability of explanation models has not been systematically investigated in the clinical context.

We aimed to address this research gap by conducting a user study on the effectiveness of current modalities of ML explainability amongst healthcare professionals for data-driven decision support. The study was performed using an explainable ML clinical decision support tool designed to help manage the admissions of patients with cancer and COVID-19: CORONET (COVID-19 Risk in ONcology Evaluation Tool [9]). In contrast to existing explainability

<sup>\*</sup>Corresponding author: oskar.wysocki@manchester.ac.uk

studies performed over simplified tasks [10, 11, 12, 13], our experiment closely mimicked a real-world clinical scenario [14], where the decision is made based on multiple factors supported by a recommendation.

As part of the contributions, we propose a novel methodology to assess the pragmatic utility of explanations, which accounts for a systematic understanding of the interaction between a clinician's mental model and the explanable ML model. Additionally, we introduce a framework for evaluating the impact of ML explanations on perceived utility. We present the results of an experiment conducted involving 23 HCPs, who were presented with 10 scenarios and had to decide whether to admit/discharge a patient with cancer and COVID-19 supported by a model firstly without an explanation (colour bar + score) and with explanation (+contributing features).

#### 2 Methods

#### 2.1 Framework for pragmatic evaluation of the model's explanation




Figure 1: Overview of the framework for pragmatic evaluation of model's explanation. It highlights key aspects of the user's mental model that affects the perceived utility of the tool a clinical setting. Pink boxes - negative effects.

We introduce a framework for pragmatic evaluation of the model's explanation, outlining the key components from the model's output (including its explanatory components) to the model's usefulness in a clinical setting (see Fig.1). It allows for investigating key aspects of the user's attitude that impact the perceived utility of the model in a clinical decision-making setting. Our end-to-end approach allows for the evaluation of the change in perception of the overall system's clinical suitability, i.e. evaluation of system-level effect, instead of item-level effect (single recommendation) [14].

First, the ease of interpretation of the tool's output is evaluated, which is a prerequisite for understanding, satisfaction and trust [15, 16]. Note, that in the clinical decision-making process, the cognitive load due to the model output is one of many disjointed contextual factors and thus should not require the full working memory of the user [17].

Then questions regarding trust, satisfaction and model understanding are systematically assessed. Trust in the model leads to its high persuasive power and *reassurance* in cases where the user is less certain about the right decision [18]. Understanding the model's reasoning process supports the critical assessment of the model (i.e. understanding when the tool provide incorrect recommendations). Satisfaction measures how well the provided output corresponds the user's expectations.

Trust may be enhanced by providing the explanation [19, 20, 21], uncertainty of the model's recommendation [21, 18] and/or by ensuring the high performance of the model at a given task [22]. The latter aspect can be characterised by the the declared performance by the model's developer in testing and validation, and the subjective performance perceived by the user during the experiment [23, 14]. However, excessive trust may have negative effects, i.e. *confirmation bias* [24] and *automation bias* [25, 26]. These are evaluated via the concordance between user's action, recommendation and correct action, and via time spent on decisions.

Finally, all aspects account for the perceived pragmatic utility of the model, evaluated via asking whether HCPs would use the model in their clinical practice.

The output of the model may consist of three main components: i) recommendation (R, i.e. prediction, prognosis or classification); ii) uncertainty of the model's recommendation (U); iii) explanation (Exp); and can be presented to the user as: R, R+U, R+Exp and R+U+Exp. In our empirical analysis, as the CORONET model does not communicate its uncertainty, we put aside this aspect and performed a 'R then R+Exp' within-subject scenario with one group of HCPs. Although one of the visual components of the explanation (i.e. scatter plot, described in 2.2) makes the user aware of the model's imperfect performance (predicted scores for the whole training cohort, including incorrect recommendations), we did not evaluate this aspect directly. We did not provide explicitly any performance metric to the HCPs. Similarly, the perceived performance of the model was not measured. Thus, we did not evaluate the contribution of the model's performance to the HCPs' attitude towards the model. Individual characteristics of the user, such as user's background, knowledge and expectations that may impact their attitude [27] were evaluated.

Guided by the framework, we investigate the overall pragmatic impact of the explanations on the clinical decision by targeting the following research questions:

- RQ1 Does the explanation improve understanding of the model's reasoning mechanisms?
- RQ2 Does the explanation lead to an increase in model satisfaction?
- RQ3 Does the explanation lead to an increase in trust?
- RQ4 Do explanations improve the HCPs' critical understanding of the model?
- RQ5 Do explanations enable improved clinical understanding?
- RQ6 How do explanations impact the automation bias?
- RQ7 Does the explanation increase the confirmation bias?

This study indicates that there is a major unaddressed communication gap between explainable ML models and healthcare professionals and elicits priority areas for investigation.


#### 2.2 Clinical setting & supporting model

**Recommendation (R):** The basic output of the model is the recommended action and the CORONET score, which defines action. The score is presented on a colour bar, scaled 0-3 with decision thresholds (Fig.2a). The score below 1.0 recommends discharge, above 1.0 admission, and above 2.3 suggests high risk of severe COVID-19 illness. Critically, this is in a group of patients who may also present to hospital with cancer or treatment related problems and not just COVID-19. This therefore adds complexity to the decision-making for the HCP. However, CORONET is only built to aid decisions regarding COVID-19 severity and requirement for admission. The CORONET model is a regression random forest, which was trained using the ordinal 0-1-2-3 scale as a dependent variable. True outcomes used as *y* are: 0-patient discharged; 1-patient admitted; 2-admitted and required supplemental oxygen; 3-admitted, required O2 and died. Thus, the model predicts the score in range 0-3 and the recommended action is delivered based on a pragmatically defined threshold [9, 28, 29]. CORONET's user interface is depicted in Fig.S.7.


**Model explanation (Exp):** The explanation of the output consists of two visual components. Firstly, it shows a *scatter plot* (Fig.2b) with the predicted score on the x axis for all patients used in model derivation (scores predicted in Leave-One-Out Cross Validation [9]. Each dot represents an individual patient and the colour corresponds to their true outcome. All patients are sorted from left to right according to their predicted score. The plot allows the user to locate the patient in question (marked by a star) in the whole cohort, considering both true outcomes and model's recommendations. The point distribution elicits the model's errors, i.e. some recommendations may be incorrect. For several 'death outcomes' the model predicted scores below admission threshold. Similarly, for some discharged patients the tool recommended admission. This reveals the performance of the model to the user.

Second, the explanation shows a bar plot with features contributing to the individual prediction (later referred to as *contribution plot*, Fig.2c). The length of the bar represents the magnitude obtained using SHAP explanation [7]. The colour shows the direction of the contribution (towards discharge or admission). Features are ordered from top to bottom depending on the contribution. The figure serves as a local explanation of the model, where the bars change for each individual recommendation.


Overall, the model's output consists of three cognitive chunks [20] organized in a three-step hierarchy: i) colour bar with the score; ii) scatter plot; iii) contribution plot. The derivation of the model and its associated source code are available at https://github.com/digital-ECMT/CORONET\_tool.



(a) Recommended action and the CORONET score, which defines action. The score is presented on a colour bar, scaled 0-3 with decision thresholds



(b) The plot shows all patients used for training of the CORONET model. Each dot represents an individual patient. The colour corresponds to their true outcome. The location on the X-axis is determined by the CORONET score based on the individual's data.



(c) The contribution plot with features contributing to the individual prediction obtained using SHAP explanation.

Figure 2: Output of the CORONET tool provided for each individual patient.

#### 2.3 Study design

The study follows a within-subject design whereby each participant uses the tool in two conditions. Firstly, the participant is provided information in the form of a recommended action and a score on a scale of 0-3 (later referred to as **CS** - CORONET score), and told that the higher the patient's score, the more severe the predicted outcome is. Secondly, the participant is then provided with the recommendation, the CORONET score and an explanation of how the tool has arrived at this score for the patient (later referred to as **CS+Exp** - CORONET score plus explanation). This approach minimises any random noise introduced by the variability on user's clinical and technological exposure and previous experience. The order of presented cases was the same for all participants. The study design is depicted in the Fig.3.

The online questionnaire consisted of six stages: i) introductory questions refering to the HCP's background and expectations ii) five artificial patient cases with only the recommendation provided; iii) questions evaluating the model's usefulness; iv) another five artificial cases this time provided with the recommendation and explanation; v) questions evaluating the model's usefulness (same as in iii) vi) overall impressions of the tool. The questions were designed specifically for this study.




Figure 3: Experimental design. The within-subject design enabled examination of the change in attitude towards the model of each participant, as each of them was first supported by CORONET Score (CS) alone and then by CORONET Score and explanation (CS+Exp).

Ten artificial patient case scenarios were constructed, reviewed and approved by a senior oncology fellow and a consultant oncologist. The decision to admit or discharge was based on clinical guidelines and best practice. The scenarios were similarly structured and comprised an introduction to the patient (including demographics, presenting complaint, and relevant past medical history - all 10 cases are detailed in the supplementary material), the patient's parameters from observations (vital signs), and blood test results.

Two of the ten patient cases (Daniel in CS, and Christine in CS+Exp scenarios) were intentionally built in a way that the model would wrongly discharge the patient, as CORONET considers features of COVID-19 severity only [30] and not oncological emergencies which might be concomitant e.g. neutropenic sepsis. In these cases, the healthcare professional should be able to apply their clinical judgement and recognise the patient would require admission for oncological reasons and subsequently override the model's decision. In Daniel's case, the admission is due to renal dysfunction and tumor lysis syndrome following chemotherapy received three days prior for his diagnosis of diffuse large B cell lymphoma. Therefore, this is a scenario where CORONET is thought to potentially 'wrongly discharge' the patient as it does not take into consideration oncological emergencies as part of it's recommendation.. In Christine's case the admission is due to her recent chemotherapy course. Again, because CORONET does not take other types of infections and treatment related presentations, this is a situation where CORONET is not adequate in making an accurate decision.

These cases were included to assess for over-reliance on the model (automation bias). We analysed the concordance between clinicians' decisions, and the model's recommendations and the correct actions approved by our team of experts.

We tracked the time spent on each part of the questionnaire ensuring that it was not quickly skipped when the user lost interest or was engaged with external tasks. It also allowed for tracking the time spent on decision of each patient case.

#### 2.4 Selection of participants

This research project was ethically approved by the research ethics committee [REC reference: 20/WA/0269]. Participants were required to be clinically active healthcare professionals (determined as having patient contact in at least 10% of their working hours) and currently working with patients with cancer presenting with COVID-19. This included senior doctors, junior doctors, physician associates, pharmacists, advanced practitioner nurses, and staff nurses. Any participants who failed to meet these inclusion criteria were excluded. Participants were recruited via email invitations to NHS Trusts, clinician networks, and social media groups.

#### 3 Results

#### 3.1 Participants

23 healthcare professionals participated in the experiment, with various level of experience and expertise (Table 1). The median knowledge regarding managing patients with COVID-19 was five (in one to seven scale), evaluated via direct question. Most of the HCPs felt very comfortable using new technologies. 35% of participants perform statistical analysis at work, and majority of participants do use online calculators or scoring systems, similar to CORONET (83% and 65%, more in Supp Table S.1).

Table 1: Characteristics	of 23 healthcare	nrofessionals	narticinated	in the experiment	

Table 1. Characteristics of 23 heatificate professionals participated	in the experiment.
Age group	n
21-30	6
31-40	10
41-50	3
51-60	4
	n
Advanced Nurse Practitioner or higher	2
Consultant / attending physician	5
Doctor (within 2-4 years of graduating / specialty trainee (ST) / fellow	2
Doctor within first year of graduating (FY1) / intern	1
General practitioner / GPST / community doctor	3
Pharmacist	1
Registered Nurse	1
Specialist Nurse	1
Specialist registrar / ST 3+ / senior resident/senior fellow	7
	Median [min,max]
How comfortable you feel when using new technology?	6 [3,7]
Knowledge on the management of patients with cancer who have developed COVID-19	5 [2,7]

# 3.2 HCPs want to know both contributing features and uncertainty. Uncertainty was considered to be more important than contributing features

The responses regarding the expectations for the ML-based DSS are depicted in Fig.4. 87% (20/23) of HCPs were interested in knowing the features contributing to the model's recommendation, three HCPs had neutral opinions. 91% (21/23) considered the model's explanation of an individual recommendation as important. Apart from knowing *why* the model recommends such action, the uncertainty behind the recommendation was even more essential for the HCPs: 96% (22/23) of them at least slightly agreed, and 43% (10/23) strongly agreed, which is the highest 'strongly agree' proportion in the study.

Knowing the mathematical framework behind the model is significantly less important than the local/global explanation and the model's uncertainty (p < 0.001, Kruskal-Wallis test, Supp Table S.5). Intriguingly, 73% of HCPs who were not

interested in the mathematics behind the model would like to see an associated model explanation and uncertainty. This points into largely unaddressed research questions on the dialogue between explanations and safety properties (such as a representation for uncertainty) as proxies for risk assessment.




Figure 4: Responses to questions related to HCP's expectations regarding a ML model. Q3-Q5 relate directly to the expectation regarding model transparency and explanation. Q3 investigates the need for global explanation, and Q4 asks about local explanation for the "individual patient".




Figure 5: Responses related to HCP's easiness of interpretation of visual output of CORONET and whether it convinces the user to accept the model's recommendation.

# 3.3 Visual explanations were easy to interpret

Explanation visualisations were easy to interpret and contributed to convincing HCPs to accept or reject the model's recommendation (Fig. 5, Supp Table S.2). For the majority of HCPs, the colour bar was easy to interpret (83%; 19/23). Interestingly, the CS (score without further explanation) presented on 0-3 colour scale convinces 48% (11/23) HCPs to accept/reject the recommendation. Among these 11 convinced by CS, the average response to questions regarding expectations are high (>5.7).

We found no significant difference between the ease of interpretation of the colour bar, scatterplot or contribution plot, as well as no difference in their persuasive power (p>0.05, Supp Table S.3). We did not find any significant correlation between questions related to the visual output and either knowledge on the management of patients with cancer who have developed COVID-19, nor to expectations for ML-based DSS (Supp Table S.4). Thus, the perception of the visual output was not affected by expectations, nor the competence in the task.

#### 3.4 Explanations did not significantly impact on the decision-making

In the pairwise comparison of responses between CS and CS+Exp scenarios, we did not find a statistically significant change in the HCPs' attitude towards the model. The results are summarized in Fig. 6, Fig.S.1, Supp Table S.6.

The explanation did not improve satisfaction (RQ2, Fig.6 question A), trust (RQ3, Fig.6 questions B,C,E) nor the understanding of how the model produced the recommendation (RQ1, Fig.6 questions D,F,G). Of note, we observe a slight but not significant positive change (p=0.056) in 'help in cases where I am less confident in the decision on how to proceed'.

The majority of HCPs replied positively (57-74% answers at least 'Slightly agree' in quest. A-C) regarding the CS output alone. We established the following associations between positive responses and HCPs' expertise and expectations:

- The lower the expertise, the more helpful the tool appeared to be, even when no explanation is provided (r = -0.482, p = 0.02, Fig.S.2).
- The higher the need for knowing the contributing features, the less helpful the CS output is likely to be: in making safe decisions (r = -0.653; p = 0.001, Fig.S.3) and in cases where HCPs are less confident (r = -0.553, p = 0.006, Fig.S.3)

Additionally, being convinced by the colour bar was correlated with satisfaction and reassurance in the CS only scenario (r = 0.484, p = 0.019 and r = 0.527, p = 0.01, Fig.S.4).

#### 3.5 Model explanations may have an adverse effect on HCPs

CS-Exp output of the model led to both positive and negative changes in satisfaction, helpfulness of the model, reassurance and understanding. 34% HCPs were more satisfied with CS+Exp than CS, but in 23% their satisfaction decreased (RQ2). Higher satisfaction was weakly correlated with ease in the interpretation of the contribution plot (r=0.436, p=0.038), but not correlated with being convinced by it (p=0.24). HCPs who were convinced by the contribution plot in the CS+Exp scenario (65%; 15/23), showed high levels of satisfaction both without (average 5.4) and an with explanation (average 5.6). This is also true regarding the scatter plot.

26% of HCPs said that the tool with CS+Exp was more helpful in making safe clinical decisions, with 26% stating the opposite. There was a positive correlation (r = 0.63, p = 0.001,  $p_{adj} = 0.135$ ) between change in satisfaction and change in help in making safe decisions. 22% of HCPs felt more reassured in the CS+Exp scenario, while 35% felt the opposite.

The provided explanations did not change the level of understanding of when and why the model produces wrong recommendations for 57% (13/23) of HCPs (RQ1). For 17% the explanation even led to lower understanding. This cannot be attributed to the time spent on the patients' analyses, nor ease in interpreting the diagrams (p > 0.05).

The change in understanding wrong recommendations was associated with the change in reassuring the utility of the tool (r = 0.762, p < 0.001,  $p_{adj} = 0.003$ ), which is the strongest correlation found between changes (Supp Table S.7).

The highest percentage - 52% of positive changes was observed to help in cases where the HCP is less confident. Among them, there were three HCPs who rated their knowledge of managing patients with COVID-19 as 2 (RQ5). However, 22% found CS+Exp less helpful.

#### 3.6 Over-reliance on the model's recommendation leads to wrong decisions

There are only three cases where all the HCPs made the correct decision, and six cases where the concordance with the correct decisions was more than 80% (Fig.7). Of note, the concordance does not depend on whether a patient should be admitted or not when the tool recommends correctly (on average 88% vs 87%, p=0.73).

However, the lowest concordance with the correct decision was observed for cases where CORONET recommended the wrong action (RQ6, Fig.7). In Daniel's case 65% (15/23) HCPs decided to discharge, as CORONET recommended, while for Christine this accounted for 48% (11/23). As the level of complexity and difficulty in decisions across the cases 1-10 was similar, we argue that the lowest concordance for Daniel and Christine is caused by the wrong recommendation provided by the model. The results strongly suggest that for both types of output (CS and CS+Ex) HCPs over-relied in the recommendation provided by the model.




Figure 6: Paired answers for the same questions among 23 Healthcare Professionals. Each dot is an individual answer. Lines: gray - no change; green - positive change up in the Likert scale; red - negative change. No statistically significant change observed in pairwise comparison.




Figure 7: Concordance between decisions made by respondents and correct action approved by our team. Higher is better. (A) - correct decision 'admit', (D) - 'discharge'. Daniel and Christine (orange bars) were cases where CORONET recommends incorrect discharge.

#### 3.7 Quicker decisions due to over-reliance on explanations

Decisions which were in agreement with the model recommendation required less time (RQ6, Table 2, Fig.S.5). For cases 1-5, where the tool provides only the CS, the median time spent on one case was 48s when the user agreed with the tool, and 61s otherwise. For cases 6-10, where the CS+Exp was provided, we observed a statistically significant

Table 2: Time spent on individual decisions for cases 1-5 (CS) and 6-10 (CS+Ex). p values from Mann Whitney-U Test; significance: ns - non significant, p > 0.05, \* - p < 0.05, \*\* - p < 0.01, \*\*\* - p < 0.001

Time [s] median [O1,O3]	Action the same as the recommendation	Action against the recommendation	p	Significance
Cases 1-5, output: CS	48 [34,74]	61 [39,85]	0.097	ns
Cases 6-10, output: CS+Exp	38 [22,64]	84 [50,116]	0.000	***
p	0.042	0.170		
Significance	*	ns		

difference: decisions required 38s for decisions aligned with and 84s (median values) for decisions opposed to the tool's recommendation (p < 0.001, CLES = 0.754, MWU test).

As described in the previous section, some respondents needed less time when deciding on cases where the explanation was delivered. Further investigation of the agreement with the model reveals that when the HCP agrees with the recommendation (RQ7), the combined CS+Exp output leads to significantly quicker decisions than CS output: 38s for CS+Exp and 48s for CS (p=0.04, CLES=0.59). Moreover, when the explanation is provided, it takes more time to decide when the model contradicts the user's decision, but the difference to CS is not significant. Median time for CS output was 61s, compared to 84s for CS+Exp (p=0.17, MWU test).

#### 3.8 Positive feedback, but no attribution to the explanatory model

General feedback regarding the CORONET model is positive (Fig.8). The model and supporting interface is considered easy to use, and the majority of respondents would recommend the tool to their colleagues.




Figure 8: Positive feedback regarding the CORONET evaluated at the end of the experiment.

We found no impact of the explanatory component on the clinical utility of the model (Fig.9), as the explanation did not improve the attitude towards whether to use the CORONET in clinical practice or not (RQ5, p > 0.05, Wilcoxon signed-rank test).

#### 3.9 Respondents' engagement

Our experiment is supported by an online questionnaire, which is to be completed without direct supervision and time limits. Although the questionnaire is estimated to require  $\approx 30 \mathrm{min}$ , the actual average completion time was  $\approx 19 \mathrm{min}$  (median 16 min, Fig.10). Of note, nine HCPs completed it in less than 15 min, spending less than six min on deciding about 10 patients. We also identified several anomalies in the time spent in particular sections which we manually curated (see details in supplementary material). Most likely they were caused by occasional interruptions, which are inherent in the experiment conducted in the clinical setting, where the respondents may be distracted by more urgent matters.

Overall, we did not find a significant difference between total time spent on deciding on patient cases between CS and CS+Exp (p=0.846, Wilcoxon signed-rank test, Fig.S.6). However, 56% (13/23) HCPs spent less time when CS+Exp was provided. We did not find any correlation between this decrease and other aspects investigated in previous paragraphs.




Figure 9: Evaluation of clinical utility of the tool. Same question was asked twice, for CS and CS+Ex. No significant change (p > 0.05).




Figure 10: Total time spent by 23 Healthcare Professionals.

# 4 Discussion: Pragmatic clinical embedding of explainable ML models

#### **CORONET** explanation among the existing taxonomies

The empirical analysis conducted in this study fits into a *domain expert experiment with the exact application task*, which is an application-grounded evaluation, with real humans and real tasks (according to the taxonomy introduced in [20]).

In designing the CORONET output, we followed the *perceptive interpretability* framework [31], emphasising an immediate interpretation: *you see, and you know*. We argue that the contribution plot follows this assumption.

According to taxonomies established in [5] and [12], the output of the tool investigated in this study delivers:

- modularity: each interpretable element of the output (cognitive chunk) appears as a separate visual component. In the contribution plot, each bar delivers a meaningful portion of information and can be interpreted independently (e.g. low contribution of albumin level to recommendation of 'admission')
- **relevance:** it provides insight for a particular audience, in this case clinicians seeking a patient's characteristics critical for the decision (contribution plot), and also how the patient in question can be located in the whole cohort (scatter plot). Additionally, the colour bar and score describe how severe the patient's outcome is expected to be.

However, CORONET's output does not deliver:

- **simulatability:** a model is simulatable when the user can reason and simulate how the model produces the output for an arbitrary input. It could be possible (assuming limited complexity of the model) for algorithms such decision tree or a set of rules. However, as CORONET uses a Random Forest regression model, with multiple (>100) decision trees used to arrive at the prediction, the user cannot precisely elicit the formal decision process.
- unambiguity: CORONET provides local explanations, focusing on individualised patient care. Based on that, it is not possible to reveal how the model behaves in various parts of the feature space. Such behavior was investigated during model derivation [9] with the support of dependency plots. However, these are not presented to the user in the context of this study.

Analogously to unambiguity, the user is not able to evaluate the *descriptive accuracy* [5], which measures how well the relationship learned by the model is reflected in the explanation. Such evaluation was performed during model development, and then intentionally excluded from the cognitive chunks presented to the user. We argue that additional cognitive load would hamper the benefit of such information [32, 16].

#### Characterising HCPs' attitude on the explanations

More than 87% of participants stated that knowing the contribution of features is important, both for the overall model and individual recommendations. The most important aspect, with the highest number of 'strongly agree' answers, is to know the model's uncertainty. This could suggest that HCPs, are self-reportedly more comfortable in assessing risks and uncertainty, rather than interpreting the feature contribution to the output of the ML model. Hence, delivering a model's uncertainty may contribute more to building trust in comparison to a break-down explanation of the features' contributions. However, this conclusion is solely based on the proportion of 'strongly Agree', as we did not find significant differences between expectations regarding uncertainty, feature contribution and individual recommendation (p>0.05, Kruskal-Wallis test, Supp Table S.5). An intuition on the mathematical principles behind the model is perceived as significantly less important (p<0.05).

Showing uncertainty increases trust and the likelihood of following the model prediction [18], under a low cognitive load setting [21]. Limited cognitive investment in the model interpretation would result in a reduced understanding of the output. While the model used in our study does not explicitly provide uncertainty of its prediction, it recommends a binary action based on a numeric score and a threshold, which are shown to the HCPs. We argue it serves for the user as a proxy of the model's recommendation confidence.

#### Unmet need to build pragmatically grounded explanatory models

Although the majority of HCPs (78%; 18/23) found the explanation component accounting for the features' contributions easy to interpret, still 17% of HCPs found the explanation model difficult (RQ2). This points to opportunities for the design and optimisation of explanatory models from the point of view of end-user interaction. Contemporary explanation methods such as SHAP or LIME focus on the calculation of a faithful explanation for the model, abstracting away the pragmatic aspects of this explanation (e.g. their interaction with domain experts). Recent work binding representation design and cognitive models [33, 34] may provide a formal avenue for designing more pragmatically efficient explanatory models.

Explanations in the form of feature importance may act as a safety feature, drawing the user's attention to features omitted by initial judgment but marked as highly relevant by the model [35]. According to [36] the feature-based explanatory models can be in one of three groups: desired (expected by included in the model), ambivalent (user is indifferent about whether they are included or not) or prohibited features (expected to be omitted, following ethical ML principles). Displaying contributing features allows the user to verify whether the model utilises expected features instead of inconsistent or irrelevant features, based on supporting domain knowledge. In our study, we asked the HCPs explicitly whether the model included all desired features. Additionally, no prohibited features were recognized in the model. One of the downsides of feature importance is that it may lead to confirmation bias, both in the model derivation phase [37] and in the post-hoc interpretation of the output. Practically, it is caused by noticing only the features that confirm previous assumptions and overlooking other, potentially incorrect features.

#### Ambiguity on the utility and perceived value of explanations

Depending on the evaluated aspect, 26-57% of HCPs did not change their attitude to the model after the explanation was provided (RQ1). This highlights previous adoption barriers towards ML based decision support tools, i.e. lack of agreement with the model, lack of knowledge of the model, HCPs' attitude toward the model [27, 38]. Despite the widespread notion on the need for safe and explainable models, we found in our study that a step towards such models remained unnoticed or negatively perceived among almost half of the participants (RQ3). For 17-35% of HCPs, a negative effect of explanation was observed. Two factors are likely to be involved: (i) The explanation could introduce further uncertainty in the clinical judgment, as it may have differed from the clinical assessment in particular cases;

(ii) Information overload (too much cognitive effort) or simply not meeting the expectations regarding the explanation model. Further research

This implies that explainable ML researchers should further move from an algorithm-centric perspective [31], to a pragmatic perspective, i.e. prioritising their pragmatic embedding within the experts' decision-making process. Feature contribution, well-received in the data science community, is not necessarily acknowledged among HCPs. Intensifying the dialog and creating feedback loops between explainable AI researchers and domain experts will be an essential part of the development of pragmatically relevant explainable ML models.

Although explainable ML is advocated as the methodological silver bullet for addressing transparency issues [1], some user studies, similarly to this paper, show evidence of a limited or even no impact of the explanation on performance in a task [39, 11, 40, 41, 42]. This highlights the direction of further research in what explanation is expected depending on the context, e.g. clinical setting. Our study shows that we should be careful about our preconceived notions about what needs to be explained.

#### For many HCPs, explanations did not deliver a critical understanding of the model

Explanations did not improve the critical understanding on why the model recommends a different action to the HCPs for 57% of the participants (RQ4). One possible reason may be a misunderstanding of the difference between local and global explanations. The barplot used to visualise feature importance shows the magnitude of the contribution of each feature to the recommendation for an individual patient. For each analysed case the bars can be reordered. This may lead to an extra complexity in interpretation and shows the simplistic design of standard explanation visualisation devices. However, when promoting more individualised patient care one should also expect higher variation in local explanations of the model. Second, the tool is oversimplifying the complexity of the patient and does not refer to the underlying deep biological processes which may explain a biomarker and its relation to the recommended action. The tool is focused on one aspect of the patient (in this scenario COVID-19) and a finite set of predictors. This is often overlooked by the HCPs, and the limitations of the model remained unidentified even with the explanation provided.

#### Explainable models help HCPs to address ambiguous cases

The highest number of positive changes between CS and CS+Exp was observed to support clinical cases where HCPs were less confident in their decisions (RQ5). Interestingly, less experienced HCPs tended to trust the model more. This confirms the findings of [43] and [44] where less experienced clinicians are more likely to rely on the recommendation, changing their initial decision. This suggests that the main function of a ML-based (data-driven) clinical decision support system, such as CORONET, is to be helpful when the user is uncertain, pragmatically positioning the model within the clinical workflow [45]. This is aligned with the direct feedback received from HCPs (see Supp 5). The recommended action, together with the explanation, may deliver new evidence and justifications for more ambiguous cases.

#### Explainable models help HCPs to acquire new domain knowledge

When the user still lacks a consolidated domain prior knowledge, the explanation may act as a guideline or a source of new evidence-based knowledge, pointing into a second function of explainable ML models (RQ5). In [13] explanations only had an impact on HCPs who felt they had insufficient knowledge to achieve their given task, which may signify that the knowledge gap was addressed by the model's output.

## Quicker decisions for explanatory models in contrast to black-box models

57% (13/23) of HCPs spent less time on decisions for cases 6-10 (the cases with explanations), when more information was delivered to end-users (RQ6). Potential reasons are: i) lost interest in the experiment; ii) gained fluency in interpreting the provided information. After the firstfivecases, HCPs familiarized with the layout of the decryption were able to decide on upcoming cases quicker; iii) provided explanations expedited the decision due to over-reliance on the tool. The questionnaire was not designed to verify points i) and ii) leaving this point of ambiguity. Cases 6-10 reflect the same level of difficulty from cases 1-5.

#### Models and explanations can increase confirmation bias

The results suggest that when the recommendation agrees with their initial decision, HCPs decide quicker, when compared to a model disagreement setting (RQ6). Furthermore, when explanations are provided, the decision is comparatively quicker. When the tool contradicts the clinician, explanations lead to a longer reflection on the decision. This highlights the risk of confirmation bias, possibly caused by the explanation that increases the reassurance on the decision (RQ7). This aligns to the results in [1] where users tend to use the explanation to support the justification of their prior decision. The authors found that users reinforced pre-existing beliefs when the explanation supported them but did not abandon them when it was the opposite. Of note, model accuracy also affects the likelihood of adjusting prior beliefs. In some cases, the explanation may appear reassuring even if it does not explain the model [46].

#### Evidence of explanations reducing automation bias

On the other hand, explanations drove users to reflect longer on the decision when disagreeing with the model outcome (RQ6). More time may indicate cognitive forcing, which is reported to reduce over-reliance on the model [16]. At the same time, to reduce the automation bias, the explanation must reduce the cognitive load [32, 26]. Working memory constraints, together with increased cognitive load, lead to higher uncertainty in decision-making processes, thus increasing automation bias. This is particularly true in clinical settings where multiple contextual factors affect the reasoning process of HCPs [19].

#### Over-reliance on the model's recommendation

Some evidence in the literature suggests models that use explainability techniques can hamper the user's ability to detect when a serious error is made [41]. The explanation may excessively increase users' confidence in an algorithmic decision, communicating a false impression of correctness and rigour and resulting in decreased vigilance and auditing of the output [47, 48]. In addition, [25] and [49] reported higher error rates when decisions relied too heavily on automation, resulting in bias towards a wrong action despite, evidence available which should have resulted in the user overriding the model. Automation bias is also linked to higher cognitive load and to consistently higher accuracy of the model [32], and familiarity with the task [13]. In our study, we identified over-reliance on the model for the cases where CORONET was intentionally wrong (RQ6). For these incorrect recommendations, the HCP should have overridden the decision and admitted the patient. Only 35% (CS scenario) and 52% (CS+Ex) HCPs took that critical perspective over the model.

#### 4.1 Assumptions & limitations of this study

**General concordance among clinical judgment.** In the assessment of AI models for decision support in healthcare, a well-known challenge is the definition of a ground truth [50], as decisions may vary across HCPs. Thus, a panel of HCPs is recommended rather than a small number of annotators. In our study, the 'correct' actions were defined by a panel of two experienced oncologists.

Lack of controlled environment. An ideal environment for HCPs to participate would be at a face-to-face setting. Due to the pandemic and social distancing measures, the study was developed in an online setup. In this context, devices used by individual HCPs could not be controlled, which may have impacted on the consistency and the quantity of the usability feedback on each type of device. Despite this, the developers of CORONET designed its interface for cross-device compatibility (e.g. phones, tablets, and computers), and therefore by undertaking this method of design, we were able to capture feedback on CORONET's usability based on a mixture of device types.

**Rush in completing the questionnaire.** We argue that attempting to understand the model and therefore building trust in the tool require a substantial time commitment for some HCPs. Designing a study which is supervised, better controlled, and which better incentiveses HCPs for a longer time commitment would improve the current study design.

**Use of simulated patient cases.** To preserve the confidentiality of the patients, artificial patient case scenarios were used instead of real-life data. The cases were constructed in a clinically consistent manner by two domain experts. However, this does not take away the fact that in real life, a HCP would undertake clinical observations of the patient and account for any visible signs of distress and illness, such as any skin changes, unusual responses, and patient examination findings, which is not possible to reproduce in an artificial setting.

#### 5 Conclusions

In addition to the acceptability of the predictive model [24, 51], a pragmatic evaluation of the user-model interaction is key to a successful deployment of ML-based recommendation tools in healthcare. This paper contributes a pragmatic evaluation framework for explainable ML models for clinical decision support. The study used a within-subject design involving 23 healthcare professionals and compared an explainable ML model with a black-box model. The study revealed a more nuanced role of ML explanation models when these are pragmatically embedded in the clinical context. HCPs acknowledged the role of explanations as a safety and trust mechanism. Communicating the uncertainty behind the model emerged as a stronger requirement, when compared to the explanations.

Despite the general positive attitude towards explanations, for a significant set of participants (17-35%), a negative effect for the explanations was observed, possibly due to an increase in cognitive effort. Moreover, explanations did not improve the critical understanding of the model for 57% of the participants (i.e. their ability to detect an error in the model). We also found that explanations can increase confirmation bias, possibly accentuating over-reliance in the model. On the other hand, explanations drove HCPs to reflect longer on the decision when disagreeing with the model outcome, evidencing explanations as a possible mechanism for reducing automation bias. There was strong evidence

that explainable models better supported HCPs to address ambiguous clinical cases (cases where HCPs were not certain about their decision). Also, explainable models helped less experienced HCPs to acquire new domain knowledge.

This work points to still open research questions in the area, including the need for further pragmatic evaluation of explainable models in complex clinical workflows, the co-development of models of explanations with domain experts and a better understanding of the dialogue between other model safety mechanisms and explanations.

# Acknowledgements

We would like to express our great gratitude to the healthcare professionals who participated in the experiment. We appreciate that they voluntary devoted their time, so limited in the time of pandemic, to complete the experiment. This paper and the research behind it would not have been possible without the significant feedback they delivered.

## **Funding**





























This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 965397

Funding for developing the CORONET online tool has been provided by The Christie Charitable fund (1049751). Dr Rebecca Lee is supported by the National Institute for Health Research.

### References

- [1] Kevin Bauer, Moritz von Zahn, and Oliver Hinz. "Expl(Ai)Ned: The Impact of Explainable Artificial Intelligence on Cognitive Processes". In: SSRN Electronic Journal (2021). ISSN: 1556-5068. DOI: 10/gm9786. URL: https://www.ssrn.com/abstract=3872711.
- [2] Zachary C. Lipton. *The Mythos of Model Interpretability*. Mar. 6, 2017. arXiv: 1606.03490 [cs, stat]. URL: http://arxiv.org/abs/1606.03490.
- [3] Leilani H. Gilpin et al. "Explaining Explanations: An Overview of Interpretability of Machine Learning". In: 2018 IEEE 5th International Conference on Data Science and Advanced Analytics (DSAA). 2018 IEEE 5th International Conference on Data Science and Advanced Analytics (DSAA). Oct. 2018, pp. 80–89. DOI: 10/ggccrr.
- [4] Wojciech Samek and Klaus-Robert Müller. "Towards Explainable Artificial Intelligence". In: *Explainable AI: Interpreting, Explaining and Visualizing Deep Learning*. Ed. by Wojciech Samek et al. Vol. 11700. Lecture Notes in Computer Science. Cham: Springer International Publishing, 2019, pp. 5–22. ISBN: 978-3-030-28953-9 978-3-030-28954-6. DOI: 10.1007/978-3-030-28954-6\_1. URL: http://link.springer.com/10.1007/978-3-030-28954-6\_1.

- [5] W. James Murdoch et al. "Definitions, Methods, and Applications in Interpretable Machine Learning". In: *Proceedings of the National Academy of Sciences* 116.44 (Oct. 29, 2019), pp. 22071–22080. ISSN: 0027-8424, 1091-6490. DOI: 10/ggbhmq. URL: http://www.pnas.org/lookup/doi/10.1073/pnas.1900654116.
- [6] Mokanarangan Thayaparan, Marco Valentino, and André Freitas. "A survey on explainability in machine reading comprehension". In: *arXiv preprint arXiv:2010.00389* (2020).
- [7] Scott M Lundberg and Su-In Lee. "A Unified Approach to Interpreting Model Predictions". In: *Advances in Neural Information Processing Systems*. Vol. 30. Curran Associates, Inc., 2017. URL: https://proceedings.neurips.cc/paper/2017/hash/8a20a8621978632d76c43dfd28b67767-Abstract.html.
- [8] Marco Tulio Ribeiro, Sameer Singh, and Carlos Guestrin. "Anchors: High-precision model-agnostic explanations". In: *Proceedings of the AAAI conference on artificial intelligence*. Vol. 32. 1. 2018.
- [9] Rebecca Lee et al. "Establishment of CORONET; COVID-19 Risk in Oncology Evaluation Tool to identify cancer patients at low versus high risk of severe complications of COVID-19 infection upon presentation to hospital". In: *JCO Clinical Cancer Informatics (in press)* (Mar. 2022).
- [10] Yasmeen Alufaisan et al. *Does Explainable Artificial Intelligence Improve Human Decision-Making?* preprint. PsyArXiv, June 18, 2020. DOI: 10.31234/osf.io/d4r9t. URL: https://osf.io/d4r9t.
- [11] Samuel Carton, Qiaozhu Mei, and Paul Resnick. "Feature-Based Explanations Don't Help People Detect Misclassifications of Online Toxicity". In: *Proceedings of the International AAAI Conference on Web and Social Media* 14 (May 26, 2020), pp. 95–106. ISSN: 2334-0770. URL: https://ojs.aaai.org/index.php/ICWSM/article/view/7282.
- [12] Himabindu Lakkaraju et al. "Faithful and Customizable Explanations of Black Box Models". In: *Proceedings of the 2019 AAAI/ACM Conference on AI, Ethics, and Society*. AIES '19: AAAI/ACM Conference on AI, Ethics, and Society. Honolulu HI USA: ACM, Jan. 27, 2019, pp. 131–138. ISBN: 978-1-4503-6324-2. DOI: 10/ggvzms. URL: https://dl.acm.org/doi/10.1145/3306618.3314229.
- [13] James Schaffer et al. "I Can Do Better than Your AI: Expertise and Explanations". In: *Proceedings of the 24th International Conference on Intelligent User Interfaces*. IUI '19. New York, NY, USA: Association for Computing Machinery, Mar. 17, 2019, pp. 240–251. ISBN: 978-1-4503-6272-6. DOI: 10/ggkkjj. URL: https://doi.org/10.1145/3301275.3302308.
- [14] Eoin M. Kenny et al. "Explaining Black-Box Classifiers Using Post-Hoc Explanations-by-Example: The Effect of Explanations and Error-Rates in XAI User Studies". In: *Artificial Intelligence* 294 (2021), p. 103459. ISSN: 0004-3702. DOI: 10.1016/j.artint.2021.103459.
- [15] Antoine Hudon et al. "Explainable Artificial Intelligence (XAI): How the Visualization of AI Predictions Affects User Cognitive Load and Confidence". In: *Information Systems and Neuroscience*. Ed. by Fred D. Davis et al. Vol. 52. Lecture Notes in Information Systems and Organisation. Cham: Springer International Publishing, 2021, pp. 237–246. ISBN: 978-3-030-88999-2 978-3-030-88900-5. DOI: 10.1007/978-3-030-88900-5\_27. URL: https://link.springer.com/10.1007/978-3-030-88900-5\_27.
- [16] Zana Buçinca, Maja Barbara Malaya, and Krzysztof Z. Gajos. "To Trust or to Think: Cognitive Forcing Functions Can Reduce Overreliance on AI in AI-assisted Decision-making". In: *Proceedings of the ACM on Human-Computer Interaction* 5 (CSCW1 Apr. 22, 2021), 188:1–188:21. DOI: 10.1145/3449287. URL: https://doi.org/10.1145/3449287.
- [17] Mark A. Musen, Blackford Middleton, and Robert A. Greenes. "Clinical Decision-Support Systems". In: *Biomedical Informatics: Computer Applications in Health Care and Biomedicine*. Ed. by Edward H. Shortliffe and James J. Cimino. Cham: Springer International Publishing, 2021, pp. 795–840. ISBN: 978-3-030-58721-5. DOI: 10.1007/978-3-030-58721-5\_24. URL: https://doi.org/10.1007/978-3-030-58721-5\_24.
- [18] Sean McGrath et al. When Does Uncertainty Matter?: Understanding the Impact of Predictive Uncertainty in ML Assisted Decision Making. Nov. 13, 2020. arXiv: 2011.06167 [cs]. URL: http://arxiv.org/abs/2011.06167.
- [19] Divya Ramani et al. "Examining the Patterns of Uncertainty across Clinical Reasoning Tasks: Effects of Contextual Factors on the Clinical Reasoning Process". In: *Diagnosis* 7.3 (Aug. 27, 2020), pp. 299–305. ISSN: 2194-802X, 2194-8011. DOI: 10/gm9785. URL: https://www.degruyter.com/document/doi/10.1515/dx-2020-0019/html.
- [20] Finale Doshi-Velez and Been Kim. "Considerations for Evaluation and Generalization in Interpretable Machine Learning". In: *Explainable and Interpretable Models in Computer Vision and Machine Learning*. Ed. by Hugo Jair Escalante et al. The Springer Series on Challenges in Machine Learning. Cham: Springer International Publishing, 2018, pp. 3–17. ISBN: 978-3-319-98130-7 978-3-319-98131-4. DOI: 10.1007/978-3-319-98131-4\_1. URL: http://link.springer.com/10.1007/978-3-319-98131-4\_1.

- [21] Jianlong Zhou et al. "Effects of Uncertainty and Cognitive Load on User Trust in Predictive Decision Making". In: *Human-Computer Interaction INTERACT 2017*. Ed. by Regina Bernhaupt et al. Vol. 10516. Cham: Springer International Publishing, 2017, pp. 23–39. ISBN: 978-3-319-68058-3 978-3-319-68059-0. DOI: 10.1007/978-3-319-68059-0\_2. URL: https://link.springer.com/10.1007/978-3-319-68059-0\_2.
- [22] Onur Asan, Alparslan Emrah Bayrak, and Avishek Choudhury. "Artificial Intelligence and Human Trust in Healthcare: Focus on Clinicians". In: *Journal of Medical Internet Research* 22.6 (June 19, 2020), e15154. ISSN: 1438-8871. DOI: 10/ghxwzf. URL: http://www.jmir.org/2020/6/e15154/.
- [23] Taehyun Ha et al. "Examining the effects of power status of an explainable artificial intelligence system on users' perceptions". In: *Behaviour & Information Technology* 0.0 (2020), pp. 1–13. DOI: 10.1080/0144929X.2020. 1846789. eprint: https://doi.org/10.1080/0144929X.2020.1846789. URL: https://doi.org/10.1080/0144929X.2020.1846789.
- [24] Marzyeh Ghassemi, Luke Oakden-Rayner, and Andrew L. Beam. "The False Hope of Current Approaches to Explainable Artificial Intelligence in Health Care". In: *The Lancet Digital Health* 3.11 (Nov. 1, 2021), e745–e750. ISSN: 2589-7500. DOI: 10/gm8bdh. pmid: 34711379. URL: https://www.thelancet.com/journals/landig/article/PIIS2589-7500(21)00208-9/fulltext.
- [25] Linda J. Skitka, Kathleen L. Mosier, and Mark Burdick. "Does Automation Bias Decision-Making?" In: *International Journal of Human-Computer Studies* 51.5 (1999), pp. 991–1006. ISSN: 1071-5819. DOI: 10/bg5rb7. URL: https://www.sciencedirect.com/science/article/pii/S1071581999902525.
- [26] Kate Goddard, Abdul Roudsari, and Jeremy Wyatt. "Automation Bias A Hidden Issue for Clinical Decision Support System Use". In: *Studies in health technology and informatics* 164 (Jan. 1, 2011), pp. 17–22. DOI: 10.3233/978-1-60750-709-3-17.
- [27] Srikant Devaraj et al. "Barriers and Facilitators to Clinical Decision Support Systems Adoption: A Systematic Review". In: *Journal of Business Administration Research* 3.2; https://web.archive.org/web/20220303231827/https://www.sciedu.ca/journal/index.php/jbar/article/view/5217 (2 July 24, 2014), p. 36. ISSN: 1927-9515. DOI: 10.5430/jbar.v3n2p36. URL: https://www.sciedu.ca/journal/index.php/jbar/article/view/5217.
- [28] R.J. Lee et al. "Establishment of CORONET; COVID-19 Risk in Oncology Evaluation Tool to identify cancer patients at low versus high risk of severe complications of COVID-19 infection upon presentation to hospital". In: *medRxiv* (2020). DOI: 10.1101/2020.11.30.20239095. URL: https://www.medrxiv.org/content/early/2020/12/03/2020.11.30.20239095.
- [29] Rebecca Lee et al. "CORONET; COVID-19 in Oncology evaluation Tool: Use of machine learning to inform management of COVID-19 in patients with cancer." In: *Journal of Clinical Oncology* 39.15\_suppl (2021), pp. 1505–1505. DOI: 10.1200/JCO.2021.39.15\\_suppl.1505. URL: https://doi.org/10.1200/JCO.2021.39.15\_suppl.1505.
- [30] R.J. Lee et al. "Longitudinal characterisation of haematological and biochemical parameters in cancer patients prior to and during COVID-19 reveals features associated with outcome". In: *ESMO Open* 6.1 (2021), p. 100005. ISSN: 2059-7029. DOI: https://doi.org/10.1016/j.esmoop.2020.100005. URL: https://www.sciencedirect.com/science/article/pii/S2059702920328647.
- [31] Erico Tjoa and Cuntai Guan. "A Survey on Explainable Artificial Intelligence (XAI): Toward Medical XAI". In: *IEEE Transactions on Neural Networks and Learning Systems* 32.11 (Nov. 2021), pp. 4793–4813. ISSN: 2162-2388. DOI: 10.1109/TNNLS.2020.3027314.
- [32] David Lyell and Enrico Coiera. "Automation Bias and Verification Complexity: A Systematic Review". In: *Journal of the American Medical Informatics Association: JAMIA* 24.2 (Mar. 1, 2017), pp. 423–431. ISSN: 1527-974X. DOI: 10/gng5ps. pmid: 27516495.
- [33] Daniel Raggi et al. *Dissecting Representations*. Springer International Publishing, Aug. 24, 2020. ISBN: 978-3-030-54248-1. DOI: 10.17863/CAM.57293; https://web.archive.org/web/20220314103546/https://www.repository.cam.ac.uk/handle/1810/310207. URL: https://www.repository.cam.ac.uk/handle/1810/310207.
- [34] Peter C.-H. Cheng et al. "Cognitive Properties of Representations: A Framework". In: *Diagrammatic Representation and Inference*. Ed. by Amrita Basu et al. Lecture Notes in Computer Science. Cham: Springer International Publishing, 2021, pp. 415–430. ISBN: 978-3-030-86062-2. DOI: 10.1007/978-3-030-86062-2\_43.
- [35] Hilde J. P. Weerts, Werner van Ipenburg, and Mykola Pechenizkiy. *A Human-Grounded Evaluation of SHAP for Alert Processing*. July 7, 2019. arXiv: 1907.03324 [cs, stat]. URL: http://arxiv.org/abs/1907.03324.

- [36] Himabindu Lakkaraju and Osbert Bastani. ""How Do I Fool You?": Manipulating User Trust via Misleading Black Box Explanations". In: *Proceedings of the AAAI/ACM Conference on AI, Ethics, and Society.* AIES '20: AAAI/ACM Conference on AI, Ethics, and Society. New York NY USA: ACM, Feb. 7, 2020, pp. 79–85. ISBN: 978-1-4503-7110-0. DOI: 10/ghj6hb. URL: https://dl.acm.org/doi/10.1145/3375627.3375833.
- [37] I. Elizabeth Kumar et al. "Problems with Shapley-value-based Explanations as Feature Importance Measures". In: *Proceedings of the 37th International Conference on Machine Learning*. International Conference on Machine Learning. PMLR, Nov. 21, 2020, pp. 5491–5500. URL: https://proceedings.mlr.press/v119/kumar20e.html.
- [38] Reed T. Sutton et al. "An Overview of Clinical Decision Support Systems: Benefits, Risks, and Strategies for Success". In: *npj Digital Medicine* 3.1 (Dec. 2020), p. 17. ISSN: 2398-6352. DOI: 10.1038/s41746-020-0221-y. URL: http://www.nature.com/articles/s41746-020-0221-y.
- [39] Jasper van der Waa et al. "Evaluating XAI: A Comparison of Rule-Based and Example-Based Explanations". In: Artificial Intelligence 291 (2021), p. 103404. ISSN: 0004-3702. DOI: 10/gjm97h. URL: https://www.sciencedirect.com/science/article/pii/S0004370220301533.
- [40] Ben Green and Yiling Chen. "The Principles and Limits of Algorithm-in-the-Loop Decision Making". In: *Proc. ACM Hum.-Comput. Interact.* 3.CSCW (Nov. 2019). DOI: 10.1145/3359152. URL: https://doi.org/10.1145/3359152.
- [41] Forough Poursabzi-Sangdeh et al. *Manipulating and Measuring Model Interpretability*. Aug. 15, 2021. arXiv: 1802.07810 [cs]. URL: http://arxiv.org/abs/1802.07810.
- [42] Yunfeng Zhang, Q. Vera Liao, and Rachel K. E. Bellamy. "Effect of Confidence and Explanation on Accuracy and Trust Calibration in AI-assisted Decision Making". In: *Proceedings of the 2020 Conference on Fairness, Accountability, and Transparency*. FAT\* '20. New York, NY, USA: Association for Computing Machinery, Jan. 27, 2020, pp. 295–305. ISBN: 978-1-4503-6936-7. DOI: 10/ggjpcr. URL: https://doi.org/10.1145/3351095.3372852.
- [43] Kate Goddard, Abdul Roudsari, and Jeremy C. Wyatt. "Automation Bias: Empirical Results Assessing Influencing Factors". In: *International Journal of Medical Informatics* 83.5 (May 1, 2014), pp. 368–375. ISSN: 1386-5056. DOI: 10/f2wz3k. URL: https://www.sciencedirect.com/science/article/pii/S1386505614000148.
- [44] Dawn Dowding et al. "Nurses' Use of Computerised Clinical Decision Support Systems: A Case Site Analysis". In: Journal of Clinical Nursing 18.8 (2009), pp. 1159–1167. ISSN: 1365-2702. DOI: 10.1111/j.1365-2702.2008.02607.x. URL: https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2702.2008.02607.x.
- [45] Qian Yang, Aaron Steinfeld, and John Zimmerman. "Unremarkable AI: Fitting Intelligent Decision Support into Critical, Clinical Decision-Making Processes". In: *Proceedings of the 2019 CHI Conference on Human Factors in Computing Systems*. CHI '19. New York, NY, USA: Association for Computing Machinery, May 2, 2019, pp. 1–11. ISBN: 978-1-4503-5970-2. DOI: 10.1145/3290605.3300468. URL: https://doi.org/10.1145/3290605.3300468.
- [46] Julius Adebayo et al. Sanity Checks for Saliency Maps. Nov. 6, 2020. arXiv: 1810.03292 [cs, stat]. URL: http://arxiv.org/abs/1810.03292.
- [47] Marzyeh Ghassemi et al. *ClinicalVis: Supporting Clinical Task-Focused Design Evaluation*. Oct. 13, 2018. arXiv: 1810.05798 [cs]. URL: http://arxiv.org/abs/1810.05798.
- [48] Malin Eiband et al. "The Impact of Placebic Explanations on Trust in Intelligent Systems". In: *Extended Abstracts of the 2019 CHI Conference on Human Factors in Computing Systems*. CHI EA '19. New York, NY, USA: Association for Computing Machinery, May 2, 2019, pp. 1–6. ISBN: 978-1-4503-5971-9. DOI: 10/gmv78z. URL: https://doi.org/10.1145/3290607.3312787.
- [49] Ericka Rovira, Kathleen McGarry, and Raja Parasuraman. "Effects of Imperfect Automation on Decision Making in a Simulated Command and Control Task". In: *Human Factors* 49.1 (Feb. 1, 2007), pp. 76–87. ISSN: 0018-7208. DOI: 10/fmh7q6. URL: https://doi.org/10.1518/001872007779598082.
- [50] Po-Hsuan Cameron Chen, Craig H Mermel, and Yun Liu. "Evaluation of Artificial Intelligence on a Reference Standard Based on Subjective Interpretation". In: *The Lancet Digital Health* 3.11 (Nov. 2021), e693–e695. ISSN: 25897500. DOI: 10/gmv7xj. URL: https://linkinghub.elsevier.com/retrieve/pii/S2589750021002168.
- [51] Alex John London. "Artificial Intelligence and Black-Box Medical Decisions: Accuracy versus Explainability". In: *Hastings Center Report* 49.1 (2019), pp. 15–21. ISSN: 1552-146X. DOI: 10/ghvzdb. URL: https://onlinelibrary.wiley.com/doi/abs/10.1002/hast.973.

# **Supplementary Information**

#### **Supplementary Methods**

# List of questions

Below we list all questions from the experiment, that are relevant in the analysis aiming to answer broader research questions from the previous section:

Questions related to demographics:

- D1. What is your job title?
- D2. What is your age group?
- D3. Which of the following tasks do you perform on computers at work?
- D4. Rate your knowledge on the management of patients with cancer who have developed COVID-19

#### Questions related to expectations:

- Q1. I feel comfortable when using new technology
- Q2. It is important for me to know the mathematics behind the model's recommendations
- Q3. It is important for me to know the features of my patient contribute to the model's recommendation
- Q4. It is important for me to know how the model makes its recommendation for my individual patient
- Q5. It is important for me to know how uncertain (in %) the model is about its recommendation

#### Questions related to the visual representation:

- Q6. The colour bar with the score is easy to interpret
- Q7. The colour bar with the score convinces me to accept or reject the model's recommendation
- Q8. The scatterplot with all patients is easy to interpret
- Q9. The scatterplot with all patients convinces me to accept or reject the model's recommendation
- Q10. The barplot with feature contribution is easy to interpret
- Q11. The barplot with feature contribution convinces me to accept or reject the model's recommendation

#### Ouestions related to the user's attitude towards the model:

- A. I am satisfied with the output information that CORONET provides
- B. CORONET helps me in making safe clinical decisions on patient management
- C. When my initial decision was the same as CORONET had recommended, I felt reassured
- D. I understand when and why CORONET may provide the wrong recommendation in some cases
- E. CORONET helps in cases where I am less confident in the decision on how to proceed
- F. Even when my initial course of action was different to what CORONET recommended, I still had full confidence in my original decision.
- G. I was surprised when CORONET recommended an action different to my own

#### Feedback from Healthcare Professionals

Below we quote received feedback regarding the usability of the tool from the Healthcare Professionals:

- This could be a very useful tool to aid the clinician's decision. With every risk stratification score/tool I'm used to using the MASCC score for febrile neutropenia, it helps your final decision. If my gut said admit, irrespective of any score, I would admit. Tools can be great to add confidence to a less obvious choice of admit/discharge.
- With all the patients I would have discharged home, I still would have asked my team to safety-net call for the next 10-14 days.

- I don't think this can be proposed in clinical practice
- decision to admit a cancer patient is complex and multifactorial often dependent on social support, patient engagement and both cancer and treatment related complications
- Really good tool. Very easy to use. It would be great to get individual hospital trusts to consider adding it to hospital guidelines so doctors felt safer using it to base decisions off.
- Easy to use but differed a lot with my clinical assessment
- Having played with the app a little, it seems like the biggest impact on score comes from NEWS, so I think for many patients it won't add much, but can be a useful adjunct.
- An interesting concept but not one that I think would be used in its current format in clinical care-too formulaic and not individualized to patient risk factors

Additionally, at the end of the experiment Healthcare Professionals were asked about aspects in which the tool could be improved in the future. Below we quote received feedback:

- Inclusion in guidelines, clarify the legal implications
- CORONET is a standalone 'App' to be useful (and used) it has to integrate with patient EPRs (Electronic Patient Record) so HCPs are not copying electronic information from EPR to CORONET this not only takes time but prone to errors This would be a major factor in my decision to recommend CORONET
- May be useful to separate high risk and low risk comorbidities. For example, some conditions such as COPD
  will cause more severe issues when combined with COVID than other conditions such as diabetes.
- It does not seem to take into account oxygen requirements independently of NEWS2; it is difficult to discharge someone who is needing 4L of oxygen!
- Take into consideration if they are active chemotherapy/immunotherapy. As even with a covid positive test, they still require iv antibiotics and monitoring. Higher risk of deterioration, especially lung cancers, if discharged home.
- The wording on the red/yellow diagram seems to be inconsistent re categories. 1st and 2nd are actions e.g. 'consider' admit/discharge, the 3rd is commenting on the disease severity. These are different categories, so the third should be more like 'consider level 2+ care' or another type of action
- The labeling of the red/yellow score is a bit hard to follow, suggest removing the bottom legend

# **Supplementary Tables**

Table S.1: Tasks performed on computers at work by the healthcare professionals participating in the experiment.

Tuble 5.1. Tubks periorified	J11 C	0111	Puu	015	ut **	OII	. 0	uic	1100	ai tii	cuic	, P.	OIC	00101	iiui	Pu	LLICI	Pui	5	111 0	110	· AP	JI 111	iciit.
Emailing	Advanced Nurse Practitioner or higher	Advanced Nurse Practitioner or higher	Consultant/attending physician	Consultant/attending physician	Consultant / attending physician	Consultant / attending physician	Consultant / attending physician	Doctor within first year of graduating (FY1) / intern	Doctor (within 2-4 years of graduating / specialty trainee (ST) / fellow	Doctor (within 2-4 years of graduating / specialty trainee (ST) / fellow	General practitioner / GPST / community doctor	General practitioner / GPST / community doctor	deneral practitioner / GPST / community doctor	Other (please enter text)	Registered Nurse	Specialist Nurse	s Specialist registrar / ST 3+/ senior resident/senior fellow	s Specialist registrar / ST 3+/ senior resident/senior fellow	s Specialist registrar / ST 3+/ senior resident/senior fellow	s Specialist registrar / ST 3+/ senior resident/senior fellow	s Specialist registrar / ST 3+/ senior resident/senior fellow	<ul> <li>Specialist registrar / ST 3+/ senior resident/senior fellow</li> </ul>	× Specialist registrar / ST 3+ / senior resident/senior fellow	96%
Accessing clinical guidelines	X X	X X	X X	X X	X X	X	X X	X X	X X	X X		X X	X X	X X	Х	X X	X X	X X	X X	X X	X X	X X	X X	91%
Viewing test results		λ.		Α.		λ.							λ.											96%
Viewing designation	X	X	X	X	X	X	X	X	Х	X	Х	Х	X	X		X	Х	X	X	X	X	X	X	74%
Viewing drug charts Online clinical calculators	X	X	X	X	Х	Х	X	X		X			X	X		Х		X		X	X	X	X	83%
		X	X	X			X	X	Х	X	Х	X	X	Х	Х		X	X	X	X	Х	X	X	
Online scoring systems and/or algorithms Statistical analysis	X	Х	X	X			X	Х		Х		Х	Х				X	Х	X	Х		X	X	65% 35%
Statistical alialysis	Х		Х				Х									Х	Х				Х	Х	Х	33%

Table S.2: Responses related to ease of interpretation and convincing capacity of the tool's output. No significant difference between answers.

	median respons in Likert scale 1-7 [Q1,Q3]
Of The sale of the decree	III Likert scale 1-7 [Q1,Q5]
Q6. The colour bar with the score	6 [5,6]
is easy to interpret	0 [0,0]
Q7. The colour bar with the score	
convinces me to accept or reject	4 [4,6]
the model's recommendation	
Q8. The scatterplot with all patients	5 [3.5,6]
is easy to interpret	5 [5.5,0]
Q9. The scatterplot with all patients	
convinces me to accept or reject	5 [4,5.5]
the model's recommendation	
Q10. The barplot with feature contribution	5 [5 6]
is easy to interpret	5 [5,6]
Q11. The barplot with feature contribution	
convinces me to accept or reject	5 [4,6]
the model's recommendation	- / <b>-</b>
	p>0.05
	Friedman test

Table S.3: Individual comparisons between reported ease of interpretation of the visuals and their convincing capacity. No significant differences; all p>0.05

			p (Kruskal–Wallis)
Q6. The diagram (colorbar) is easy to interpret	vs.	Q8. The diagram (dotplot) is easy to interpret	0.131
Q10. The diagram (barplot) is easy to interpret	vs.	Q6. The diagram (colour bar) is easy to interpret	0.716
Q10. The diagram (barplot) is easy to interpret	vs.	Q8. The diagram (dotplot) is easy to interpret	0.244
Q11. The diagram (barplot) convinces me to accept or reject the model's recommendation	vs.	Q7. The diagram (colour bar) convinces me to accept or reject the model's recommendation	0.387
Q11. The diagram (barplot) convinces me to accept or reject the model's recommendation	vs.	Q9. The diagram (dotplot) convinces me to accept or reject the model's recommendation	0.281
Q7. The diagram (colour bar) convinces me to accept or reject the model's recommendation	vs.	Q9. The diagram (dotplot) convinces me to accept or reject the model's recommendation	0.740

Table S.4: Spearman correlations between questions regarding ease of interpretation of the visuals or their convincing capacity (Q6-Q11) and questions regarding user competence in the task (D4) or expectations regarding the output (Q1-Q5). No significant correlations were found (Holm adjusted p>0.05).

Spearman correlation r	D4. Rate your knowledge on the management of patients with cancer who have developed COVID-19	Q1. I feel comfortable when using new technology	Q2. It is important for me to know the mathematics behind the model's recommendations	Q3. It is important for me to know the features of my patient contribute to the model's recommendation	Q4. It is important for me to know how the model makes its recommendation for my individual patient	Q5. It is important for me to know how uncertain (in %) the model is about its recommendation
Q6. The colour bar with the score is easy to interpret	0.026	-0.101	-0.199	-0.109	-0.003	-0.187
Q7. The colour bar with the score convinces me to accept or reject the model's recommendation	-0.214	0.062	-0.195	0.031	0	-0.011
Q8. The scatterplot with all patients is easy to interpret	-0.105	0.333	-0.122	-0.309	0.085	-0.107
Q9. The scatterplot with all patients convinces me to accept or reject the model's recommendation	-0.156	0.433	-0.054	-0.283	0.059	-0.167
Q10. The barplot with feature contribution is easy to interpret	-0.368	0.077	-0.288	-0.06	-0.164	-0.044
Q11. The barplot with feature contribution convinces me to accept or reject the model's recommendation	-0.255	0.388	-0.268	-0.124	-0.038	-0.081

Table S.5: Responses to questions Q2-Q5 related to user expectation regarding the output of the recommendation tool. Answers to Q2 are significantly lower than to other questions. No significant difference between Q3,Q4,Q5.

	median response in Likert scale 1-7 [Q1,Q3]	
Q2. It is important for me to know the mathematics behind the model's recommendations	4 [2,5]	b
Q3. It is important for me to know the features of my patient contribute to the model's recommendation	6 [5,6.5]	a
Q4. It is important for me to know how the model makes its recommendation for my individual patient	6 [5,6]	a
Q5. It is important for me to know how uncertain (in %) the model is about its recommendation	6 [5.5, 7]	a
	p<0.001 Friedman test	p<0.001 posthoc Kruskal-Wallis

Supplementary Tables 23

Table S.6: Responses to questions A-G after deciding on patient cases when supported by recommendation only (CS) and after decisions supported by recommendation and its explanation (CS+Exp). No statistically significant change observed in pairwise comparison (Wilcoxon signed-rank test).

Index	Question	Median in Likert scale	Pairwise comparison		
		CORONET Score (CS)	CORONET Score + Explanation (CS+Exp)	p	significance*
A	I am satisfied with the output information that CORONET provides	5 [4.5,6]	5 [5,6]	0.976	ns
В	CORONET helps me in making safe clinical decisions on patient management	5 [3.5,5]	5 [4,5]	0.425	ns
C	When my initial decision was the same as CORONET had recommended, I felt reassured	5 [4,6]	5 [4,6]	0.746	ns
D	I understand when and why CORONET may provide the wrong recommendation in some cases	5 [3,6]	5 [4,5]	0.275	ns
E	CORONET helps in cases where I am less confident in the decision on how to proceed	4 [3,5]	5 [4,5.5]	0.056	ns
F	Even when my initial course of action was different to what CORONET recommended, I still had full confidence in my original decision	5 [5,6]	5 [5,6]	0.890	ns
G	I was surprised when CORONET recommended an action different to my own	4 [3,5]	4 [3,5]	0.821	ns

Table S.7: Statistically significant correlations between changes in responses between CS and CS+Exp scenarios. Only one after adjustment for multiple comparison (using Sidak correction).

Change in:	Change in:	r (spearman)	p	p-adjusted (Sidak)	significance
Reassurance	Understanding wrong recommendation	0.762	0.000	0.003	**
Satisfaction with the output	Help in making safe decisions	0.632	0.001	0.135	ns
Reassurance	Help when HCP is less confident	0.549	0.007	0.553	ns
Satisfaction with the output	Reassurance	0.513	0.012	0.771	ns
Help in making safe decisions	Help when HCP is less confident	0.418	0.047	0.997	ns




Figure S.1: Responses to questions A-G which were asked twice: after deciding on patients 1-5, where only the CORONET score (CS) was presented, and after deciding on the patients 6-10, when additional model's explanation was delivered (CS+Exp).




Figure S.2: Correlation indicating positive feedback to the model's recommendation, which is not supported by an explanation (CS). The lower the expertise, the more helpful the tool appeared to be, even when no explanation is provided. Answers on X axis are from CS scenario; points are scattered for visibility.




Figure S.3: Correlation indicating positive feedback to the model's recommendation, which is not supported by an explanation (CS). The higher the need for knowing the contributing features, the less helpful CS output is likely to be. Answers on X axis are from CS scenario; points are scattered for visibility.




Figure S.4: Correlation indicating positive feedback to the model's recommendation, which is not supported by an explanation (CS). Satisfaction with the CS output, and reassurance when the tool recommended the same action as their own were correlated with being convinced by the colour bar. Answers on X axis are from CS scenario; points are scattered for visibility.

# Agree or disagree with the recommendation?




Figure S.5: Time spent on individual decisions for cases 1-5 (CS) and 6-10 (CS+Exp) stratified by the concordance between the user's final decision and the model's recommendation.





Figure S.6: Total time spent on making decisions about patients: cases 1-5 where only the score with colour bar were provided; cases 6-10 enhanced with two additional figures with explanation. Each line is an individual clinician. Green - time increased, red - decreased. No significant difference (p=0.846, Wilcoxon sign rank test).











# COVID-19 Risk in ONcology Evaluation Tool

CORONET is an online tool to support decisions regarding hospital admissions or discharge in cancer patients presenting with symptoms of COVID-19 and the likely severity of illness. It is based on real world patient data. This is an early version that could possibly change. The model is currently undergoing peer review and we would be grateful for feedback as to utility of this tool (email the-christie.coronet@nhs.net or tweet @beckilee @CORONET). It should not be used for patient management currently (pending peer review). Detailed information as to how version 2 of the tool was created can be found in the pre-print on Preprints with THE LANCET. Version 1 is available at medRxiv. Version 2 uses data from many countries and new analysis has resulted in an update of the features important for the model.

What does CORONET do? Who is CORONET for? Limitations

CORONET asks for some details about the patient, their cancer and blood test results on presentation to hospital with symptoms of COVID-19. It then uses data about the admission, requirement for oxygen and survival of similar patients in the past to show likely outcome of the patient.




Figure S.7: User's interface of the CORONET available at https://coronet.manchester.ac.uk