

Timed Contact Algebras

Ivo Düntsch, Michael Winter

Department of Computer Science
Brock University
St. Catharines, ON, Canada
Email: {duentsch|mwinter}@brocku.ca

Abstract

Timed contact algebras constitute an approach to a temporal version of a region based theory of space. The general theory does not provide a notion of an underlying static world, i.e. it does not explicitly contain a set of non moving regions. Furthermore, the model of time does not have any structure, i.e. time is neither ordered nor required to be discrete or continuous. In this paper we want to investigate two extensions of the basic theory. The first extension considers grounded timed contact algebras that make the underlying static world explicit. In this context we introduce the Axiom of Construction that relates the existence of certain regions and the time structure for the first time. The second addition is given by a betweenness relation on the set of time. In this context we introduce the Axiom of Continuity (CONT), ensuring “smooth” movement of regions through time. Last but not least, we show that both axioms together do not allow finite models.

I. Introduction

Traditionally, space has been considered using point-based theories such as geometric (e.g. Euclidean geometry) or topological representations (point-set topology) of space. In recent years region-based theories have been of interest in AI and, in particular, Knowledge Representation (KR). These theories of qualitative spatial reasoning (QSR) are often based on mereotopology. This “pointless” approach considers regions as first order elements and introduces points as second order definable entities using sets of

regions, similar to the representation of Boolean algebras, where elements can be recovered as ultrafilters. The origins of mereotopology go back to the works of [9] on mereology and, on the other hand, the works of [6], [11], and [13] who used regions instead of points as the basic entity of geometry. The mereological structures of Leśniewski are, basically, complete Boolean algebras B without a smallest element. Whitehead’s addition consists of a “connection” (or “contact”) relation C among nonempty regions, which, in its simplest form is a reflexive and symmetric relation satisfying an additional extensionality axiom. The standard model of a mereotopological structures is the set of regular closed (or regular open) sets of topological spaces $\langle X, \tau \rangle$ with the *standard (Whiteheadian) contact* among regions defined by

$$uCv \iff u \cap v \neq \emptyset.$$

The primary example is the collection of all nonempty regular closed sets of the Euclidean plane.

The simplest algebraic theory of mereotopology is given by Boolean contact algebras, which are Boolean algebras extended with a contact relation C satisfying some axioms. They appear in different papers under various names and are closely related to the Region Connection Calculus RCC of [2], and, therefore, also the basis of relational approaches such as the relation algebra RCC8 and its extensions [4], [10]. For a recent overview of the topic we invite the reader to consult [12].

In [5] timed contact algebras were introduced. They add a notion of time to Boolean contact algebras, i.e. they constitute an adequate theory for regions moving in time. In this approach regions are modeled by a collection of snapshots in a static world, i.e. given a set of points in time T a moving region is just a function from T to the static world. In other words, a region is described by a collection of static regions at any point in time t - its (spatial) extent at

Both authors gratefully acknowledge support from the Natural Sciences and Engineering Research Council of Canada.

time t . In terms of [7] this theory falls into the category of snapshot spatial-temporal models. However, timed contact algebras are much weaker than any theory studied in [7]. In the general theory, the set of points in time does not have any structure and the underlying static world is hidden and cannot necessarily be identified. It is even possible that the static world differs at any point in time. We want to demonstrate this by an example.

Example 1. In general, a region at rest, i.e. a region that does not move (or change its shape), is modeled by a function that maps every point in time to the same static region. In this example we want to demonstrate the static world can be different at different points in time. In particular, there might be static regions available at some point in time that do not exist at others. Therefore, let $\mathbf{2} = \{0, 1\}$ and $\mathbf{4} = \{0, a, a^*, 1\}$ be the (finite) Boolean algebras with two and four elements, respectively. The contact relation on both algebras is given by overlap, i.e. regions a and b are in contact if they have a nonzero meet or, more formally, aCb if and only if $a \cdot b \neq 0$. As model of time we choose the discrete set $\{0, 1\}$. Now, consider all function $f : \{0, 1\} \rightarrow \mathbf{2} \cup \mathbf{4}$ with $f(0) \in \mathbf{2}$ and $f(1) \in \mathbf{4}$. In this case there is no non-moving region that has a as its extent at time 1.

In this paper we want to investigate grounded timed contact algebras that make the static world explicit. This theory will motivate a new axiom, the Axiom of Construction (CONS), relating the existence of regions and the time structure for the first time.

The second addition to the general theory of timed contact algebras in this paper is given by a betweenness relation on the set of time, i.e. we introduce some structure on time. We are using a betweenness relation instead of some order structure because of its generality. The structure induced by a betweenness relation includes all ordered time structures but is not restricted to those. However, this additional structure motivates a new axiom, the Axiom of Continuity (CONT), ensuring “smooth” movement of regions through time. To motivate this axiom we want to give an example of a region that does not move “smoothly” in the sense considered in this paper.

Example 2. We choose the Euclidean plane as static world and the set $\mathbb{R}^{\geq 0}$ of the non-negative real numbers as the model of time. We denote by $B(p, r)$ the regular closed disc with radius r and center at point p . In this model a moving region is a function from $\mathbb{R}^{\geq 0}$ to the regular closed sets of the plane. We consider two regions r_1 and r_2 . r_1 is the region so that $r_1(x) = B((0, 0), 1)$ is the closed disc with radius 1 at the origin for all $x \in \mathbb{R}^{\geq 0}$. This region does neither move nor change its shape, i.e. is a region at

rest. r_2 is defined by

$$r_2(x) = \begin{cases} B((1, 0), 1 - x), & 0 \leq x < 1; \\ B((3, 0), 1), & \text{otherwise.} \end{cases}$$

For any $0 \leq x < 1$ the regions r_1 and r_2 are in contact (they share the common point $(1, 0)$). At any other point in time they are not in contact. However, even though none of the regions disappears at any time they are never in external contact, i.e. sharing just a boundary point.

Finally, we want to relate both extension of the general theory. We will show that both axioms together (CONS and CONT) do not allow finite models. In particular, they imply that the set of points in time must be infinite.

II. Contact Algebras

For any set X and $Y \subseteq X$ we denote by $X - Y$ the complement of Y in X . If X is clear from the context we simply write $-Y$. For a binary relation R on D , and $x \in D$, we let $R(x) = \{y : xRy\}$, the *range of x with respect to R* .

If \sim is an equivalence relation on a set A , we will denote the equivalence class of an element $a \in A$ by $[a]$, i.e. $[a] = \{x \in A \mid a \sim x\}$.

A. Contact/Pre-Contact Relations

Throughout this paper, $(B, 0, 1, +, \cdot, *)$ is a Boolean algebra; if no confusion can arise we sometimes denote algebras by their base set. For details on Boolean algebras not explicitly mentioned in this paper we refer to [8].

In the remainder we will also use difference $a \setminus b := a \cdot b^*$ and the symmetric difference $a \triangle b := (a \setminus b) + (b \setminus a)$ of two elements a and b .

A binary relation C on a Boolean algebra B is called a *contact relation* (CR) if it satisfies:

- C0) $(\forall a) 0(-C)a$;
- C1) $(\forall a)[a \neq 0 \Rightarrow aCa]$;
- C2) $(\forall a)(\forall b)[aCb \Rightarrow bCa]$;
- C3) $(\forall a)(\forall b)(\forall c)[(aCb \text{ and } b \leq c) \Rightarrow aCc]$;
- C4) $(\forall a)(\forall b)(\forall c)[aC(b + c) \Rightarrow (aCb \text{ or } aCc)]$.

The pair $\langle B, C \rangle$ is called a *Boolean Contact Algebra* (BCA).

Axioms C0 and C1 imply that 0 is the only element with $C(0) = \emptyset$, i.e. that is not connected to any element. This axiom is motivated in the static world, i.e. in contact structures without any notion of time, by the fact that we

are not able to distinguish the empty region and a region that is not in contact to anything. In a timed environment this might not longer be true. A region in time might disappear, i.e. in our interpretation it is equal to the empty region, for a certain amount of time but still be in contact to other regions for the rest of the time. To model this we use relations that are weaker than contact relations: A relation C on a Boolean algebra B is called a *pre-contact relation* if it satisfies C0, C2–C4 and

$$P1) (\forall a)[(\exists b) aCb \Rightarrow aCa];$$

Any contact relation is a pre-contact relation. Conversely, the set $I := \{a \mid C(a) = \emptyset\}$ is a Boolean ideal for every pre-contact relation. We denote the congruence induced by I by \sim_C . The following lemma was shown in [5].

Lemma 1: Let C be a pre-contact relation on a Boolean algebra B . Then:

- 1) I is an ideal and the relation C_I on B/I defined by

$$[a]C_I[b] :\Longleftrightarrow aCb$$

is a contact relation.

- 2) $a \sim_C b$ if and only if $C(a \triangle b) = \emptyset$.

Notice that the range of C distributes over $+$ for any (pre-) contact relation, i.e.

$$\begin{aligned} x \in C(a + b) &\Longleftrightarrow (a + b)Cx \\ &\Longleftrightarrow aCx \text{ or } bCx && \text{by C2-4} \\ &\Longleftrightarrow x \in C(a) \cup C(b). \end{aligned}$$

The following property will be used later.

Lemma 2: Let C be a pre-contact relation on a Boolean algebra B . Then $a \sim_C b$ implies $C(a) = C(a \cdot b) = C(b)$.

Proof: Let $a \sim_C b$. Then $C(a \setminus b) \subseteq C(a \triangle b) = \emptyset$, and therefore,

$$\begin{aligned} C(a) &= C(a \cdot b + (a \setminus b)) \\ &= C(a \cdot b) \cup C(a \setminus b) \\ &= C(a \cdot b). \end{aligned}$$

The property $C(b) = C(a \cdot b)$ is shown analogously. ■

A (pre-)contact relation is called *extensional* if and only if it satisfies the following axiom:

$$C5) (\forall a)(\forall b)[C(a) = C(b) \Rightarrow a = b].$$

Notice that an extensional pre-contact relation is already a contact relation.

B. Timed Contact Algebras

A timed contact structure was first defined in [5], and is given by a set modeling time, a Boolean algebra of regions, and a family of pre-contact relations.

Definition 1: We call $\langle T, B, (C_t)_{t \in T} \rangle$ a timed contact algebra if T is a non-empty set, B Boolean algebra, and C_t is a pre-contact relation on B for every $t \in T$.

The relationship aC_tb denotes the fact that the regions a and b are in contact at time t .

Notice that the family of pre-contact relations can be seen as a ternary relation $C \subseteq T \times B \times B$ defined by $C = \bigcup_{t \in T} \{t\} \times C_t$. In the remainder of the paper we will adapt that view even though we will not change the notation.

It is easy to verify that the set of function B^T from an arbitrary set T to a Boolean (pre-)contact algebra B is a timed contact algebra. In general, we cannot guarantee that the contact structures at each point in time are the same (or are isomorphic). We will show below that every faithful timed contact algebra can be represented by a subdirect product.

For every $t \in T$ the pre-contact relation C_t induces a congruence relation on B which we denote by \sim_t . Furthermore, we denote the quotient algebra by B_t and the natural homomorphism by $\pi_t : B \rightarrow B_t$.

We call a timed contact algebra *faithful* if and only if it satisfies the following:

$$\text{FAT. } (\forall a)(\forall b)[(\forall t)(a \sim_t b) \Rightarrow a = b].$$

The next lemma provides an algebraic characterization of faithfulness:

Lemma 3: A timed contact algebra $\langle T, B, (C_t)_{t \in T} \rangle$ is faithful if and only if the mapping $\pi : B \rightarrow \prod_{t \in T} B_t$ defined by $\pi(b)_t = \pi_t(b)$, i.e. the t -component of $\pi(b)$ is $\pi_t(b)$, is a subdirect embedding.

Proof: It is well known that π is an embedding if and only if the family $(\pi_t)_{t \in T}$ separates points (see e.g. [1]), and the latter property is just FAT. Since each B_t is a quotient of B , π is a subdirect embedding. ■

As a straightforward generalization of C5 we call a timed contact algebra *extensional* if and only if it satisfies the following condition:

$$\text{EXT. } (\forall a)(\forall b)[(\forall t)(C_t(a) = C_t(b)) \Rightarrow a = b].$$

From Lemma 2 we immediately conclude that every extensional timed contact algebra is also faithful.

C. Topological models

First we recall some notions from topology. By a topological space (X, τ) we mean a set X provided with a family τ of subsets, called open sets, which contains the empty set and the whole set X , and is closed with respect to arbitrary unions and finite intersections. A subset $a \subseteq X$ is called *closed* if it is the complement of an open set.

In every topological space one can define the following operations on subsets $a \subseteq X$:

- 1) $Int(a) = \bigcup \{o \in \tau \mid o \subseteq a\}$ (the interior of a), i.e., the union of all open sets contained in a .
- 2) $Cl(a) = \bigcap \{c \text{ is closed} \mid a \subseteq c\}$ (the closure of a), i.e., the intersection of all closed sets containing a .

Cl and Int are interdefinable, i.e. $Cl(a) = -Int(-a)$ and $Int(a) = -Cl(-a)$.

A subset a of X is called *regular closed* if $Cl(Int(a)) = a$. We denote by $RC(X)$ the family of regular closed sets of X . It is a well known fact that $RC(X)$ is a Boolean algebra with respect to the following operations and constants:

$$0 = \emptyset, 1 = X, a + b = a \cup b \text{ and } a \cdot b = Cl(Int(a \cap b)).$$

$RC(X)$ naturally provides a contact relation C defined by aCb if and only if $a \cap b \neq \emptyset$. C is called *the standard* (or *Whiteheadian*) *contact relation* on $RC(X)$.

We regard a region in time as a function $f : T \rightarrow RC(X)$, i.e. the function providing the spatial extent of the region for every point in time.

We denote the set of all such function by X_T and define $C_t \subseteq X_T \times X_T$ by

$$fC_tg : \iff f(t) \cap g(t) \neq \emptyset.$$

Notice that for each $t \in T$ the relation C_t is defined by the standard contact relation on B , i.e. fC_tg if and only if $f(t)C^wg(t)$.

The following theorem was shown in [5].

Theorem 1: Let $\langle T, B, (C_t)_{t \in T} \rangle$ be a timed contact structure. Then there is a compact and semi-regular T_0 space $\langle X, \tau \rangle$ and an embedding $k : B \rightarrow X_T$ with aC_tb if and only if $k(a)(t) \cap k(b)(t) \neq \emptyset$.

III. Basis of Regions At Rest

In a concrete topological model we may call the constant functions *regions at rest* since they neither move nor change their shape. Since the elements in the abstract

theory need not to be functions we cannot immediately transfer this idea. Our approach will be based on a binary relation and maximal cliques thereof.

We say that a region a is *at rest with respect to a region* b (or *a and b are at rest* or *a is not moving with respect to b*), denoted by aRb , if and only if

$$(\exists t)(aC_tb) \Rightarrow (\forall t)(aC_tb).$$

Definition 2: Let $\langle T, B, (C_t)_{t \in T} \rangle$ be a timed contact structure. A subalgebra S of B is called *static* if and only if the elements of S are pairwise at rest, i.e. aRb for all $a, b \in S$.

Lemma 4: Each timed contact structure $\langle T, B, (C_t)_{t \in T} \rangle$ permits a non-empty and maximal static subalgebra.

Proof: First, it is easy to verify that the subalgebra $\{0, 1\}$ is indeed static because $0(-C_t)1$ for any $t \in T$. The existence of a maximal subalgebra now follows from Zorn's Lemma: In order to apply that lemma we show that $S = \bigcup_{i \in I} S_i$ of a chain $(S_i)_{i \in I}$ (with respect to inclusion) of static subalgebras is again a static subalgebra. It is easy to verify that S is a subalgebra of B . Now, assume $a, b \in S$ and that there is a $t \in T$ with aC_tb . Then there is a $j \in I$ with $a, b \in S_j$ since $(S_i)_{i \in I}$ is a chain. From the fact that S_j is a static subalgebra of B we conclude aC_tb for all $t \in T$. ■

Maximal static subalgebras are usually not unique. Furthermore, such a maximal static subalgebra might not be rich enough to characterize the extent of every region at any point in time. To illustrate this, consider Example 1 again. The set of constant functions, i.e. the functions $\bar{0}(x) = 0$ and $\bar{1}(x) = 1$, is the universe of a maximal static subalgebra, but none of those functions satisfies $f(1) = a$.

Definition 3: Let $\langle T, B, (C_t)_{t \in T} \rangle$ be a timed contact structure. A maximal static subalgebra S is called a *basis of regions at rest* (or simply *basis*) if and only if for all $b \in B$ and all $t \in T$ there is $a \in S$ with $b \sim_t a$. In this situation the structure $\langle T, B, (C_t)_{t \in T}, S \rangle$ is called a *grounded (timed) contact algebra*.

Since elements of a basis are pairwise at rest, the restriction of any of the relations C_t to S yields the same relation. We will denote this relation by C_S , i.e. for all $a, b \in S$ we have

$$aC_Sb \iff (\exists t \in T)aC_tb \iff (\forall t \in T)aC_tb.$$

Theorem 2: Let $\langle T, B, (C_t)_{t \in T}, S \rangle$ be a grounded contact algebra. Then $\langle S, C_S \rangle$ is

- 1) a BCA if B is faithful,
- 2) an extensional BCA if B is extensional.

Proof: 1) It is sufficient to show the ideal $I = \{a \in S \mid C_S(a) = \emptyset\}$ just contains 0. Assume there is $a \neq 0$

with $C_S(a) = \emptyset$. We want to show that $C_t(a) = \emptyset$ for all $t \in T$ since this implies $a \sim_t 0$ for all $t \in T$, and, hence, $a = 0$ using FAT, a contradiction. Therefore, assume that there is a $c \in B$ and a $t \in T$ with $a C_t c$. Then there is an element $b \in S$ with $b \sim_t c$ because S is a basis. We obtain $a \in C_t(c) = C_t(b)$ by Lemma 2, and, hence, $a C_S b$, a contradiction.

2) Since extensionality implies injectivity S is a BCA by 1. Now, assume $C_S(a) = C_S(b)$ for $a, b \in S$. We want to show that $C_t(a) = C_t(b)$ for all $t \in T$ since this implies $a = b$ by EXT. Therefore, assume $a C_t c$. Then there is a $d \in S$ with $c \sim_t d$ since S is a basis. We get $a \in C_t(c) = C_t(d)$, i.e. $a C_t d$. From $d \in S$ we conclude $a C_S d$, and, hence, $d \in C_S(a) = C_S(b)$. This immediately implies $b \in C_t(d) = C_t(c)$, i.e. $b C_t c$. ■

In the next theorem we want to establish that a basis really captures regions at rest, i.e. that the elements in S can be regarded as constant functions.

Theorem 3: Let $\langle T, B, (C_t)_{t \in T}, S \rangle$ be a faithful grounded contact algebra. Then B can be embedded into the algebra of functions S^T . Furthermore, the basis S corresponds to the set of constant functions in S^T .

Proof: By Theorem 2 we know that S together with C_S is a BCA. We want to show that $c \sim_t a$ and $c \sim_t b$ for a $t \in T$ with $a, b \in S$ implies $a = b$, i.e. that the mapping $h_t : B \rightarrow S$ defined by $h_t(c) = a$ if and only if $c \sim_t a$ and $a \in S$ is well defined. Since \sim_t is an equivalence relation we obtain $a \sim_t b$. From $a \sim_t b$ is equivalent to $C(a \Delta b) = \emptyset$ and $a, b \in S$ we get $a \sim_t b$ for all $t \in T$. Injectivity implies $a = b$. Now, define $h(c)(t) := h_t(c)$, and we have to show that $c C_t d$ if and only if $h_t(c) C_S h_t(d)$. This follows immediately from $C_t(h_t(c)) = C_t(c) \ni d$ and $c \in C_t(d) = C_t(h_t(d))$.

It remains to show that h maps the elements of S to constant functions. Since \sim_t is reflexive we have $a \sim_t a$, and, hence, $h_t(a) = a$ for all $t \in T$. Therefore, $h(a)(t) = a$ for all $a \in S$. ■

Notice that the previous theorem can also be stated in terms of a subdirect product. It shows that in the current case all components of the product B_t are isomorphic to S .

IV. Axiom of Construction

We are interested in ensuring the existence of a moving region if two snapshots are provided. This is of particular interest if we have a notion of static regions, i.e. a grounded contact structure. Assume one fixes two points in time, and for both points a static region. Is there a region that moves through those static regions at those points in time? The

following axiom of construction ensures the existence of such a region:

$$\text{CONS. } (\forall a)(\forall b)(\forall t_1)(\forall t_2)[t_1 \neq t_2 \Rightarrow (\exists c)(c \sim_{t_1} a \text{ and } c \sim_{t_2} b)].$$

Even though the Axiom of Construction is motivated by regions at rest, the axiom does not require the existence of a basis.

V. Betweenness Relation

Betweenness relations were studied in detail in [3]. A ternary relation Btw on a set U is called a betweenness relation if it satisfies

$$\begin{aligned} \text{BT0) } & (\forall a) \text{ Btw}(a, a, a); \\ \text{BT1) } & (\forall a)(\forall b)(\forall c)[\text{Btw}(a, b, c) \Rightarrow \text{Btw}(c, b, a)]; \\ \text{BT2) } & (\forall a)(\forall b)(\forall c)[\text{Btw}(a, b, c) \Rightarrow \text{Btw}(a, a, b)]; \\ \text{BT3) } & (\forall a)(\forall b)(\forall c)[(\text{Btw}(a, b, c) \text{ and } \text{Btw}(a, c, b)) \Rightarrow b = c]. \end{aligned}$$

Every partial order on a set induces a betweenness relation on that set but not necessarily vice versa. It was shown in [3] that a betweenness relation is generated by a partial order if and only if it satisfies the additional axioms:

$$\begin{aligned} \text{BT4) } & \text{There are no odd cycles in the graph of the relation } \text{Btw}(a, a, b); \\ \text{BT5) } & (\forall a)(\forall b)(\forall c)[(\text{Btw}(a, b, c) \text{ and } \text{Btw}(b, c, d) \text{ and } b \neq c) \Rightarrow \text{Btw}(a, b, d)]. \end{aligned}$$

Notice that BT4 can neither be expressed by a finite set of first-order axioms nor by using a finite number of variables. Axiom BT5 is also called “outer transitivity axiom”.

We call a betweenness relation Btw on U *dense* if and only if for all $u_1, u_2 \in U$ with $u_1 \neq u_2$ there is a $u \in U$ with $u \neq u_1, u \neq u_2$ and $\text{Btw}(u_1, u, u_2)$.

As already mentioned in the introduction we want to stay as general as possible. Therefore, we are going to use the more general concept of betweenness on the set of time instead of a partial order.

Definition 4: Let $\langle T, B, (C_t)_{t \in T} \rangle$ be a timed contact structure, and Btw be a betweenness relation on T . Then $\langle T, \text{Btw}, B, (C_t)_{t \in T} \rangle$ is called a BLT-algebra.

VI. Axiom of Continuity

In order to state the axiom we need two more predicates. The ternary relation NZ defined by

$$NZ(t_1, t_2, a) : \Longleftrightarrow (\forall t)(\text{Btw}(t_1, t, t_2) \Rightarrow a \not\sim_t 0)$$

indicates that the region a does not disappear at any time between t_1 and t_2 . Furthermore, we are interested in timed version of external contact. In the static world external contact $aECb$ can be defined by $aCt b$ and $a \cdot b = 0$. Using the relation \sim_t we define external contact at t by

$$aEC_t b : \Longleftrightarrow aC_t b \text{ and } a \cdot b \sim_t 0.$$

The Axiom of Continuity requires that two regions that

- 1) are in contact at t_1 ,
- 2) are not in contact at t_2 , and
- 3) do not disappear between t_1 and t_2

must be in external contact at some point in time between t_1 and t_2 . Formally, we define

$$\begin{aligned} \text{CONT. } (\forall a)(\forall b)(\forall t_1)(\forall t_2)(\exists t)[(NZ(t_1, t_2, a) \\ \text{and } NZ(t_1, t_2, b) \text{ and } aC_{t_1} b \text{ and } a(-C_{t_2})b) \\ \Rightarrow (\text{Btw}(t_1, t, t_2) \text{ and } a(EC_t b))]. \end{aligned}$$

Example 2 shows that not every BLT-algebra satisfies CONT. However, we obtain the following relationship between the temporal structure and the Axiom of Construction:

Lemma 5: Let $\langle T, \text{Btw}, B, (C_t)_{t \in T}, S \rangle$ be an extensional grounded BLT algebra satisfying CONT, CONS and $|S| > 2$. Then Btw is dense.

Proof: Since $|S| > 2$ there is an element $a \in S$ with $a \neq 0, 1$. By Theorem 2 $\langle S, C_S \rangle$ is an extensional BCA, i.e. there is $b \in S$ with $b \neq 0$ and $a(-C_S)b$.

From $a \neq 0$ we conclude that $C_S(a) \neq \emptyset$ using the extensionality of C_S . This is equivalent to $aC_t a$ for all $t \in T$. The property $bC_t b$ for all $t \in T$ follows analogously.

Now, assume $t_1, t_2 \in T$ with $t_1 \neq t_2$. CONS implies that there is a region $c \in B$ with $c \sim_{t_1} a$ and $c \sim_{t_2} b$. We distinguish two cases:

$-NZ(t_1, t_2, c)$: In this case there is a $t \in T$ with $\text{Btw}(t_1, t, t_2)$ and $c \sim_t 0$, i.e. $c(-C_t)c$. From $a \sim_{t_1} c$ and $aC_{t_1} a$ we conclude $cC_{t_1} a$. This implies $c \not\sim_{t_1} 0$, and, hence, $t \neq t_1$. The property $t \neq t_2$ is shown analogously using $c \sim_{t_2} b$ and $bC_{t_2} b$.

$NZ(t_1, t_2, c)$: Since $aC_t a$ for all $t \in T$ we get $NZ(t_1, t_2, a)$. Furthermore, we have $a \in C_{t_1}(a) = C_{t_1}(c)$, i.e. $aC_{t_1} c$, and $a \notin C_{t_2}(b) = C_{t_2}(c)$, i.e. $a(-C_{t_2})c$. From CONT we conclude that there is a $t \in T$ with $\text{Btw}(t_1, t, t_2)$ and $aEC_t c$. Since $a \in C_{t_1}(a) = C_{t_1}(a \cdot c)$ we get $a(-EC_{t_1})c$. This verifies $t \neq t_1$. Since $a(-C_{t_2})b$ and $c \sim_{t_2} b$ we have $a(-C_{t_2})c$. This shows $t \neq t_2$. ■

VII. Conclusion and Outlook

In this paper we started to investigate axioms that relate the existence of regions and the temporal structure of timed contact algebras. Our axiom of continuity serves as a first approximation of a 'smooth' movement of regions. Future work will concentrate on this aspect.

Furthermore, the spatial component of a timed contact algebra is based on a binary contact relation at each point in time. A temporal/modal logic based on a temporal and spatial modality for this kind of theory is of interest.

References

- [1] Burris, S. and Sankappanavar, H.P. (1981). A Course in Universal Algebra. *Springer Verlag*, New York, 1981.
- [2] Clarke, B.L. (1981) A calculus of individuals based on 'connection'. *Notre Dame J. Formal Logic* 22, 204-218.
- [3] Düntsch, I. and Urquhart, A. (2006). Betweenness and Comparability Obtained from Binary Relations. *Lecture Notes in Computer Science* vol. 4136 (2006), 148-161.
- [4] Düntsch, I., Schmidt, G. and Winter, M. (2001). A necessary Relation Algebra for Mereotopology. *Studia Logica* 69, 381-409.
- [5] Düntsch, I. and Winter, M. (2008). Moving Spaces. In: S. Demri, C.S. Jensen (Eds.), *TIME 2008 - 15th International Symposium on Temporal Representation and Reasoning*, IEEE Computer Society (2008), 59-63.
- [6] de Laguna, T. (1922). Point, line and surface as sets of solids. *The Journal of Philosophy*, 19:449-461.
- [7] Kontchakov, R., Kurucz, A., Wolter, F. and Zakharyashev M. (2007) Spatial Logic+Temporal Logic = ?. In Aiello, M., Pratt-Hartmann I. and van Benthem, J. *Handbook of Spatial Logics*, pp. 497-564 Springer
- [8] Koppelberg, S. (1989). *General Theory of Boolean Algebras*, volume 1 of *Handbook on Boolean Algebras*. North Holland.
- [9] Leśniewski, S. (1927 – 1931). O podstawach matematyki. 30–34.
- [10] Renz, J. and Nebel B. (2007) Qualitative Spatial Reasoning using Constraint Calculi. In Aiello, M., Pratt-Hartmann I. and van Benthem, J. *Handbook of Spatial Logics*, pp. 161-215 Springer
- [11] Tarski, A. (1956). Foundation of the geometry of solids. In Woodger, J. H., editor, *Logic, Semantics, Metamathematics*, pages 24–29. Clarendon Press, Oxford. Translation of the summary of an address given by A. Tarski to the First Polish Mathematical Congress, Lwów, 1927.
- [12] Vakarelov, D. (2007) Region-Based Theory of space: Algebras of Regions, Representation Theory, and Logics. In: Dov Gabbay et al. (Eds.) *Mathematical Problems from Applied Logic II. Logics for the XXIst Century*, pages 267-348. Springer.
- [13] Whitehead, A. N. (1929). *Process and reality*. MacMillan, New York.