

NB: The graded, first version of the report must be returned if you hand in a second time!

H1b: MD simulation – dynamic properties

Andréas Sundström and Linnea Hesslow

November 29, 2018

Task N ^o	Points	Avail. points
Σ		

Introduction

The velocity Verlet algorithm is a semi-implicit and efficient method to simulate an ensemble of particles whose trajectories are governed by Newton's equation of motion. Accordingly, it is a suitable algorithm to study molecular dynamics and to determine statistical properties of a system. Here, we use the velocity Verlet algorithm to study a system of aluminum atoms in a face centered cubic (FCC) lattice. By scaling the positions, momenta and lattice parameter, we can equilibrate the temperature and pressure to prescribed values. We study the aluminum system at 500 °C and 700 °C, which correspond to the solid and liquid state respectively, and compute two dynamic quantities: mean squared displacements and the velocity correlation function.

For this report, we used scripts provided by Anders Lindman for initializing FCC lattices and calculating lattice energies, lattice forces and virials of an aluminium lattice. All simulations were done in a simulation box containing $4^3 = 64$ unit FCC cells, and a total of $4 \cdot 64$ aluminum atoms placed at the corresponding FCC lattice points. The simulation box had periodic boundary conditions.

The velocity Verlet algorithm

The main idea behind the velocity Verlet algorithm is to split up the time steps of the velocity, in order to make the update process of the state more symmetric. The position, velocity and acceleration, x_i , v_i and a_i ¹ respectively, are updated according to

$$\begin{aligned}v_{i+\frac{1}{2}} &= v_i + \frac{1}{2}a_i dt, \\x_{i+1} &= x_i + v_{i+\frac{1}{2}} dt, \\a_{i+1} &= \text{get_acceleration}(x_{i+1}), \\v_{i+1} &= v_{i+\frac{1}{2}} + \frac{1}{2}a_{i+1} dt.\end{aligned}\tag{1}$$

By effectively using an average of the old and new acceleration, $(a_{i+1} + a_i)/2$, for updating the velocity, $v_i \rightarrow v_{i+1}$, the velocity Verlet algorithm becomes semi-implicit; this also results in better energy-conservation properties of the algorithm, compared to, e.g., a fully explicit algorithm ($v_{i+1} = v_i + a_i dt$). However, in contrast to a fully implicit algorithm, there is no need for a computationally costly matrix inversion for each time step, and the velocity Verlet algorithm is also self-starting on an initial condition of $x_{i=0} = x_0$, $v_{i=0} = v_0$, and $a_{i=0} = \text{get_acceleration}(x_0)$.

Task 1: potential energy

The theoretical, minimum energy lattice parameter for aluminum can be determined by calculating the minimum potential energy per unit cell in a lattice with zero initial momenta for all particles.

Figure 1 shows the potential energy as a function of the lattice parameter. We used a quadratic fit to find the minimum energy², and obtained $V_{\text{eq}} \approx 65.38 \text{ \AA}^3$. This corresponds to the equilibrium lattice parameter $a_{\text{eq}} \approx 4.029 \text{ \AA}$ at 0 K, which we took as the initial lattice parameter for the following tasks. We find that figure 1 looks similar to the figure 1 in the homework problem file, which is encouraging.

Task 2: determine the time step

In this task, we use a lattice with the equilibrium lattice constant $a_{\text{eq}} \approx 4.029 \text{ \AA}$, found in the previous task, but then we added a random perturbation, uniformly distributed in the interval $\pm 0.065a_{\text{eq}}$, to each atom position. This creates a non-equilibrium system, which has a non-trivial time evolution. To determine the time evolution, we used the velocity Verlet algorithm, as described in the introduction.

¹In most situations the acceleration need not be saved for each time step, which might be insinuated by the index on a_i . The index is just used for notational convenience.

²We performed the quadratic fit in the volume V , which to a small error corresponds to a quadratic fit in the lattice parameter a , since $E \approx \alpha(V - V_0)^2 \approx \alpha a_0^4(a - a_0)^2$ in a close vicinity of the minimum $a \approx a_0$.




Figure 1: The potential energy per unit cell for aluminum as a function of the lattice parameter cubed.

The first step when doing simulations of this kind is to determine a suitable timestep. Even though the velocity Verlet algorithm have good energy conservation properties, it only gives an approximation to the “true” continuous solution; an approximation which gets better the smaller dt we choose. However, choosing dt too small will result in unnecessary computational costs for the same total simulation time. We are therefore interested in finding the largest dt we can get away with without losing energy conservation. From figure 2, we see that $dt = 2 \cdot 10^{-2}$ ps clearly does not conserve energy, while $dt = 1 \cdot 10^{-1}$ ps dose conserve energy. To be on the safe side, we chose $dt = 5 \cdot 10^{-3}$ ps = 5 fs as our time step. This is in line with the lecture notes, where it is stated that a suitable time step would normally be a few femtoseconds, or somewhat larger for heavy atoms.

The total energy of the simulated system at each time step is determined by sum of the kinetic energy of each particle, $E_{\text{kin}}^{(\text{atom})} = m_{\text{Al}} v^2 / 2$, and the total lattice energy of the system. Then, to calculate the temperature, we can use the *equipartition theorem* stating that $\langle E_{\text{kin}}^{(\text{atom})} \rangle = 3k_B T / 2$, or equivalently that $T = 2 \langle E_{\text{kin}}^{(\text{atom})} \rangle / (3k_B)$. We can therefore define an instantaneous temperature

$$\mathcal{T}(t) = \sum_{\text{all atoms}} 2E_{\text{kin}}^{(\text{atom})}(t) / (3N_{\text{atoms}}k_B). \quad (2)$$




Figure 2: The temperature and kinetic energy per unit cell as a function of time for four different time steps.

With the random noise, the temperature and the energy differ between runs, but are in the same order of magnitude. We note that the temperature in several cases is higher than desired value of 600-800 K from the problem sheet. The temperatures and energies up to one standard deviation are quantified in table 1.

Table 1: Energies and temperatures with one standard deviation uncertainties for four different values of the time steps.

dt [ps]	T [K]	E_{tot} [eV/unit cell]
$2 \cdot 10^{-2}$	unstable	unstable
$1 \cdot 10^{-2}$	$783 \pm 3.5\%$	$-12.61 \pm 9.2 \cdot 10^{-3}\%$
$5 \cdot 10^{-3}$	$793 \pm 3.7\%$	$-12.62 \pm 2.2 \cdot 10^{-3}\%$
$2 \cdot 10^{-3}$	$769 \pm 3.7\%$	$-12.65 \pm 5.2 \cdot 10^{-4}\%$

Tasks 3 and 4: temperature and pressure equilibration

There is no simple method to initialize a system to a specified temperature and pressure, but prescribed values can be obtained by scaling the velocities and the positions of the particles in the system. Given that the temperature of the system is given by the average kinetic energy of the atoms, we can change the temperature by scaling the velocities of all atoms. A scheme for this temperature scaling is to first calculate the instantaneous temperature \mathcal{T} according to equation (2), at that time step, and then scale it by a scaling factor

$$\alpha_T = 1 - \frac{dt}{\tau_T} \frac{\mathcal{T} - T_{\text{eq}}}{\mathcal{T}}, \quad (3)$$

where T_{eq} is the desired equilibrium temperature, and τ_T turns out to be a typical time scale on which the system equilibrates. However, as we only have direct control of the particle velocities, we equilibrate the temperature by scaling the velocities by $v_i \rightarrow \sqrt{\alpha_T} v_i$, since the temperature depends quadratically on the velocities.

A similar scheme for pressure equilibration is to instead scale the particle positions and simulation-box volume, using a scaling factor of

$$\alpha_P = 1 - \kappa \frac{dt}{\tau_P} (\mathcal{P} - P_{\text{eq}}), \quad (4)$$

where \mathcal{P} is the instantaneous pressure, P_{eq} is the desired equilibrium pressure, τ_P is the characteristic equilibration time, and κ is the isothermal compressibility of the material simulated³. This time the positions are scaled according to $x_i \rightarrow \alpha_P^{1/3} x_i$, and similarly the simulation box volume is scaled by scaling its side lengths by $L \rightarrow \alpha_P^{1/3} L$.

We set $\tau_P = \tau_T = 100dt$, where $dt = 5 \cdot 10^{-3}$ ps, and equilibrated the temperature and pressure by scaling the particle momenta and positions (and box size) respectively. Choosing a slower equilibration time did not affect the results qualitatively. Both temperature and pressure were equilibrated in the same Verlet loop, but for the higher temperature the system was first melted by increasing the temperature to 1100 °C. To determine the isothermal compressibility κ , the values of Young's modulus Y and shear modulus G were taken from Physics Handbook, table T 1.1. From F 1.15 in Physics Handbook, the bulk modulus can then be calculated as

$$B = \frac{YG}{9G - 3Y} \quad \kappa_{\text{AI}} = \frac{1}{B} \approx (6.6444 \cdot 10^5 \text{ bar})^{-1}, \quad (5)$$

where 1 bar = $6.2415 \cdot 10^{-7} \text{ eV/\AA}^3$ in atomic units.

The results are shown in figure 3, where we overlay the instantaneous values of \mathcal{T} and \mathcal{P} with a moving average using 100 time steps. The desired temperatures and pressures were approximately obtained in the equilibration process. Although the average pressure was slightly below zero, this is within the fluctuation error bars and is in line with the figure 2 in the homework problem document.

Due to the large fluctuations, it is difficult to determine if the system has reached its equilibrium directly from the instantaneous pressure. Instead, we consider the time-evolution of the lattice parameter a_0 , as shown in figure 4. After $t = 15\tau_{\text{eq}}$, the lattice parameter has reached a constant which means that the equilibrium is reached. The equilibrium values of the lattice parameter were found to be

$$a_0 \approx 4.09 \text{ \AA}, \quad T = 500 \text{ °C}, \quad (6)$$

$$a_0 \approx 4.26 \text{ \AA}, \quad T = 700 \text{ °C}. \quad (7)$$

³The isothermal compressibility is defined according to $\kappa = -(V \partial P / \partial V)_T^{-1}$. This compressibility should therefore be used when scaling the volume of the box to change the pressure.




Figure 3: The instantaneous values of \mathcal{T} and \mathcal{P} overlaid with with a moving average using 100 time steps, which corresponds to $\Delta t = \tau_P/2$. Left panel: $T = 500^\circ\text{C}$, right panel: $T = 700^\circ\text{C}$.

These value are larger than the zero-temperature constants, and it is reasonable that the higher 700°C case corresponds to a larger lattice parameter at constant pressure.




Figure 4: The time evolution of the lattice parameter a_0 as the system equilibrates.

Tasks 3-5: particle trajectories

Starting with the temperature- and pressure-equilibrated systems from the previous section, we study the particle trajectories for both systems. Here, we decrease the time step to $dt = 1 \cdot 10^{-3}$ ps and the simulation length to $t_{\text{end}} = 10$ ps to get better statistics. This was mostly motivated by increasing the resolution in tasks 6-7.

First, we note that the cumulative averages of the instantaneous temperatures and pressures stayed close to their initial values. This is shown in figure 5. Numerically, we obtained

$$T = 506 \pm 28^\circ\text{C}, \quad P = 403 \pm 1244 \text{ bar}, \quad (8)$$

and

$$T = 699 \pm 36^\circ\text{C}, \quad P = -640 \pm 1860 \text{ bar}. \quad (9)$$

From equation (82) in MD lecture notes, the mean squared displacement can be




Figure 5: The instantaneous values, and the cumulative averages, of the temperature and the pressure in the production runs. Left panel: $T = 500\text{ }^{\circ}\text{C}$, right panel: $T = 700\text{ }^{\circ}\text{C}$

calculated as

$$\Delta_{\text{MSD}}(t) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt' \frac{1}{N_{\text{atoms}}} \sum_{i=0}^{N_{\text{atoms}}-1} [\mathbf{r}_i(t+t') - \mathbf{r}_i(t')]^2 \quad (10)$$

\Rightarrow

$$\Delta_{\text{MSD}}(t_k) \approx \frac{1}{N_T - k} \frac{1}{N_{\text{atoms}}} \sum_{j=0}^{N_T-k-1} \sum_{i=0}^{N_{\text{atoms}}-1} [\mathbf{r}_i(t_{k+j}) - \mathbf{r}_i(t_j)]^2 \quad (11)$$

COMMENT ON THE LARGE FLUCTUATIONS IN P!!!!!!!!!!!!!!!!!!!!




Figure 6: Five individual particle trajectories (gray thin lines), overlaid with the mean squared displacement $\Delta_{\text{MSD}} \approx 0.16\text{ }^{\circ}\text{Å}^2$ (thick black line). In the top panel, $T = 500\text{ }^{\circ}\text{C}$, the system is in a solid state. In the bottom panel, $T = 700\text{ }^{\circ}\text{C}$, the system is in a liquid state, where $\Delta_{\text{MSD}} \approx 6tD_s$ (dotted red line).

We now consider the particle trajectories. Figure 6 shows the trajectories of five individual particles along with the mean squared displacement as determined in equation (11). We can clearly see that the particle trajectories are bounded in the left figure 6,

while for the high-temperature case in the right panel, they increase as square root of time ($\Delta_{\text{MSD}} \propto t$). Consequently, the former is in a solid state while the latter is in a liquid state. In the solid, the obtained mean squared displacement was $\Delta_{\text{MSD}} \approx 0.16 \text{ \AA}^2$. We determined the self-diffusion coefficient as average slope of the mean squared displacement for $1 \text{ ps} \leq t \leq 9 \text{ ps}$, yielding $D_s \approx 0.42 \text{ \AA}^2/\text{ps}$.

Tasks 6-7: velocity correlation and power spectrum

We calculate the discrete auto-correlation function similarly to the MSD,

$$\Phi_j = \frac{1}{N-j} \sum_{i=0}^{N-j-1} \langle v_{i+j} v_i \rangle, \quad (12)$$

where $j = 0, 1, \dots, N-1$ and the average is taken over all atoms. Figure 7(left) shows that it is noticeably different between the solid and the liquid states: while the solid state remains non-zero at longer time, presumably because of oscillations around lattice points, the liquid velocity correlation quickly decays to zero after some initial oscillations.




Figure 7: Top panel: The velocity correlation function, and (bottom panel) its spectrum, calculated both directly from the velocity correlation (solid line) and from the power spectrum of the particle velocity (dotted line). Blue lines show $T = 500 \text{ °C}$ and red lines $T = 700 \text{ °C}$.

We now proceed to numerically approximate the integral

$$\hat{\Phi}(f) = 2 \int_0^\infty dt \Phi(t) \cos(2\pi f t) \approx 2 \int_0^{T_s} dt \Phi(t) \cos(2\pi f t) \quad (13)$$

using a trapezoidal method in MATLAB, with a frequency range $f = 0$ to $f = 1/(2\Delta t) = f_{\text{Nyquist}}$, and frequency steps $\Delta f = 1/T_s$, where T_s is a time at about half the simulation end time. This is to avoid including noisy data in $\Phi(t)$ at later times, where the statistics are poor.

We may then calculate the power spectrum according to

$$\begin{aligned}
 \hat{P}(\omega) &= \lim_{T \rightarrow \infty} \frac{1}{T} \left\langle \left| \int_0^T dt v(t) e^{i\omega t} \right|^2 \right\rangle \\
 &\approx \frac{1}{T} \left\langle \left| \int_0^T dt v(t) e^{i\omega t} \right|^2 \right\rangle \\
 \Rightarrow \hat{P}_k &= \frac{1}{T} \left\langle \left| \frac{T}{N} \sum_{i=0}^{N-1} v_i \exp\left(i2\pi \frac{ik}{N}\right) \right|^2 \right\rangle = \frac{T}{N} \langle |\hat{v}_k|^2 \rangle
 \end{aligned} \tag{14}$$

where the averages is taken over all atoms, and

$$\hat{v}_k = \sqrt{N} \sum_{i=0}^{N-1} v_i \exp\left(i2\pi \frac{ik}{N}\right) \tag{15}$$

is the discrete Fourier transform of v_i . Note that the factor of T was not included in the C scripts, and we therefore multiplied by this factor in the MATLAB plotting scripts. When we compare $\hat{\Phi}_k$ and \hat{P}_k in figure 7 (right), we find that they are very similar, as, indeed, they should be according to the Wiener-Khinchine theorem. If we instead take $T_s = T$ (using the full time evolution of $\Phi(t)$, we get a more noisy signal. This is because there is less statistics at high values of j in equation (12). Therefore, even though the results are similar, the power spectrum method in equation (14), should be considered more accurate since it includes more statistics of the data points.

The self-diffusion coefficient as determined by the power spectral density at $f = 0$, was found to be $D_s = 0.44 \text{ \AA}^2/\text{ps}$ and $D_s = 0.42 \text{ \AA}^2/\text{ps}$ determined from $\hat{\Phi}$, and \hat{P} respectively. (Since the data was noisy, we took the mean of the three first data points to obtain these values). As expected, these values are close to the values obtained from the mean squared displacement.

Concluding discussion

Using the velocity Verlet algorithm, we study a system of aluminum atoms at 500° C and 700° C , which correspond to the solid and liquid state respectively.

From both the mean squared displacements and the velocity correlation function, the solid state is clearly distinguishable from the liquid state. The mean squared displacement reaches a constant value in the solid state, whereas it grows linearly with time in the liquid state, which is characteristic of diffusion in a random walk process. Similarly, the spectrum of the velocity correlation function vanishes at zero frequency which means that the average velocity correlation is zero and hence there is no net movement of the particles; in contrast for the liquid state, the zero-frequency value of the spectrum is finite and proportional to the diffusion coefficient.

A Source Code

A.1 Main program task 1: main.T1.c

```
1  /*
2  main_T1.c Task 1 H1b
3  In this task, we scan over a range of lattice parameters, a0, to determine
4  which one results in the lowest potential energy stored in the lattice.
5
6  System of units:
7  Energy   - eV
8  Time     - ps
9  Length   - Angstrom
10 Temp     - K
11 Mass     - eV (ps)^2 A^(-2)
12 Pressure - eV A^(-3)
13 */
14 #include <stdio.h>
15 #include <math.h>
16 #include <stdlib.h>
17
18 #include "initfcc.h"
19 #include "alpotential.h"
20
21 #define N_cells 4
22 #define N_lattice_params 25
23
24 /* Main program */
25 int main()
26 {
27     int N_atoms = 4*N_cells*N_cells*N_cells;
28     double a0;
29     double a0_min = 4.0;
30     double a0_max = 4.2;
31     double da0 = (a0_max - a0_min)/N_lattice_params;
32
33     double (*pos)[3] = malloc(sizeof(double[N_atoms][3]));
34     double *energy = malloc(sizeof(double[N_lattice_params]));
35
36     FILE *file_pointer;
37
38     for (int i=0; i<N_lattice_params; i++){
39         a0 = a0_min + i*da0; // The lattice constant of this iteration
40         init_fcc(pos, N_cells, a0); // Init, FCC cells with lattice constant `a0`
41         // energy per unit cell
42         energy[i] = get_energy_AL(pos, N_cells*a0, N_atoms )*4/N_atoms;
43     }
44
45     // Write to files
46     file_pointer = fopen("../data/lattice_energies.tsv", "w");
47     for (int i=0; i<N_lattice_params; i++){
48         a0 = a0_min + i*da0;
49         fprintf(file_pointer, "%.8f \t %.8f \n", a0, energy[i]);
50     }
51     fclose(file_pointer);
52
53     free(pos); pos = NULL;
54     free(energy); energy = NULL;
55     return 0;
56 }
```

A.2 Main program Task 2: main.T2.c

```
1  /*
2  main_T2.c, Task 2, H1b
3  In this task, we add random noise to the particle positions and see how the
4  system evolves in time. Using the kinetic energy of the particles, we can
5  derive an instantaneous temperature of the system.
6
7  System of units:
8  Energy   - eV
9  Time     - ps
10 Length   - Angstrom
11 Temp     - K
12 Mass     - eV (ps)^2 A^(-2)
13 Pressure - eV A^(-3)
14 */
15
16 #include <stdio.h>
17 #include <math.h>
18 #include <stdlib.h>
19 #include <time.h>
20
```

```

21 #include "initfcc.h"
22 #include "alpotential.h"
23 #include "funcs.h"
24
25 #define N_cells 4
26 #define AMU 1.0364e-4
27 #define kB 8.6173303e-5
28
29 /* Main program */
30 int main()
31 {
32     int N_atoms = 4*N_cells*N_cells*N_cells;
33     double m_Al = 27*AMU;
34
35     double a_eq = 4.03; // Min potential energy lattice constant
36
37     double noise_amplitude = 6.5e-2 * a_eq;
38     double t_max=10; //
39     double dt = 2e-3;
40     int N_timesteps = t_max/dt;
41     double t, E_kin;
42
43     double (*pos)[3] = malloc(sizeof(double[N_atoms][3]));
44     double (*momentum)[3] = malloc(sizeof(double[N_atoms][3]));
45     double (*forces)[3] = malloc(sizeof(double[N_atoms][3]));
46     double *temperature = malloc(sizeof(double[N_timesteps]));
47     double *E_tot = malloc(sizeof(double[N_timesteps]));
48
49     FILE *file_pointer;
50
51     init_fcc(pos, N_cells, a_eq); // initialize fcc lattice
52     add_noise( N_atoms, 3, pos, noise_amplitude ); // adds random noise to pos
53     set_zero( N_atoms, 3, momentum); // set momentum to 0
54     get_forces_AL( forces, pos, a_eq*N_cells, N_atoms); //initial cond forces
55
56     for (int i=0; i<N_timesteps; i++){
57         /*
58          * The loop over the timesteps first takes a timestep according to the
59          * Verlet algorithm, then calculates the energies and temeperature.
60          */
61         timestep_Verlet (N_atoms, pos, momentum, forces, m_Al, dt, a_eq*N_cells);
62
63         E_kin = get_kin_energy(N_atoms, momentum, m_Al );
64         E_tot[i] = (E_kin + get_energy_AL(pos, a_eq*N_cells, N_atoms))*4/N_atoms;
65
66         /*  $3N \cdot k_B \cdot T/2 = 1/(2m) \cdot \sum_{i=1}^N p_i^2 = p_{sq}/(2m)$  */
67         temperature[i] = E_kin * 2/(3*N_atoms*kB);
68     }
69
70     /* Write tempertaure to file */
71     char file_name[100];
72     sprintf(file_name, "../data/temperature_dt-%0.0e_Task2.tsv", dt);
73     file_pointer = fopen(file_name, "w");
74     for (int i=0; i<N_timesteps; i++){
75         t = i*dt; // time at step i
76         fprintf(file_pointer, "%.4f \t %.8f \n", t, temperature[i]);
77     }
78     fclose(file_pointer);
79
80     /* Write total energy to file */
81     sprintf(file_name, "../data/total_energy_dt-%0.0e_Task2.tsv", dt);
82     file_pointer = fopen(file_name, "w");
83     for (int i=0; i<N_timesteps; i++){
84         t = i*dt; // time at step i
85         fprintf(file_pointer, "%.4f \t %.8f \n", t, E_tot[i]);
86     }
87     fclose(file_pointer);
88
89     free(pos); pos = NULL;
90     free(momentum); momentum = NULL;
91     free(forces); forces = NULL;
92     free(temperature); temperature = NULL;
93     free(E_tot); E_tot = NULL;
94     return 0;
95 }

```

A.3 Temperature and pressure equilibration for tasks 3-7: main_T3.c

```

1 /*
2  * main_T3.c, Tasks 3 and 4, Hib. Also used as input in Tasks 5-7.
3  * In this task, we use an equilibration scheme, based on scaling particle momenta
4  * and positions, to equilibrate the temperature and pressure in the system. We do
5  * this for T=500 degC and T=700 degC and P=1 bar. The difference between the two
6  * temperatures are that the higher temperature results in a melted system. (To

```

```

7 ensure that the system is melted properly, we first raise the temperature to
8 900 degC and then lower it back to 700 degC.)
9
10 After the system has equilibrated, we save the full phase space (all particle
11 positions and momenta) as well as the equilibrated lattice parameter to a
12 binary file which then can be read in for a production run.
13
14 System of units:
15 Energy - eV
16 Time - ps
17 Length - Angstrom
18 Temp - K
19 Mass - eV (ps)^2 A^(-2)
20 Pressure - eV A^(-3)
21 */
22
23 #include <stdio.h>
24 #include <math.h>
25 #include <stdlib.h>
26 #include <time.h>
27
28 #include "initfcc.h"
29 #include "alpotential.h"
30 #include "funcs.h"
31
32 #define N_cells 4
33 /* define constants in atomic units: eV, , ps, K */
34 #define AMU 1.0364e-4
35 #define degC_to_K 273.15
36 #define bar 6.2415e-07
37 #define kB 8.61733e-5
38
39 /* Main program */
40 int main()
41 {
42     char file_name[100];
43
44     int N_atoms = 4*N_cells*N_cells*N_cells;
45     double m_Al = 27*AMU;
46     /*
47      Values of Young's and shear modulus, Y and G resp., taken from
48      Physics Handbook, table T 1.1. Bulk modulus then calculated as
49      B = Y*G / (9*G - 3*Y) [F 1.15, Physics Handbook]
50      kappa = 1/B
51     */
52     double kappa_Al = 1/(6.6444e+05 * bar);
53     double a_eq = 4.03;
54     double cell_length = a_eq*N_cells;
55     double inv_volume = pow(cell_length, -3);
56     double noise_amplitude = 6.5e-2 * a_eq;
57
58     double T_final_C = 500;
59     int nRuns = 1; //2 if melt, 1 otherwise
60     double T_melt_C = 1100;
61
62     double P_final_bar = 1;
63
64     double T_eq;
65     double P_eq = P_final_bar*bar;
66     double dt = 5e-3;
67     double tau_T = 100*dt;
68     double tau_P = 100*dt;
69     double t_eq = 15*tau_P; //equilibration times
70     int N_timesteps = t_eq/dt;
71
72     double alpha_T, alpha_P, alpha_P_cube_root;
73     double t, E_kin, virial;
74
75     double (*pos)[3] = malloc(sizeof(double[N_atoms][3]));
76     double (*momentum)[3] = malloc(sizeof(double[N_atoms][3]));
77     double (*forces)[3] = malloc(sizeof(double[N_atoms][3]));
78     double *temperature = malloc(sizeof(double[N_timesteps]));
79     double *pressure = malloc(sizeof(double[N_timesteps]));
80     double *a0 = malloc(sizeof(double[N_timesteps]));
81
82     FILE *file_pointer;
83
84
85     init_fcc(pos, N_cells, a_eq); // initialize fcc lattice
86     add_noise( N_atoms, 3, pos, noise_amplitude ); // adds random noise to pos
87     set_zero( N_atoms, 3, momentum); // set momentum to 0
88     get_forces_AL( forces, pos, cell_length, N_atoms); //initial cond forces
89
90     for (int irun=0; irun < nRuns; irun++){// last run: final, irun = 0
91         if (irun == nRuns - 1){ // final run
92             T_eq = T_final_C + degC_to_K;
93         }else{
94             T_eq = T_melt_C + degC_to_K;
95         }
96         for (int i=0; i<N_timesteps; i++){
97             /*

```

```

98     The loop over the timesteps first takes a timestep according to the
99     Verlet algorithm, then calculates the energies and temeperature.
100    */
101    timestep_Verlet(N_atoms, pos, momentum, forces, m_Al, dt, cell_length);
102
103    E_kin = get_kin_energy(N_atoms, momentum, m_Al );
104    virial = get_virial_AL(pos, cell_length, N_atoms);
105
106    /* 3N*kB*T/2 = 1/(2m) * \sum_{i=1}^N p_i^2 = p_sq/(2m) */
107    temperature[i] = E_kin * 1/(1.5*N_atoms*kB);
108    /* PV = NkT + virial */
109    pressure[i] = inv_volume * (E_kin/1.5 + virial);
110
111    /* Equilibrate temperature by scaling momentum by a factor sqrt(alpha_T).
112       N.B. It is equally valid to scale the momentum instead of the velocity←
113       since they only differ by a constant factor m.
114    */
115    alpha_T = 1 + 2*dt*(T_eq - temperature[i]) / (tau_T * temperature[i]);
116    scale_mat(N_atoms, 3, momentum, sqrt(alpha_T));
117
118    // Equilibrate pressure by scaling the posistions by a factor of
119    // alpha_P^(1/3)
120    alpha_P = 1 - kappa_Al* dt*(P_eq - pressure[i])/tau_P;
121    alpha_P_cube_root = pow(alpha_P, 1.0/3.0);
122    scale_mat(N_atoms, 3, pos, alpha_P_cube_root);
123
124    cell_length*=alpha_P_cube_root;
125    inv_volume*=1/alpha_P;
126
127    //temperature[i]*=alpha_T;
128    //pressure[i]*=alpha_P;
129    a0[i] = cell_length/N_cells;
130  }
131 }
132
133 printf("equilibrium a0 = %.4f A\n", cell_length/N_cells);
134
135 /* Write tempertaure, pressure and cell size to file */
136 sprintf(file_name, "../data/temp-%d_pres-%d_Task3.tsv",
137         (int) T_final_C, (int) P_final_bar);
138 file_pointer = fopen(file_name, "w");
139 for (int i=0; i<N_timesteps; i++){
140     t = i*dt; // time at step i
141     fprintf(file_pointer, "%.4f \t %.8f \t %.8f \t %.8f \n",
142             t, temperature[i], pressure[i], a0[i]);
143 }
144 fclose(file_pointer);
145
146 /* Write phase space coordinates to file */
147 sprintf(file_name, "../data/phase-space-temp-%d_pres-%d.tsv",
148         (int) T_final_C, (int) P_final_bar);
149 file_pointer = fopen(file_name, "w");
150 for (int i=0; i<N_atoms; i++){
151     for (int j=0; j<3; j++){
152         fprintf(file_pointer, " %.16e \t", pos[i][j]);
153     }
154     for (int j=0; j<3; j++){
155         fprintf(file_pointer, " %.16e \t", momentum[i][j]);
156     }
157     fprintf(file_pointer, "\n");
158 }
159 fclose(file_pointer);
160
161 /* save equilibrated position and momentum as a binary file */
162 sprintf(file_name, "../data/INIDATA-temp-%d_pres-%d.bin",
163         (int) T_final_C, (int) P_final_bar);
164 file_pointer = fopen(file_name, "wb");
165 fwrite(pos, sizeof(double), 3*N_atoms, file_pointer);
166 fwrite(momentum, sizeof(double), 3*N_atoms, file_pointer);
167 fwrite(&cell_length, sizeof(double), 1, file_pointer);
168 fclose(file_pointer);
169
170 free(pos); pos = NULL;
171 free(momentum); momentum = NULL;
172 free(forces); forces = NULL;
173 free(temperature); temperature = NULL;
174 free(pressure); pressure = NULL;
175 free(a0); a0 = NULL;
176 return 0;
177 }

```

A.4 Production runs for tasks 3-7 : main_Prod.c

```

1  /*
2  main_Prod.c, Production runs, H1b

```

```

3 In this program, we use the equilibrated micro-states from Tasks 3-4 to study
4 dynamical properties, such as mean squared displacement (MSD), velocity
5 auto-correlation function, and the power spectral density of the atom
6 movements.
7
8 System of units:
9 Energy - eV
10 Time - ps
11 Length - Angstrom
12 Temp - K
13 Mass - eV (ps)2 A(-2)
14 Pressure - eV A(-3)
15 */
16
17 #include <stdio.h>
18 #include <math.h>
19 #include <stdlib.h>
20 #include <time.h>
21
22 #include "initfcc.h"
23 #include "alpotential.h"
24 #include "funcs.h"
25
26 #define N_cells 4
27 /* define constants in atomic units: eV, , ps, K */
28 #define AMU 1.0364e-4
29 #define degC_to_K 273.15
30 #define bar 6.2415e-07
31 #define kB 8.61733e-5
32
33 /* Main program */
34 int main()
35 {
36     char file_name[100];
37
38     int N_atoms = 4*N_cells*N_cells*N_cells;
39     double m_Al = 27*AMU;
40     double cell_length;
41     double inv_volume;
42
43     double T_eq_C = 700;
44     double P_eq_bar = 1;
45
46     double dt = 1e-3; // higher res for spectral function
47     double t_end = 10;
48     int N_timesteps = t_end/dt;
49     int N_between_steps = 1; // save all steps for max res in spectral function
50     int N_save_timesteps = N_timesteps / N_between_steps;
51     int N_save_atoms = 5;
52
53     double t, E_kin, virial;
54
55     double (*pos)[3] = malloc(sizeof(double[N_atoms][3]));
56     double (*pos_0)[3] = malloc(sizeof(double[N_atoms][3])); //for displacements
57     double (*momentum)[3] = malloc(sizeof(double[N_atoms][3]));
58     double (*forces)[3] = malloc(sizeof(double[N_atoms][3]));
59     double (*displacements)[N_save_atoms] =
60         malloc(sizeof(double[N_save_timesteps][N_save_atoms]));
61     double (*pos_all)[N_atoms][3] =
62         malloc(sizeof(double[N_save_timesteps][N_atoms][3]));
63     double (*vel_all)[N_atoms][3] =
64         malloc(sizeof(double[N_save_timesteps][N_atoms][3]));
65     double *temperature = malloc(sizeof(double[N_timesteps]));
66     double *pressure = malloc(sizeof(double[N_timesteps]));
67     double *msd = malloc(sizeof(double[N_save_timesteps]));
68     double *vel_corr = malloc(sizeof(double[N_save_timesteps]));
69     double *pow_spec = malloc(sizeof(double[N_save_timesteps]));
70     double *freq = malloc(sizeof(double[N_save_timesteps]));
71
72     // Initialize to 0
73     for (int i = 0; i < N_save_timesteps; i++){
74         msd[i] = 0;
75         pow_spec[i] = 0;
76         vel_corr[i] = 0;
77     }
78     FILE *file_pointer;
79
80     // read positions, momenta and cell_length
81     sprintf(file_name, "../data/INIDATA-temp-%d-pres-%d.bin",
82         (int) T_eq_C, (int) P_eq_bar);
83     file_pointer = fopen(file_name, "rb");
84     fread(pos, sizeof(double), 3*N_atoms, file_pointer);
85     fread(momentum, sizeof(double), 3*N_atoms, file_pointer);
86     fread(&cell_length, sizeof(double), 1, file_pointer);
87     fclose(file_pointer);
88
89     for (int i=0; i<N_atoms; i++){
90         for (int j=0; j<3; j++){
91             pos_0[i][j]=pos[i][j];
92         }
93     }

```

```

94 inv_volume = pow(cell_length, -3);
95 get_forces_AL( forces, pos, cell_length, N_atoms); //initial cond forces
96
97 printf("Initialized. Starting with Verlet timestepping.\n");
98 for (int i=0; i<N_timesteps; i++){
99     /*
100     The loop over the timesteps first takes a timestep according to the
101     Verlet algorithm, then calculates the energies and temeperature.
102     */
103     timestep_Verlet(N_atoms, pos, momentum, forces, m_Al, dt, cell_length);
104
105     E_kin = get_kin_energy(N_atoms, momentum, m_Al );
106     virial = get_virial_AL(pos, cell_length, N_atoms);
107     /* PV = NkT + virial */
108     pressure[i] = inv_volume * (E_kin/1.5 + virial);
109     /* 3N*kB*T/2 = 1/(2m) * \sum_{i=1}^N p_i^2 = p_sq/(2m) */
110     temperature[i] = E_kin * 1/(1.5*N_atoms*kB);
111
112     if (i % N_between_steps == 0){
113         int k = i/N_between_steps; // number of saved timesteps so far
114         // Saves the displacements of some atoms into 'displacements'
115         get_displacements (N_save_atoms, pos, pos_0, displacements[k]);
116
117         // Saves all the positions
118         copy_mat(N_atoms, 3, pos, pos_all[k]);
119
120         // Saves all the velocities
121         copy_mat(N_atoms, 3, momentum, vel_all[k]);
122         //But we need to scale the momenta to get the velocities
123         scale_mat(N_atoms, 3, vel_all[k], 1/m_Al);
124     }
125
126     if ((i*10) % N_timesteps == 0){ //Print out progress at every 10%
127         printf("done %d%% of Verlet timestepping\n", (i*10)/N_timesteps);
128     }
129 }
130 printf("done 100%% of Verlet timestepping\n");
131
132 //Calculating MSD
133 printf("calculating MSD\n");
134 get_MSD(N_atoms, N_save_timesteps, pos_all, msd);
135
136 //Calculating the velocity correlation function
137 printf("calculating velocity correlation\n");
138 get_vel_corr(N_atoms, N_save_timesteps, vel_all, vel_corr);
139
140 //Calculating the velocity power spectrum
141 printf("calculating power spectrum\n");
142 get_powerspectrum(N_atoms, N_save_timesteps, vel_all, pow_spec);
143 fft_freq(freq, dt, N_save_timesteps);
144
145 printf("writing to file\n");
146 /* Write temperture to file */
147 sprintf(file_name, "../data/temp-%d_pres-%d_Prod-test.tsv",
148         (int) T_eq_C, (int) P_eq_bar);
149 file_pointer = fopen(file_name, "w");
150 for (int i=0; i<N_timesteps; i++){
151     t = i*dt; // time at step i
152     fprintf(file_pointer, "%.4f \t %.8f \t %.8f \n",
153             t, temperature[i], pressure[i]);
154 }
155 fclose(file_pointer);
156
157 /* Write displacements to file */
158 sprintf(file_name, "../data/temp-%d_pres-%d_displacements.tsv",
159         (int) T_eq_C, (int) P_eq_bar);
160 file_pointer = fopen(file_name, "w");
161 for (int i=0; i<N_save_timesteps; i++){
162     t = i*dt*N_between_steps; // time at step i
163     fprintf(file_pointer, "%.4f", t);
164     for (int j=0; j<N_save_atoms; j++){
165         fprintf(file_pointer, "\t %.8f", displacements[i][j]);
166     }
167     fprintf(file_pointer, "\n");
168 }
169 fclose(file_pointer);
170
171 /* Write MSD to file */
172 sprintf(file_name, "../data/temp-%d_pres-%d_dynamicProperties.tsv",
173         (int) T_eq_C, (int) P_eq_bar);
174 file_pointer = fopen(file_name, "w");
175 // write header
176 fprintf(file_pointer, "%s t[ps] \t MSD[A^2] \t vel_corr [A/ps]^2 \n");
177 for (int i=0; i<N_save_timesteps; i++){
178     t = i*dt*N_between_steps; // time at step i
179     fprintf(file_pointer, "%.4f \t %.8f \t %.8f \n", t, msd[i], vel_corr[i]);
180 }
181 fclose(file_pointer);
182
183 /* Write power spectrum to file */
184 sprintf(file_name, "../data/temp-%d_pres-%d_power-spectrum.tsv",

```

```

185     (int) T_eq_C, (int) P_eq_bar);
186     file_pointer = fopen(file_name, "w");
187     // write header
188     fprintf(file_pointer, "%% f[1/ps] \t P[A/ps]^2 \n");
189     for (int i=0; i<N_save_timesteps/2; i++){ // only save from f=0 to f_crit
190         fprintf(file_pointer, "%.4f \t %.8f \n", freq[i], pow_spec[i]);
191     }
192     fclose(file_pointer);
193
194     // Freeing all the memory
195     free(pos);          pos = NULL;
196     free(pos_0);        pos_0 = NULL;
197     free(momentum);     momentum = NULL;
198     free(forces);       forces = NULL;
199     free(temperature);  temperature = NULL;
200     free(pressure);     pressure = NULL;
201     free(displacements); displacements = NULL;
202     free(pos_all);      pos_all = NULL;
203     free(vel_all);      vel_all = NULL;
204     free(msd);          msd = NULL;
205     free(vel_corr);     vel_corr = NULL;
206     free(pow_spec);     pow_spec = NULL;
207     free(freq);         freq = NULL;
208
209     return 0;
210 }

```

A.5 Production runs for tasks 3-7 : main_Prod.c

```

1  /*
2   main_Prod.c, Production runs, H1b
3   In this program, we use the equilibrated micro-states from Tasks 3-4 to study
4   dynamical properties, such as mean squared displacement (MSD), velocity
5   auto-correlation function, and the power spectral density of the atom
6   movements.
7
8   System of units:
9   Energy   - eV
10  Time     - ps
11  Length   - Angstrom
12  Temp     - K
13  Mass     - eV (ps)^2 A^(-2)
14  Pressure - eV A^(-3)
15  */
16
17  #include <stdio.h>
18  #include <math.h>
19  #include <stdlib.h>
20  #include <time.h>
21
22  #include "initfcc.h"
23  #include "alpotential.h"
24  #include "funcs.h"
25
26  #define N_cells 4
27  /* define constants in atomic units: eV, , ps, K */
28  #define AMU 1.0364e-4
29  #define degC_to_K 273.15
30  #define bar 6.2415e-07
31  #define kB 8.61733e-5
32
33  /* Main program */
34  int main()
35  {
36      char file_name[100];
37
38      int N_atoms = 4*N_cells*N_cells*N_cells;
39      double m_Al = 27*AMU;
40      double cell_length;
41      double inv_volume;
42
43      double T_eq_C = 700;
44      double P_eq_bar = 1;
45
46      double dt = 1e-3; // higher res for spectral function
47      double t_end = 10;
48      int N_timesteps = t_end/dt;
49      int N_between_steps = 1; // save all steps for max res in spectral function
50      int N_save_timesteps = N_timesteps / N_between_steps;
51      int N_save_atoms = 5;
52
53      double t, E_kin, virial;
54
55      double (*pos)[3] = malloc(sizeof(double[N_atoms][3]));
56      double (*pos_0)[3] = malloc(sizeof(double[N_atoms][3])); //for displacements
57      double (*momentum)[3] = malloc(sizeof(double[N_atoms][3]));

```

```

58 double (*forces)[3] = malloc(sizeof(double)[N_atoms][3]));
59 double (*displacements)[N_save_atoms] =
60     malloc(sizeof(double)[N_save_timesteps][N_save_atoms]));
61 double (*pos_all)[N_atoms][3] =
62     malloc(sizeof(double)[N_save_timesteps][N_atoms][3]));
63 double (*vel_all)[N_atoms][3] =
64     malloc(sizeof(double)[N_save_timesteps][N_atoms][3]));
65 double *temperature = malloc(sizeof(double)[N_timesteps]);
66 double *pressure = malloc(sizeof(double)[N_timesteps]);
67 double *msd = malloc(sizeof(double)[N_save_timesteps]);
68 double *vel_corr = malloc(sizeof(double)[N_save_timesteps]);
69 double *pow_spec = malloc(sizeof(double)[N_save_timesteps]);
70 double *freq = malloc(sizeof(double)[N_save_timesteps]);
71
72 // Initialize to 0
73 for (int i = 0; i < N_save_timesteps; i++){
74     msd[i] = 0;
75     pow_spec[i] = 0;
76     vel_corr[i] = 0;
77 }
78 FILE *file_pointer;
79
80 // read positions, momenta and cell_length
81 sprintf(file_name, "../data/INIDATA-temp-%d-pres-%d.bin",
82         (int) T_eq_C, (int) P_eq_bar);
83 file_pointer = fopen(file_name, "rb");
84 fread(pos, sizeof(double), 3*N_atoms, file_pointer);
85 fread(momentum, sizeof(double), 3*N_atoms, file_pointer);
86 fread(&cell_length, sizeof(double), 1, file_pointer);
87 fclose(file_pointer);
88
89 for (int i=0; i < N_atoms; i++){
90     for (int j=0; j < 3; j++){
91         pos_0[i][j]=pos[i][j];
92     }
93 }
94 inv_volume = pow(cell_length, -3);
95 get_forces_AL( forces, pos, cell_length, N_atoms); //initial cond forces
96
97 printf("Initialized. Starting with Verlet timestepping.\n");
98 for (int i=0; i < N_timesteps; i++){
99     /*
100      The loop over the timesteps first takes a timestep according to the
101      Verlet algorithm, then calculates the energies and temeperature.
102     */
103     timestep_Verlet(N_atoms, pos, momentum, forces, m_AL, dt, cell_length);
104
105     E_kin = get_kin_energy(N_atoms, momentum, m_AL );
106     virial = get_virial_AL(pos, cell_length, N_atoms);
107     /* PV = NkT + virial */
108     pressure[i] = inv_volume * (E_kin/1.5 + virial);
109     /* 3N*kB*T/2 = 1/(2m) * \sum_{i=1}^N p_i^2 = p_sq/(2m) */
110     temperature[i] = E_kin * 1/(1.5*N_atoms*kB);
111
112     if (i % N_between_steps == 0){
113         int k = i/N_between_steps; // number of saved timesteps so far
114         // Saves the displacements of some atoms into `displacements`
115         get_displacements (N_save_atoms, pos, pos_0, displacements[k]);
116
117         // Saves all the positions
118         copy_mat(N_atoms, 3, pos, pos_all[k]);
119
120         // Saves all the velocities
121         copy_mat(N_atoms, 3, momentum, vel_all[k]);
122         //But we need to scale the momenta to get the velocities
123         scale_mat(N_atoms, 3, vel_all[k], 1/m_AL);
124     }
125
126     if ((i*10) % N_timesteps == 0){ //Print out progress at every 10%
127         printf("done %d%% of Verlet timestepping\n", (i*10)/N_timesteps);
128     }
129 }
130 printf("done 100%% of Verlet timestepping\n");
131
132 //Calculating MSD
133 printf("calculating MSD\n");
134 get_MSD(N_atoms, N_save_timesteps, pos_all, msd);
135
136 //Calculating the velocity correlation function
137 printf("calculating velocity correlation\n");
138 get_vel_corr(N_atoms, N_save_timesteps, vel_all, vel_corr);
139
140 //Calculating the velocity power spectrum
141 printf("calculating power spectrum\n");
142 get_powerspectrum(N_atoms, N_save_timesteps, vel_all, pow_spec);
143 fft_freq(freq, dt, N_save_timesteps);
144
145 printf("writing to file\n");
146 /* Write temperture to file */
147 sprintf(file_name, "../data/temp-%d-pres-%d-Prod-test.tsv",
148         (int) T_eq_C, (int) P_eq_bar);

```



```

149 file_pointer = fopen(file_name, "w");
150 for (int i=0; i<N_timesteps; i++){
151     t = i*dt; // time at step i
152     fprintf(file_pointer, "%.4f \t %.8f \t %.8f \n",
153         t, temperature[i], pressure[i]);
154 }
155 fclose(file_pointer);
156
157 /* Write displacements to file */
158 sprintf(file_name, "../data/temp-%d_pres-%d_displacements.tsv",
159     (int) T_eq_C, (int) P_eq_bar);
160 file_pointer = fopen(file_name, "w");
161 for (int i=0; i<N_save_timesteps; i++){
162     t = i*dt*N_between_steps; // time at step i
163     fprintf(file_pointer, "%.4f", t);
164     for (int j=0; j<N_save_atoms; j++){
165         fprintf(file_pointer, "\t %.8f", displacements[i][j]);
166     }
167     fprintf(file_pointer, "\n");
168 }
169 fclose(file_pointer);
170
171 /* Write MSD to file */
172 sprintf(file_name, "../data/temp-%d_pres-%d_dynamicProperties.tsv",
173     (int) T_eq_C, (int) P_eq_bar);
174 file_pointer = fopen(file_name, "w");
175 // write header
176 fprintf(file_pointer, "%s t[ps] \t MSD[A^2] \t vel_corr [A/ps]^2 \n");
177 for (int i=0; i<N_save_timesteps; i++){
178     t = i*dt*N_between_steps; // time at step i
179     fprintf(file_pointer, "%.4f \t %.8f \t %.8f \n", t, msd[i], vel_corr[i]);
180 }
181 fclose(file_pointer);
182
183 /* Write power spectrum to file */
184 sprintf(file_name, "../data/temp-%d_pres-%d_power-spectrum.tsv",
185     (int) T_eq_C, (int) P_eq_bar);
186 file_pointer = fopen(file_name, "w");
187 // write header
188 fprintf(file_pointer, "%s f[1/ps] \t P[A/ps]^2 \n");
189 for (int i=0; i<N_save_timesteps/2; i++){ // only save from f=0 to f_crit
190     fprintf(file_pointer, "%.4f \t %.8f \n", freq[i], pow_spec[i]);
191 }
192 fclose(file_pointer);
193
194 // Freeing all the memory
195 free(pos); pos = NULL;
196 free(pos_0); pos_0 = NULL;
197 free(momentum); momentum = NULL;
198 free(forces); forces = NULL;
199 free(temperature); temperature = NULL;
200 free(pressure); pressure = NULL;
201 free(displacements); displacements = NULL;
202 free(pos_all); pos_all = NULL;
203 free(vel_all); vel_all = NULL;
204 free(msd); msd = NULL;
205 free(vel_corr); vel_corr = NULL;
206 free(pow_spec); pow_spec = NULL;
207 free(freq); freq = NULL;
208
209 return 0;
210 }

```

A.6 Misc functions : funcs.c

```

1 #include "funcs.h"
2
3 void add_noise(int M, int N, double mat[M][N], double noise_amplitude )
4 {
5     const gsl_rng_type *T; // static info about rngs
6     gsl_rng *q; // rng instance
7     gsl_rng_env_setup (); // setup the rngs
8     T = gsl_rng_default; // specify default rng
9     q = gsl_rng_alloc(T); // allocate default rng
10    gsl_rng_set(q, time(NULL)); // Initialize rng
11
12    // Loops over all the elemtns in the matrix, to which we want to add noise
13    for (int i=0; i<M; i++){
14        for (int j=0; j<N; j++){
15            // adds uniformly distributed random noise in range +/-`noise_amplitude`
16            mat[i][j] += noise_amplitude * (2*gsl_rng_uniform(q)-1);
17        }
18    }
19    gsl_rng_free(q); // deallocate rng
20 }
21

```

```

22 void timestep_Verlet ( int N_atoms, double (*pos)[3], double (*momentum)[3],
23                       double (*forces)[3], double m, double dt,
24                       double cell_length){
25     for (int i = 0; i < N_atoms; i++) {
26       // Half-steps the momentum, then steps the position
27       for (int j = 0; j < 3; j++){
28         // p(t+dt/2)
29         momentum[i][j] += dt * 0.5 * forces[i][j];
30         // q(t+dt)
31         pos[i][j] += dt * momentum[i][j] / m;
32       }
33     }
34     // Updates the forces, based on the new positions
35     // F(t+dt)
36     get_forces_AL( forces, pos, cell_length, N_atoms);
37     // Another half-step in the momenta
38     for (int i = 0; i < N_atoms; i++) {
39       for (int j = 0; j < 3; j++) {
40         // p(t+dt/2)
41         momentum[i][j] += dt * 0.5 * forces[i][j];
42       }
43     }
44 }
45
46 double get_kin_energy ( int N_atoms, double (*momentum)[3], double m ) {
47   double p_sq=0; // momentum squared
48   for (int i = 0; i < N_atoms; i++) {
49     for (int j = 0; j < 3; j++) {
50       p_sq += momentum[i][j] * momentum[i][j];
51     }
52   }
53   // E_kin = p^2/(2m)
54   return p_sq / (2*m);
55 }
56
57 void get_displacements ( int N_atoms, double (*positions)[3],
58                         double (*initial_positions)[3], double disp[]) {
59   for (int i = 0; i < N_atoms; i++) {
60     for (int j = 0; j < 3; j++) {
61       disp[i] += (positions[i][j] - initial_positions[i][j])
62         *(positions[i][j] - initial_positions[i][j]);
63     }
64     disp[i] = sqrt(disp[i]);
65   }
66 }
67
68
69 void get_MSD ( int N_atoms, int N_times, double all_pos[N_times][N_atoms][3],
70              double MSD[N_times]) {
71   // all_pos = positions of all particles at all (saved) times
72   // outer time index it starts at outer it = 1, since MSD[0] = 0
73   for (int it = 1; it < N_times; it++) { //
74     for (int jt = 0; jt < N_times-it; jt++) { // summed time index
75       for (int kn = 0; kn < N_atoms; kn++) { // particle index
76         for (int kd = 0; kd < 3; kd++) { // three dimensions
77           MSD[it] += (all_pos[it+jt][kn][kd] - all_pos[jt][kn][kd])
78             *(all_pos[it+jt][kn][kd] - all_pos[jt][kn][kd]);
79         }
80       }
81     }
82     MSD[it] *= 1/( (double)N_atoms * (N_times-it));
83   }
84 }
85
86 void get_vel_corr ( int N_atoms, int N_times,
87                   double all_vel[N_times][N_atoms][3],
88                   double vel_corr[N_times]) {
89   /* all_vel = velocity of all particles at all (saved) times */
90   for (int it = 0; it < N_times; it++) { //
91     for (int jt = 0; jt < N_times-it; jt++) { // summed time index
92       for (int kn = 0; kn < N_atoms; kn++) { // particle index
93         for (int kd = 0; kd < 3; kd++) { // three dimensions
94           vel_corr[it] += (all_vel[it+jt][kn][kd] * all_vel[jt][kn][kd]);
95         }
96       }
97     }
98     vel_corr[it] *= 1/( (double)N_atoms * (N_times-it));
99   }
100 }
101
102 void get_powerspectrum ( int N_atoms, int N_times,
103                         double all_vel[N_times][N_atoms][3],
104                         double pow_spec[N_times]) {
105   /* all_vel = velocity of all particles at all (saved) times */
106
107   double vel_component[N_times]; // temp. var. "all_vel[:,i][j]"
108   double pow_spec_component[N_times]; // temp. var. one component F-transf.
109   double normalization_factor = 1/((double) N_atoms*N_times);
110
111   for (int kn = 0; kn < N_atoms; kn++){ // for particle index
112     for (int kd = 0; kd < 3; kd++){ // for 3D

```

```

113 // Copies the velocity component of one particle, into temporary variable
114 for (int it = 0; it < N_times; it++){
115     vel_component[it] = all_vel[it][kn][kd];
116 }
117 //Calculates the power spect. of this one velocity component
118 powerspectrum(vel_component, pow_spec_component, N_times);
119 //Adds the powerspectrum to the "output" variable
120 for (int iw = 0; iw < N_times; iw++) { // for all frequencies
121     pow_spec[iw] += pow_spec_component[iw];
122 } // end for all frequencies
123 } // end for 3D
124 } // end for particle index
125 for (int iw = 0; iw < N_times; iw++) { // for all frequencies
126     pow_spec[iw] *= normalization_factor;
127 }
128 }
129
130
131
132 void copy_mat (int M, int N, double mat_from[M][N], double mat_to[M][N]){
133     /* Copies all matrix elements of `mat_from` to `mat_to` */
134     //loops over all indices
135     for (int i = 0; i < M; i++) {
136         for (int j = 0; j < N; j++) {
137             mat_to[i][j] = mat_from[i][j];
138         }
139     }
140 }
141
142 void set_zero (int M, int N, double mat[M][N]){
143     /* Sets all the elements of matrix `mat` to zero */
144     //loops over all indices
145     for (int i = 0; i < M; i++) {
146         for (int j = 0; j < N; j++) {
147             mat[i][j] = 0;
148         }
149     }
150 }
151
152 void scale_mat (int M, int N, double mat[M][N], double alpha){
153     /* Scales the matrix `mat` by factor `alpha` */
154     //loops over all indices
155     for (int i = 0; i < M; i++) {
156         for (int j = 0; j < N; j++) {
157             mat[i][j] *= alpha;
158         }
159     }
160 }

```

B Auxiliary

B.1 Makefile

```

1
2 CC = gcc
3 CFLAGS = -O3 -Wall -Wno-unused-result
4
5 LIBS = -lm -lgsl -lgslcblas
6
7 HEADERS = initfcc.h alpotential.h funcs.h fft_func.h
8 OBJECTS = initfcc.o alpotential.o funcs.o fft_func.o
9
10
11 %.o: %.c $(HEADERS)
12     $(CC) -c -o $@ $< $(CFLAGS)
13
14 all: Task1 Task2 Task3 main_Prod.c
15
16 Task1: $(OBJECTS) main_T1.c
17     $(CC) -o $@ $^ $(CFLAGS) $(LIBS)
18
19 Task2: $(OBJECTS) main_T2.c
20     $(CC) -o $@ $^ $(CFLAGS) $(LIBS)
21
22 Task3: $(OBJECTS) main_T3.c
23     $(CC) -o $@ $^ $(CFLAGS) $(LIBS)
24
25 Prod: $(OBJECTS) main_Prod.c
26     $(CC) -o $@ $^ $(CFLAGS) $(LIBS)
27
28 # $(PROGRAMS): $(OBJECTS) main_T1.c
29 #     $(CC) -o $@ $^ $(CFLAGS) $(LIBS)
30
31 clean:

```

```

32 rm -f *.o
33 touch *.c

```

C MATLAB scripts

C.1 Analysis scripts for tasks 3-7: Al_energies.m

```

1 %% initial
2
3 tmp = matlab.desktop.editor.getActive; %% cd to current path
4 cd(fileparts(tmp.Filename));
5 set(0,'DefaultFigureWindowStyle','docked');
6 warning('off','MATLAB:handle_graphics:exceptions:SceneNode'); % interpreter
7 GRAY = 0.7*[0.9 0.9 1];
8 AMU = 1.0364e-4;
9 m_Al = 27*AMU;
10 %% task 1: lattice energies
11 clc
12
13 energy_data = load('../data/lattice_energies.tsv');
14 a0 = energy_data(:,1);
15 v0 = a0.^3;
16
17 energy = energy_data(:,2);
18 figure(1);clf;
19 plot(v0,energy, 'xk');
20
21 start_v = 64;
22 end_v = 68;
23 indToInclude = (v0 > start_v) & (v0 < end_v);
24 p = polyfit(v0(indToInclude),energy(indToInclude),2);
25 hold on;
26
27 vvec = linspace(start_v, end_v);
28 plot(vvec, p(1)*vvec.^2 + p(2)*vvec + p(3), '-r');
29 xlim([64 68]);
30
31 v_min = -p(2)/(2*p(1));
32 a_min = v_min^(1/3);
33 omega_res = sqrt(2*p(1)*a_min^4/m_Al);
34 f_res = omega_res/(2*pi); % rough estimation of resonance frequency (?)
35
36 ax = gca; ax.YLim = [-13.45 -13.42];
37 h1 = plot(v_min*[1 1], ax.YLim, '--k'); % plot vertical line at v_min
38
39 ax.YTick = (-13.45:0.01:-13.42);
40 ylabel('$E_{\rm pot}$ [eV/unit cell]');
41 xlabel('$a_0^3$ [\AA^3]');
42 legend('data', 'quadratic fit', ['$V_{\rm eq} \approx \backslash, $' ...
43       num2str(round(v_min,2)) '\, \AA^3$'], ...
44       'location', 'southeast')
45 ax = gca; ax.Children = ax.Children(3:-1:1);
46 ImproveFigureCompPhys(gcf); h1.LineWidth = 2; setFigureSize(gcf, 300, 600);
47 saveas(gcf, '../figures/potential_energy.eps', 'epsc')
48 %% task 2: find a suitable timestep
49 clc;
50
51 dt=[2e-2,1e-2,5e-3,2e-3];
52 t_eq=0.5;
53
54 figure(1);clf;figure(2);clf;
55
56 for i=1:numel(dt)
57     T_data = load(sprintf('../data/temperature_dt-%0.0e_Task2.tsv',dt(i)));
58     E_data = load(sprintf('../data/total_energy_dt-%0.0e_Task2.tsv',dt(i)));
59     t = T_data(:,1);
60     T = T_data(:,2);
61     E = E_data(:,2);
62
63     fprintf('dt = %0.0e\n',dt(i));
64
65     T_avg=mean(T(t>t_eq));
66     T_std=std(T(t>t_eq));
67     fprintf('\tT = %0.2f +- %0.1f %%\n', T_avg, abs(T_std/T_avg)*100);
68
69     E_avg=mean(E(t>t_eq));
70     E_std=std(E(t>t_eq));
71     fprintf('\tE = %0.2f +- %0.1e %%\n', E_avg, abs(E_std/E_avg)*100);
72
73     figure(1)
74     plot(t, T); hold on;
75
76     figure(2)
77     plot(t, E);hold on;

```

```

78 end
79 for ifig = 1:2
80     figure(ifig);
81     h = legend(strcat({'$dt = $ '}, num2str(round(dt',4)) , ' ps'));
82     xlabel('$t$ [ps]');
83     ax = gca;
84     if ifig ==1
85         ylabel('$\mathcal{T}$ [K]');
86         ax.YLim = [400 1800];
87     else
88         ylabel('$E_{\rm tot}$ [eV/unit cell]');
89         ax.YTick = (-13:0.1:-10);
90         ax.YLim = [-12.75 -12.5];
91     end
92     ImproveFigureCompPhys(gcf,'Linewidth', 2);setFigureSize(gcf, 400, 400);
93 end
94 saveas(1, '../figures/dt-scan-temperature.eps', 'epsc')
95 saveas(2, '../figures/dt-scan-energy.eps', 'epsc')
96 %% task 3: temperature and pressure equilibration,
97 % and task4: test production pressure and temperature
98
99 clf; figure(10);clf;
100 temps = [500 700 500 700];
101 temperatures_str = num2str([500;700]);
102 FILENAMES = [strcat({'../data/temp-'}, temperatures_str,...
103     '_pres-1_Task3.tsv');
104     strcat({'../data/temp-'}, temperatures_str, '_pres-1_Prod-test.tsv')];
105 bar = 6.2415e-07;
106 Kelvin_to_degC = -273.15;
107 t_eqs = [1 1 0.5 0.5]; % approximate equilibration time
108 N_average_points = 100;
109 dt = 5e-3;
110 tau_eqilibration = 100*dt;
111
112 for iFile = 1:numel(FILENAMES)
113     figure(iFile);clf;
114     data = load(FILENAMES{iFile});
115
116     t = data(:,1);
117     T = data(:,2)+Kelvin_to_degC;
118     P = data(:,3)/bar/1e4; %GPa
119
120     t_eq=t_eqs(iFile);
121
122     T_avg=mean(T(t>t_eq));
123     T_std=std(T(t>t_eq));
124     fprintf('\tT = %0.2f +- %0.1f C\n', T_avg, abs(T_std));
125
126     P_avg=mean(P(t>t_eq));
127     P_std=std(P(t>t_eq));
128     fprintf('\tP = %0.2f +- %0.1f bar\n', P_avg*1e4, abs(P_std*1e4)); % convert ←
129         to bar
130
131     yyaxis left
132     %subplot(2,1,1)
133
134     if iFile <=2 % equilibration run, otherwise production
135         plot(t./tau_eqilibration,T, 'color', GRAY),hold on;
136         plot(t./tau_eqilibration, movmean(T,N_average_points),'-k')
137     else
138         plot(t,T, 'color', GRAY),hold on;
139         plot(t, cumsum(T)./(1:length(t)),'-k')
140     end
141     ylabel('$T \, [\circ \rm C]$')
142
143     if iFile <=2 % equilibration run, otherwise production
144         ylim(temps(iFile)*(1+ 0.3*[-3*0.6,0.6]))
145         yyaxis right
146         plot(t./tau_eqilibration,P),hold on;
147         plot(t./tau_eqilibration, movmean(P,N_average_points),'-k')
148
149         legend('$\mathcal{T}$', 'mov avg', '$\mathcal{P}$', 'mov avg', 'location'←
150             , 'west');
151         xlabel('$t/\tau_{\rm eq}$')
152         %xlim([0 5])
153     else
154         ylim(temps(iFile)+ 100*[-3,1])
155         yyaxis right
156         plot(t,P),hold on;
157         plot(t, cumsum(P)./(1:length(t)),'-k')
158         legend('$\mathcal{T}$', 'cum avg', '$\mathcal{P}$', 'cum avg', 'location'←
159             , 'west');
160         xlabel('$t$, [ps]')
161     end
162     ylabel('$P \, [\rm GPa]$')
163     ylim([-0.6,0.6*3])
164     ImproveFigureCompPhys(gcf, 'linewidth', 3, 'LineColor', ...
165         {'MYORANGE', GRAY, 'MYBLUE', GRAY});
166     %ImproveFigureCompPhys(gcf, 'linewidth', 3, 'LineColor', ...
167         {'k', GRAY});
168     setFigureSize(gcf, 400, 400);

```

```

166     if iFile <=2 % plot a0 too
167         figure(10)
168         a0 = data(:,4);
169         plot(t./tau_eqilibration, a0);hold on;
170     end
171 end
172
173
174 figure(10);
175 xlabel('$t/\tau_{\rm eq}$');
176 ylabel('$a_0$');
177 legend('$T = 500^\circ\text{C}$', '$T = 700^\circ\text{C}$')
178 setFigureSize(gcf, 300, 600);
179 ImproveFigureCompPhys(gcf, 'LineColor', {'MYRED', 'MYLIGHTBLUE'}, 'LineStyle', {'-','-.'});
180
181 saveas(1, '../figures/TP-eq-500.eps', 'epsc')
182 saveas(2, '../figures/TP-eq-700.eps', 'epsc')
183 saveas(3, '../figures/TP-prod-500.eps', 'epsc')
184 saveas(4, '../figures/TP-prod-700.eps', 'epsc')
185 saveas(10, '../figures/a0.eps', 'epsc')
186 %% determine displacements and MSD
187 temperatures_str = num2str([500;700]);
188 clc; clf;
189 figure(10); clf;
190 FILENAMES = strcat({'../data/temp-'}, temperatures_str, ...
191 '_pres-1_displacements.tsv');
192 FILENAMES_Dyn = strcat({'../data/temp-'}, temperatures_str, ...
193 '_pres-1_dynamicProperties.tsv');
194 FILENAMES_Pow = strcat({'../data/temp-'}, temperatures_str, ...
195 '_pres-1_power-spectrum.tsv');
196 for iFile = 1:numel(FILENAMES)
197     figure(iFile); clf;
198     data = load(FILENAMES{iFile});
199     t = data(:,1);
200     dx = data(:,2:end);
201
202     data = load(FILENAMES_Dyn{iFile});
203     MSD = data(:,2);
204     vel_corr = data(:,3);
205     plot(t, MSD, 'k'); hold on;
206
207     if iFile ==2 % liquid
208         tStart = 1; tEnd = t(end)-tStart;
209         goodInd = (t>tStart & t<tEnd);
210
211         D = MSD(goodInd)./(6*t(goodInd));
212
213         selfDiffusionCoeff = mean(D); % in  $\text{\AA}^2/\text{ps}$ 
214         plot(t, 6*t*selfDiffusionCoeff, 'r');
215     end
216
217     plot(t, dx.^2, 'color', GRAY); hold on;
218
219     xlabel('$t$ [ps]')
220     ylabel('$\Delta x^2$, [\rm \AA^2]$')
221     if iFile ==1
222         ylim([0 0.5]);
223         leg = legend('$\Delta_{\rm MSD}$', 'individual trajectories');
224     else
225         ylim([0 30]);
226         leg = legend('$\Delta_{\rm MSD}$', '$6 t D_{\rm s}$', ...
227             'individual trajectories');
228     end
229     xlim([0 6])
230
231     leg.Location='northwest';
232     ImproveFigureCompPhys(gcf, 'Linewidth', 2);
233     ax = gca; [ax.Children(6:end).LineWidth] = deal(5);
234     ax.Children = ax.Children([6:end 1:5]);
235     setFigureSize(gcf, 300, 600);
236 end
237
238 % velocity correlation
239 figure(10);clf; figure(11);clf;
240 n_average_points = 1;%30;
241 for iFile = 1:numel(FILENAMES)
242     data = load(FILENAMES_Dyn{iFile});
243     t = data(:,1);
244     vel_corr = data(:,3);
245
246     data = load(FILENAMES_Pow{iFile});
247     freq = data(:,1);
248     pow_spec = data(:,2);
249
250     figure(10);
251     plot(t, vel_corr/vel_corr(1)); hold on;
252
253     dt = t(2)-t(1);
254     N_times = round(length(t)/1.5);% we have bad statistics at later times.
255     deltaf = 1/(N_times * dt);

```

```

256     freqvec = 0:deltaf:(1/(2*dt));
257     PhiHat = 2 * trapz(t(1:N_times), ...
258         (vel_corr(1:N_times) * ones(size(freqvec))) .* ...
259         cos(2*pi*t(1:N_times) * freqvec), 1);
260
261     figure(11);
262     plot(freqvec, PhiHat); hold on;
263     plot(freq, pow_spec*t(end), '-.'); hold on;
264     if iFile ==2 % liquid
265         tStart = 1;
266         selfDiffusionCoeff_spectral1 = mean(PhiHat(1:3))/6; % in  $\text{\AA}^2/\text{ps}$ 
267         selfDiffusionCoeff_spectral2 = mean(pow_spec(1:3)*t(end))/6; % in  $\text{\AA}^2/\text{ps}$ 
268     end
269
270 end
271
272 disp([selfDiffusionCoeff selfDiffusionCoeff_spectral1 ←
    selfDiffusionCoeff_spectral2]);
273
274 figure(10)
275 xlim([0 1]);
276 leg = legend(strcat({'$T='}, num2str([500;700]), '\, '\circ $C'));
277 leg.Location='northeast';
278 xlabel('$t$ [ps]')
279 ylabel('$\Phi(t)/\Phi(0)$')
280 ImproveFigureCompPhys(gcf,'LineColor', {'MYRED', 'MYLIGHTBLUE'});
281 setFigureSize(gcf, 300, 600);
282
283 figure(11)
284 leg = legend('$T= 500 \, \circ $C, $ \hat{\Phi}$', ...
285     '$T= 500 \, \circ $C, $ \hat{P}$',...
286     '$T= 700 \, \circ $C, $ \hat{\Phi}$', ...
287     '$T= 700 \, \circ $C, $ \hat{P}$');
288 xlim([0 20])
289 ylim([0 Inf])
290 xlabel('$f$ [THz]')
291 ylabel('$\hat{P}$ [\AA$^2$/ps]')
292 setFigureSize(gcf, 300, 600);
293
294 ImproveFigureCompPhys(gcf,'LineColor', ...
295     {'r', 'MYRED', 'GERIBLUE','MYLIGHTBLUE'}, 'linewidth', 3);
296 saveas(1, '../figures/MSD-500.eps', 'epsc')
297 saveas(2, '../figures/MSD-700.eps', 'epsc')
298 saveas(10, '../figures/Phi-t.eps', 'epsc')
299 saveas(11, '../figures/P-freq.eps', 'epsc')

```

C.2 Improve figure appearance: ImproveFigureCompPhys.m

```

1 function ImproveFigureCompPhys(varargin)
2 %ImproveFigureCompPhys Improves the figures of supplied handles
3 % Input:
4 % - none (improve all figures) or handles to figures to improve
5 % - optional:
6 %     LineWidth int
7 %     LineStyle column vector cell, e.g. {'-','-','-'},
8 %     LineColor column vector cell, e.g. {'k',[0 1 1], 'MYBLUE'}
9 %             colors: MYBLUE,MYORANGE,MYGREEN,MYPURPLE, MYYELLOW,
10 %             MYLIGHTBLUE, MYRED
11 %     Marker column vector cell, e.g. {'.', 'o', 'x'}
12
13 % ImproveFigure was originally written by Adam Stahl, but has been heavily
14 % modified by Linnea Hesslow
15
16
17 %%% Handle inputs
18 % If no inputs or if the first argument is a string (a property rather than
19 % a handle), use all open figures
20 if nargin == 0 || ischar(varargin{1})
21     %Get all open figures
22     figHs = findobj('Type','figure');
23     nFigs = length(figHs);
24 else
25     % Check the supplied figure handles
26     figHs = varargin{1};
27     figHs = figHs(ishandle(figHs) == 1); %Keep only those handles that are ←
        proper graphics handles
28     nFigs = length(figHs);
29 end
30
31 % Define desired properties
32 titleSize = 24;
33 interpreter = 'latex';
34 lineWidth = 4;
35 axesWidth = 1.5;
36 labelSize = 22;

```

```

37 textSize = 20;
38 legTextSize = 18;
39 tickLabelSize = 18;
40 LineColor = {};
41 LineStyle = {};
42 Marker = {};
43
44 % define colors
45 co = [ 0      0.4470  0.7410
46       0.8500  0.3250  0.0980
47       0.9290  0.6940  0.1250
48       0.4940  0.1840  0.5560
49       0.4660  0.6740  0.1880
50       0.3010  0.7450  0.9330
51       0.6350  0.0780  0.1840 ];
52 colors = struct('MYBLUE', co(1,:),...
53               'MYORANGE', co(2,:),...
54               'MYYELLOW', co(3,:),...
55               'MYPURPLE', co(4,:),...
56               'MYGREEN', co(5,:),...
57               'MYLIGHTBLUE', co(6,:),...
58               'MYRED', co(7,:),...
59               'GERIBLUE', [0.3000  0.1500  0.7500],...
60               'GERIRED', [1.0000  0.2500  0.1500],...
61               'GERIYELLOW', [0.9000  0.7500  0.1000],...
62               'LIGHTGREEN', [0.4  0.85  0.4],...
63               'LINNEAGREEN', [7 184 4]/255);
64
65 % Loop through the supplied arguments and check for properties to set.
66 for i = 1:nargin
67     if ischar(varargin{i})
68         switch lower(varargin{i}) %Compare lower case strings
69             case 'linewidth'
70                 lineWidth = varargin{i+1};
71             case 'linestyle'
72                 LineStyle = varargin{i+1};
73             case 'linecolor'
74                 LineColor = varargin{i+1};
75                 for iLineColor = 1:numel(LineColor)
76                     if isfield(colors, LineColor{iLineColor})
77                         LineColor{iLineColor} = colors.(LineColor{iLineColor});
78                     end
79                 end
80             case 'marker'
81                 Marker = varargin{i+1};
82         end
83     end
84 end
85 %%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
86
87 %%% Improve the figure(s)
88
89 for iFig = 1:nFigs
90     fig = figHs(iFig);
91
92     lineObjects = findall(fig, 'Type', 'line');
93     textObjects = findall(fig, 'Type', 'text');
94     axesObjects = findall(fig, 'Type', 'axes');
95     legObjects = findall(fig, 'Type', 'legend');
96     contourObjects = findall(fig, 'Type', 'contour'); % not counted as lines
97
98     %%% TEXT APPEARANCE: first set all to textSize and then change the ones
99     %%% that need to be changed again
100
101     %Change size of any text objects in the plot
102     set(textObjects, 'FontSize', textSize);
103     set(legObjects, 'FontSize', legTextSize);
104
105     %%% FIX LINESTYLE, COLOR ETC. FOR EACH PLOT SEPARATELY
106     for iAx = 1:numel(axesObjects)
107         lineObjInAx = findall(axesObjects(iAx), 'Type', 'line');
108
109         %set line style and color style (only works if all figs have some
110         %number of line plots..)
111         if ~isempty(LineStyle)
112             set(lineObjInAx, {'LineStyle'}, LineStyle)
113             set(contourObjects, {'LineStyle'}, LineStyle); %%%%%%%%%
114         end
115         if ~isempty(LineColor)
116             set(lineObjInAx, {'Color'}, LineColor)
117             set(contourObjects, {'LineColor'}, LineColor); %%%%%%%%%
118         end
119         if ~isempty(Marker)
120             set(lineObjInAx, {'Marker'}, Marker)
121             set(lineObjInAx, {'Markersize'}, num2cell(10+22*strcmp(Marker, '.'))←
122                 )
123         end
124
125         %%% change font sizes.
126         % Tick label size

```



```

127     xLim = axesObjects(iAx).XLim;
128     axesObjects(iAx).FontSize = tickLabelSize;
129     axesObjects(iAx).XLim = xLim;
130     %Change label size
131     axesObjects(iAx).XLabel.FontSize = labelSize;
132     axesObjects(iAx).YLabel.FontSize = labelSize;
133
134     %Change title size
135     axesObjects(iAx).Title.FontSize = titleSize;
136 end
137
138 %%% LINE APPEARANCE
139 %Change line thicknesses
140 set(lineObjects,'LineWidth',lineWidth);
141 set(contourObjects, 'LineWidth', lineWidth);
142 set(axesObjects, 'LineWidth',axesWidth)
143
144 % set interpreter: latex or tex
145 set(textObjects, 'interpreter', interpreter)
146 set(legObjects, 'Interpreter', interpreter)
147 set(axesObjects, 'TickLabelInterpreter', interpreter);
148 end
149 end

```

C.3 Change size of figures: setFigureSize.m

```

1 function [ fig ] = setFigureSize( fig, H, W )
2 fig.Units = 'points';
3 fig.WindowStyle = 'normal'; % undock
4 fig.Position(3:4) = [W H];
5 end

```