THIS IS FULL TITLE

AUTHOR A¹ AND AUTHOR B^{2,*}

ABSTRACT. This is an example LATEX article. This can be used as a template for new articles. Abstracts must be able to stand alone and so cannot contain citations to the paper's references, equations, etc. An abstract must consist of a single paragraph and be concise. Because of online formatting, abstracts must appear as plain as possible. Any equations should be inline.

1. Introduction

The introduction introduces the context and summarizes the manuscript. It is important to clearly state the contributions of this piece of work.

In this paper we present a new method for solving the model equation

$$\begin{cases}
\partial_t u - \varepsilon^2 \Delta u + u^3 - u = 0, & \text{in } \Omega \times \mathcal{T}, \\
u(x, y, t) = g(t), & \text{on } \partial \Omega, \\
u(x, y, 0) = \varphi(x, y), & \text{on } \Omega.
\end{cases}$$
(1.1)

where ε is a small parameter.

This is an example of quoting an equation (1.1).

The merits of our method are as follows:

- item one
- item two
- item three
- (1) item one
- (2) item two
- (3) item three

The outline is not required, but we show an example here.

The paper is organized as follows. Our main results are in 3, our new algorithm is in 4, experimental results are in 5, discussion is in 6 and the conclusions follow in 7.

2. Preliminaries

2.1. This is subsection.

2.1.1. This is subsubsection.

Date: August 31, 2024.

 $2000\ Mathematics\ Subject\ Classification.\ 65M60,\ 65M12.$

Key words and phrases. Keyword 1, keyword 2, error analysis.

^{*}Corresponding author (E-mail: xyz@math.univ.edu).

 $^{^{1}\}mathrm{Department}$ of Computer Science, LATEX University.

 $^{^2 \}mbox{Department}$ of Mechanical Engineering, LATEX University.

This work is supported in part by NSFC grants No. 12345678 and \dots

3. Main results

We interleave text filler with some example theorems and theorem-like items.

Definition 3.1. This is a definition environment.

Lemma 3.1. This is a lemma environment.

Theorem 3.1. This is a theorem environment.

Proof. This is a proof environment.

Lemma 3.2. This is a lemma environment

- (i) item A
- (ii) item B

$$\lim_{n \to \infty} \left(1 + \frac{1}{n} \right)^n = e. \tag{3.1}$$

This is a citation example [1,5]. This statement requires citations [3–5].

Here we state our main result as 3.2.

Theorem 3.2 (LDL^T Factorization [2]). If $A \in \mathbb{R}^{n \times n}$ is symmetric and the principal submatrix A(1:k,1:k) is nonsingular for k=1:n-1, then there exists a unit lower triangular matrix L and a diagonal matrix

$$D = \operatorname{diag}(d_1, \dots, d_n),$$

such that $A = LDL^T$. The factorization is unique.

Theorem 3.3 (Mean Value Theorem). Suppose f is a function that is continuous on the closed interval [a,b]. and differentiable on the open interval (a,b). Then there exists a number c such that a < c < b and

$$f'(c) = \frac{f(b) - f(a)}{b - a}.$$

In other words,

$$f(b) - f(a) = f'(c)(b - a).$$

Remark 3.1. Observe that 3.2, 3.3 correctly mix references to multiple labels.

Corollary 3.1. Let f(x) be continuous and differentiable everywhere. If f(x) has at least two roots, then f'(x) must have at least one root.

Proof. Let a and b be two distinct roots of f. By 3.3, there exists a number c such that

$$f'(c) = \frac{f(b) - f(a)}{b - a} = \frac{0 - 0}{b - a} = 0.$$

Note that it may require two LATEX compilations for the proof marks to show.

Display matrices can be rendered using environments from amsmath:

$$S = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad C = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \tag{3.2}$$

Equation 3.2 shows some example matrices.

We calculate the Fréchet derivative of F as follows:

$$F'(U,V)(H,K) = \langle R(U,V), H\Sigma V^T + U\Sigma K^T - P(H\Sigma V^T + U\Sigma K^T) \rangle$$
(3.3a)

SHORT TITLE 3

$$= \langle R(U, V), H\Sigma V^T + U\Sigma K^T \rangle$$

= $\langle R(U, V)V\Sigma^T, H \rangle + \langle \Sigma^T U^T R(U, V), K^T \rangle.$ (3.3b)

3.3a is the first line, and 3.3b is the last line.

4. Algorithm

Our analysis leads to the algorithm in 1.

Algorithm 1 Build tree

```
Define P := T := \{\{1\}, \dots, \{d\}\}

while \#P > 1 do

Choose C' \in \mathcal{C}_p(P) with C' := \operatorname{argmin}_{C \in \mathcal{C}_p(P)} \varrho(C)

Find an optimal partition tree T_{C'}

Update P := (P \setminus C') \cup \{\bigcup_{t \in C'} t\}

Update T := T \cup \{\bigcup_{t \in \tau} t : \tau \in T_{C'} \setminus \mathcal{L}(T_{C'})\}

end while

return T
```

Adjust the width of the algorithm environment

```
Algorithm 2 Euclid's algorithm
                                                                                    \triangleright The g.c.d. of a and b
 1: procedure Euclid(a, b)
         r \leftarrow a \bmod b
         while r \neq 0 do
                                                                           \triangleright We have the answer if r is 0
 3:
             a \leftarrow b
 4:
             b \leftarrow r
 5:
 6:
             r \leftarrow a \bmod b
         end while
 7:
                                                                                               \triangleright The gcd is b
         return b
 9: end procedure
```

5. Experimental results

Some experimental results here.

Example 1. This is example environment.



FIGURE 5.1. This is an example of a figure caption with text.

Some figures and tables here.

Figure 5.2 shows some example results.




FIGURE 5.2. Example figure using external image files.

The two figures are placed side by side, sharing one title, as shown in Figure 5.3.




FIGURE 5.3. Left: Caption 1, Right: Caption 2.

Use the tabular environment to generate Table 5.1. The commands \tabcolsep and \tabcolsep control the spacing between columns and the vertical space between rows, respectively.


Date Location **Event Name** Remarks New York 2023-02Academic Symposium Topic: Artificial Intelligence 2023-05 London Academic Exchange Focus: Mathematical Modeling 2023-09 Shanghai Technology Exhibition Display of New Chips 2024-02 Tokyo Research Workshop Subject: Quantum Computing 2024-06 Paris International Conference Theme: Climate Change

Table 5.1. Event Schedule

Some description and explanation of pictures and tables.

SHORT TITLE 5

Use minipage package to set images side-by-side, each with its own title, as shown in Figure 5.4 and Figure 5.5.



Use subfig package to set subfigure, each with its subcaption, as shown in Figure 5.6.




FIGURE 5.6. Two subfigures

Use the tabularx environment to generate Table 5.2.

Table 5.2. Height and weight sample of students from a school

Serial No.	Age	Height (cm)	Weight (kg)
001	15	156	42
002	16	158	45
003	14	162	48
004	15	163	50
Average	15	159.75	46.25

Additional results are available in the supplement in Table 5.3.

degree	step-size h	L^2 -errors	order	H^1 -errors	order	L^{∞} -errors	order
1	1/128	9.18E-06	2.02	7.70E-03	1.01	6.46E-07	2.02
	1/256	2.29E-06	2.01	1.92E-03	1.00	1.61E-07	2.01
	1/512	5.70E-07	2.00	9.56E-04	1.00	4.01E-08	2.00
2	1/128	1.39E-08	3.01	1.15E-05	2.01	3.48E-12	4.02
	1/256	1.73E-09	3.01	2.88E-06	2.01	3.27E-13	3.94
	1/512	2.17E-10	3.00	7.24E-06	2.00	6.66E-13	1.55
3	1/32	2.28E-09	4.05	6.92E-07	3.04	1.45E-15	8.21
	1/64	1.42E-10	4.03	8.65E-08	3.02	2.06E-14	3.85
	1/128	8.91E-12	4.01	1.08E-08	3.01	3.86E-14	0.91

Table 5.3. Numerical error

6. Discussion of $Z = X \cup Y$

Some discussions here. Some discussions here.

7. Conclusions

Some conclusions here. Some conclusions here.

APPENDIX A. AN EXAMPLE APPENDIX

The contents of the appendix are here.

Lemma A.1. Test Lemma.

This is a equation in appendix.

$$a^2 + b^2 = c^2. (A.1)$$

ACKNOWLEDGMENTS

We would like to acknowledge the assistance of volunteers in putting together this example manuscript and supplement.

References

- [1] Robert A. Adams and John J. F. Fournier. Sobolev spaces. Elsevier, Amsterdam, 2nd edition, 2003.
- [2] Gene H. Golub and Charles F. Van Loan. Matrix computations. The Johns Hopkins University Press, Baltimore, 4th edition, 2013.
- [3] Jie Shen. Efficient spectral-Galerkin method I. Direct solvers of second- and fourth-order equations using Legendre polynomials. SIAM J. Sci. Comput., 15(6):1489–1505, 1994.
- [4] Eitan Tadmor. A review of numerical methods for nonlinear partial differential equations. *Bull. Amer. Math. Soc.*, 49(4):507–554, 2012.
- [5] L. N. Trefethen and J. A. C. Weideman. The exponentially convergent trapezoidal rule. SIAM Rev., 56(3):385–458, 2014.