# Assignment #1

Course: Numerical methods of differential equation Name: Student name Major: XXX Student ID: 123002584

Date: April 2, 2023

**Problem 1.** Here is a question.

**Answer.** Here is the answer of the question.

**Problem 2.** Here is a proof question.

**Proof.** Here is the proof of the problem.

**Definition 1.** This is a definition.

**Proposition 1.** This is a proposition.

Lemma 1 (Lemma). This is a lemma.

**Theorem 1** (Theorem). This is a Theorem.

**Proof.** This is the proof environment.

Corollary 1. This is a corollary.

**Proposition 2** (Proposition). This is a proposition.

Remark 1. This is a remark.

Example 1. This is an example.

#### **Table**

This is an example of table, such as Table 1

Table	1	$T_2h$	اما	name
таше	- 1	12(1)	œ	паше

A	N = 3	N = 5	N = 7	N = 9	N = 11	N = 13
В	1.5789	1.3478	1.0645	0.8780	0.7222	0.5942
$\mathbf{C}$	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
D	7.2632	14.3913	21.0323	27.3171	30.9630	34.0870

## Figures

This is an example of figure, as shown in the Figure 2.




Figure 1 Image of function  $y = \sin(x)$ 

Two pictures placed side by side, as shown in Figure 2 and Figure 3.




Figure 2 Description of image 1




Figure 3 Description of image 2

### Code highlighting

This is MATLAB code highlighting environment

#### MATLAB code

```
1\, % Euler method for the ODE model
u'(x) = x^2 + x - u, x in [0.1]
3 % Initial condition: u(0) = 0.
4 clear all; clf
5 h = 0.1;
6 x=0:h:1;
7 \text{ N=length(x)-1};
8 u(1)=0;
9 fun=@(t,u) t.^2+t-u;
10 for n=1:N
       u(n+1)=u(n)+h.*fun(x(n),u(n));
11
13 ue=-exp(-x)+x.^2-x+1; % exact solution
14 plot(x,ue,'b-',x,u,'r+','LineWidth',1)
15 legend('Exact', 'Numerical', 'location', 'North')
16 xlabel('x'), ylabel('u')
```

This is Python code highlighting environment

#### Python code

```
1 #PythonDraw.py
2 import turtle as t
3 t.setup(650, 350, 200, 200)
4 t.penup()
5 t.fd(-250)
6 t.pendown()
7 t.pensize(25)
8 t.pencolor("purple color")
9 t.seth(-40)
10 for i in range (4):
11
       t.circle(40, 80)
       t.circle(-40, 80)
12
13 t.circle(40, 80/2)
14 t.fd(40)
15 t.circle(16, 180)
16 \text{ t.fd}(40 * 2/3)
17 t.done()
```

## References

- [1] E. Tadmor. A review of numerical methods for nonlinear partial differential equations. Bull. Amer. Math. Soc., 2012, 49(4): 507-554.
- [2] R. A. Adams, Fournier J J F. Sobolev spaces. Elsevier, 2003.
- [3] L. N. Trefethen, J. A. C. Weideman. The exponentially convergent trapezoidal rule. SIAM Rev., 2014, 56(3): 385-458.
- [4] J. Shen. Efficient spectral-Galerkin method I. Direct solvers of second- and fourth-order equations using Legendre polynomials. SIAM J. Sci. Comput., 1994, 15(6): 1489-1505.