


Lambda-Calculus (cont):

Fixpoints, Naming

Lecture 10

CS 565



133

Recursion and Divergence



Consider the application:

$$\Omega \equiv ((\lambda x. (x x)) (\lambda x. (x x)))$$

Ω evaluates to itself in one step.

It has no normal form.

A lambda term is in normal form if it does not contain any redex (i.e., a term that is subject to β -reduction)

Now, consider: $Y \equiv ((\lambda x. (f (x x))) (\lambda x. (f (x x))))$

$Y \rightarrow$

$$(f ((\lambda x. (f (x x))) (\lambda x. (f (x x)))) \rightarrow$$


$$(f (f (\lambda x. (f (x x))) (\lambda x. (f (x x)))) \rightarrow$$

...

$$(f (f (\dots (f (\lambda x. (f (x x))) (\lambda x. (f (x x))) \dots)))$$

134

Normal forms and order of evaluation




No expression can be converted to two distinct normal forms (Church-Rosser Theorem 1)

Is there an order of evaluation guaranteed to terminate whenever a particular expression is reducible to normal form?

- Normal-order: leftmost, outermost reduction: no expression in the argument position of a redex is reduced until the redex is reduced
- If there is a reduction from A to B and B is in normal form, then there exists a normal order reduction from A to B (Church-Rosser Theorem 2)

135

Recursion



The previous definition applies f an infinite number of times

Basis for iterated application

But, how can we slow its rate of unfolding?

Consider:

$$\Omega v \equiv (\lambda y. ((\lambda x. (\lambda y. (x x y))) \\ (\lambda x. (\lambda y. (x x y))) \\ y))$$

Ωv is in normal form. However, if it is applied to an argument it diverges.

136

Recursion (cont)



$(\Omega v \ v) \rightarrow$

$((\lambda y. ((\lambda x. (\lambda y. (x \ x \ y)))$
 $(\lambda x. (\lambda y. (x \ x \ y)))$
 $y) \ v) \rightarrow$

$((\lambda y. ((\lambda x. (\lambda y. (x \ x \ y)))$
 $(\lambda x. (\lambda y. (x \ x \ y)))$
 $y) \ v) \rightarrow$
...

137

Recursion (cont)



Now, consider

$Zf \equiv (\lambda y. ((\lambda x. (f (\lambda y. (x \ x \ y)))$
 $(\lambda x. (f (\lambda y. (x \ x \ y)))$
 $y))$


If we apply Zf to an argument:

$((\lambda y. ((\lambda x. (f (\lambda y. (x \ x \ y)))$
 $(\lambda x. (f (\lambda y. (x \ x \ y)))$
 $y) \ v) \rightarrow$
 $(f (\lambda y. ((\lambda x. (f (\lambda y. (x \ x \ y)))$
 $(\lambda x. (f (\lambda y. (x \ x \ y)))$
 $y)) \ v) \rightarrow$

Since the arguments to f are all values, this expression is equivalent to: $f \ Zf \ v$

138

Recursion (cont)



How do we apply these insights?

```
f ≡ λ fact.  
    λ n. if n = 0  
        then 1  
        else n * (fact (n - 1))
```

We can use Zf to turn f into a real factorial function:

139

Fixpoints




```
Zf 3 →  
f Zf 3 →  
(λ fact. λ n. ...) Zf 3 →  
if 3 = 0 then 1 else 3 * (Zf 2) →  
3 * (f Zf 2)  
...
```

We'll stop when $n = 0$

140

Fixpoints



Define $z = \lambda f. z_f$

Now, z defines a fixpoint for any f :

$$z \equiv \lambda f. (\lambda y. ((\lambda x. (f (\lambda y. (x x y)))) \\ (\lambda x. (f (\lambda y. (x x y)))) \\ y))$$

z computes the least fixpoint of a function.

141

Fixpoints and order of evaluation



Consider an alternative definition:

$$Y \equiv \lambda h. (\lambda x. h(x x)) (\lambda x. h(x x))$$


- What happens if we apply Y to f (the factorial functional) with argument 3?
- Under normal-order evaluation:

$$Y f 3 \equiv (\lambda x. f(x x)) (\lambda x. f(x x)) 3 \rightarrow \\ f ((\lambda x. f(x x)) (\lambda x. f(x x))) 3$$

- What happens under applicative-order?

142

Naming and substitution



Although we claimed that lambda calculus essentially manipulates functions (it does), we've spent a lot of time thinking about variables

- substitutions
- free variables
- equivalence upto renaming

Implementations must consider these issues seriously

- Rename bound variables when performing substitutions with “fresh” names.
- Impose a condition that all bound variables be distinct from each other, and other free variables.
- Derive a canonical representation that does not require renaming at all.

143

Terms and Contexts



De Bruijn indices:

- Have variable occurrences “point” directly to their binders rather than referring to them by name.
- Do so by replacing variable occurrences with numbers:
number k stands for “the variable bound by the k th enclosing λ -term

Example: $\lambda x. \lambda y. x (y x) \equiv \lambda. \lambda. 1 (0 \ 1)$

Similar to static offsets in an activation record or display.

144

Examples



$\text{identity} \equiv \lambda x. x \equiv \lambda . 0$

$\text{true} \equiv \lambda x. \lambda y. x \equiv \lambda . \lambda . 1$

$\text{false} \equiv \lambda x. \lambda y. y \equiv \lambda . \lambda . 0$

$\text{two} \equiv \lambda s. \lambda z. s (s z) \equiv \lambda . \lambda . (1 (1 0))$

145

Contexts



How do we replace free variables with their binders?

Assume an ordered context listing all free variables that can occur, and map free variables to their index in this context (counting right to left)

Context: a, b

$a \mapsto 1, b \mapsto 0$


$\lambda x. a \equiv \lambda . 2$

$\lambda x. b \equiv \lambda . 1$

$\lambda x. b (\lambda y. a) \equiv \lambda . 1 (\lambda . 3)$

146

Shifting and substitution



When substituting into a λ term, indices must be adjusted:

$\lambda y. x [z/x]$ in context x, y, z

$[2 \mapsto 0] \lambda. 2 \equiv \lambda. [3 \mapsto 1] 3 \equiv \lambda. 1$

Key point: context becomes longer when substituting inside an abstraction. Need to be careful to adjust free variables, not bound ones.

$\text{shift}(d, c)(k) = k$ if $k < c$

$k + d$ if $k \geq c$


$\text{shift}(d, c)(\lambda. t) = (\lambda. \text{shift}(d, c+1)(t))$

$\text{shift}(d, c)(t \ t') =$

$(\text{shift}(d, c)(t)) \ (\text{shift}(d, c)(t'))$

147

Example



$\text{shift}(2, 0)(\lambda. \lambda. 1 \ (0 \ 2))$

$\lambda. \lambda. 1 \ (0 \ 4)$

$\text{shift}(2, 0)(\lambda. 0 \ 1 \ (\lambda. 0 \ 1 \ 2))$

$\lambda. 0 \ 3 \ (\lambda. 0 \ 1 \ 4)$

148

Substitution



$$[j \mapsto s] k = s \text{ if } k = j \\ k \text{ otherwise}$$

$$[j \mapsto s](\lambda . t) = \lambda . [j+1 \mapsto \text{shift}(1,0)s] t$$

$$[j \mapsto s](t1 \ t2) = ([j \mapsto s] t1) ([j \mapsto s] t2)$$

Beta-reduction:

$$(\lambda t) v \rightarrow \text{shift}(-1,0)([0 \mapsto \text{shift}(1,0)(v)] t)$$

149

Examples



Assume context $\langle a, b \rangle$


► Then, $a \mapsto 1, b \mapsto 0$

$$[a / b] b \ \lambda x. \ \lambda y. \ b \\ [0 \mapsto 1] \ 0 \ \lambda . \ \lambda . \ 2 \\ 1 \ \lambda . \ \lambda . \ 3 \equiv a \ \lambda x. \ \lambda y. \ a$$

$$[(a \ (\lambda z. \ a)) / b] \ (b \ (\lambda x. \ b)) \\ [0 \mapsto (1 \ (\lambda . \ 2))] \ (0 \ \lambda . \ 1) \\ 1 \ (\lambda . \ 2) \ (\lambda . \ (2 \ (\lambda . \ 3))) \\ (a \ (\lambda z. \ a)) \ (\lambda x. \ (a \ (\lambda z. \ a)))$$

150

Examples



$[a/b] (\lambda b. (b a))$
 $[0 \mapsto 1] (\lambda . (0 2))$
 $(\lambda . (0 2))$
 $(\lambda b. (b a))$
 $[a/b] (\lambda a. (b a))$
 $[0 \mapsto 1] \lambda . (1 0)$
 $\lambda . (2 0)$
 $(\lambda a'. a a')$