

## Announcements:

Final exam: Tues. 5/7 8:00am-11:00am,  
1047 Sidney Lu Mech. E. Bldg.

(email ASAP w/ any issues)

Exam will be cumulative

## Schedule:

Today, Monday: projective space

Tuesday: problem session

Wednesday: review

Should we have another review session? When?

HW10 posted (due Wed. 4/31)

Midterm 3 graded

Q1: 68%

Median: 53 / 80

Q2: 19%

Mean: 54.9 / 80

Q3: 60%

Std. dev: 18.1

Q4: 68%

Gradelines:

A- / A: to

B+ / B / B-: to -E

C+ / C / C-: to -E

D+ / D / D-: to -E

Solns posted to website


"Where do I stand" spreadsheet updated

# Projective space (see Cox, Little, O'Shea: Ideals, Varieties, and Algorithms, Ch 8.)

Motivation: recall

Bézout's Thm: The "usual" situation is that two poly. in  $\mathbb{C}[x,y]$  of degrees  $m$  and  $n$  have  $m \cdot n$  intersection points in  $\mathbb{C}$

But what about parallel lines?



$(\deg L)(\deg M) = 1 \cdot 1 = 1$ , but  $L$  and  $M$  don't intersect

Fix: add pts. "at  $\infty$ " where parallel lines meet

Consider equiv. classes of parallel lines

Def (version 1): The (complex) projective plane is the set

$\nearrow \widetilde{\mathbb{P}^2(\mathbb{C})} = \mathbb{C}^2 \cup \{\text{one pt "at } \infty" \text{ for each equiv. class of parallel lines}\}$

for now, to distinguish from def. 2 class of parallel lines }  $H_\infty$

Works, but kind of a weird def'n

For a nicer one, let's define homogenous coords. in  $\mathbb{C}^3$

We say that  $\underbrace{(a_0, a_1, a_2)}_{\in \mathbb{C}^3} \sim (b_0, b_1, b_2)$

if  $(b_0, b_1, b_2) = (\lambda a_0, \lambda a_1, \lambda a_2)$  for some  $\lambda \in \mathbb{C} \setminus \{0\}$

i.e. if all the ratios are the same:  $\frac{a_0}{a_1} = \frac{b_0}{b_1}, \frac{a_0}{a_2} = \frac{b_0}{b_2}, \frac{a_1}{a_2} = \frac{b_1}{b_2}$

i.e. if  $a, b \neq 0, a \sim b \iff a$  and  $b$  are on the same line thru. origin in  $\mathbb{C}^3$

Denote equiv. classes  $[a_0 : a_1 : a_2]$

Def (version 2): The complex proj. plane is the set of equivalence classes

$$\mathbb{P}^2(\mathbb{C}) = (\mathbb{C}^3 \setminus \{0\}) / \sim$$

i.e. the set of 1D subspaces of  $\mathbb{C}^3$

Prop: There is a (nice) bijection

$$\begin{array}{ccc} \mathbb{P}^2(\mathbb{C}) & \longrightarrow & \widetilde{\mathbb{P}^2(\mathbb{C})} \\ \text{def 2} & & \text{def 1} \end{array}$$

$$\text{Pf: } \mathbb{P}^2(\mathbb{C}) = \underbrace{\{[1:x:y] \mid x, y \in \mathbb{C}\}}_{S_1} \cup \underbrace{\{[0:1:y] \mid y \in \mathbb{C}\}}_{S_2} \cup \underbrace{\{[0:0:1]\}}_{S_3}$$

$$[1:x:y] \mapsto (x, y) \text{ is a bij. } S_1 \rightarrow \mathbb{C}^2$$

Let  $a_m \in H_\infty$ ,  $m \in \mathbb{C} \cup \{\infty\}$  be the equiv. class of lines in  $\mathbb{C}^2$  of slope  $m$

$$\text{Then } [0:1:m] \mapsto a_m$$

$$[0:0:1] \mapsto a_\infty$$

gives a bijection  $S_2 \cup S_3 \rightarrow H_\infty$

□

Def: (complex) projective space is the set

$$\mathbb{P}^n(\mathbb{C}) = \{\text{lines thru. origin in } \mathbb{C}^{n+1}\}$$

$$= \{a = (a_0, \dots, a_{n+1}) \in \mathbb{C}^{n+1} \setminus \{0\}\} / (a \sim \lambda a, \lambda \in \mathbb{C})$$

$$= \{[a_0 : \dots : a_n]\}$$

$$\text{Cor: } \mathbb{P}^n(\mathbb{C}) = \mathbb{C}^n \cup \mathbb{P}^{n-1}(\mathbb{C})$$

Pf: Use the maps from the previous prop:


$$[1 : a_1 : \dots : a_n] \mapsto (a_1, \dots, a_n) \in \mathbb{C}^n$$

$$[0 : a_1 : \dots : a_n] \mapsto \underbrace{[a_1 : \dots : a_n]}_{\text{not all 0}} \in \mathbb{P}^{n-1}(\mathbb{C})$$

□

$$\text{Ex: } \mathbb{P}^1(\mathbb{C}) = \{\text{lines in } \mathbb{C}^2\} = \{[x : y]\}$$

$$= \{[1 : m] \mid m \in \mathbb{C}\} \cup \{[0 : 1]\}$$



Also called the Riemann sphere

Want to define projective varieties in  $\mathbb{P}^n(\mathbb{C})$

$$\text{Let } f(x, y, z) = xy - z$$

$$\text{Then } f(1, 1, 1) = 0$$

$$f(2, 2, 2) = 2$$

So what does  $f([1:1:1])$  mean?

Problem: When we scaled the variables, we doubled  $z$  but quadrupled  $xy$

Fix:

Def:  $f(x_0, \dots, x_n) \in \mathbb{C}^{n+1}$  is homogeneous of degree  $d$  if every term has degree  $d$

If  $f$  homog. of degree  $d$

$$f(\lambda a_0, \dots, \lambda a_n) = \lambda^d f(a_0, \dots, a_n)$$

$$\text{If } \lambda \neq 0, f(\lambda a_0, \dots, \lambda a_n) = 0 \iff f(a_0, \dots, a_n) = 0$$

Def: If  $f \in \mathbb{C}[x_0, \dots, x_n]$  homog.,

$$V(f) := \{[a_0 : \dots : a_n] \in \mathbb{P}^n(\mathbb{C}) \mid f(a_0, \dots, a_n) = 0\}$$

is the projective variety assoc. to  $f$ .

Next time:  $V(I)$  for "homog. ideal"  $I$