

Announcements

HW1 posted

due Wed. 2/11 @ 9:00am via Gradescope

make sure you've joined the Gradescope course!

course code: email me

Last time: Schur's Lemma & Maschke's Thm

Today: reps of abelian gps.

intro to character theory [F-H §2.1, Serre §2.1]

Proposition 6: If G is abelian, then every G -irrep. is one-dimensional.

PF: First we claim that $\rho(g) \in \text{End}_G(V) \ \forall g \in G$.

Indeed, if $h \in G$, $v \in V$, then since G is abelian,

$$\rho(g)\rho(h)v = \rho(h)\rho(g)v,$$

so $\rho(g)$ is G -equivariant.

By Schur's Lemma, every $\rho(g)$ is a scalar mult. of the identity. This means that every subspace of V is G -invariant, so V must be 1-dim. \square

$$\text{Ex: } G = \mathbb{Z}/n\mathbb{Z} = \langle g \rangle.$$

Irreps of G are of the form

$$\rho_g: G \rightarrow \mathbb{C}^*$$

g : n th root of 1

$$g^a = g^a$$

So there are exactly n nonisom. irreps. of G !

$G = \mathbb{Z}/4\mathbb{Z}$	$\rho(1)$	$\rho(g)$	$\rho(g^2)$	$\rho(g^3)$
ρ_1	1	1	1	1
ρ_i	1	i	-1	$-i$
ρ_{-1}	1	-1	1	-1
ρ_{-i}	1	$-i$	-1	i

Very nice looking table! (Square, roots of unity,
orthog. rows/cols.)

And tells us everything we'd want to know
about the repn theory of G

Goal: get as close as we can to
this for all finite gps.

Def 7: The character of a repn. (ρ, V) is the function $\chi_V: G \rightarrow \mathbb{C}$ given by the trace of ρ :

$$\chi_V(g) = \text{Tr}(\rho(g)).$$

Prop 8:

a) The character is a class function:

$$\chi_V(hgh^{-1}) = \chi_V(g)$$

$$b) \chi_V(1) = \dim V$$

$$c) \chi_{V \oplus W} = \chi_V + \chi_W$$

$$d) \chi_{V \otimes W} = \chi_V \chi_W$$

$$e) \chi_{V^*} = \overline{\chi_V}$$

$$f) \chi_{\text{sym}^2 V}(g) = \frac{1}{2} [\chi_V(g)^2 + \chi_V(g^2)]$$

$$g) \chi_{\wedge^2 V}(g) = \frac{1}{2} [\chi_V(g)^2 - \chi_V(g^2)]$$

Pf: a) Trace is a class function

$$b) \rho(1) = \text{Id}_V$$

c), d), e) Let $g \in G$, and let $\lambda_1, \dots, \lambda_k$ be the e-values of $\rho_V(g)$, and μ_1, \dots, μ_l be the e-values of $\rho_W(g)$. Then,

c) The e-values of $\rho_{V \oplus W}(g) = \rho_V(g) \oplus \rho_W(g)$ are $\lambda_1, \dots, \lambda_k, \mu_1, \dots, \mu_l$

d) The e-values of $\rho_{V \otimes W}(g) = \rho_V(g) \otimes \rho_W(g)$ are $\lambda_i \mu_j$, $1 \leq i \leq k, 1 \leq j \leq l$

e) The e-values of $\rho_{V^*}(g) = \rho_V(g^{-1})^T$ are $\lambda_1^{-1}, \dots, \lambda_k^{-1}$. Since $\rho_V(g)$ has finite order, its e-values are roots of unity, so $\lambda_i^{-1} = \overline{\lambda_i}$, and so

$$\chi_{V^*}(g) = \overline{\lambda_1} + \dots + \overline{\lambda_k} = \overline{\chi_V(g)}$$

f), g) See HW1

□

We collect the characters of irreps. of G into the character table

- Rows indexed by irreps.
- Cols indexed by conjugacy classes in G
- Entries are the character value for the irrep. at (any elt. of) the conj. class.

Ex: Recall from Lecture 1 the three irreps of S_3 :
(we'll see soon that there are no more)

$$\rho_{\text{triv}}: w \mapsto [1]$$

$$\rho_{\text{sgn}}: w \mapsto [(-1)^w]$$

$$\rho_{\text{ref}}: (1) \mapsto \begin{bmatrix} 1 & \\ & 1 \end{bmatrix} \quad (23) \mapsto \begin{bmatrix} 1 & -1 \\ & -1 \end{bmatrix}$$

$$(12) \mapsto \begin{bmatrix} & 1 \\ 1 & \end{bmatrix} \quad (123) \mapsto \begin{bmatrix} & -1 \\ 1 & -1 \end{bmatrix}$$

$$(13) \mapsto \begin{bmatrix} -1 & \\ -1 & 1 \end{bmatrix} \quad (132) \mapsto \begin{bmatrix} -1 & 1 \\ & -1 \end{bmatrix}$$

Conj. classes are det'd by cycle type

So the character table is:

S_3	1elt ()	3elts (12)	2elts (123)
χ_{triv}	1	1	1
χ_{sgn}	1	-1	1
χ_{ref}	2	0	-1

Class activity: Compute the characters of the following reps? Can they be written as sums of irred. chars.?

a) $\rho_{\text{triv}} \oplus \rho_{\text{sgn}}$

b) ρ_{ref}^*

c) $\rho_{\text{ref}}^{\otimes 2}$

d) $\text{Sym}^2 \rho_{\text{ref}}$

e) $\wedge^2 \rho_{\text{ref}}$

f) The regular repn $V_{\text{reg}} = \langle v_g \mid g \in S_3 \rangle$

w/ the action $\omega \cdot v_u := v_{\omega u}$
 $\omega \in S_3$