

A factorial analogue of the boson-fermion correspondence

Andrew Hardt

University of Illinois Urbana-Champaign

Misra Birthday Conference, October 2024

arXiv:2410.06582 (double factorial); in preparation (factorial, algebraic proofs)

Joint with Daniel Bump and Travis Scrimshaw

Some (brief) background

Isomorphism between two representations of the infinite Heisenberg algebra, the *fermionic* and *bosonic* Fock spaces

Some (brief) background

Isomorphism between two representations of the infinite Heisenberg algebra, the *fermionic* and *bosonic* Fock spaces

Connections to KP hierarchy of differential equations, symmetric functions, integrable systems, vertex algebras, and representation theory of affine Lie algebras

Some (brief) background

Isomorphism between two representations of the infinite Heisenberg algebra, the *fermionic* and *bosonic* Fock spaces

Connections to KP hierarchy of differential equations, symmetric functions, integrable systems, vertex algebras, and representation theory of affine Lie algebras

- Skyrme
- Kyoto school: Sato, Jimbo, Miwa, Date, Kashiwara etc.
- Frenkel, Kac, Peterson, etc.

Some (brief) background

Isomorphism between two representations of the infinite Heisenberg algebra, the *fermionic* and *bosonic* Fock spaces

Connections to KP hierarchy of differential equations, symmetric functions, integrable systems, vertex algebras, and representation theory of affine Lie algebras


- Skyrme
- Kyoto school: Sato, Jimbo, Miwa, Date, Kashiwara etc.
- Frenkel, Kac, Peterson, etc.

Various generalizations:

- Jing (Hall-Littlewood)
- Kashiwara-Miwa-Stern (quantum Fock space)


The classical boson-fermion correspondence

Fermionic Fock space



$$|\lambda\rangle = \dots \quad \begin{array}{ccccccccc} \bullet & \bullet & \circ & \bullet & \circ & | & \circ & \bullet & \circ & \bullet & \circ \\ -4 & -3 & -2 & -1 & 0 & & 1 & 2 & 3 & 4 & 5 \end{array} \quad \dots$$

Fermionic Fock space




$$|\lambda\rangle = \cdots \quad \bullet \quad \bullet \quad \circ \quad \bullet \quad \circ \quad | \quad \circ \quad \bullet \quad \circ \quad \bullet \quad \circ \quad \cdots$$

-4 -3 -2 -1 0 1 2 3 4 5

$$\mathfrak{F}^{(0)} := \mathbb{C}\text{-span of } \{|\lambda\rangle\},$$

Fermionic Fock space



$$|\lambda\rangle = \cdots \quad \bullet \quad \bullet \quad \circ \quad \bullet \quad \circ \quad | \quad \circ \quad \bullet \quad \circ \quad \bullet \quad \circ \quad \cdots$$

-4 -3 -2 -1 0 1 2 3 4 5

$$\mathfrak{F}^{(0)} := \mathbb{C}\text{-span of } \{|\lambda\rangle\}, \quad \mathfrak{F} = \bigoplus_{m \in \mathbb{Z}} \mathfrak{F}^{(m)}$$

$$\mathfrak{F}^{(m)} := \mathbb{C}\text{-span of } \{\Sigma^m |\lambda\rangle\}, \text{ where } \Sigma \text{ shifts all particles one unit right}$$

Fermionic Fock space

ψ_i : *creation* at position i ,

$$\begin{array}{c} - \quad \dots \quad \bullet \quad \bullet \quad \circ \quad \bullet \quad \bullet \quad | \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \dots \quad \in \mathfrak{F}^{(0)} \\ \psi_{-2} \downarrow \\ \dots \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad | \quad \circ \quad \circ \quad \circ \quad \bullet \quad \circ \quad \dots \quad \in \mathfrak{F}^{(1)} \end{array}$$

Fermionic Fock space

ψ_i : *creation* at position i , ψ_i^* : *deletion* at position i

— ... ● ● ○ ● ● | ○ ○ ○ ● ○ ... $\in \mathfrak{F}^{(0)}$

$\psi_{-2} \downarrow$


... ● ● ● ● ● | ○ ○ ○ ● ○ ... $\in \mathfrak{F}^{(1)}$

$\psi_3^* \downarrow$

... ● ● ● ● ● | ○ ○ ○ ○ ○ ... $\in \mathfrak{F}^{(0)}$

Fermionic Fock space

ψ_i : *creation* at position i , ψ_i^* : *deletion* at position i




These operators form a Clifford algebra:

$$\psi_i \psi_j + \psi_j \psi_i = \psi_i^* \psi_j^* + \psi_j^* \psi_i^* = 0, \quad \psi_i \psi_j^* + \psi_j^* \psi_i = \delta_{ij}$$

Fermionic Fock space

ψ_i : *creation* at position i , ψ_i^* : *deletion* at position i



These operators form a Clifford algebra:

$$\psi_i \psi_j + \psi_j \psi_i = \psi_i^* \psi_j^* + \psi_j^* \psi_i^* = 0, \quad \psi_i \psi_j^* + \psi_j^* \psi_i = \delta_{ij}$$

Fermion fields:

$$\psi(z) = \sum_{i \in \mathbb{Z}} \psi_i z^i, \quad \psi^*(w) = \sum_{j \in \mathbb{Z}} \psi_j^* w^{-j},$$

\mathfrak{gl}_∞ -representation

$$\mathfrak{gl}_\infty = \left\{ \sum_{i,j} a_{ij} E_{ij} \mid a_{ij} = 0 \text{ for } |i-j| \gg 0 \right\}$$

$\mathfrak{gl}_\infty = \left\{ \sum_{i,j} a_{ij} E_{ij} \mid a_{ij} = 0 \text{ for } |i-j| \gg 0 \right\}$ acts on $\mathfrak{F}^{(m)}$ via

$$E_{ij} := i \begin{bmatrix} & j \\ & 1 \\ & \end{bmatrix} \mapsto \psi_i \psi_j^*$$

$\mathfrak{gl}_\infty = \left\{ \sum_{i,j} a_{ij} E_{ij} \mid a_{ij} = 0 \text{ for } |i - j| \gg 0 \right\}$ acts on $\mathfrak{F}^{(m)}$ via

$$E_{ij} := i \begin{bmatrix} & j \\ & 1 \\ & \end{bmatrix} \mapsto \psi_i \psi_j^*$$

The *current operators* $J_k := \sum_{i \in \mathbb{Z}} : \psi_{i-k} \psi_i^* :$ form a Heisenberg algebra:

$$[J_k, J_\ell] = k \delta_{k, -\ell} \cdot 1$$

\mathfrak{gl}_∞ -representation

$\mathfrak{gl}_\infty = \left\{ \sum_{i,j} a_{ij} E_{ij} \mid a_{ij} = 0 \text{ for } |i-j| \gg 0 \right\}$ acts on $\mathfrak{F}^{(m)}$ via

$$E_{ij} := i \begin{bmatrix} & j \\ & 1 \\ & \end{bmatrix} \mapsto \psi_i \psi_j^*$$

The *current operators* $J_k := \sum_{i \in \mathbb{Z}} : \psi_{i-k} \psi_i^* :$ form a Heisenberg algebra:

$$[J_k, J_\ell] = k \delta_{k, -\ell} \cdot 1$$

Half-vertex operator:

$$e^{H_\pm(\mathbf{p})} := \exp \left(\sum_{k \geq 1} \frac{1}{k} p_k J_{\pm k} \right) \quad \left(\text{Think: } p_k = \sum_{i \geq 1} x_i^k \right)$$

Bosonic Fock Space

Bosonic Fock space:

$$B^{(m)} = \mathbb{C}[p_1, p_2, \dots] \cdot s^m, \quad B = \bigoplus_{m \in \mathbb{Z}} B^{(m)}$$

(Think: symmetric functions in x_1, x_2, \dots)

Bosonic Fock Space

Bosonic Fock space:

$$B^{(m)} = \mathbb{C}[p_1, p_2, \dots] \cdot s^m, \quad B = \bigoplus_{m \in \mathbb{Z}} B^{(m)}$$

(Think: symmetric functions in x_1, x_2, \dots)

Heisenberg algebra acts via

$$J_k \mapsto \begin{cases} k \frac{\partial}{\partial p_k}, & \text{if } k > 0 \\ p_{|k|}, & \text{if } k < 0 \end{cases}$$

Boson-fermion correspondence

Theorem (Boson-Fermion Correspondence, first part)

There exist vertex operators

$$\psi(z)|_{\mathfrak{F}(m)} = z^m e^{H_-(z)} \Sigma e^{-H_+(z^{-1})}$$

and

$$\psi^*(z)|_{\mathfrak{F}(m)} = z^{-m} e^{-H_-(z)} \Sigma^{-1} e^{H_+(z^{-1})}$$

Boson-fermion correspondence

Theorem (Boson-Fermion Correspondence, first part)

There exist vertex operators

$$\psi(z)|_{\mathfrak{F}(m)} = z^m e^{H_-(z)} \Sigma e^{-H_+(z^{-1})}$$

and

$$\psi^*(z)|_{\mathfrak{F}(m)} = z^{-m} e^{-H_-(z)} \Sigma^{-1} e^{H_+(z^{-1})}$$

Theorem (Boson-Fermion Correspondence, second part)

The map $\mathfrak{F}^{(0)} \rightarrow B^{(0)}$,

$$|\lambda\rangle \mapsto \langle \emptyset | e^{H_+(\mathbf{p})} |\lambda\rangle$$

is an isomorphism of Heisenberg algebra modules. Under this map,

$$|\lambda\rangle \mapsto s_\lambda.$$

The factorial boson-fermion correspondence

(Supersymmetric) factorial Schur functions

Factorial Schur function (Biedenharn-Louck, Macdonald 6th variation):

$$s_{\lambda}(x|\alpha) = \frac{\det \left((x_i - \alpha_1) \cdots (x_i - \alpha_{\lambda_j + j}) \right)_{1 \leq i, j \leq n}}{\prod_{i < j} (x_i - x_j)}.$$

(Supersymmetric) factorial Schur functions

Factorial Schur function (Biedenharn-Louck, Macdonald 6th variation):

$$s_{\lambda}(x|\alpha) = \frac{\det \left((x_i - \alpha_1) \cdots (x_i - \alpha_{\lambda_j + j}) \right)_{1 \leq i, j \leq n}}{\prod_{i < j} (x_i - x_j)}.$$

$$s_{\lambda}(x|\alpha) \xrightarrow{\alpha_i \mapsto 0} s_{\lambda}(x)$$

factorial
Schur

Schur

(Supersymmetric) factorial Schur functions

Factorial Schur function (Biedenharn-Louck, Macdonald 6th variation):

$$s_{\lambda}(x|\alpha) = \frac{\det \left((x_i - \alpha_1) \cdots (x_i - \alpha_{\lambda_j + j}) \right)_{1 \leq i, j \leq n}}{\prod_{i < j} (x_i - x_j)}.$$

$$s_{\lambda}(x/y|\alpha) \xrightarrow{y_i \mapsto \alpha_i} s_{\lambda}(x|\alpha) \xrightarrow{\alpha_i \mapsto 0} s_{\lambda}(x)$$

supersymmetric
factorial
Schur
(Molev)

factorial
Schur

Schur

(Supersymmetric) factorial Schur functions

Factorial Schur function (Biedenharn-Louck, Macdonald 6th variation):

$$s_{\lambda}(x|\alpha) = \frac{\det \left((x_i - \alpha_1) \cdots (x_i - \alpha_{\lambda_j + j}) \right)_{1 \leq i, j \leq n}}{\prod_{i < j} (x_i - x_j)}.$$

$$s_{\lambda}(\mathbf{p}|\alpha) \xrightarrow{p_k \mapsto p_k(x/y)} s_{\lambda}(x/y|\alpha) \xrightarrow{y_i \mapsto \alpha_i} s_{\lambda}(x|\alpha) \xrightarrow{\alpha_i \mapsto 0} s_{\lambda}(x)$$

supersymmetric
factorial
Schur
(Molev)

factorial
Schur

Schur

Deformed fermion fields

Deform the fermion fields:

$$\psi(z|\alpha) = \sum_{i \in \mathbb{Z}} \frac{z^i}{(1 - z\alpha_1) \cdots (1 - z\alpha_i)} \psi_i$$

$$\psi^*(w|\alpha) = \sum_{j \in \mathbb{Z}} \frac{(1 - w\alpha_1) \cdots (1 - w\alpha_{j-1})}{w^j} \psi_j^*.$$

Deformed fermion fields

Deform the fermion fields:

$$\psi(z|\alpha) = \sum_{i \in \mathbb{Z}} \frac{z^i}{(1 - z\alpha_1) \cdots (1 - z\alpha_i)} \psi_i$$

$$\psi^*(w|\alpha) = \sum_{j \in \mathbb{Z}} \frac{(1 - w\alpha_1) \cdots (1 - w\alpha_{j-1})}{w^j} \psi_j^*.$$

This induces an automorphism of the Clifford algebra

Deformed current operators

$$J_k^{(\alpha)} := \sum_{i,j} A_{ij}^k E_{ij}, \text{ where } A_{ij}^k = \begin{cases} e_{j-i-k}(-\alpha_{i+1}, \dots, -\alpha_{j-1}) & \text{if } j \geq i+k \text{ and } k > 0, \\ h_{j-i-k}(\alpha_j, \dots, \alpha_i) & \text{if } j \leq i \leq j-k \text{ and } k \leq 0, \\ 0 & \text{otherwise.} \end{cases}$$

Deformed current operators

$$J_k^{(\alpha)} := \sum_{i,j} A_{ij}^k E_{ij}, \text{ where } A_{ij}^k = \begin{cases} e_{j-i-k}(-\alpha_{i+1}, \dots, -\alpha_{j-1}) & \text{if } j \geq i+k \text{ and } k > 0, \\ h_{j-i-k}(\alpha_j, \dots, \alpha_i) & \text{if } j \leq i \leq j-k \text{ and } k \leq 0, \\ 0 & \text{otherwise.} \end{cases}$$

These live in the space

$$\mathfrak{gl}_{\infty}^+ = \left\{ \sum_{i,j} a_{ij} E_{ij} \mid a_{ij} = 0 \text{ for } i - j \gg 0 \right\}$$

Deformed current operators

$$J_k^{(\alpha)} := \sum_{i,j} A_{ij}^k E_{ij}, \text{ where } A_{ij}^k = \begin{cases} e_{j-i-k}(-\alpha_{i+1}, \dots, -\alpha_{j-1}) & \text{if } j \geq i+k \text{ and } k > 0, \\ h_{j-i-k}(\alpha_j, \dots, \alpha_i) & \text{if } j \leq i \leq j-k \text{ and } k \leq 0, \\ 0 & \text{otherwise.} \end{cases}$$

These live in the space

$$\mathfrak{gl}_{\infty}^+ = \left\{ \sum_{i,j} a_{ij} E_{ij} \mid a_{ij} = 0 \text{ for } i-j \gg 0 \right\}$$

The operators $J_k^{(\alpha)}$ still satisfy the (undeformed) Heisenberg relations

$$[J_k^{(\alpha)}, J_{\ell}^{(\alpha)}] = k\delta_{k,-\ell} \cdot 1$$

Deformed boson-fermion correspondence

Theorem (Deformed Boson-Fermion Correspondence, first part)

There exist vertex operators

$$\psi(z|\alpha)|_{\mathfrak{F}(m)} = z^m e^{H_-(z|\alpha)} \Sigma_{(\alpha)} e^{-H_+(z^{-1}|\alpha)}$$

and

$$\psi^*(z|\alpha)|_{\mathfrak{F}(m)} = z^{-m} e^{-H_-(z|\alpha)} \Sigma_{(\alpha)}^{-1} e^{H_+(z^{-1}|\alpha)},$$

Deformed boson-fermion correspondence

Theorem (Deformed Boson-Fermion Correspondence, first part)

There exist vertex operators

$$\psi(z|\alpha)|_{\mathfrak{F}(m)} = z^m e^{H_-(z|\alpha)} \Sigma_{(\alpha)} e^{-H_+(z^{-1}|\alpha)}$$

and

$$\psi^*(z|\alpha)|_{\mathfrak{F}(m)} = z^{-m} e^{-H_-(z|\alpha)} \Sigma_{(\alpha)}^{-1} e^{H_+(z^{-1}|\alpha)},$$

Theorem (Deformed Boson-Fermion Correspondence, second part)

The map $\mathfrak{F}^{(0)} \rightarrow B^{(0)}$,

$$|\lambda\rangle \mapsto \langle \emptyset | e^{H_+^{(\mathbf{p}|\alpha)}} |\lambda\rangle$$

is an isomorphism of Heisenberg algebra modules. Under this map,

$$|\lambda\rangle \mapsto s_\lambda(\mathbf{p}|\alpha).$$

Consequences

The structure of the boson-fermion correspondence allows us to turn the crank and prove various identities for the (skew) factorial Schur functions:

Consequences

The structure of the boson-fermion correspondence allows us to turn the crank and prove various identities for the (skew) factorial Schur functions:

- Symmetry

Consequences

The structure of the boson-fermion correspondence allows us to turn the crank and prove various identities for the (skew) factorial Schur functions:

- Symmetry
- branching

Consequences

The structure of the boson-fermion correspondence allows us to turn the crank and prove various identities for the (skew) factorial Schur functions:

- Symmetry
- branching
- Giambelli

Consequences

The structure of the boson-fermion correspondence allows us to turn the crank and prove various identities for the (skew) factorial Schur functions:

- Symmetry
- branching
- Giambelli
- Jacobi-Trudi, Nägelsbach–Kostka

Consequences

The structure of the boson-fermion correspondence allows us to turn the crank and prove various identities for the (skew) factorial Schur functions:

- Symmetry
- branching
- Giambelli
- Jacobi-Trudi, Nägelsbach–Kostka
- skew-Cauchy

Consequences

The structure of the boson-fermion correspondence allows us to turn the crank and prove various identities for the (skew) factorial Schur functions:

- Symmetry
- branching
- Giambelli
- Jacobi-Trudi, Nögelbach–Kostka
- skew-Cauchy
- action of symmetric function involution

Consequences

The structure of the boson-fermion correspondence allows us to turn the crank and prove various identities for the (skew) factorial Schur functions:

- Symmetry
- branching
- Giambelli
- Jacobi-Trudi, Nögelbach–Kostka
- skew-Cauchy
- action of symmetric function involution
- Murnaghan-Nakayama

Consequences

The structure of the boson-fermion correspondence allows us to turn the crank and prove various identities for the (skew) factorial Schur functions:

- Symmetry
- branching
- Giambelli
- Jacobi-Trudi, Nägelsbach–Kostka
- skew-Cauchy
- action of symmetric function involution
- Murnaghan-Nakayama

In addition,

- Our half vertex operator matches the transfer matrix of a solvable, free fermionic lattice model of Bump-McNamara-Nakasuji

Consequences

The structure of the boson-fermion correspondence allows us to turn the crank and prove various identities for the (skew) factorial Schur functions:

- Symmetry
- branching
- Giambelli
- Jacobi-Trudi, Nägelsbach–Kostka
- skew-Cauchy
- action of symmetric function involution
- Murnaghan-Nakayama

In addition,

- Our half vertex operator matches the transfer matrix of a solvable, free fermionic lattice model of Bump-McNamara-Nakasuji
- The factorial Schur functions are tau-functions of the KP-hierarchy

Murnaghan-Nakayama rule

Theorem (Factorial Murnaghan–Nakayama rule)

$$p_k s_\lambda(\mathbf{p}|\alpha) = * \cdot s_\lambda(\mathbf{p}|\alpha) + \sum_{\nu} (-1)^{ht(\nu/\lambda)-1} A_{j,i}^{-k} s_\nu(\mathbf{p}|\alpha),$$

where the sum is over all ν such that ν/λ is a nonzero ribbon of size at most k and the contents of the boxes in ν/λ are $i, i+1, \dots, j-1$

(Similar rule for differentiation)

Murnaghan-Nakayama rule



Theorem (Factorial Murnaghan–Nakayama rule)

$$p_k s_\lambda(\mathbf{p}|\alpha) = * \cdot s_\lambda(\mathbf{p}|\alpha) + \sum_{\nu} (-1)^{ht(\nu/\lambda)-1} A_{j,i}^{-k} s_{\nu}(\mathbf{p}|\alpha),$$


where the sum is over all ν such that ν/λ is a nonzero ribbon of size at most k and the contents of the boxes in ν/λ are $i, i+1, \dots, j-1$






(Similar rule for differentiation)

Example: $\lambda = (8, 3, 1)$. Compute $p_3 s_\lambda(\mathbf{p}|\alpha)$:



Example: $\lambda = (8, 3, 1)$. Compute $p_3 s_\lambda(\mathbf{p}|\alpha)$:





Thank You



Happy Birthday, Kailash Misra!