

**Measurement of  $D\bar{D}$  Decays from the  $\psi(3770)$  Resonance**

A DISSERTATION  
SUBMITTED TO THE FACULTY OF THE GRADUATE SCHOOL  
OF THE UNIVERSITY OF MINNESOTA  
BY

Andy Jarod Julin

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS  
FOR THE DEGREE OF  
DOCTOR OF PHILOSOPHY

Prof. Ron Poling

May, 2017

© Andy Jarod Julin 2017



The text of this work is licensed under a Creative Commons  
Attribution-ShareAlike 4.0 International license.

# Acknowledgements

This is where the Acknowledgements go!

# Dedication

This is where the Dedications go!

## Abstract

We measure the production cross section of  $e^+e^- \rightarrow \psi(3770) \rightarrow D\bar{D}$  near the peak of the  $\psi(3770)$  resonance. The  $(69.80 \pm 0.03)\text{ pb}^{-1}$  of  $e^+e^-$  annihilation data used were collected in 2010 over a center-of-mass energy range of 3.735 GeV to 3.870 GeV. From previously observed  $e^+e^- \rightarrow D\bar{D}$  cross section measurements, this shape cannot be explained by a single Breit-Wigner. Instead, this analysis fitted using interference effects from non-resonant  $D\bar{D}$  production, and measured the mass and width of the  $\psi(3770)$  more precisely than previous results.

# Contents

<b>Acknowledgements</b>	<b>i</b>
<b>Dedication</b>	<b>ii</b>
<b>Abstract</b>	<b>iii</b>
<b>List of Tables</b>	<b>viii</b>
<b>List of Figures</b>	<b>xii</b>
<b>1 Introduction</b>	<b>1</b>
1.1 Procedure . . . . .	3
<b>2 Theoretical Background</b>	<b>5</b>
2.1 Standard Model . . . . .	5
2.1.1 Electromagnetic Force . . . . .	5
2.1.2 Weak Force . . . . .	6
2.1.3 Strong Force . . . . .	7
2.1.4 Elementary Particles . . . . .	8
2.2 Charmonium . . . . .	11
<b>3 Detector and Related Systems</b>	<b>14</b>
3.1 BEPCII Accelerator . . . . .	15
3.2 BESIII Detector . . . . .	15
3.2.1 Multi-Layer Drift Chamber . . . . .	17
3.2.2 Time-of-Flight System . . . . .	19

3.2.3	Electromagnetic Calorimeter . . . . .	20
3.2.4	Muon Identifier . . . . .	22
3.3	Triggering Systems . . . . .	22
<b>4</b>	<b>Analysis Software</b>	<b>26</b>
4.1	BESIII Offline Software System . . . . .	26
4.1.1	Framework . . . . .	26
4.1.2	Simulation . . . . .	27
4.1.3	Reconstruction . . . . .	28
4.1.4	Calibration . . . . .	29
4.2	Detector Simulation . . . . .	29
4.2.1	Multi-Layer Drift Chamber . . . . .	29
4.2.2	Time-of-Flight System . . . . .	30
4.2.3	Electromagnetic Calorimeter . . . . .	31
4.2.4	Muon Identifier . . . . .	31
4.3	<i>D</i> -Tagging . . . . .	32
4.3.1	Selection Cuts . . . . .	33
4.3.2	Reconstruction . . . . .	35
<b>5</b>	<b>Measurement of <math>\sigma_{\psi(3770) \rightarrow D\bar{D}}</math> near <math>\psi(3770)</math></b>	<b>37</b>
5.1	Derivation of $\sigma_{\psi(3770) \rightarrow D\bar{D}}$ . . . . .	37
5.2	Form Factors . . . . .	40
5.3	Data and Monte Carlo Samples . . . . .	41
5.3.1	Data Samples . . . . .	41
5.3.2	Center-of-Mass Energy Measurement . . . . .	42
5.3.3	Luminosity Measurement . . . . .	44
5.3.4	Monte Carlo Generation . . . . .	47
5.3.5	Software Packages . . . . .	47
5.4	Signal Determination . . . . .	48
5.5	Efficiency Correction . . . . .	49
5.5.1	CP Violation Correction . . . . .	49
5.6	Fitting Procedure . . . . .	55
5.6.1	Coulomb Correction . . . . .	59

5.7	Systematics . . . . .	61
5.7.1	$\psi(3770)$ Parameter Systematics . . . . .	61
5.7.2	Cross Section Systematics . . . . .	67
5.8	Results . . . . .	70
<b>6</b>	<b>Measurement of Hadronic Production and <math>\Gamma(\psi(3770) \rightarrow \text{non-}D\bar{D})</math></b>	<b>72</b>
6.1	Data and Monte Carlo Samples . . . . .	72
6.1.1	Data Samples . . . . .	72
6.1.2	Center-of-Mass Energy Measurement . . . . .	73
6.2	Event Selection . . . . .	74
6.3	Hadron Counting . . . . .	75
6.4	Background Subtraction . . . . .	76
6.5	Efficiency Extrapolation . . . . .	78
6.6	Results for $\psi(3770)$ Data . . . . .	86
6.6.1	$D\bar{D}$ Correction . . . . .	87
6.6.2	Reconstruction Efficiencies . . . . .	90
6.6.3	Signal Amounts . . . . .	91
6.6.4	Non- $D\bar{D}$ Branching Fraction (Breit-Wigner) . . . . .	92
6.7	$\psi(2S)$ Background Investigation . . . . .	93
6.7.1	Non- $D\bar{D}$ Branching Fraction ( $\psi(2S)$ from Data) . . . . .	94
6.7.2	Non- $D\bar{D}$ Branching Fraction ( $\psi(2S)$ Excluded) . . . . .	98
6.8	Results for Scan Data . . . . .	102
<b>7</b>	<b>Conclusion</b>	<b>107</b>
<b>References</b>		<b>108</b>
<b>Appendix A.</b>	<b>Glossary and Acronyms</b>	<b>112</b>
A.1	Glossary . . . . .	112
A.2	Acronyms / Initialisms . . . . .	112
<b>Appendix B.</b>	<b><math>D^0</math> Signal Fits</b>	<b>114</b>
<b>Appendix C.</b>	<b><math>D^+</math> Signal Fits</b>	<b>126</b>

Appendix D. Scan Data Hadronic Counting Fits	137
Appendix E. Scan Data Reconstruction Efficiency	149
Appendix F. Scan Data Signal Counts	159

# List of Tables

1.1	Previous experimental results for the mass of the $\psi(3770)$ . . . . .	2
4.1	The reconstructed $D$ -tag modes used in this analysis. . . . .	33
4.2	The required cuts to identify charged tracks as $\pi^\pm$ or $K^\pm$ . . . . .	34
4.3	The required cuts to identify neutral showers as a $\gamma$ . . . . .	34
4.4	The required cuts to reconstruct $\gamma\gamma$ pairs as a $\pi^0$ . . . . .	34
4.5	The required cuts to reconstruct $\pi^+\pi^-$ pairs as a $K_S^0$ . . . . .	35
4.6	Cuts to prevent cosmic ray and lepton backgrounds in $D^0 \rightarrow K^-\pi^+$ . . .	35
5.1	Selection cuts on muon tracks used to determine the center-of-mass energy.	42
5.2	Selection cuts on electron tracks used to determine the luminosity. . . .	44
5.3	Measured luminosities for each energy bin. . . . .	46
5.4	Number of events contained in each generated sample and the scan data.	47
5.5	The software packages used in this analysis and their version numbers. .	48
5.6	Number of proper and generated particles for $D^0$ . . . . .	50
5.7	Number of proper and generated particles for $D^+$ (part 1). . . . .	51
5.8	Number of proper and generated particles for $D^+$ (part 2). . . . .	52
5.9	Mode-by-mode reconstruction efficiencies for $D^0$ and $D^+$ . . . . .	53
5.11	The quantum correlated factors for the $D^0$ modes. . . . .	53
5.10	The overall reconstruction efficiency of $D^0$ and $D^+$ for each energy bin.	54
5.12	The $DD$ cross section at each $E_{\text{cm}}$ point. . . . .	56
5.13	Single-tag fitting differences by mode. . . . .	63
5.14	Comparison of input and output fit parameters. . . . .	64
5.15	Comparison of output fit parameters between ISR generators. . . . .	65
5.16	Systematic uncertainties relative to the measured parameters of the $\psi(3770)$ .	67
5.17	Systematics shifts affecting the cross section measurements. . . . .	68

5.18	Measurements of the $D^0 / D^+$ cross sections.	69
5.19	Final results for the $\psi(3770)$ parameters.	70
5.20	Fit results compared to the KEDR results and the PDG.	70
5.21	Comparison of cross section calculations at $E_{\text{cm}} = 3.7732 \text{ GeV}$ .	71
6.1	Data samples used for the inclusive measurement.	73
6.2	Selection cuts on charged tracks used to count hadronic events.	74
6.3	Selection cuts on neutral tracks used to count hadronic events.	74
6.4	Selection cuts to remove Bhabha and two-photon backgrounds.	74
6.5	Standard selection cuts (SHAD) for counting hadronic events.	75
6.6	Loose selection cuts (LHAD) for counting hadronic events.	75
6.7	Tight selection cuts (THAD) for counting hadronic events.	75
6.8	Reconstruction of background samples for the old continuum data.	77
6.9	Hadronic events selected in the old continuum data.	77
6.10	Reconstruction of background samples for the new continuum data.	80
6.11	Hadronic events selected in the new continuum data.	81
6.12	Selection methods for the $D\bar{D}$ efficiency correction.	87
6.13	Efficiency corrections for the $\psi(3770)$ samples.	87
6.14	Reconstruction of background samples for the $\psi(3770)$ data.	90
6.15	Hadronic events selected in the $\psi(3770)$ data.	91
6.16	Cross sections for $\psi(3770) \rightarrow \text{non-}D\bar{D}$ found using the $\psi(3770)$ data.	92
6.17	Branching fractions for $\psi(3770) \rightarrow \text{non-}D\bar{D}$ found using the $\psi(3770)$ data.	92
6.18	Cross sections of $\psi(2S)$ calculated using two different methods.	93
6.19	Cross sections for $\psi(3770) \rightarrow \text{non-}D\bar{D}$ found using the $\psi(3770)$ data.	94
6.20	Branching fractions for $\psi(3770) \rightarrow \text{non-}D\bar{D}$ found using the $\psi(3770)$ data.	94
6.21	Cross sections for $\psi(3770) \rightarrow \text{non-}D\bar{D}$ found using the $\psi(3770)$ data.	98
6.22	Branching fractions for $\psi(3770) \rightarrow \text{non-}D\bar{D}$ found using the $\psi(3770)$ data.	98
6.23	Energy values and background cross sections for the scan data.	103
A.1	Acronyms and Initialisms	112
E.1	Reconstruction of background samples for the 3734 (Scan) data.	149
E.2	Reconstruction of background samples for the 3736 (Scan) data.	149
E.3	Reconstruction of background samples for the 3744 (Scan) data.	150
E.4	Reconstruction of background samples for the 3748 (Scan) data.	150

E.5	Reconstruction of background samples for the 3750 (Scan) data.	150
E.6	Reconstruction of background samples for the 3751 (Scan) data.	150
E.7	Reconstruction of background samples for the 3753 (Scan) data.	151
E.8	Reconstruction of background samples for the 3755 (Scan) data.	151
E.9	Reconstruction of background samples for the 3756 (Scan) data.	151
E.10	Reconstruction of background samples for the 3759 (Scan) data.	151
E.11	Reconstruction of background samples for the 3762 (Scan) data.	152
E.12	Reconstruction of background samples for the 3765 (Scan) data.	152
E.13	Reconstruction of background samples for the 3767 (Scan) data.	152
E.14	Reconstruction of background samples for the 3771 (Scan) data.	152
E.15	Reconstruction of background samples for the 3774 (Scan) data.	153
E.16	Reconstruction of background samples for the 3777 (Scan) data.	153
E.17	Reconstruction of background samples for the 3780 (Scan) data.	153
E.18	Reconstruction of background samples for the 3782 (Scan) data.	153
E.19	Reconstruction of background samples for the 3786 (Scan) data.	154
E.20	Reconstruction of background samples for the 3789 (Scan) data.	154
E.21	Reconstruction of background samples for the 3792 (Scan) data.	154
E.22	Reconstruction of background samples for the 3797 (Scan) data.	154
E.23	Reconstruction of background samples for the 3800 (Scan) data.	155
E.24	Reconstruction of background samples for the 3802 (Scan) data.	155
E.25	Reconstruction of background samples for the 3807 (Scan) data.	155
E.26	Reconstruction of background samples for the 3809 (Scan) data.	155
E.27	Reconstruction of background samples for the 3813 (Scan) data.	156
E.28	Reconstruction of background samples for the 3815 (Scan) data.	156
E.29	Reconstruction of background samples for the 3822 (Scan) data.	156
E.30	Reconstruction of background samples for the 3832 (Scan) data.	156
E.31	Reconstruction of background samples for the 3839 (Scan) data.	157
E.32	Reconstruction of background samples for the 3849 (Scan) data.	157
E.33	Reconstruction of background samples for the 3855 (Scan) data.	157
E.34	Reconstruction of background samples for the 3863 (Scan) data.	157
F.1	Hadronic events selected in the 3734 (Scan) data.	159
F.2	Hadronic events selected in the 3736 (Scan) data.	159

F.3	Hadronic events selected in the 3744 (Scan) data.	160
F.4	Hadronic events selected in the 3748 (Scan) data.	160
F.5	Hadronic events selected in the 3750 (Scan) data.	160
F.6	Hadronic events selected in the 3751 (Scan) data.	161
F.7	Hadronic events selected in the 3753 (Scan) data.	161
F.8	Hadronic events selected in the 3755 (Scan) data.	161
F.9	Hadronic events selected in the 3756 (Scan) data.	162
F.10	Hadronic events selected in the 3759 (Scan) data.	162
F.11	Hadronic events selected in the 3762 (Scan) data.	162
F.12	Hadronic events selected in the 3765 (Scan) data.	163
F.13	Hadronic events selected in the 3767 (Scan) data.	163
F.14	Hadronic events selected in the 3771 (Scan) data.	163
F.15	Hadronic events selected in the 3774 (Scan) data.	164
F.16	Hadronic events selected in the 3777 (Scan) data.	164
F.17	Hadronic events selected in the 3780 (Scan) data.	164
F.18	Hadronic events selected in the 3782 (Scan) data.	165
F.19	Hadronic events selected in the 3786 (Scan) data.	165
F.20	Hadronic events selected in the 3789 (Scan) data.	165
F.21	Hadronic events selected in the 3792 (Scan) data.	166
F.22	Hadronic events selected in the 3797 (Scan) data.	166
F.23	Hadronic events selected in the 3800 (Scan) data.	166
F.24	Hadronic events selected in the 3802 (Scan) data.	167
F.25	Hadronic events selected in the 3807 (Scan) data.	167
F.26	Hadronic events selected in the 3809 (Scan) data.	167
F.27	Hadronic events selected in the 3813 (Scan) data.	168
F.28	Hadronic events selected in the 3815 (Scan) data.	168
F.29	Hadronic events selected in the 3822 (Scan) data.	168
F.30	Hadronic events selected in the 3832 (Scan) data.	169
F.31	Hadronic events selected in the 3839 (Scan) data.	169
F.32	Hadronic events selected in the 3849 (Scan) data.	169
F.33	Hadronic events selected in the 3855 (Scan) data.	170
F.34	Hadronic events selected in the 3863 (Scan) data.	170

# List of Figures

1.1	Measurements of $R = \sigma(e^+e^- \rightarrow \text{hadrons})/\sigma(e^+e^- \rightarrow \mu^+\mu^-)$ . . . . .	1
1.2	Measured and predicted charmonium resonances. . . . .	2
2.1	The standard model of particle physics. . . . .	8
2.2	The Cabibbo-Kobayashi-Maskawa (CKM) Matrix. . . . .	9
2.3	The standard form of the Cabibbo-Kobayashi-Maskawa (CKM) Matrix. . . . .	10
2.4	An example Feynman diagram for the decay of $\psi(3686)$ . . . . .	12
2.5	An example Feynman diagram for the decay of $\psi(3770)$ . . . . .	13
3.1	A schematic of the BESIII detector. . . . .	16
3.2	MDC energy deposition for various particles as a function of momenta. .	19
3.3	ToF measurements for various particles as a function of momenta. . .	20
3.4	EMC energy deposition for various particles as a function of momentum. .	21
3.5	Triggering systems for event filtering at BESIII. . . . .	24
5.1	The differences measured for scan data center-of-mass energies. . . . .	43
5.2	The values measured for on-peak data center-of-mass energies. . . . .	44
5.3	An example 2D ( $\Delta E$ vs. $m_{\text{BC}}$ ) signal fit. . . . .	48
5.4	Mode-by-mode MC efficiencies for $D^0$ and $D^+$ . . . . .	55
5.5	The measured $e^+e^- \rightarrow D\bar{D}$ cross sections. . . . .	57
5.6	The Exponential Model fit results. . . . .	58
5.7	The Vector Dominance Model fit results. . . . .	59
5.8	The Vector Dominance Model fit results with Coulomb interactions. .	60
5.9	The ratio of measured $D^+$ to $D^0$ cross sections. . . . .	61
5.10	The $K\pi$ vs. $\pi\pi$ invariant masses for the mode $D^+ \rightarrow K^-\pi^+\pi^+$ . . . . .	66
6.1	The number of hadrons found in the 3650 (Old) data sample. . . . .	76
6.2	The number of hadrons found in the 3500 (New) data sample. . . . .	78

6.3	The number of hadrons found in the 3542 (New) data sample. . . . .	78
6.4	The number of hadrons found in the 3600 (New) data sample. . . . .	79
6.5	The number of hadrons found in the 3650 (New) data sample. . . . .	79
6.6	The number of hadrons found in the 3671 (New) data sample. . . . .	79
6.7	The extrapolation for SHAD events using the new continuum data. . . . .	83
6.8	The extrapolation for LHAD events using the new continuum data. . . . .	84
6.9	The extrapolation for THAD events using the new continuum data. . . . .	85
6.10	The number of hadrons found in the $\psi(3770)$ (R1) data sample. . . . .	86
6.11	The number of hadrons found in the $\psi(3770)$ (R2) data sample. . . . .	86
6.12	The other-side $D^0$ tracks and corresponding $D^0\overline{D}^0$ multiplicities for R1. . . . .	88
6.13	The other-side $D^+$ tracks and corresponding $D^+D^-$ multiplicities for R1. . . . .	88
6.14	The other-side $D^0$ tracks and corresponding $D^0\overline{D}^0$ multiplicities for R2. . . . .	89
6.15	The other-side $D^+$ tracks and corresponding $D^+D^-$ multiplicities for R2. . . . .	89
6.16	The extrapolation for SHAD events using $\psi(2S)$ data. . . . .	95
6.17	The extrapolation for LHAD events using $\psi(2S)$ data. . . . .	96
6.18	The extrapolation for THAD events using $\psi(2S)$ data. . . . .	97
6.19	The extrapolation for SHAD events without including $\psi(2S)$ events. . . . .	99
6.20	The extrapolation for LHAD events without including $\psi(2S)$ events. . . . .	100
6.21	The extrapolation for THAD events without including $\psi(2S)$ events. . . . .	101
6.22	The inclusive cross sections measured for the scan data region. . . . .	104
6.23	The non- $D\overline{D}$ cross sections measured for the scan data region. . . . .	105
6.24	The non- $D\overline{D}$ branching fractions measured for the scan data region. . . . .	106
B.1	The 2D signal fit for $D^0$ in Bin 0. . . . .	114
B.2	The 2D signal fit for $D^0$ in Bin 1. . . . .	114
B.3	The 2D signal fit for $D^0$ in Bin 2. . . . .	115
B.4	The 2D signal fit for $D^0$ in Bin 3. . . . .	115
B.5	The 2D signal fit for $D^0$ in Bin 4. . . . .	115
B.6	The 2D signal fit for $D^0$ in Bin 5. . . . .	116
B.7	The 2D signal fit for $D^0$ in Bin 6. . . . .	116
B.8	The 2D signal fit for $D^0$ in Bin 7. . . . .	116
B.9	The 2D signal fit for $D^0$ in Bin 8. . . . .	117
B.10	The 2D signal fit for $D^0$ in Bin 9. . . . .	117

B.11	The 2D signal fit for $D^0$ in Bin 10.	117
B.12	The 2D signal fit for $D^0$ in Bin 11.	118
B.13	The 2D signal fit for $D^0$ in Bin 12.	118
B.14	The 2D signal fit for $D^0$ in Bin 13.	118
B.15	The 2D signal fit for $D^0$ in Bin 14.	119
B.16	The 2D signal fit for $D^0$ in Bin 15.	119
B.17	The 2D signal fit for $D^0$ in Bin 16.	119
B.18	The 2D signal fit for $D^0$ in Bin 17.	120
B.19	The 2D signal fit for $D^0$ in Bin 18.	120
B.20	The 2D signal fit for $D^0$ in Bin 19.	120
B.21	The 2D signal fit for $D^0$ in Bin 20.	121
B.22	The 2D signal fit for $D^0$ in Bin 21.	121
B.23	The 2D signal fit for $D^0$ in Bin 22.	121
B.24	The 2D signal fit for $D^0$ in Bin 23.	122
B.25	The 2D signal fit for $D^0$ in Bin 24.	122
B.26	The 2D signal fit for $D^0$ in Bin 25.	122
B.27	The 2D signal fit for $D^0$ in Bin 26.	123
B.28	The 2D signal fit for $D^0$ in Bin 27.	123
B.29	The 2D signal fit for $D^0$ in Bin 28.	123
B.30	The 2D signal fit for $D^0$ in Bin 29.	124
B.31	The 2D signal fit for $D^0$ in Bin 30.	124
B.32	The 2D signal fit for $D^0$ in Bin 31.	124
B.33	The 2D signal fit for $D^0$ in Bin 32.	125
B.34	The 2D signal fit for $D^0$ in Bin 33.	125
C.1	The 2D signal fit for $D^+$ in Bin 2.	126
C.2	The 2D signal fit for $D^+$ in Bin 3.	126
C.3	The 2D signal fit for $D^+$ in Bin 4.	127
C.4	The 2D signal fit for $D^+$ in Bin 5.	127
C.5	The 2D signal fit for $D^+$ in Bin 6.	127
C.6	The 2D signal fit for $D^+$ in Bin 7.	128
C.7	The 2D signal fit for $D^+$ in Bin 8.	128
C.8	The 2D signal fit for $D^+$ in Bin 9.	128

C.9	The 2D signal fit for $D^+$ in Bin 10.	129
C.10	The 2D signal fit for $D^+$ in Bin 11.	129
C.11	The 2D signal fit for $D^+$ in Bin 12.	129
C.12	The 2D signal fit for $D^+$ in Bin 13.	130
C.13	The 2D signal fit for $D^+$ in Bin 14.	130
C.14	The 2D signal fit for $D^+$ in Bin 15.	130
C.15	The 2D signal fit for $D^+$ in Bin 16.	131
C.16	The 2D signal fit for $D^+$ in Bin 17.	131
C.17	The 2D signal fit for $D^+$ in Bin 18.	131
C.18	The 2D signal fit for $D^+$ in Bin 19.	132
C.19	The 2D signal fit for $D^+$ in Bin 20.	132
C.20	The 2D signal fit for $D^+$ in Bin 21.	132
C.21	The 2D signal fit for $D^+$ in Bin 22.	133
C.22	The 2D signal fit for $D^+$ in Bin 23.	133
C.23	The 2D signal fit for $D^+$ in Bin 24.	133
C.24	The 2D signal fit for $D^+$ in Bin 25.	134
C.25	The 2D signal fit for $D^+$ in Bin 26.	134
C.26	The 2D signal fit for $D^+$ in Bin 27.	134
C.27	The 2D signal fit for $D^+$ in Bin 28.	135
C.28	The 2D signal fit for $D^+$ in Bin 29.	135
C.29	The 2D signal fit for $D^+$ in Bin 30.	135
C.30	The 2D signal fit for $D^+$ in Bin 31.	136
C.31	The 2D signal fit for $D^+$ in Bin 32.	136
C.32	The 2D signal fit for $D^+$ in Bin 33.	136
D.1	The number of hadrons found in the 3734 (Scan) data sample.	137
D.2	The number of hadrons found in the 3736 (Scan) data sample.	138
D.3	The number of hadrons found in the 3744 (Scan) data sample.	138
D.4	The number of hadrons found in the 3748 (Scan) data sample.	138
D.5	The number of hadrons found in the 3750 (Scan) data sample.	139
D.6	The number of hadrons found in the 3751 (Scan) data sample.	139
D.7	The number of hadrons found in the 3753 (Scan) data sample.	139
D.8	The number of hadrons found in the 3755 (Scan) data sample.	140

D.9	The number of hadrons found in the 3756 (Scan) data sample.	140
D.10	The number of hadrons found in the 3759 (Scan) data sample.	140
D.11	The number of hadrons found in the 3762 (Scan) data sample.	141
D.12	The number of hadrons found in the 3765 (Scan) data sample.	141
D.13	The number of hadrons found in the 3767 (Scan) data sample.	141
D.14	The number of hadrons found in the 3771 (Scan) data sample.	142
D.15	The number of hadrons found in the 3774 (Scan) data sample.	142
D.16	The number of hadrons found in the 3777 (Scan) data sample.	142
D.17	The number of hadrons found in the 3780 (Scan) data sample.	143
D.18	The number of hadrons found in the 3782 (Scan) data sample.	143
D.19	The number of hadrons found in the 3786 (Scan) data sample.	143
D.20	The number of hadrons found in the 3789 (Scan) data sample.	144
D.21	The number of hadrons found in the 3792 (Scan) data sample.	144
D.22	The number of hadrons found in the 3797 (Scan) data sample.	144
D.23	The number of hadrons found in the 3800 (Scan) data sample.	145
D.24	The number of hadrons found in the 3802 (Scan) data sample.	145
D.25	The number of hadrons found in the 3807 (Scan) data sample.	145
D.26	The number of hadrons found in the 3809 (Scan) data sample.	146
D.27	The number of hadrons found in the 3813 (Scan) data sample.	146
D.28	The number of hadrons found in the 3815 (Scan) data sample.	146
D.29	The number of hadrons found in the 3822 (Scan) data sample.	147
D.30	The number of hadrons found in the 3832 (Scan) data sample.	147
D.31	The number of hadrons found in the 3839 (Scan) data sample.	147
D.32	The number of hadrons found in the 3849 (Scan) data sample.	148
D.33	The number of hadrons found in the 3855 (Scan) data sample.	148
D.34	The number of hadrons found in the 3863 (Scan) data sample.	148

# Chapter 1

## Introduction

Since the discovery of the  $J/\psi$  particle in 1974, the charm energy range (3.0 GeV to 4.5 GeV) has been one of the most precisely studied regions in particle physics. This has led to the further discovery of many more resonances, as shown in Figure 1.1.




Figure 1.1: Measurements of  $R = \sigma(e^+e^- \rightarrow \text{hadrons})/\sigma(e^+e^- \rightarrow \mu^+\mu^-)$ .

Many of the lower mass resonances are predictable within the context of the quark model. However, there are a number of states which have been predicted, but not yet discovered. There are also states which were discovered experimentally without any corresponding predictions. A variety of these particles are shown in Figure 1.2.




Figure 1.2: Measured and predicted charmonium resonances.  
 Solid black lines represent measurements while the dashed red lines are predictions.

The resonances below the  $D\bar{D}$  threshold, like the  $J/\psi$  and the  $\psi(2S)$ , show solid agreement with their theoretical predictions. However, many of the ones above, such as the  $\psi(3770)$ , still show some disagreement. This is likely due to the more complicated interactions introduced from  $D\bar{D}$  decays. Several experiments have attempted to measure the shape of the  $\psi(3770)$  based on different assumptions. The most prominent of these is the assumption of interference. This can have notable effects on the resultant parameters of the  $\psi(3770)$ , such as the mass, shown in Table 1.1.

$M^{\psi(3770)}$ [MeV] (No Interference)	$M^{\psi(3770)}$ [MeV] (With Interference)
BES-II [1] $3772.0 \pm 1.9$	BaBar [4] $3778.8 \pm 1.9 \pm 0.9$
Belle [2] $3776.0 \pm 5.0 \pm 4.0$	KEDR [5] $3779.2^{+1.8+0.5+0.3}_{-1.7-0.7-0.3}$
BaBar [3] $3775.5 \pm 2.4 \pm 0.5$	

Table 1.1: Previous experimental results for the mass of the  $\psi(3770)$ . Where applicable, the first errors are statistical, the second are systematic, and the third are model-dependent.

Both BaBar [4] and KEDR [5] found it necessary to include interference effects for fitting the  $D\bar{D}$  spectrum. However, the statistics of the KEDR sample were insufficient to fully resolve the discrepancies seen with other experiments that ignored interference. **Using the larger data sample available at BESIII, we have precisely measured and analyzed the shape of the  $D\bar{D}$  spectrum around the  $\psi(3770)$  resonance.** We have also used this measurement to probe the branching fraction of non- $D\bar{D}$  decays in this region.

## 1.1 Procedure

The basis of this measurement involves identifying  $D\bar{D}$  pairs which have decayed from  $\psi(3770)$ . All data used in this analysis was collected from  $e^+e^-$  collisions analyzed by the BESIII detector. As  $e^-$  and  $e^+$  are fundamental particles, the total energy of each is transferred in the collision. Additionally, the ability to precisely tune these two beam energies allows for the targeted production of specific particles, such as the  $\psi(3770)$ . However, resonances like this will decay almost instantaneously. For example, the  $\psi(3770)$  has an average lifetime of  $\sim 10^{-23}$  s. Even if it were traveling at  $c$ , the farthest distance this could travel would be  $(3 \times 10^8 \text{ m/s}) \times (\sim 10^{-23} \text{ s}) \approx 1 \text{ fm}$ . This distance, approximately the radius of a proton, is far too small to be resolved by our detectors. Even the initial decay into  $D\bar{D}$  pairs is too small to precisely measure, as their lifetime of  $4 \times 10^{-13}$  s corresponds to a maximum distance of  $\sim 1 \text{ mm}$  if it were traveling at  $c$ . However, the limited available energy in these decays ( $m_{\psi(3770)} - 2 \times m_D \approx 40 \text{ MeV}$ ) means the velocity is much, much slower.

Instead, the reconstruction of candidate  $D$  particles relies on measuring their decays into other known modes. While there are dozens of possible decays modes, we focus on those which have high branching fractions, and are comprised of particles identifiable by the BESIII detector. For our purposes, these are  $\pi^\pm, K^\pm, \pi^0$ , and  $K_S^0$ . Each of these particles leave ‘tracks’ within the detector; charged particles travel along curved paths and interact electromagnetically with charged wires along their trajectory, while neutral particles travel along straight paths and deposit bursts of energy after contacting crystals located along the inner walls of the detector. The various components of the BESIII detector analyze these tracks to determine the type of particle, as well as its momentum

and energy. By analyzing sets of particles corresponding to the chosen decay modes, we can reconstruct the combination most likely to have originated from a  $D$  based on their total energy.

To determine the  $\psi(3770) \rightarrow D\bar{D}$  cross section, we need to count the number of  $D\bar{D}$  pairs produced, as well as measure the integrated luminosity, at each energy point in our data sample. This counting is done not only for the actual collision data collected, but also for computed generated Monte Carlo (MC) background samples. Each of these backgrounds corresponds to a particular event type which may be mistaken as signal, such as  $e^+e^- \rightarrow \tau^+\tau^-$ . We subtract these misidentified contributions from the total amount found in data to determine the actual number of reconstructed  $D\bar{D}$  events. Also from MC, we calculate the efficiency of reconstructing  $D\bar{D}$  decays based on our selection criteria. By dividing the actual reconstructed events by this efficiency, we identify the true number of  $D\bar{D}$  events produced. Then, using the measured luminosity for each energy point, we determine the cross section as a function of center-of-mass energy.

The specific details of this analysis start with background on relevant theoretical concepts. Next, we list the specifications for the collider and detector which collected the data used for these measurements along with their related analysis software and reconstruction methods. From here, we further describe the procedure for determining the  $\psi(3770) \rightarrow D\bar{D}$  cross section and show the results with systematic uncertainties. Finally, we examine the current progress of measuring the non- $D\bar{D}$  branching fraction.

# Chapter 2

## Theoretical Background

### 2.1 Standard Model

Developed throughout the 1960s and 1970s, the Standard Model provides the most complete description of observable matter in the universe to date. It is a classification of all confirmed subatomic particles currently known, and predicts the most accurate results of any scientific theory ever measured. Each of the electromagnetic, weak, and strong fundamental forces are well described by this formulation. These three are described by an  $SU(3) \times SU(2) \times U(1)$  group, where the  $SU(3)$  corresponds to the strong force, the  $SU(2)$  corresponds to the weak force, and the  $U(1)$  corresponds to the electromagnetic force. The remaining fundamental force, gravity, is not included in the Standard Model. It is negligible on the scale of the masses of fundamental particles, and will be ignored in the discussions that follow.

#### 2.1.1 Electromagnetic Force

The electromagnetic force is responsible for the forces between objects with electric charge, most notably binding together electrons and protons to form atoms and the structures they comprise. The theory of electromagnetic interactions is known as Quantum Electrodynamics (QED). Within this theory, the mediator of this force is the photon, a massless vector boson. As there is only a single mediator, and a single conserved quantity (electric charge), the formulation of QED is relatively simple compared to

the other forces. Still, the predictions it makes show astounding consistency with experiment, such as correctly calculating the anomalous magnetic dipole moment of the electron to more than 10 significant figures. Much of this success is due to QED being calculable through perturbation theory, where corrections are applied in terms of higher order factors of the coupling constant,  $\alpha$ . This is possible due to its relatively small value ( $\alpha \approx 1/137$ ), as the terms are convergent below very high orders of  $\alpha$ .

### 2.1.2 Weak Force

The weak force is responsible for radioactive decay and other subatomic phenomena. This is distinct from the electromagnetic and strong interactions, where the constituent particles cannot change their types (or flavors). The mediators of this force are the  $W$  and  $Z$ , which are massive vector bosons. Not only are each of their masses non-zero, they are extremely heavy particles at  $80.4\text{ GeV}$  and  $90.2\text{ GeV}$ , respectively [6]. These large masses not only limit the interaction distance of the weak force, but also minimize the interaction strength (which is inversely proportion to mass). Furthermore, the large  $W$  and  $Z$  masses also lead to much slower interaction times, further reducing the effects of the weak force in comparison to the strong and electromagnetic forces.

In addition to transforming particle flavor, the weak force is also unique in its violation of various symmetries. The first discovery of symmetry violation came in 1957, when Wu and others [8] discovered the weak force did not behave identically under parity (P) transformations (i.e., mirror reflection). To account for this, a new theory conserving a compound symmetry was proposed. This combined charge conjugation (C), the swapping of particles with their antiparticles, with parity to form CP parity. However, in 1964, evidence of CP violation was also discovered by Cronin and Fitch [9]. The resolution to this symmetry conservation involves yet a third symmetry, time reversal (T), in which time is replaced with its negative ( $t \rightarrow -t$ ). While the weak force violates these symmetries individually, the application of all three (CPT) is conserved across all known processes, and is known as the CPT Theorem.

At higher energy scales, the electromagnetic and weak forces unify into the electroweak force. In this theory, there are initially four massless gauge bosons mediating the interactions. As a result of the Higgs mechanism, the initial gauge symmetry is broken at lower energies, and three of these bosons acquire a mass. These three bosons

are the  $W^\pm$  and  $Z$ , while the remaining massless boson is the  $\gamma$ . The energies scales required for this unification were only present in the early universe. Before this, it is also believed there was an epoch of even higher energy, in which the electroweak force merged with the strong force.

### 2.1.3 Strong Force

The strong force is responsible for binding together particles known as hadrons. The theory of strong interactions is known as Quantum Chromodynamics (QCD). Like the electromagnetic force, the mediator of the strong force is also a massless vector boson, the gluon. However, while massless particles typically correspond to an infinite interaction range, the strong potential becomes very large at higher separations. This prevents particles which interact through the strong force, known as quarks (see Section 2.1.4), from existing as isolated entities in a process known as confinement. The typical interaction range is on the order of the proton radius, around  $10^{-15}$  m. QCD calculations face serious challenges, however, as the coupling constant is not small ( $\alpha_S \gtrsim 1$ ). This excludes the use of perturbation theory for most cases, as the higher order terms do not converge.

Strong interactions are associated with a corresponding conserved quantity known as the color charge. Despite its name, however, the term 'color' has no association with light, which is a purely electromagnetic phenomena. There are three colors associated with this charge, red ( $r$ ), green ( $g$ ), and blue ( $b$ ). For anti-particles, there are oppositely charged values ( $\bar{r}$ ,  $\bar{g}$ , and  $\bar{b}$ ). In order for hadrons to be formed, the total color values of the constituents must be colorless. This means the total sum must involve all three colors ( $rgb$  or  $\bar{r}\bar{g}\bar{b}$ ) or pairs of opposite colors ( $r\bar{r}$ ,  $g\bar{g}$ , or  $b\bar{b}$ ). However, these individual colors are not observable in nature. Because particles with different color values are distinct, this effectively triples the number of possible particle combinations, due to combinatorics.

Unlike the photon, which does not carry an electric charge, gluons do possess a color charge. There are eight possible color combinations which a gluon may possess, which are typically expressed using the Gell-Mann representation of  $SU(3)$ . With this basis, each gluon is linearly independent, and no combination of gluons can be used to form a color singlet state. This non-zero charge of the force carrier makes QCD significantly

more complex than QED. In fact, carrying color charge means gluons can also interact with each other directly, leading to certain theoretical states such as glueballs.

#### 2.1.4 Elementary Particles

There are two primary groups contained in the Standard Model, fermions and bosons. This division is based on the Spin Statistics theorem, where fermions (bosons) have half-integer (integer) spins. As described by the Pauli Exclusion principle, nature restricts fermions from occupying the same quantum state. Bosons, however, do not have this restriction, and can have any number occupying the same state.




Figure 2.1: The standard model of particle physics.

It is comprised of two main groups: fermions, which includes the quarks and leptons, and bosons, which includes the gauge bosons and the Higgs boson. Image reproduced courtesy of [7].

## Fermions

The fermions are divided by their interaction types into two major groups, quarks ( $q$ ) and leptons ( $l$ ). Each of these groups contains six particles with their corresponding antiparticles. These can be grouped into three generations, which aligns particles with the same electric charges, but greatly differing masses. As an example, the up ( $u$ ), charm ( $c$ ), and top ( $t$ ) quarks all have an electric charge of  $+2/3$  (in terms of the electron charge,  $e$ ), but  $t$  is approximately five orders of magnitude more massive than  $u$ . For the quarks and fermions in Figure 2.1, rows indicate particles with the same electric charge, while columns represent each generation of particles.

Although all fermions interact both electromagnetically and weakly, only the quarks interact strongly. Because of confinement, quarks cannot exist as isolated particles, and are only found in nature as groups of particles called hadrons. The most common types of hadrons exist as quark-antiquark pairs, known as mesons, or as groups of three quarks (or antiquarks), known as baryons. There are, however, indications of more exotic combinations of quarks, such as tetra- ( $qq\bar{q}\bar{q}$ ) or penta-quark ( $qqqq\bar{q}$ ) states seen by recent experiments [10, 11, 12].

While the negatively charged quarks ( $d$ ,  $s$ , and  $b$ ) are labeled as definite states, the quarks are actually mixed states. Through weak interactions, each of these quarks can transform into other quarks. The probabilities for these transformations are expressed by the Cabibbo-Kobayashi-Maskawa (CKM) Matrix [13], shown in Figure 2.2. From the experimentally measured values [6], it is evident the matrix is nearly diagonal. It is also clear that correlations are strongest within each generation, as the off-diagonal terms are generally smaller than the diagonal ones. Additionally, though the convention splits the negatively charged quarks into mixed states (leaving the positively charged quarks fixed), this choice has no physical basis. The reverse choice of having mixed positively charged quarks is equivalent.

$$\begin{bmatrix} |V_{ud}| & |V_{us}| & |V_{ub}| \\ |V_{cd}| & |V_{cs}| & |V_{cb}| \\ |V_{td}| & |V_{ts}| & |V_{tb}| \end{bmatrix} = \begin{bmatrix} 0.97427 \pm 0.00014 & 0.22536 \pm 0.00061 & 0.00355 \pm 0.00015 \\ 0.22522 \pm 0.00061 & 0.97343 \pm 0.00015 & 0.0414 \pm 0.0012 \\ 0.00886^{+0.00033}_{-0.00032} & 0.0405^{+0.0011}_{-0.0012} & 0.99914 \pm 0.00005 \end{bmatrix}$$

Figure 2.2: The Cabibbo-Kobayashi-Maskawa (CKM) Matrix.

The values of the CKM matrix are typically parameterized using three Euler angles

$(\theta_{12}, \theta_{23}, \theta_{13})$  and a CP-violating phase parameter  $(\delta_{13})$ , where the indices represent the three generations of quarks. This formulation allows the matrix to be cast in the “standard” parametrization, shown in Figure 2.3. The form with three separated matrices clearly shows the connections between each generation of quarks. Namely, the third shows the original formulation in terms of a single rotation, the Cabibbo angle  $(\theta_{12})$ . This theory is known as the Glashow-Iliopoulos-Maiani (GIM) mechanism [[14] and was used to explain the suppression of flavor-changing neutral currents (FCNC) before the discovery of the charm quark.

$$\begin{aligned} & \begin{bmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{bmatrix} \begin{bmatrix} c_{13} & 0 & s_{13}e^{-i\delta_{13}} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta_{13}} & 0 & c_{13} \end{bmatrix} \begin{bmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta_{13}} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta_{13}} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta_{13}} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta_{13}} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta_{13}} & c_{23}c_{13} \end{bmatrix} \end{aligned}$$

Figure 2.3: The standard form of the Cabibbo-Kobayashi-Maskawa (CKM) Matrix. The parameterization is in terms of three angles  $(\theta_{12}, \theta_{23}, \theta_{13})$  and a phase angle  $(\delta_{13})$ .

Here,  $c_{ij} = \cos \theta_{ij}$  and  $s_{ij} = \sin \theta_{ij}$ .

The leptons are also organized into generations consisting of particles with two distinct charges. The electron ( $e^-$ ), muon ( $\mu^-$ ), and tau ( $\tau^-$ ) are all negatively charged particles. There is also a neutral particle, a neutrino ( $\nu$ ), corresponding to each of the charged leptons ( $\nu_e, \nu_\mu, \nu_\tau$ ). These are very small mass ( $< 1$  eV) particles with extremely low interactions. With the exception of mass, the interaction properties of each flavor is very similar. However, the three flavors themselves are treated as separate conserved quantities.

The Standard Model assumes neutrinos to be massless particles. However, this was violated by the discovery of neutrino oscillations, where transformations occur between neutrino flavor states due to differences in their masses. As with the quarks, the flavor states,  $\nu_e, \nu_\mu$ , and  $\nu_\tau$ , are not the states observed in nature. Rather, the states with definite mass, labeled  $\nu_1, \nu_2$ , and  $\nu_3$ , are linear combinations of the three flavor states. This can be expressed in a rotation of bases called the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix [15, 16]. Its formulation is analogous to the CKM Matrix for quarks.

## Bosons

For each of the three forces included in the Standard Model, there are accompanying gauge bosons. These are the photon ( $\gamma$ ) for electromagnetic force, the  $W^\pm$  and  $Z$  for the weak force, and the gluon ( $g$ ) for the strong force. Each of the gauge bosons is a spin-1 vector boson, which means there are three available polarization states (-1, 0, +1). However, since the photon and gluon are both massive, gauge invariance requires these to have transverse polarizations. This means the spin-0 state is eliminated, and there are only two polarization states for each. There is also the Higgs boson ( $H$ ), which unifies the electromagnetic and weak forces, and whose interactions with other particles is responsible for their mass. This is the only known fundamental spin-0 particle, which means it has only one polarization state.

Even with the amazing success of the Standard Model, the theory is not complete. Along with neutrino oscillations, other effects, such as dark matter and dark energy, remain major obstacles to constructing a unified theory. Such a theory must also include gravity, but there remain significant difficulties in explaining its effects through a quantum field theory. There also remains no conclusive explanation for various constants, such as the masses of each fundamental particle. Still, the Standard Model remains the most precise description of the universe to date, and continues to provide the basis for much current and future experimental and theoretical work.

## 2.2 Charmonium

The majority of this analysis focuses on a specific group of particles known as Charmonium. These particles are resonances formed by a  $c\bar{c}$  pair, and can be treated analogously to the hydrogen atom. Namely, there is a spectrum of various excited states in the Charmonium region, just as the spectrum of states associated with the emission lines of hydrogen. The first three charmonium states to be discovered were the  $J/\psi$ ,  $\psi'$ , and  $\psi''$ . The ' and '' marks indicate these are the first and second excited states of the  $J/\psi$ , respectively. More commonly, the  $\psi'$  is denoted  $\psi(3686)$  and the  $\psi''$  is denoted  $\psi(3770)$ . The numbers in parentheses represent the mass of the particle in MeV.

An alternative labeling scheme for these states uses the quantum numbers for each particle. This is written in the form  $N^{2s+1}L_J$ , where  $N$  refers to the principal quantum

number,  $s$  refers to the total spin angular momentum of the particle,  $L$  refers to the orbital angular momentum, and  $J$  refers to the total angular momentum. Here, the values of  $L$  are in spectroscopic notation, where  $L = 1, 2, 3, 4 \dots$  is denoted  $S, P, D, F \dots$ , and higher values follow alphabetically (excluding  $J$ ). As each of these states is comprised of two spin- $\frac{1}{2}$  particles, the value of  $s$  in this case can only be 0 (opposite) or 1 (aligned). With this, the  $J/\psi$ ,  $\psi(3686)$ , and  $\psi(3770)$  can be denoted  $1^3S_1$ ,  $2^3S_1$ , and  $1^3D_1$ . The values of  $n$  and  $L$  are used for the alternate notation of  $\psi(2S)$  representing  $\psi(3686)$ . However, the notation of  $\psi(1D)$  is not often used for  $\psi(3770)$ . This is due to evidence of mixing in  $\psi(3770)$  between the  $2^3S_1$  and  $1^3D_1$  states that suggests more complicated underlying interactions [17, 18].

In fact, while the comparisons from this model work well for states less massive than the  $\psi(3770)$ , the predictions made above this often break down. This is likely based on the energy required to produce open-charm  $D$  mesons, such as  $D^+(c\bar{u})$  and  $D^0(c\bar{d})$ . The  $D\bar{D}$  threshold (twice the mass of the  $D^0$ ) is just above the  $\psi(2S)$  mass, and just slightly below the  $\psi(3770)$  mass. Therefore, the decay products of the two particles are drastically different, even while the available phase space is similar. Example Feynman diagrams for these two particles can be seen in Figures 2.4 and 2.5.




Figure 2.4: An example Feynman diagram for the decay of  $\psi(3686)$ . Without sufficient energy to produce  $D$  mesons, the decays of  $\psi(3686)$  must be mediated by three hard gluons and are suppressed, as described by the OZI rule.




Figure 2.5: An example Feynman diagram for the decay of  $\psi(3770)$ . With sufficient energy to produce  $D$  mesons, the open-charm decays of  $\psi(3770)$  are allowed to proceed, greatly increasing the total decay width.

The difference is also clearly seen in the total decay widths, where the most recent experimental averages [6] are  $\Gamma^{\psi(2S)} = 286 \text{ keV}$  and  $\Gamma^{\psi(3770)} = 27.5 \text{ MeV}$ . An explanation for this discrepancy was proposed independently in the 1960s by Okubo [19], Zweig [20], and Iizuka [21], and is named the OZI rule. This states that any Feynman Diagram where the initial and final particles are separated at some point by only gluons represents a suppressed decay. Such behavior requires that the momentum transfer from the initial particles must occur entirely through these gluons. Because of the decreasing strength of the strong interaction with higher momentum transfer, the rate of these decays is thereby inhibited. This is further compounded by the need for three gluons in such an interaction, as one gluon could not conserve color charge, and two could not converse C-parity. Once above the  $D\bar{D}$  threshold, the allowed open-charm decays dominate, and the total width is greatly increased. This dominance points to a high branching fraction expected for decays of the type  $\psi(3770) \rightarrow D\bar{D}$ .

## Chapter 3

# Detector and Related Systems

All data used for this analysis were collected at the third Beijing Spectrometer (BESIII), located in Beijing, China, at the Institute of High Energy Physics (IHEP) campus. This detector records  $e^+e^-$  collision events provided by the second Beijing Electron-Positron Collider (BEPCII). The center-of-mass energy range for this facility was selected to concentrate on  $\tau^+\tau^-$  and  $c\bar{c}$  production, from about 2.0 GeV to 4.6 GeV. Both of these machines are upgrades from previous versions built on the same sites. The first BEPC and BES were originally constructed in 1989, while the upgrade to BESII occurred in 1996. Their operation was terminated in 2004 to prepare for the upgrades to the current systems.

In 2009, BEPCII and BESIII began operation with the goal of utilizing greatly increased luminosity. For example, instead of the single-bunch electron collisions of BEPC, the new design utilized multiple bunch collisions. This creates many groups of electrons and positrons which are tightly packed in each ring during run time. BEPCII also utilizes a dual-storage ring for the electrons and positrons, compared to the single-ring available at BEPC. The improvements provide BEPCII with a design luminosity of  $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ , two orders of magnitude larger than the previous installation. This luminosity is optimized for energies near the  $\psi(3770)$  resonance, as BESIII conducts many precision measurements and rare decay searches around this region. A detailed description of the BESIII detector can be found in Ref. [24].

### 3.1 BEPCII Accelerator

The setup for collisions in BEPCII begins with bombarding a fixed target with electrons in a linear accelerator. This generates high energy photons which interact with the target material to form  $e^+e^-$  pairs. The positrons from these pairs are then separated magnetically. These positrons are then injected into the desired storage ring until they reach the desired beam current. As electrons do not need to be created in this manner, they are instead directly accelerated in the linear accelerator and injected into the opposite storage ring. These injections occur at a rate of 50 mA/min for positrons and 200 mA/min for electrons.

To achieve the necessary high beam currents, many bunches of electrons and positrons are packed into the evacuated rings. During operation, each ring contains 93 bunches of length 1.5 cm separated by 8 ns (2.4 m). These provide a maximum beam current of 0.91 A while operating in collision mode. At the interaction point, each beam is focused using super-conducting quadrupole magnets to compress the beam size to about 5.7  $\mu\text{m}$  vertically, while the horizontal beam size is about 380  $\mu\text{m}$ . For collisions, each beam is also angled towards the center of the storage rings with an angle of 11 mrad. This crossing angle introduces a net momentum in the radial direction to help prevent against tracks which may continue in the direction of the beam pipe, thereby avoiding detection.

For a normal run, collisions continue occurring until the instantaneous luminosity falls below useful levels. While this is typically depleted due to the collisions between the  $e^+e^-$  particles, other unwanted interactions (such as those with residual gas in the storage rings) also reduce these currents. When this happens, BEPCII can replenish the beams using top-off injections. This allows the collider to continue utilizing the remaining particles within the storage rings without dumping the beams completely. Recycling these leftover electrons and positrons saves considerable time, and allows for more efficient data taking.

### 3.2 BESIII Detector

Centered around the interaction point of BEPCII, the BESIII detector records information about the particles produced by the  $e^-e^+$  collisions. Each collision occurs

within the beam-pipe of the detector, which has an inner and outer radii of 31.5 mm and 57.0 mm, and is evacuated to  $5 \times 10^{-10}$  Torr. Most of the apparatus is a region of uniform 1.0 T magnetic field provided by a super-conducting solenoid with a mean radius of 1.482 m and a length of 3.53 m. The field points in the  $z$ -direction, which is along the direction of the  $e^+$  beam. The  $x$ -direction points towards the center of the storage rings, while the  $y$ -direction is vertically upwards. This magnetic field is used to allow measurement of the momenta of charged particles based on the curvature of their trajectories. It allows typical charged particles to sufficiently interact with much of the tracking volume, while minimizing those which curl too much to reach all layers of the detector.




Figure 3.1: A schematic of the BESIII detector.

There are four main layers surrounding the beam-pipe: the Main Drift Chamber (MDC), the Time-of-Flight (ToF), the Electromagnetic Calorimeter (EMC), and the Muon Chamber (MUC). All but the MUC are surrounded by a uniform 1.0 T magnetic field produced by the super conducting solenoidal magnet.

The BESIII detector, shown in Figure 3.1, consists of four main components which address different aspects of measuring and identifying particles. Starting from the most

interior, these layers are the Multi-Layer Drift Chamber (MDC), the Time-of-Flight System (ToF), the Electromagnetic Calorimeter (EMC), and the Muon Identifier (MUC). Using the information provided by each of these, the tracks seen in the detector are given a particle hypothesis for their most likely identity. Only charged particles stable enough to traverse the detector volume are identifiable in this way. The charged particles identified at BESIII are electrons ( $e$ ), muons ( $\mu$ ), pions ( $\pi$ ), kaons ( $K$ ), and protons ( $p$ ). Short-lived particles, such as  $D^0$  and  $D^+$ , must be reconstructed from their decays into these constituents, as well as neutral shower energy from photons ( $\gamma$ ).

### 3.2.1 Multi-Layer Drift Chamber

The purpose of the Multi-Layer Drift Chamber is to determine the momenta and trajectories of charged particles. Because of the magnetic field, charged particles travel in helical trajectories. The direction of curvature is used to determine the charge, while the radius of curvature of the track is used to determine the momentum.

The MDC is comprised of many layers of tungsten sense wires to detect the ionization of particles which pass through its gas-filled volume. The sense and field wires create an electric field which causes ionization electrons to drift towards the sense wires. This field is tuned to a strength which minimizes secondary ionization, except in the immediate vicinity of the sense wire. This produces an avalanche of secondary ionizations which creates a measurable current pulse in the sense wire. The amount of energy deposited by this process is proportional to the primary ionization produced by the track. Tracing the path of energy depositions over time allows for the reconstruction of each charged particle trajectory.

The main design of the MDC focuses on cylindrical layers of drift cells comprised of sense wires running coaxial to the beam pipe. The inner and outer radii of the MDC are 59 mm and 810 mm, respectively. There are 43 layers of sense wires which cover 93% of the  $4\pi$  solid angle in the detector. These include 8 layers in the inner chamber, and 35 in the outer chamber. Each of the layers in the inner chamber are stereo layers (for measuring along the  $z$ -axis), while the 16 stereo layers and 19 axial layers (for measuring along the  $x-y$  plane) are interleaved in the outer chamber. This arrangement provides position resolutions of 130  $\mu\text{m}$  and 2 mm in the  $r-\phi$  plane and beam direction, respectively, for each hit. The uncertainty in the transverse plane measurements is

dominated by electron diffusion and the readout time uncertainty for the electronics. For the transverse momentum, the resolution is about 0.5 % for tracks with momenta of 1 GeV, with uncertainties coming mainly from wire position measurements, and multiple scattering from material in the MDC.

The gas used for ionization is a mixture of 60% helium (He) and 40% methane ( $C_3H_8$ ). Helium, because of its low atomic number, and thus, long radiation length, also minimizes multiple scattering which degrades the momentum resolution. Methane, with extra rotational and vibrational degrees of freedom not accessible to helium, quenches the ionization energy. Without this effect, electrons would be accelerated by the electric field, produced secondary ionization energy, and lead to electric breakdown.

In addition to trajectory, the MDC also measures the rate of energy loss over distance for a particle traveling through a material [25], as described by the Bethe-Bloch equation,

$$-\frac{dE}{dx} = 4\pi N \frac{z^2 e^4}{m_e \beta^2} \left[ \log \left( \frac{2m_e \beta^2}{I(1 - \beta^2)} \right) - \beta^2 \right], \quad (3.1)$$

where  $N$  is the electron number density of the material,  $z$  is the charge of the particle in terms of  $e$ , the charge of the electron,  $m_e$  is the mass of the electron,  $\beta$  is the velocity of the particle, and  $I$  is the mean excitation potential for electrons in the material being traversed. The resolution of  $dE/dx$  in the MDC is about 6 % for particles incident 90° to the beam-axis. The uncertainty is due to fluctuations in the number of primary ionizations along the flight path, fluctuations in the avalanche process, as well as from edge effects on each cell.




Figure 3.2: MDC energy deposition for various particles as a function of momenta. Distinction between  $K$  and  $\pi$  tracks is easier for low momenta, but becomes very difficult for higher values.

The energy deposition provides a method of determining particle identity, as this quantity depends on the velocity of the particle. An example of this behavior for multiple types of particles can be seen in Figure 3.2. To identify a particle from the various candidates, the measured energy deposition ( $dE/dx_{\text{meas}}$ ) is compared against the expected value ( $dE/dx_{\text{exp}}$ ) of each hit used to reconstruct the particle's trajectory ( $i$ ):

$$\chi^2 = \sum_i \chi_i^2 = \left( \frac{dE/dx_{\text{meas}} - dE/dx_{\text{exp}}}{\sigma} \right)_i^2, \quad (3.2)$$

where  $\sigma$  represents the uncertainty on the measured energy deposition. This process provides a separation of  $3\sigma$  between  $K$  and  $\pi$  tracks with momenta up to 770 MeV.

### 3.2.2 Time-of-Flight System

The purpose of the Time-of-Flight System is to determine the velocity of charged particles. This is useful for distinguishing particles with similar momenta, but different masses, as shown in Figure 3.3. It uses information provided by the MDC to determine the probability for each charged track to match the possible particle hypotheses. Namely, this includes the measured momentum, the expected time interval based on its

trajectory, and the mass for each particle hypothesis. This process provides a separation of  $3\sigma$  between  $K$  and  $\pi$  tracks with momenta up to 900 MeV.




Figure 3.3: ToF measurements for various particles as a function of momenta. Distinction between  $K$  and  $\pi$  tracks is very easy for low momenta, but becomes more difficult for higher values.

The ToF is comprised of two bands of staggered plastic scintillators attached to photomultiplier tubes (PMTs). These two bands, located at 0.81 m and 0.86 m from the beam-pipe, measure a time difference used to determine the speed of each charged particle. The resolution is about 100 ps, and is largely limited by the scintillation light rise time, as well as fluctuations associated with the PMTs. The layer is split into two regions, barrel and endcap, which cover the ranges  $|\cos \theta| < 0.82$  and  $0.85 < |\cos \theta| < 0.95$ , respectively. The former is dual-layer with each containing 88 scintillators of 5 cm thickness arranged in a trapezoidal cross section, while the latter contains two single layers of 48 fan-shaped scintillators. Between the two are support structures for the MDC as well as other service lines.

### 3.2.3 Electromagnetic Calorimeter

The purpose of the Electromagnetic Calorimeter is to measure the energy deposited by photons and electrons. Since most of the candidates identified in the detector are

relativistic, they are minimum ionizing particles. This causes each to deposit a relatively constant value of energy, independent of the measured momenta. However, electrons, with their extremely small mass, will deposit significant amounts of energy due to Bremsstrahlung radiation and subsequent secondary  $e^+e^-$  pair production. This provides a clear distinction in the detector between electron and muon tracks above 200 MeV, as seen in Figure 3.4. Energy measurements from the EMC are also useful for identifying neutral particles which decay only to photons, such as  $\pi^0$ .




Figure 3.4: EMC energy deposition for various particles as a function of momentum. As the mass of  $e^-$  is so small, it deposits virtually all of its energy in the EMC. Both  $\mu^-$  and  $\pi$  are generally minimum ionizing particles in this momenta range, and their similar masses make them difficult to distinguish with the EMC measurements alone.

The EMC is comprised of tellurium-doped cesium iodide ( $\text{CsI}(\text{Tl})$ ) crystals with square front faces attached to a photodiode. Each of the 6240 crystals is 5.2 cm long on the square edges and 28 cm (15 radiation lengths) deep. To prevent photons from aligning with the gaps between each crystal, each one is offset with a tilt of  $1.5^\circ$  in the  $\phi$ -direction and  $1.5^\circ$  to  $3^\circ$  in the  $\theta$ -direction. These crystals provide an energy resolution ( $\sigma/E$ ) of 2.5% at 1 GeV and 4% down to 100 MeV. This is limited by energy not deposited over the length of the crystal, the gaps between crystals, and by non-uniform light production. Additionally, only measurements of energy above 20 MeV are considered, because of the irreducible noise level in the detector. The position resolution is  $\sigma = 0.6 \text{ cm}/\sqrt{E \text{ [GeV]}}$ , and is primarily limited by the crystal segmentation. The

EMC has an inner radius of 94 cm and a total weight of approximately 24 tons. It covers the regions  $|\cos\theta| < 0.83$  (barrel) and  $0.85 < |\cos\theta| < 0.93$  (endcap), with a gap in between that does not provide reliable energy measurements.

### 3.2.4 Muon Identifier

The purpose of the Muon Identifier is to determine the likelihood of a charged particle being a muon. Since electrons are significantly lower mass, they deposit virtually all of their remaining energy in the EMC. Additionally, since muons do not interact strongly, they will penetrate notably further than will pions, kaons, or protons. This provides a clear indication of a muon when a particle penetrates the MUC. However, due to the magnetic field, only muons with  $p > 0.4 \text{ GeV}$  can traverse deep enough to be identifiable.

The MUC is comprised of resistive plate counters (RPC) which are interspersed between the steel plates of the super-conducting solenoid's flux-return iron. Each of the steel layers increase in thickness working outwards from the center: 3 cm, 3 cm, 3 cm, 4 cm, 4 cm, 8 cm, 8 cm, 8 cm, and 15 cm. Like the other components, the MUC is split into a barrel and an endcap region. The barrel has nine RPC layers of 4 cm thickness. In the endcap, the first RPC layer is after the first steel layer, leaving only eight RPC layers. Each of these layers has RPC strips oriented along only one direction. For the barrel, the  $z$  ( $\phi$ ) orientation is read out for only the odd (even) layers. Conversely, the endcap only reads out the  $x$  ( $y$ ) orientation in the odd (even) layers.

## 3.3 Triggering Systems

In order to maintain a high efficiency for selecting physics events, many backgrounds must be filtered out. At BESIII, this is done through a triggering system with two-tiers: a hardware trigger (L1) and a software event filter (L3). This process is illustrated in Figure 3.5. The filtered background events are primarily from beam-related sources, such as beam-gas or beam-wall interactions, and occur at a rate of about 13 MHz. To assist with this process, collimators and masks are used to prevent lost electrons from interacting with the detector. However, there are also other sources of backgrounds, such as cosmic rays, which occur at a rate of about 1.5 kHz. The total backgrounds must be suppressed to a rate which does not overwhelm the recording of events by

the readout systems. This rate is roughly 2 kHz at the  $J/\psi$  peak, and 600 Hz for the  $\psi(2S)$  when running near peak luminosity. For Bhabha events ( $e^+e^- \rightarrow e^+e^-$ ), which are used for calibration and luminosity measurements, this rate is 800 Hz within the detector acceptance.




Figure 3.5: Triggering systems for event filtering at BESIII.

To prevent overloading the event recording system, non-physics background events must be eliminated from the data stream. This processing uses simple read-out information from each of the layer of the detector to quickly and efficiently determine whether or not events should be further considered for analysis.

The first step (L1) reads out every clock cycle (24 ns) at a rate of 41.65 MHz. It uses information from the MDC, ToF, and EMC collectively to reduce the rates of beam-related backgrounds to 1.84 kHz and cosmic rays to about 200 Hz. However, L1 has a maximum rate of about 4 kHz. Because of this, when the buffer holding the subdetector data is around 80 % full, L1 triggering is halted until the buffer drops below 10 % full.

From the MDC, L1 gathers information about each charged track. The main parameter examined is the number of superlayers a track passed through. Here, a superlayer is the collection of wires at the same radial distance away from the center of the detector. Tracks are defined as ‘short’ if they deposit energy in segments of superlayers 3–5, or ‘long’ for superlayers 3–5 and 10. To ensure a sufficient momentum to reach the outer superlayers while originating at the interaction point, a minimum transverse momentum cut is applied to each track. This cut is 90 MeV and 120 MeV for short and long tracks, respectively. In addition to the numbers of short and long tracks for an event, the information about back-to-back tracks is also used.

From the ToF, L1 gathers information about the number of hits in the barrel and end-cap regions. It also examines the number of back-to-back hits in each of the two regions. Here, ‘back-to-back’ is defined as having hits within a range of 9 counters on the opposite side of the detector.

From the EMC, L1 gathers information about the clustering of energies around a local maximum-energy crystal. This includes the number of isolated clusters, as well as the information about back-to-back hits in the barrel and end-cap. Additionally, the balance of energy in the  $\phi$ -direction (barrel) and in the  $z$ -direction (endcap) is also used.

The subdetector information gathered during L1 is then passed off to an online computer farm (L3) where the event is assembled. This step reduces backgrounds from a rate of about 2 kHz to about 1 kHz. Combined with the signal rate at the  $J/\psi$  peak (2 kHz), this corresponds to a total maximum event rate of 3 kHz, or a tape write speed of 40 MB/s.

# Chapter 4

## Analysis Software

### 4.1 BESIII Offline Software System

Reconstructing and processing event data gathered by the BESIII detector is done using the BESIII Offline Software System (BOSS) [26]. This is an analysis software distribution written using the C++ language and running primarily on the Scientific Linux CERN operating system [27]. There are five main parts to BOSS: framework, simulation, reconstruction, calibration, and analysis.

#### 4.1.1 Framework

The framework is built on the Gaudi software architecture [28], which provides a standard interface and utilities for things such as event simulation, data processing, and physics analysis. The software is managed using the Configuration Management Tool [29], which provides a method for creating packages, handling package dependencies, and producing executables from source code. There are three main filetypes for data stored by the framework: raw data (`.raw`), reconstructed data (`.rec`), and Data-Summary-Tape (`.dst`). The latter two of these file types are derived from the ROOT [30] format (`.root`) for easy management and usage in various analyses.

### 4.1.2 Simulation

There are four main parts to the simulation process: event generation, detector description, particle tracking, and detector response. Event generation is primarily handled by the Monte Carlo (MC) generators KKMC, BesEvtGen, and Babayaga, which are described below. To model its geometry and materials, a unique description of the detector has been created using a format based on XML. This allows both simulation and reconstruction packages to appropriately model the behavior of events within the specific environment of BESIII. For particle tracking, interactions with detector materials are handled by GEANT4 [31]. Lastly, detector responses are modeled by the so-called ‘digitization code’. This takes into account each detector component, as well as readout electronics, and realistic situations such as noise or dead channels. There is also a simulation of the triggering system implemented.

#### **KKMC**

Originally developed for the LEP and SLC colliders, KKMC [32] is a generator used to model electroweak interactions. Namely, the processes generated are of the form  $e^+e^- \rightarrow f\bar{f} + (n)\gamma$ , where  $f = \{\mu, \tau, u, d, s, c, b\}$ , and  $(n)\gamma$  represents any number of additional photons. These are modeled taking into account second-order sub-leading corrections, as well as initial-state radiation (ISR), and interference between initial- and final-state radiation (FSR). The effects of beam energy spread, typically on the order of 1 MeV near the  $\psi(3770)$ , can also be included.

After generation, the  $f\bar{f}$  pair is decayed by models depending on the fermions involved. The TAUOLA library [33] is used to decay  $\tau^+\tau^-$  pairs, and takes into account spin-polarization effects. The PYTHIA model [34] is used to hadronize final-state  $q\bar{q}$  continuum production using the parton shower model. For resonances like the  $\psi(3770)$ , the only action performed by KKMC is the generation of ISR. After this, the virtual photon produced is handed off to BesEvtGen.

#### **BesEvtGen**

Originally developed for the CLEO and BaBar collaborations, EvtGen [35] is another widely used generator. It is the basis for BesEvtGen [36], which incorporates many

different decay models into a single utility. Over 30 exclusive decay models are available in BesEvtGen, as well as the capability to incorporate user-created models.

The simulation process occurs sequentially using dynamic information from decay amplitude probabilities and forward / backward spin-density matrices. From this, final state radiation is handled by the PHOTOS model [37]. To generate unknown decays of charmonium resonances, the LundCharm model [38] is used, while other unknown hadronic decays are handled by PYTHIA. For radiative processes, such as radiative return to  $J/\psi$  or  $\psi(2S)$ , the VECTORISR model [39] is used. This occurs when one particle in the initial  $e^+e^-$  pair radiates a photon of high enough energy that only lower mass resonances can be produced from the reduced center-of-mass energy. When the radiation is less intense, the  $\psi(3770)$  resonance is directly produced through the combination of KKMC and BesEvtGen.

### Babayaga

Production of QED processes is done using the Babayaga generator [40]. This includes  $e^+e^- \rightarrow \{e^+e^-, \mu^+\mu^-, \gamma\gamma\}$ . The results are of very high precision, with an estimated theoretical accuracy of 0.1 %. It also matches exact next-leading-order corrections from the parton shower algorithm. The high precision also helps determine the efficiencies and acceptances required to precisely measure the integrated luminosities of each data sample.

#### 4.1.3 Reconstruction

Reconstruction primarily involves information about specific types of particles from each of the four main detector layers. These sources of information are as follows:

- a charged track finding algorithm and a Kalman-filter-based track-fitter
- a particle identifying algorithm based on  $dE/dx$  and time-of-flight measurements
- a shower- and cluster-finding algorithm for EMC energy and position
- a muon track finding algorithm

Further descriptions about each of these processes can be found in Sec. 4.2. Additionally, algorithms for determining the corresponding beam bunch crossing, as well as for secondary vertex and track refitting, are also utilized.

#### 4.1.4 Calibration

To maintain consistent production and analysis of datasets, a centralized source of run-dependent information is maintained by BOSS. This includes algorithms which determine the calibration constants for each sub-detector, as well as a centralized database to store the results. Each of the calibration outputs are stored in a ROOT file along with other details such as the beam energy, luminosity, magnetic field information, trigger conditions, and hardware / software versions. While all of this information is stored by a central MySQL [41] server at IHEP, databases for other institutions in BESIII regularly synchronize with this server to create a mirrored copy of these values.

### 4.2 Detector Simulation

The following sections detail the simulation, calibration, and reconstruction processes for each layer. Each of these relies on a geometry description created using GEANT4.

#### 4.2.1 Multi-Layer Drift Chamber

Simulating events in the MDC accounts for the axial layers and endplates, the stereo layers, and the stereo cells. The simulation also heavily relies on the calibration parameters to determine things such as wire efficiency and resolution as a function of drift distance for each wire, noise in each layer, and other misalignment issues.

Calibration of the MDC uses  $J/\psi \rightarrow \mu^+ \mu^-$  events to determine position and  $dE/dx$  measurements. Using  $J/\psi$  events allows for quickly obtaining sufficient statistics due to the very large production cross section at the peak. The information determined includes constants such as  $x - t$  relations, timing, alignment, and absolute wire efficiency. Each of these values are stored in the database for each run. Additionally, turning off the magnetic field could allow for precise determination of wire positions.

Reconstructing MDC events starts by finding axial track segments using raw hits. Each of these are found by searching for pre-calculated patterns. Next, these segments

are linked to circular tracks by a circular fit using the least-squares method. Stereo segments are then added using an iterative helix fit. Lastly, additional hits which were possibly missed from the initial reconstruction are applied using a Kalman-filter process. This process also examines the tracks over multiple particle hypotheses. The reconstruction is remarkably efficient, with over 98 % of tracks with  $p_T > 150 \text{ MeV}$  being reconstructed, even amidst high backgrounds. From this, the charge, momentum, and trajectory can be determined for each track.

In addition to tracking, the MDC also analyzes  $dE/dx$  for each particle. This calculates the energy deposition of each track as it passes through the chamber, and corrects the measured charge amplitudes to determine a probability for each particle hypothesis. Corrections applied account for things such as multiple scatterings, magnetic deflections, and ionization. This likelihood from  $dE/dx$  is used in conjunction with information from the ToF to identify the actual type of particle for each track.

#### 4.2.2 Time-of-Flight System

Simulating events in the ToF accounts for the scintillator, wrapping materials, and photomultiplier tubes (PMTs). The process converts the energy deposited in the scintillator into photons, then propagates the shape of a photon pulse (rather than individual photons) to the PMTs in order to generate an electronic signal. A discriminator is applied to each pulse to determine the analog-to-digital conversion (ADC) and time-to-digital conversion (TDC) outputs. The algorithm is tested using the results of beam tests, however, subsequently gathered data sets require updated tuning. A full simulation tracing each optical photon can also be used for further details on the timing measurement.

Calibration of the ToF also uses  $J/\psi$  decays to leptons to determine both timing and energy measurements. The information determined includes effective velocity, attenuation length, and muon energy loss. The status and performance of the ToF are also regularly monitored by a laser-fiberoptics pulsing system.

Reconstructing ToF events starts by using tracks with trajectories extrapolated from the MDC. Each track is then matched with a particular ToF module; either the two layers of the barrel, or the single-layer endcap. The travel time for each hypothesis is then calculated using a weighted average of results from PMTs at both ends of the scintillator. Corrections are also applied to account for aspects like the effective light

velocity in the scintillator and the light attenuation length. Measurements of  $dE/dx$  are obtained for both charged and neutral particles, and are added back into the EMC in order to improve the shower energy resolution.

#### 4.2.3 Electromagnetic Calorimeter

Simulating events in the EMC accounts for the crystals, casing, silicon photodiodes, preamplifier boxes, cables, and the support system. For each of the crystals and photodiodes, hit information is recorded, and the deposited energy is summed. From this, photon statistics are computed, and the resulting photodiode response is converted into electronic signals. To obtain the waveform in the time domain, an inverse Laplace transform is applied. Then, a sampling and peak searching process is simulated to yield energy and time information. For each bin, Gaussian-type electronic noise is added, and the background is produced by summing over the waveforms.

Calibration of the EMC uses Bhabha electrons with  $E > 1.55 \text{ GeV}$  for the high energy response, and  $\pi^0 \rightarrow \gamma\gamma$  decays for the low energy response. The responses for individual crystals must be analyzed separately, due to their potential intrinsic variations. As a result, they are monitored frequently by a LED light pulser, and periodically recalibrated. Corrections due to temperature variations can also be applied.

Reconstructing EMC events starts by converting the ADC value of each crystal into energy based on the calibration constants. After this, clusters in both the barrel and endcaps are formed by analyzing local maximum energy deposits, called seeds. A clustering algorithm then relates hits around these seeds and sums the values for a particular shower. The positions of each shower is then calculated using energy-weighted first moments. If multiple seeds are found in one cluster, a splitting algorithm is invoked to split the cluster into multiple showers. Additionally, matching energy deposits from the ToF are also added back into the total shower energy. This improves the energy resolution, particularly for low energy photons.

#### 4.2.4 Muon Identifier

Simulating events in the MUC accounts for forming each RPC, creating sets of strips to form each read-out plane, combining each of these with aluminum boxes to form a

muon counter module, and placing the modules alongside iron slabs. The digitization from the read-out planes is selected to fire based off the distance away of each track. Noise is simulated using Poisson distributions initially determined from measurements made during the construction of the chamber, and updated with the taking of actual data.

Calibration of the MUC analyzes RPC detection efficiencies over a function of area. The cluster size and noise levels are also studied.

Reconstructing MUC events starts by searching for collected hits in each of the barrel and endcap orientations. The two collections are then combined with reconstructed tracks from the MDC. Since low momentum muons may cause only a few layers to fire, a subsequent search is performed over unused hits based on the extrapolated trajectories of MDC tracks. The reconstruction process primarily analyzes the depth of the track in the MUC, the maximum number of hits in the layers fired by a track, and the matching between a MDC track with the MUC stand-alone track. These parameters, along with the track momentum and MDC exit angle, are input into an Artificial Neural Network in order to distinguish between hadron and muon tracks. The distinction process is quite effective, generally removing  $\sim 96\%$  of pions and keeping around  $\sim 90\%$  of muons.

### 4.3 *D*-Tagging

From the decay of the  $\psi(3770)$ , the most commonly produced particles are  $D^0\overline{D}^0$  or  $D^+D^-$  pairs. Since the  $D\overline{D}$  pairs produced are two-body decays, the energy available to each  $D$  is half of the center-of-mass energy (in the center-of-mass reference frame). Each of these  $D$  mesons then quickly decays to certain sets of particles. Reconstructing one of these decays requires assembling the right combination of such particles. From energy and momentum conservation, the total 4-momentum of the constituents must match the total energy of the  $D$  meson.

This reconstruction technique is known as ‘*D*-Tagging’, and was pioneered by the MARK-III collaboration [42, 43]. For our analysis, the particles analyzed in the detector include  $\pi^\pm$ ,  $K^\pm$ ,  $\pi^0$ , and  $K_S^0$ , and the decay modes used are shown in Table 4.1. There are three  $D^0$  modes and six  $D^+$  modes, where charge conjugation (converting all particles to their anti-particles) is implied throughout the analysis. The modes used are chosen for

their effectiveness of reconstruction in the detector; they generally have higher branching fractions and minimize multiplicity (the number of tracks used). Additionally, for the neutral modes, the doubly-Cabbibo suppressed decays (DCSD), such as  $D^0 \rightarrow K^+ \pi^-$ , are also included by this procedure.

(0) $D^0 \rightarrow K^- \pi^+$	(200) $D^+ \rightarrow K^- \pi^+ \pi^+$	(203) $D^+ \rightarrow K_S^0 \pi^+ \pi^0$
(1) $D^0 \rightarrow K^- \pi^+ \pi^0$	(201) $D^+ \rightarrow K^- \pi^+ \pi^+ \pi^0$	(204) $D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-$
(3) $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$	(202) $D^+ \rightarrow K_S^0 \pi^+$	(205) $D^+ \rightarrow K^+ K^- \pi^+$

Table 4.1: The reconstructed  $D$ -tag modes used in this analysis.

This process occurs for each event and searches over each decay mode for each charm (i.e., both  $D^+$  and  $D^-$ ). The combinations chosen for reconstruction are those with the smallest energy difference from the expected value. More than one  $D$  combination can be extracted from a given event, as long as it satisfies all other requirements (see Section 4.3.1). While this may sound like it overestimates the number of actual  $D$  particles found, the process is also used to calculate reconstruction efficiency, thereby cancelling out this effect.

### 4.3.1 Selection Cuts

Before being considered as potential reconstruction candidates, each track in the detector must also pass other cuts specific to its identified particle type. The following describes the necessary criteria required for the particles in the decay modes we are using.

#### $\pi^\pm/K^\pm$ Selection

Each of the reconstructed charged tracks must pass vertex cuts in both the transverse ( $x - y$ ) and beam ( $z$ ) directions relative to the interaction point. This requires tracks originate sufficiently close to the interaction point to ensure they are not other backgrounds, such as cosmic rays, or other tracks which have decayed in flight. There is also a cut on the angle measured within the MDC ( $\theta$ ) to ensure the track has not disappeared down the beam-line (where detection cannot occur). Lastly, from the particle identification process, the probability of being a pion (kaon) must be more likely than being a kaon (pion). The values for each of these requirements can be found in Table 4.2.

Vertex ( $xy$ )	$V_{xy} < 1 \text{ cm}$
Vertex ( $z$ )	$ Vz  < 10 \text{ cm}$
MDC Angle	$ \cos \theta  < 0.93$
Pion Probability	$P(\pi) > 0, P(\pi) > P(K)$
Kaon Probability	$P(K) > 0, P(K) > P(\pi)$

Table 4.2: The required cuts to identify charged tracks as  $\pi^\pm$  or  $K^\pm$ .

### $\gamma$ Selection

To distinguish photon energy from noise, each shower in the EMC is required to have a certain amount of deposited energy. These cuts are different for the barrel ( $|\cos \theta| < 0.80$ ) and endcap ( $0.84 < |\cos \theta| < 0.92$ ) regions. Each photon must also pass a TDC timing cut to ensure they are consistent with actual physics events, and not originating at other times. The values for each of these requirements can be found in Table 4.3.

Minimum Energy (Barrel)	$E_{\text{EMC}} > 25 \text{ MeV}$	$( \cos \theta  < 0.80)$
Minimum Energy (Endcap)	$E_{\text{EMC}} > 50 \text{ MeV}$	$(0.84 <  \cos \theta  < 0.92)$
TDC Timing	$(0 \leq t \leq 14) \times 50 \text{ ns}$	

Table 4.3: The required cuts to identify neutral showers as a  $\gamma$ .

### $\pi^0$ Selection

Reconstructing  $\pi^0$  mesons involves finding  $\gamma\gamma$  pairs, as this is its most dominant decay ( $\sim 99\%$ ). Each of the  $\gamma$  showers used must pass the cuts described above. Additionally, at least one photon in the pair must be found in the barrel region. Each of the two photons are then kinematically fit to compare with the invariant mass of the  $\pi^0$ , and must also pass a proper fit cut. The resulting momentum from this fit is used for reconstructing  $D$ -tag candidates. The values used for each of these requirements are shown in Table 4.4.

Nominal Mass	$115 \text{ MeV} < m_{\pi^0} < 150 \text{ MeV}$
Fit Quality	$\chi^2 < 200, \text{Converged}$

Table 4.4: The required cuts to reconstruct  $\gamma\gamma$  pairs as a  $\pi^0$ .

## $K_S^0$ Selection

Reconstructing  $K_S^0$  mesons involves finding  $\pi^+\pi^-$  pairs, as this is its most common decay ( $\sim 70\%$ ). While  $\pi^0\pi^0$  pairs are also a substantial decay mode ( $\sim 30\%$ ), these are not considered for reconstruction due to the increased difficulty of finding correct  $4\gamma$  sets. To account for the  $K_S^0$  decaying in flight, each of the charged pions considered are not subjected to the vertex or probability cuts in Table 4.2. Instead, the two found pions are kinematically constrained to a common vertex. The results must pass a nominal mass cut ( $\sim 3\sigma$ ) and a proper fit cut to be deemed a  $K_S^0$ . From this, the resulting momentum from the vertex fit is used for reconstructing  $D$ -tag candidates. The values used for each of these requirements are shown in Table 4.5.

Nominal Mass	$487 \text{ MeV} < m_{K_S^0} < 511 \text{ MeV}$
Fit Quality	$\chi^2 < 100$ , Converged

Table 4.5: The required cuts to reconstruct  $\pi^+\pi^-$  pairs as a  $K_S^0$ .

## Cosmic Ray and Lepton Veto

Lastly, when reconstructing the mode  $D^0 \rightarrow K^-\pi^+$ , an additional veto is used. Since the mode has only two charged tracks, it is common to misidentify particles which come from cosmic ray and two-lepton backgrounds. To prevent this, cuts on the timing difference between the tracks, as well as on the particle identification process, are enacted. The values used for each of these requirements are shown in Table 4.6.

Timing (TDC)	$ t_1 - t_2  < 5 \times 50 \text{ ns}$
Particle Identification	$(\chi_{e^-}^2 + \chi_{e^+}^2) - (\chi_{K^-}^2 + \chi_{\pi^+}^2) > 10$

Table 4.6: Cuts to prevent cosmic ray and lepton backgrounds in  $D^0 \rightarrow K^-\pi^+$ .

### 4.3.2 Reconstruction

After each of the constituent particles are identified, a reconstructed  $D$  candidate can be characterized by two main properties:

$$\Delta E = |E_{\text{beam}} - E_{\text{tag}}|, \quad m_{\text{BC}} = \sqrt{E_{\text{beam}}^2 - |\vec{p}_{\text{tag}}|^2}. \quad (4.1)$$

These are the energy difference ( $\Delta E$ ) and the beam-constrained mass ( $m_{\text{BC}}$ ), and effectively represent the energy and momentum of the  $D$ -tag, respectively. As the candidate with the smallest  $\Delta E$  for each decay mode in each event is selected, the values will typically peak near 0 MeV. Distributions of  $m_{\text{BC}}$ , meanwhile, typically peak near  $m_{D^0} = 1.865 \text{ GeV}$  or  $m_{D^+} = 1.870 \text{ GeV}$ . Additionally, while invariant mass ( $m_{\text{inv}} = \sqrt{E_{\text{tag}}^2 - |\vec{p}_{\text{tag}}|^2}$ ) can also be examined,  $m_{\text{BC}}$  is generally preferred due to a higher precision for the beam energy than of the individual particles comprising the  $D$  candidate.

## Chapter 5

# Measurement of $\sigma_{\psi(3770) \rightarrow D\bar{D}}$ near $\psi(3770)$

The main objective of this analysis is to measure the  $\sigma_{\psi(3770) \rightarrow D\bar{D}}$  cross section at center-of-mass energies around the  $\psi(3770)$ . While the cross section very near the  $\psi(3770)$  mass has been measured with excellent precision [44], the shape over a wider energy range has not been measured to a precise degree. This has left open the possibility of multiple functional shapes which could describe the cross section, and asks the question of whether or not to include interference effects. Fits to this region with a single Breit-Wigner shape (i.e., no interference) have provided a unsatisfactory description of the  $\psi(3770)$ , and further necessitate the need for interference. The most prominent candidate for shapes with interference assume a contribution from the lower mass  $\psi(2S)$  state, and is the form considered here. The procedure for this analysis largely follows the methodology of the KEDR collaboration [5]. With the larger statistics available at BESIII, however, an inclusive cross section measurement (such as performed by KEDR) is not required. Instead, we measure the production of  $\psi(3770) \rightarrow D\bar{D}$  events directly.

### 5.1 Derivation of $\sigma_{\psi(3770) \rightarrow D\bar{D}}$

The production rate for a pair of  $D$  mesons coming from  $\psi(3770)$  at a given center-of-mass energy can be calculated following an approach of Kuraev and Fadin [22] applied by the KEDR collaboration [5]. This method also corrects for Initial State Radiation

(ISR), affecting particles accelerated in a collider, and is given by the following:

$$\sigma_{D\bar{D}}^{RC}(W) = \int z_{D\bar{D}}(W\sqrt{1-x}) \sigma_{D\bar{D}}(W\sqrt{1-x}) \mathcal{F}(x, W^2) dx. \quad (5.1)$$

Here,  $W$  is the given center-of-mass energy,  $x$  is an approximation for the fraction of radiated energy, and  $\mathcal{F}(x, W^2)$  is the probability of losing this energy from ISR:

$$\begin{aligned} \mathcal{F}(x, W^2) &= \beta x^{\beta-1} \left[ 1 + \frac{\alpha}{\pi} \left( \frac{\pi^2}{3} - \frac{1}{2} \right) + \frac{3}{4}\beta + \beta^2 \left( \frac{37}{96} - \frac{\pi^2}{12} - \frac{L}{72} \right) \right] = \beta x^{\beta-1} F(W^2), \\ \beta &= \frac{2\alpha}{\pi}(L-1), \quad L = \log \left( \frac{W^2}{m_e^2} \right). \end{aligned} \quad (5.2)$$

The factor  $z_{D\bar{D}}$  includes the expected Coulomb interaction between the mesons in of the charged mode ( $D^+D^-$ ),

$$z_{D^+D^-} = \frac{\pi\alpha/\beta_{D^+}}{1 - \exp(-\pi\alpha/\beta_{D^+})} \times \theta(W - 2m_{D^+}), \quad (5.3)$$

but only accounts for the  $D\bar{D}$  energy threshold in the neutral mode ( $D^0\bar{D}^0$ ),

$$z_{D^0\bar{D}^0} = \theta(W - 2m_{D^0}). \quad (5.4)$$

The theta function imposes the step in the cross section at the production threshold.

The integral in Eq. 5.1 can be simplified by taking advantage of the relatively constant values of  $z_{D\bar{D}}$  and  $\sigma_{D\bar{D}}$  over sufficiently small intervals. By splitting the full  $W$  range into such intervals and integrating over each, this becomes

$$\int \mathcal{F}(x, W^2) dx \approx \sum_{n=0}^N F(W^2) \int_{\frac{n}{N}}^{\frac{n+1}{N}} \beta x^{\beta-1} dx = \sum_n F(W^2) \left[ x_{\text{upper}}^\beta - x_{\text{lower}}^\beta \right]. \quad (5.5)$$

The upper, lower, and mid-point values are given by

$$x_i = \left[ 1 - \left( \frac{2m_D}{W} \right)^2 \right] \left( \frac{n_i}{N} \right), \quad n_i : \begin{cases} n_{\text{lower}} & = n \\ n_{\text{mid}} & = n + \frac{1}{2} \\ n_{\text{upper}} & = n + 1 \end{cases}. \quad (5.6)$$

The bracketed expression in Eq. 5.6 represents the maximum value of  $x$  determined by the theta functions of Eqs. 5.3 and 5.4. To maintain sufficient precision with this interval approximation, the value of  $N = 1024$  is used. Combining this with the other factors in Eq. 5.1, the cross section including the effect of ISR becomes

$$\sigma_{D\bar{D}}^{RC}(W) = \sum_{n=0}^N z_{D\bar{D}}(W') \sigma_{D\bar{D}}(W') F(W^2) \left[ 1 - \left( \frac{2m_D}{W} \right)^2 \right]^\beta \left[ \frac{[(n+1)^\beta - n^\beta]}{N^\beta} \right], \quad (5.7)$$

where  $W' = W\sqrt{1-x_{\text{mid}}}$ . The Born level  $D\bar{D}$  cross section is given theoretically by

$$\sigma_{D\bar{D}} = \frac{\pi\alpha^2}{3W^2} \beta_D^3 |F_D(W)|^2, \quad \beta_D = \sqrt{1 - \frac{4m_D^2}{W^2}}. \quad (5.8)$$

Here,  $\beta_D$  is the velocity of the  $D$  meson in the center-of-mass system. The form factor  $F_D$  represents the contribution of each individual resonant ( $R$ ) component and the total non-resonant ( $NR$ ) component. Each resonant piece is parametrized with a phase angle relative to the non-resonant contribution:

$$F_D(W) = F_D^{NR}(W) + \sum_r F_D^{R_r}(W) e^{i\phi_r}. \quad (5.9)$$

Each resonant contribution to the form factor is modeled by a Breit-Wigner amplitude,

$$F_D^R(W) = \frac{6W\sqrt{(\Gamma_{ee}/\alpha^2)(\Gamma_{D\bar{D}}(W)/\beta_D^3)}}{M^2 - W^2 - iM\Gamma(W)}, \quad (5.10)$$

where  $\Gamma_{ee}$  is the electron partial width, and  $\Gamma(W)$  represents the total width of the resonance with mass  $M$ :

$$\Gamma(W) = \left( \frac{M}{W} \right) \left[ \frac{z_{D\bar{D}}(W) d_{D\bar{D}}(W)}{z_{D^0\bar{D}^0}(M) d_{D^0\bar{D}^0}(M) + z_{D^+D^-}(M) d_{D^+D^-}(M)} \right] \Gamma(M). \quad (5.11)$$

The value of  $\Gamma(M)$  represents the total width at the nominal mass of the resonance. The factors  $d_{D^+D^-}$  and  $d_{D^0\bar{D}^0}$  are the Blatt-Weisskopf damping factors [23] for a vector resonance:

$$d_{D\bar{D}} = \frac{\rho_{D\bar{D}}^3}{\rho_{D\bar{D}}^2 + 1}, \quad \rho_{D\bar{D}} = q_D R_0 = \left( \frac{\beta_D W}{2} \right) R_0. \quad (5.12)$$

Here,  $q_D$  is the  $D$  momentum in the center-of-mass frame, while  $R_0$  represents the radius of the parent particle. The  $D\bar{D}$  partial width listed in Eq. 5.10 is simply the total width rescaled according to  $\mathcal{B}_{nD\bar{D}}$ , the sum of all non- $D\bar{D}$  decay modes of  $\psi(3770)$ :

$$\Gamma_{D\bar{D}}(W) = \Gamma(W) \times (1 - \mathcal{B}_{nD\bar{D}}). \quad (5.13)$$

However, as a simplifying assumption, we use  $\mathcal{B}_{nD\bar{D}} = 0$  throughout this part of the analysis.

## 5.2 Form Factors

With the cross section formula calculated, the description of the form factor contribution (Equation (5.9)) must now be determined. Since we assume the  $\psi(2S)$  resonant contribution is negligible in the energy range of our measurements, the only major resonant contribution is from the  $\psi(3770)$ :

$$F_D(W) = F_D^{\text{NR}}(W) + F_D^{\psi(3770)}(W) e^{i\phi^{\psi(3770)}}. \quad (5.14)$$

Currently, there is no definitive model for the non-resonant term, so we use two alternative parameterizations. The first is a simple exponential model:

$$F_D^{\text{NR}} = F_{\text{NR}} \exp(-q_D^2/a_{\text{NR}}^2), \quad (5.15)$$

where both  $F_{\text{NR}}$  and  $a_{\text{NR}}$  are parameters determined through fitting. The second treatment implements a Vector Dominance Model (VDM). This assumes the interference effects are due to the  $\psi(2S)$  mediating  $D\bar{D}$  production above threshold,

$$F_D^{\text{NR}}(W) = F_D^{\psi(2S)}(W) + F_0, \quad (5.16)$$

and that the effective properties of the  $\psi(2S)$  are similar to those of the  $\psi(3770)$ . The real constant  $F_0$  represents the potential effect of higher resonances, like the  $\psi(4040)$ .

The first term is similar to Equation (5.10), but with a modification to the total width:

$$\Gamma^{\psi(2S)}(W) = \left( \frac{M^{\psi(2S)}}{W} \right) \left[ \frac{z_{D^0\bar{D}^0}(W) d_{D^0\bar{D}^0}(W) + z_{D^+D^-}(W) d_{D^+D^-}(W)}{z_{D^0\bar{D}^0}(M^{\psi''}) d_{D^0\bar{D}^0}(M^{\psi''}) + z_{D^+D^-}(M^{\psi''}) d_{D^+D^-}(M^{\psi''})} \right] \Gamma^{\psi(2S)}(M). \quad (5.17)$$

Without this modification, the mass of the  $\psi(2S)$  would be below the  $D\bar{D}$  threshold, and thus, the vanishing  $z_{D\bar{D}}$  terms would cause a singularity in the width. Therefore, we use the mass of the  $\psi(3770)$  in its place to estimate the effects in this region. While it may behave like the total width in Equation (5.17), the true physical meaning of the parameter  $\Gamma^{\psi(2S)}(W)$  is uncertain. For the radii in Equation (5.12), however, the values used are distinct for each meson:  $R_{\psi(2S)} = 0.75$  fm and  $R_{\psi(3770)} = 1.00$  fm.

## 5.3 Data and Monte Carlo Samples

### 5.3.1 Data Samples

This analysis primarily uses scan data produced by BEPCII and collected by BESIII in 2010 over an energy range of 3.643 GeV to 3.890 GeV. These data are partitioned into 34 center-of-mass energy ( $E_{\text{cm}}$ ) bins of variable size over a range of 3.735 GeV to 3.870 GeV. The bin partitioning was determined from measurements of center-of-mass energies (Section 5.3.2). The total range was chosen to be above the  $D^0\bar{D}^0$  threshold (3.730 GeV) and below the  $D^{*0}\bar{D}^0$  threshold (3.872 GeV). This range includes bins which are below the threshold for  $D^+D^-$  (3.739 GeV), with production beginning in the third bin.

Additionally, there are three higher statistics points used for comparison. These include an ‘On-Peak  $\psi(3770)$ ’ sample of  $2.93 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 3.7732 \text{ GeV}$ , an ‘XYZ-scan’ sample of  $50.54 \text{ pb}^{-1}$  at  $E_{\text{cm}} = 3.810 \text{ GeV}$ , and an ‘R-scan’ sample of  $7.95 \text{ pb}^{-1}$  at  $E_{\text{cm}} = 3.850 \text{ GeV}$ . The first of these is the high statistics measurement very near the  $\psi(3770)$  mass using double-tag reconstruction (both  $D$  and  $\bar{D}$  in a single event) performed by Derrick Toth. The other two samples were analyzed using the same procedure as for the scan data (See Sections 5.4 to 5.6). None of these points are used to determine the final results, as the differences between samples introduce additional systematics which overshadow any statistical improvement. However, these provide

useful comparisons at important energy points along the cross section shape.

### 5.3.2 Center-of-Mass Energy Measurement

To precisely determine the values of center-of-mass energy along the scan data region, events of  $e^+e^- \rightarrow \mu^+\mu^-$  collisions were analyzed to determine their invariant mass ( $m_{\text{inv}}$ ). Assuming minimal radiating energy from the muons, this value is effectively equal to the energy produced by the collider. The selection process for such events is shown in Table 5.1. The statistics of the scan data are too small to precisely determine the energy for individual runs. Therefore, multiple runs were combined together into bins based off the assumption each was taken at nearly identical energies. A comparison of the measured values to the online beam energies is shown in Figure 5.1.

Vertex ( $xy$ )	$V_{xy} < 1 \text{ cm}$
Vertex ( $z$ )	$ Vz  < 10 \text{ cm}$
MDC Angle	$ \cos \theta ^{\mu^\pm} < 0.80$
Muon Opening Angle	$\cos(179.64^\circ) < \cos \theta_{\mu\mu} < \cos(178.60^\circ)$
Energy-Momentum Ratio	$0.05 \leq (E/p)^{\mu^\pm} < 0.15$
Shower Energy	$0.0 \text{ GeV} < E_{EMC}^{\mu^\pm} < 0.3 \text{ GeV}$

Table 5.1: Selection cuts on muon tracks used to determine the center-of-mass energy.




Figure 5.1: The differences measured for scan data center-of-mass energies. The average difference (green) is about 1.29 MeV higher than the recorded energy values provided by the collider at run time (red).

The use of muons to determine energy is subject to an overall scale shift due to potentially miscalibrated magnetic field effects. This requires a point of reference in order to ensure the measured values are correctly determined. We use the first round of the on-peak  $\psi(3770)$  sample for this comparison, as its center-of-mass energy has been alternatively measured to a very precise degree [45]. The results of this method compared to our own for the on-peak data are shown in Figure 5.2. It is evident the official procedure using  $D\bar{D}$  events provides more stability over the run range than using muons. Still, the difference between average values of each measurement is taken as the offset for shifting the measured scan data points. From this, the measured center-of-mass energies for the scan data are increased by 1.86 MeV to determine the final values. Compared to the initial measurements of scan data energy, this indicates the values were lower by  $\sim 0.7$  MeV on average. The final center-of-mass energies of the scan data, along with the luminosity of each bin, are shown in Table 5.3.




Figure 5.2: The values measured for on-peak data center-of-mass energies. The difference between our measured values (blue) and the official values (red) is used to shift the results of the measured scan data energy values.

### 5.3.3 Luminosity Measurement

To precisely determine the values of luminosity for each bin, radiative Bhabha events ( $e^+e^- \rightarrow e^+e^-(\gamma)$ ) were analyzed to determine the total number of collisions produced. The procedure used is described in Ref. [46]. For each run,  $1.4 \times 10^6$  events were generated using Babayaga 3.5. We select events with only two (oppositely) charged tracks, satisfying the event selection criteria shown in Table 5.2.

Vertex ( $xy$ )	$V_{xy} < 1 \text{ cm}$
Vertex ( $z$ )	$ Vz  < 10 \text{ cm}$
MDC Angle	$ \cos \theta  < 0.80$
Exclude $e^+e^- \rightarrow \mu^+\mu^-(\gamma)$ Events	$E_{\text{EMC}} > 0.73 \times E_{\text{beam}}$
Exclude $e^+e^- \rightarrow \gamma J/\psi, J/\psi \rightarrow e^+e^-$ Events	$p > 0.93 \times E_{\text{beam}}$

Table 5.2: Selection cuts on electron tracks used to determine the luminosity.

Applying these cuts to both data and MC identically, we use the resulting number

of events found in the MC divided by the number of total generated events to determine the efficiency ( $\epsilon_{MC}$ ). From this, and using the cross section provided by the generator ( $\sigma_{Bhabha}$ ), we can take the events found in data ( $N_{\text{data}}$ ) to determine the integrated luminosity ( $\mathcal{L}$ ) of each run:

$$\mathcal{L} = \frac{N_{\text{data}}}{\sigma_{\text{Bhabha}} \epsilon_{MC}} \quad (5.18)$$

The integrated luminosity for each bin is shown in Table 5.3. The total luminosity for the scan data used is  $(69.80 \pm 0.03) \text{ pb}^{-1}$ , where the error listed is statistical.

Bin	Run Range	$E_{\text{cm}}$ Range [GeV]	$\mathcal{L}$ [pb $^{-1}$ ]
0	14041 - 14046	3.734 - 3.736	0.8293(30)
1	14360 - 14360	3.736 - 3.744	0.3287(19)
2	14047 - 14048	3.744 - 3.748	0.9524(32)
3	14049 - 14053	3.748 - 3.750	1.4055(39)
4	14057 - 14067	3.750 - 3.751	2.2717(50)
5	14068 - 14077	3.751 - 3.753	2.9702(57)
6	14078 - 14086	3.753 - 3.755	3.3080(60)
7	14087 - 14101	3.755 - 3.756	3.4162(61)
8	14103 - 14109	3.756 - 3.759	3.8712(65)
9	14110 - 14123	3.759 - 3.762	4.4382(70)
10	14129 - 14144	3.762 - 3.765	4.4896(70)
11	14145 - 14154	3.765 - 3.767	3.2828(60)
12	14155 - 14160	3.767 - 3.771	2.4418(52)
13	14161 - 14164	3.771 - 3.774	2.0151(47)
14	14165 - 14168	3.774 - 3.777	1.8261(45)
15	14169 - 14174	3.777 - 3.780	1.8237(45)
16	14200 - 14203	3.780 - 3.782	1.9505(46)
17	14209 - 14217	3.782 - 3.787	2.1500(49)
18	14221 - 14226	3.787 - 3.789	2.5488(53)
19	14230 - 14238	3.789 - 3.792	2.8320(56)
20	14239 - 14246	3.792 - 3.797	3.5310(63)
21	14247 - 14258	3.797 - 3.800	4.0479(67)
22	14259 - 14268	3.800 - 3.802	3.9284(66)
23	14269 - 14274	3.802 - 3.807	2.6929(55)
24	14275 - 14279	3.807 - 3.809	1.7604(44)
25	14280 - 14290	3.809 - 3.813	1.2539(38)
26	14291 - 14292	3.813 - 3.815	0.8969(32)
27	14293 - 14294	3.815 - 3.823	0.6803(28)
28	14295 - 14296	3.823 - 3.832	0.3997(21)
29	14297 - 14297	3.832 - 3.839	0.2846(18)
30	14298 - 14298	3.839 - 3.849	0.2802(18)
31	14299 - 14299	3.849 - 3.855	0.2764(18)
32	14300 - 14301	3.855 - 3.863	0.3188(19)
33	14302 - 14303	3.863 - 3.870	0.3002(19)

Table 5.3: Measured luminosities for each energy bin.  
The uncertainties listed are statistical errors from the data selection, as uncertainties from the MC statistics are negligible.

### 5.3.4 Monte Carlo Generation

To compare to the scan data sample, several Monte Carlo (MC) samples were produced. For the main signal determination, samples of generic  $D^0\overline{D}^0$  from  $\psi(3770)$  and generic  $D^+D^-$  from  $\psi(3770)$  were generated with  $2 \times 10^5$  events per bin. In addition,  $100 \times$  data-size samples were produced for  $q\bar{q}$ ,  $\tau^+\tau^-$ , radiative return to  $J/\psi$  (denoted  $\gamma J/\psi$ ), and radiative return to  $\psi(2S)$  (denoted  $\gamma\psi'$ ). Each of these samples was generated at the University of Minnesota in July of 2014 using BOSS 6.6.4.p02. The  $D^0\overline{D}^0$ ,  $D^+D^-$ ,  $q\bar{q}$ , and  $\tau^+\tau^-$  states were generated using KKMC, while the  $\gamma J/\psi$  and  $\gamma\psi'$  were generated with BesEvtGen. All except  $q\bar{q}$  were then decayed with BesEvtGen. The total numbers of events in each sample can be found in Table 5.4.

Sample	Number of Events
$\psi(3770) \rightarrow D^0\overline{D}^0$	$6.800 \times 10^6$
$\psi(3770) \rightarrow D^+D^-$	$6.400 \times 10^6$
$q\bar{q}$	$8.916 \times 10^7$
$\gamma J/\psi$	$7.307 \times 10^6$
$\gamma\psi'$	$2.457 \times 10^7$
$\tau^+\tau^-$	$2.164 \times 10^7$
Data	$4.844 \times 10^8$

Table 5.4: Number of events contained in each generated sample and the scan data.

In general, all MC samples were generated based off communal decay card information used within BESIII. However, the  $D\overline{D}$  samples were generated by implementing the Born-level cross section shape from the final fit into KKMC. This procedure was repeated over five iterations in order to provide a more data-driven basis for the ISR corrections. The effects of this process are examined in Section 5.7.

### 5.3.5 Software Packages

The main packages used for this analysis are shown in Table 5.5. The package `DtagRecAlg` is a custom package used to extract variables from `DTagAlg` relevant to the analysis.

Package	Version
KKMC	v00-00-51
DTagAlg	v00-00-51
DTagRecAlg	v00-00-01

Table 5.5: The software packages used in this analysis and their version numbers.

## 5.4 Signal Determination

We measure the yields of both  $D^0\overline{D}^0$  and  $D^+D^-$  events with two-dimensional fits to  $\Delta E$  and  $m_{BC}$ . MC samples are partitioned into the following four groups: proper  $D$ -tags ( $N_{D\bar{D}}$ ), misreconstructed  $D$ -tags ( $N_{\text{misrec}}$ ), continuum ( $N_{q\bar{q}}$ ), and other ( $N_{\text{other}}$ ). The first two groups are obtained using truth information from the  $D\bar{D}$  samples, while the last group is a combination of the  $\tau^+\tau^-$ ,  $\gamma J/\psi$ , and  $\gamma\psi'$  samples. These groups are fitted to data using the RooFit package to perform a negative log-likelihood minimization for each energy bin ( $E_i$ ) separately for both  $D^0$  and  $D^+$ . For each fit, the four MC sample groups are used to construct 2D ( $\Delta E$  vs.  $m_{BC}$ ) PDF functions that are fitted against the corresponding data histograms. The proper  $D\bar{D}$  shape is treated as signal, and its integral after fitting ( $N_D$ ) is used for determining the signal yields and cross sections. An example fit is shown in Figure 5.3, while the complete set of these plots can be found in Appendices B and C.




Figure 5.3: An example 2D ( $\Delta E$  vs.  $m_{BC}$ ) signal fit.

The covers  $D^0$  events in the region  $3.774 \text{ GeV} < E_{\text{cm}} \leq 3.777 \text{ GeV}$ . The data points (black) are fitted by the total MC shape (blue), which is the sum of the signal (green) and background (red) components.

## 5.5 Efficiency Correction

In addition to the parameters gathered by `DTagAlg`, truth information was taken from the generic  $D\bar{D}$  samples in order to determine the mode-by-mode reconstruction efficiency. To be deemed proper, a reconstruction must pass not only the standard  $D$ -tag cuts, but also match the generator information for the event. This process removes peaking backgrounds from modes with similar constituents. The total number of proper  $D$ -tag reconstructions is then divided by the number of  $D$  particles generated for each mode, and the mode-by-mode efficiencies are weighted by the PDG branching ratios [6] to determine the overall efficiency ( $\epsilon_D$ ) for each of  $D^0$  and  $D^+$ :

$$\epsilon_D = \sum_i \epsilon_i \text{ rec } \mathcal{B}_i = \sum_i \left( \frac{N_i \text{ prop}}{N_i \text{ gen}} \right) \mathcal{B}_i. \quad (5.19)$$

Each  $D^0$  efficiency also includes the corresponding DCSD terms for its decay (See Section 4.3).

To determine the cross section at each point, this procedure was applied separately for each energy bin. The total number of proper and generated particles for each energy bin are shown in Table 5.6 for  $D^0$  and Tables 5.7 and 5.8 for  $D^+$ . The efficiencies for  $D^+$  and  $D^0$  were also calculated across the total sample, and are shown for each mode in Table 5.9 and Figure 5.4. Total efficiencies of each bin for both  $D^0$  and  $D^+$  are shown in Table 5.10.

### 5.5.1 CP Violation Correction

Due to CP violation in the  $D^0\bar{D}^0$  system, each of the neutral decay modes must be corrected to account for quantum correlations. This is done by applying scaling factors to the efficiency for each of the three modes used in reconstruction. The corrections are parameterized for each mode ( $m$ ) by the following form [47]:

$$\alpha_{D^0 \rightarrow m} = 1 + r_m^2 + 2 \times y \times r_m \times R_m \times \cos(\delta_m). \quad (5.20)$$

Here,  $r_m$  and  $\delta_m$  represent the relative magnitudes and phases between the Cabibbo-favored and doubly-Cabibbo-suppressed modes, respectively, while the factor of  $R_m$

Bin	$D^0 \rightarrow K^- \pi^+$			$D^0 \rightarrow K^- \pi^+ \pi^0$			$D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$		
	$N_{\text{prop}}$	$N_{\text{gen}}$	$\epsilon_{\text{rec}}$	$N_{\text{prop}}$	$N_{\text{gen}}$	$\epsilon_{\text{rec}}$	$N_{\text{prop}}$	$N_{\text{gen}}$	$\epsilon_{\text{rec}}$
0	11197	15757	0.711	21600	55815	0.387	13592	33714	0.403
1	11266	15713	0.717	21735	56189	0.387	13651	33621	0.406
2	11065	15557	0.711	21683	55791	0.389	13984	33431	0.418
3	10748	15501	0.693	21114	55767	0.379	13168	33505	0.393
4	11176	15779	0.708	21922	55873	0.392	13613	33410	0.407
5	10984	15722	0.699	21366	56027	0.381	13483	33748	0.400
6	10837	15507	0.699	21144	55551	0.381	13738	33639	0.408
7	10707	15645	0.684	20889	56193	0.372	13074	33494	0.390
8	11031	15585	0.708	21788	55891	0.390	13967	33826	0.413
9	10994	15485	0.710	21978	56012	0.392	14020	33601	0.417
10	10952	15497	0.707	21400	56126	0.381	13727	33591	0.409
11	11025	15535	0.710	21700	55919	0.388	13980	33706	0.415
12	10909	15593	0.700	21713	55876	0.389	13787	33335	0.414
13	11088	15826	0.701	21696	55758	0.389	14013	33805	0.415
14	10975	15620	0.703	21576	55591	0.388	14039	33581	0.418
15	10965	15672	0.700	21501	55647	0.386	13777	33644	0.409
16	10803	15438	0.700	21505	55667	0.386	13832	33822	0.409
17	10913	15473	0.705	21574	56023	0.385	14163	34131	0.415
18	11156	15856	0.704	21461	55706	0.385	13725	33321	0.412
19	11124	15728	0.707	21775	56192	0.388	13930	33935	0.410
20	10837	15530	0.698	21149	55737	0.379	13883	33703	0.412
21	10756	15397	0.699	21212	55815	0.380	13577	33907	0.400
22	10986	15582	0.705	21178	56343	0.376	13352	33773	0.395
23	11147	15861	0.703	21088	56167	0.375	13200	33680	0.392
24	10785	15633	0.690	20763	55952	0.371	12651	33660	0.376
25	10972	15490	0.708	20687	55957	0.370	12841	33512	0.383
26	11016	15645	0.704	20707	55785	0.371	12563	33615	0.374
27	10800	15420	0.700	20244	55748	0.363	12362	33359	0.371
28	10939	15694	0.697	20077	55550	0.361	12507	33497	0.373
29	10839	15689	0.691	20685	56386	0.367	12575	33551	0.375
30	10798	15766	0.685	20238	55433	0.365	12526	33885	0.370
31	10568	15438	0.685	20533	55825	0.368	12901	33745	0.382
32	10820	15679	0.690	20706	55823	0.371	13155	33810	0.389
33	10643	15736	0.676	20488	55672	0.368	12875	33547	0.384

Table 5.6: Number of proper and generated particles for  $D^0$ .  
The mode-by-mode numbers of particles used in the efficiency calculations for  
 $D^0 \rightarrow K^- \pi^+$ ,  $D^0 \rightarrow K^- \pi^+ \pi^0$ , and  $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$ .

Bin	$D^+ \rightarrow K^- \pi^+ \pi^+$			$D^+ \rightarrow K^- \pi^+ \pi^+ \pi^0$			$D^+ \rightarrow K_S^0 \pi^+$		
	$N_{\text{prop}}$	$N_{\text{gen}}$	$\epsilon_{\text{rec}}$	$N_{\text{prop}}$	$N_{\text{gen}}$	$\epsilon_{\text{rec}}$	$N_{\text{prop}}$	$N_{\text{gen}}$	$\epsilon_{\text{rec}}$
2	20643	37609	0.549	6853	24250	0.283	2302	5900	0.390
3	19935	37582	0.530	6569	24235	0.271	2335	6114	0.382
4	20859	37915	0.550	6653	24216	0.275	2291	5879	0.390
5	20193	37647	0.536	6611	24051	0.275	2292	5965	0.384
6	20484	37814	0.542	6616	24184	0.274	2292	5924	0.387
7	19635	37462	0.524	6411	24708	0.259	2127	5815	0.366
8	20876	37758	0.553	6867	24343	0.282	2328	5841	0.399
9	20873	37739	0.553	6754	24284	0.278	2377	5911	0.402
10	20571	37487	0.549	6604	24249	0.272	2360	6010	0.393
11	20669	37468	0.552	6818	24268	0.281	2296	5904	0.389
12	20843	37838	0.551	6783	24247	0.280	2401	6074	0.395
13	20486	37286	0.549	6942	24582	0.282	2335	6184	0.378
14	20935	37961	0.551	6836	24467	0.279	2282	5963	0.383
15	20543	37458	0.548	6769	24135	0.280	2295	5889	0.390
16	20713	37543	0.552	6758	24376	0.277	2356	5921	0.398
17	21016	37757	0.557	6949	24470	0.284	2302	5903	0.390
18	21123	38024	0.556	6635	24111	0.275	2320	5889	0.394
19	20708	37357	0.554	6662	24240	0.275	2204	5855	0.376
20	20760	37761	0.550	6768	24395	0.277	2263	5883	0.385
21	20860	37893	0.550	6668	24216	0.275	2296	5996	0.383
22	20827	37820	0.551	6673	24421	0.273	2353	5997	0.392
23	20796	37554	0.554	6723	24377	0.276	2290	5957	0.384
24	20058	37410	0.536	6415	24310	0.264	2341	5966	0.392
25	20677	37552	0.551	6462	24245	0.267	2364	5987	0.395
26	20300	37491	0.541	6539	24405	0.268	2356	5993	0.393
27	20754	37704	0.550	6530	24295	0.269	2288	5969	0.383
28	20213	37428	0.540	6362	24146	0.263	2341	5988	0.391
29	20565	37545	0.548	6495	24257	0.268	2262	5916	0.382
30	20313	37773	0.538	6535	24167	0.270	2323	6075	0.382
31	20322	37280	0.545	6566	24389	0.269	2378	5927	0.401
32	20883	37992	0.550	6594	24249	0.272	2306	5855	0.394
33	20645	37738	0.547	6560	24195	0.271	2269	5945	0.382

Table 5.7: Number of proper and generated particles for  $D^+$  (part 1).  
The mode-by-mode numbers of particles used in the efficiency calculations for  
 $D^+ \rightarrow K^- \pi^+ \pi^+$ ,  $D^+ \rightarrow K^- \pi^+ \pi^+ \pi^0$ , and  $D^+ \rightarrow K_S^0 \pi^+$ .

Bin	$D^+ \rightarrow K_S^0 \pi^+ \pi^0$			$D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-$			$D^+ \rightarrow K^+ K^- \pi^+$		
	$N_{\text{prop}}$	$N_{\text{gen}}$	$\epsilon_{\text{rec}}$	$N_{\text{prop}}$	$N_{\text{gen}}$	$\epsilon_{\text{rec}}$	$N_{\text{prop}}$	$N_{\text{gen}}$	$\epsilon_{\text{rec}}$
2	5742	27235	0.211	3562	15019	0.237	1720	4019	0.428
3	5668	27699	0.205	3169	14877	0.213	1637	3970	0.412
4	5710	27527	0.207	3358	14750	0.228	1791	4016	0.446
5	5614	27654	0.203	3282	14696	0.223	1748	3981	0.439
6	5656	27475	0.206	3392	14823	0.229	1771	4075	0.435
7	5481	27597	0.199	3293	14860	0.222	1593	3951	0.403
8	5827	27975	0.208	3462	15035	0.230	1736	4042	0.429
9	6003	27589	0.218	3427	14906	0.230	1716	3940	0.436
10	5722	27751	0.206	3327	14854	0.224	1690	3941	0.429
11	5888	27649	0.213	3385	14837	0.228	1672	3930	0.425
12	5731	27651	0.207	3370	14926	0.226	1718	3889	0.442
13	5792	27618	0.210	3336	14669	0.227	1696	3958	0.428
14	5745	27608	0.208	3288	14702	0.224	1712	3904	0.439
15	5832	27480	0.212	3384	14808	0.229	1649	3855	0.428
16	5891	27758	0.212	3443	14845	0.232	1783	3947	0.452
17	5954	27639	0.215	3484	14910	0.234	1804	3941	0.458
18	5875	27669	0.212	3366	14913	0.226	1700	3968	0.428
19	5773	27838	0.207	3371	14782	0.228	1766	4037	0.437
20	6024	28019	0.215	3367	14801	0.227	1709	3903	0.438
21	5804	27645	0.210	3306	14819	0.223	1688	3863	0.437
22	5893	27692	0.213	3304	14949	0.221	1741	3975	0.438
23	5780	27865	0.207	3313	14981	0.221	1670	3891	0.429
24	5623	27811	0.202	3161	14867	0.213	1667	3954	0.422
25	5684	27490	0.207	3250	14794	0.220	1711	3966	0.431
26	5704	27829	0.205	3137	14811	0.212	1641	3958	0.415
27	5630	27497	0.205	3132	14881	0.210	1678	3972	0.422
28	5724	28006	0.204	3271	14828	0.221	1690	3931	0.430
29	5693	27554	0.207	3272	15009	0.218	1772	3994	0.444
30	5574	27717	0.201	3308	14730	0.225	1673	4026	0.416
31	5556	27556	0.202	3288	15049	0.218	1746	3965	0.440
32	5785	27634	0.209	3259	14840	0.220	1680	3923	0.428
33	5653	27528	0.205	3287	14959	0.220	1734	4031	0.430

Table 5.8: Number of proper and generated particles for  $D^+$  (part 2).  
The mode-by-mode numbers of particles used in the efficiency calculations for  
 $D^+ \rightarrow K_S^0 \pi^+ \pi^0$ ,  $D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-$ , and  $D^+ \rightarrow K^+ K^- \pi^+$ .

Decay Mode ( $i$ )	PDG $\mathcal{B}_i$ [%]	MC Efficiency $\epsilon_i$
$D^0 \rightarrow K^- \pi^+$	$3.89 \pm 0.05$	$0.7002 \pm 0.0011$
$D^0 \rightarrow K^- \pi^+ \pi^0$	$13.93 \pm 0.50$	$0.3794 \pm 0.0004$
$D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$	$8.11 \pm 0.21$	$0.3988 \pm 0.0006$
$\epsilon_{D^0} = (11.245 \pm 0.020)\%$		
$D^+ \rightarrow K^- \pi^+ \pi^+$	$9.13 \pm 0.19$	$0.5471 \pm 0.0007$
$D^+ \rightarrow K^- \pi^+ \pi^+ \pi^0$	$5.99 \pm 0.18$	$0.2739 \pm 0.0006$
$D^+ \rightarrow K_S^0 \pi^+$	$1.47 \pm 0.07$	$0.3883 \pm 0.0014$
$D^+ \rightarrow K_S^0 \pi^+ \pi^0$	$6.99 \pm 0.27$	$0.2079 \pm 0.0005$
$D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-$	$3.12 \pm 0.11$	$0.2237 \pm 0.0007$
$D^+ \rightarrow K^+ K^- \pi^+$	$0.95 \pm 0.03$	$0.4317 \pm 0.0018$
$\epsilon_{D^+} = (9.770 \pm 0.063)\%$		

Table 5.9: Mode-by-mode reconstruction efficiencies for  $D^0$  and  $D^+$ .

The values shown are over the entire data sample, while calculations for the cross sections use the values of each energy point individually (see Table 5.10). The errors listed on the PDG branching fractions are shown for reference, and are not included in the efficiency errors (see Section 5.7.1).

represents a coherence factor characterizing the variation of  $\delta_m$  over phase space. Note, there is no such variation for a two-body decay (like  $D^0 \rightarrow K^- \pi^+$ ), so  $R_{D^0 \rightarrow K^- \pi^+} = 1$ . The value of  $y$  represents the difference in total width components of the  $D^0 \bar{D}^0$  system,  $y = (\Gamma_2 - \Gamma_1)/(\Gamma_2 + \Gamma_1)$ , where 1 and 2 represent the CP-odd and CP-even states, respectively.

The mode-dependent values for these factors are listed in Table 5.11. These are taken from the *CPV*-allowed values in [48] for  $D^0 \rightarrow K^- \pi^+$ , and from [49] for  $D^0 \rightarrow K^- \pi^+ \pi^0$  and  $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$ . The value of  $y = 0.0066^{+0.0007}_{-0.0010}$  is also from [48], and is the same for all modes. After applying each of the mode-dependent corrections, the efficiency for the full sample changes from  $\epsilon_{D^0} = (11.320 \pm 0.213)\%$  to  $\epsilon_{D^0} = (11.352 \pm 0.213)\%$ , and similarly for the efficiencies of each  $E_{\text{cm}}$  bin.

Mode	$r_m$	$R_m$	$\delta_m$ [°]	$\alpha_m$
$D^0 \rightarrow K^- \pi^+$	$0.0591 \pm 0.0063$	1	$11.8^{+9.5}_{-14.7}$	$1.00426 \pm 0.00083$
$D^0 \rightarrow K^- \pi^+ \pi^0$	$0.0447 \pm 0.0012$	$0.81 \pm 0.06$	$18^{+14}_{-15}$	$1.00248 \pm 0.00014$
$D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$	$0.0549 \pm 0.0006$	$0.43^{+0.17}_{-0.13}$	$-52^{+28}_{-17}$	$1.00270^{+0.00014}_{-0.00012}$

Table 5.11: The quantum correlated factors for the  $D^0$  modes.

$E_{\text{bin}}$	$\epsilon_{D^0}$	$\epsilon_{D^+}$
0	$0.1146 \pm 0.0005$	-
1	$0.1151 \pm 0.0005$	-
2	$0.1161 \pm 0.0005$	$0.0990 \pm 0.0005$
3	$0.1119 \pm 0.0005$	$0.0952 \pm 0.0005$
4	$0.1156 \pm 0.0005$	$0.0983 \pm 0.0005$
5	$0.1131 \pm 0.0005$	$0.0964 \pm 0.0005$
6	$0.1137 \pm 0.0005$	$0.0972 \pm 0.0005$
7	$0.1104 \pm 0.0005$	$0.0934 \pm 0.0005$
8	$0.1157 \pm 0.0005$	$0.0991 \pm 0.0005$
9	$0.1165 \pm 0.0005$	$0.0996 \pm 0.0005$
10	$0.1141 \pm 0.0005$	$0.0977 \pm 0.0005$
11	$0.1157 \pm 0.0005$	$0.0990 \pm 0.0005$
12	$0.1152 \pm 0.0005$	$0.0986 \pm 0.0005$
13	$0.1154 \pm 0.0005$	$0.0985 \pm 0.0005$
14	$0.1157 \pm 0.0005$	$0.0984 \pm 0.0005$
15	$0.1146 \pm 0.0005$	$0.0986 \pm 0.0005$
16	$0.1146 \pm 0.0005$	$0.0992 \pm 0.0005$
17	$0.1151 \pm 0.0005$	$0.1003 \pm 0.0005$
18	$0.1148 \pm 0.0005$	$0.0990 \pm 0.0005$
19	$0.1151 \pm 0.0005$	$0.0984 \pm 0.0005$
20	$0.1138 \pm 0.0005$	$0.0988 \pm 0.0005$
21	$0.1129 \pm 0.0005$	$0.0982 \pm 0.0005$
22	$0.1122 \pm 0.0005$	$0.0984 \pm 0.0005$
23	$0.1118 \pm 0.0005$	$0.0982 \pm 0.0005$
24	$0.1094 \pm 0.0005$	$0.0953 \pm 0.0005$
25	$0.1105 \pm 0.0005$	$0.0975 \pm 0.0005$
26	$0.1098 \pm 0.0005$	$0.0962 \pm 0.0005$
27	$0.1082 \pm 0.0005$	$0.0969 \pm 0.0005$
28	$0.1081 \pm 0.0005$	$0.0961 \pm 0.0005$
29	$0.1087 \pm 0.0005$	$0.0971 \pm 0.0005$
30	$0.1078 \pm 0.0005$	$0.0959 \pm 0.0005$
31	$0.1092 \pm 0.0005$	$0.0969 \pm 0.0005$
32	$0.1104 \pm 0.0005$	$0.0978 \pm 0.0005$
33	$0.1090 \pm 0.0005$	$0.0971 \pm 0.0005$

Table 5.10: The overall reconstruction efficiency of  $D^0$  and  $D^+$  for each energy bin. These values are used to calculate the corresponding cross sections at each energy point. The listed errors are statistical only.




Figure 5.4: Mode-by-mode MC efficiencies for  $D^0$  and  $D^+$ .  
The error bars are negligible on the scale shown.

## 5.6 Fitting Procedure

After applying the correction in Section 5.5.1, the efficiency values (Table 5.10) were combined with the luminosity (Table 5.3) and the signal values from each 2D fit (Table 5.12) to determine the cross section at each energy point. Since each  $\psi(3770)$  produces a  $D\bar{D}$  pair, a factor of 2 is included to avoid double counting:

$$\sigma_{D\bar{D}}^{RC}(E_i) = \frac{N_D(E_i)}{2 \epsilon_D(E_i) \mathcal{L}(E_i)}. \quad (5.21)$$

The resulting cross section values for  $D^0$  and  $D^+$  are also listed in Table 5.12, and shown in Figure 5.5.

Using the measured  $D\bar{D}$  cross sections, we fit to Equation (5.7) using each of the form factor choices described in Section 5.2. In each case, there are four common fit parameters:  $M^{\psi(3770)}$ ,  $\Gamma^{\psi(3770)}$ ,  $\Gamma_{ee}^{\psi(3770)}$ , and  $\phi^{\psi(3770)}$ . These represent the mass, total

$E_{\text{mid}}$	$N_{D^0\overline{D}^0}$	$\sigma_{D^0\overline{D}^0}^{RC}$ [nb]	$N_{D^+D^-}$	$\sigma_{D^+D^-}^{RC}$ [nb]
3.7342	$32 \pm 9$	$0.169 \pm 0.046$	-	-
3.7368	$16 \pm 5$	$0.218 \pm 0.068$	-	-
3.7447	$169 \pm 16$	$0.765 \pm 0.070$	$28 \pm 7$	$0.150 \pm 0.038$
3.7483	$264 \pm 20$	$0.838 \pm 0.064$	$96 \pm 13$	$0.360 \pm 0.048$
3.7501	$509 \pm 28$	$0.968 \pm 0.053$	$196 \pm 18$	$0.439 \pm 0.040$
3.7517	$831 \pm 34$	$1.237 \pm 0.052$	$329 \pm 23$	$0.574 \pm 0.040$
3.7534	$1036 \pm 38$	$1.377 \pm 0.051$	$481 \pm 27$	$0.748 \pm 0.042$
3.7556	$1182 \pm 41$	$1.566 \pm 0.055$	$508 \pm 28$	$0.797 \pm 0.045$
3.7562	$1459 \pm 45$	$1.629 \pm 0.051$	$701 \pm 32$	$0.914 \pm 0.042$
3.7592	$2014 \pm 52$	$1.948 \pm 0.052$	$1054 \pm 39$	$1.192 \pm 0.045$
3.7624	$2444 \pm 58$	$2.385 \pm 0.058$	$1346 \pm 44$	$1.534 \pm 0.051$
3.7650	$2062 \pm 53$	$2.715 \pm 0.071$	$1223 \pm 42$	$1.882 \pm 0.066$
3.7676	$1746 \pm 48$	$3.102 \pm 0.087$	$1059 \pm 39$	$2.200 \pm 0.081$
3.7713	$1585 \pm 46$	$3.406 \pm 0.101$	$1045 \pm 38$	$2.634 \pm 0.098$
3.7742	$1569 \pm 46$	$3.714 \pm 0.111$	$1103 \pm 40$	$3.067 \pm 0.111$
3.7775	$1543 \pm 46$	$3.692 \pm 0.111$	$1108 \pm 39$	$3.078 \pm 0.111$
3.7802	$1539 \pm 46$	$3.444 \pm 0.104$	$1006 \pm 38$	$2.599 \pm 0.100$
3.7829	$1381 \pm 44$	$2.791 \pm 0.090$	$951 \pm 38$	$2.206 \pm 0.088$
3.7869	$1167 \pm 42$	$1.995 \pm 0.072$	$821 \pm 36$	$1.627 \pm 0.073$
3.7891	$888 \pm 38$	$1.361 \pm 0.058$	$656 \pm 34$	$1.178 \pm 0.062$
3.7926	$739 \pm 36$	$0.920 \pm 0.045$	$475 \pm 32$	$0.680 \pm 0.045$
3.7970	$514 \pm 34$	$0.562 \pm 0.037$	$329 \pm 31$	$0.414 \pm 0.039$
3.8003	$374 \pm 31$	$0.424 \pm 0.035$	$186 \pm 28$	$0.241 \pm 0.037$
3.8024	$196 \pm 23$	$0.325 \pm 0.038$	$125 \pm 23$	$0.236 \pm 0.043$
3.8070	$136 \pm 19$	$0.352 \pm 0.050$	$97 \pm 17$	$0.289 \pm 0.051$
3.8093	$65 \pm 14$	$0.233 \pm 0.050$	$32 \pm 13$	$0.132 \pm 0.055$
3.8135	$12 \pm 10$	$0.060 \pm 0.049$	$26 \pm 11$	$0.153 \pm 0.066$
3.8153	$8 \pm 7$	$0.056 \pm 0.051$	$12 \pm 8$	$0.089 \pm 0.063$
3.8229	$12 \pm 7$	$0.140 \pm 0.083$	$15 \pm 8$	$0.197 \pm 0.107$
3.8320	$4 \pm 5$	$0.069 \pm 0.086$	$3 \pm 5$	$0.046 \pm 0.099$
3.8390	$14 \pm 6$	$0.237 \pm 0.105$	$14 \pm 7$	$0.254 \pm 0.124$
3.8494	$11 \pm 6$	$0.186 \pm 0.104$	$10 \pm 6$	$0.186 \pm 0.118$
3.8555	$24 \pm 8$	$0.337 \pm 0.111$	$17 \pm 6$	$0.273 \pm 0.104$
3.8632	$22 \pm 8$	$0.340 \pm 0.127$	$6 \pm 5$	$0.099 \pm 0.091$

Table 5.12: The  $D\overline{D}$  cross section at each  $E_{\text{cm}}$  point.

The number of data events observed in each  $E_{\text{cm}}$  bin are also shown. The uncertainties on the cross sections are statistical only and come from the signal fitting ( $N_D$ ), PDG values (Table 5.9), and MC reconstruction efficiency (Table 5.10). The values for  $N_{D^0\overline{D}^0}$  and  $N_{D^+D^-}$  are taken from the signal fits shown in Appendices B and C.




Figure 5.5: The measured  $e^+e^- \rightarrow D\bar{D}$  cross sections.  
The  $D^0\bar{D}^0$  cross section is shown on the left and  $D^+D^-$  is shown on the right.

width, electron partial width, and relative phase to the non-resonant contribution for the  $\psi(3770)$ , respectively. This notation for the total width corresponds to  $\Gamma(M)$  in Section 5.1. Two additional fitting parameters are dependent on which form factor is being used:  $F_{NR}$  and  $a_{NR}$  for the exponential, or  $\Gamma^{\psi(2S)}$  and  $F_0$  for the VDM. For the former, these represent the amplitude and exponent normalization for the non-resonant contribution. In the latter case, these represent the modified total width for the  $\psi(2S)$  above resonance (see Section 5.2) and the constant contribution of resonances above the  $\psi(3770)$ . The fitting is done simultaneously for  $D^0$  and  $D^+$  with identical parameters using TMinuit with a  $\chi^2$  minimization. The value minimized is the sum of the  $\chi^2$  values from each of the  $D^0$  and  $D^+$  cross section functions. Results for the Exponential and VDM choices are shown in Figures 5.6 and 5.7, respectively.




Figure 5.6: The Exponential Model fit results.

The results for  $D^0$  are shown on the top and the results for  $D^+$  are shown on the bottom. The fit shape (blue) is calculated from Equation (5.1) using the non-resonant component from Equation (5.15).




Figure 5.7: The Vector Dominance Model fit results.

The results for  $D^0$  are shown on the top and the results for  $D^+$  are shown on the bottom. The fit shape (blue) is calculated from Equation (5.1) using the non-resonant component from Equation (5.16).

### 5.6.1 Coulomb Correction

In the development of this analysis, it was discovered that the theoretical formulation in Section 5.1 did not lead to a successful fit of the  $D\bar{D}$  cross sections, shown in Figure 5.8. Namely, including the Coulomb effect pulls the  $D^0\bar{D}^0$  and  $D^+\bar{D}^-$  cross sections in opposite directions. We found the best fits were achieved by altering Equation (5.3) to set the Coulomb factor to 1. While this disagrees with conventional theoretical wisdom, it is consistent with studies of  $\Upsilon(4S) \rightarrow B\bar{B}$  where applying a Coulomb correction for the charged final state also leads to inconsistency with data.




Figure 5.8: The Vector Dominance Model fit results with Coulomb interactions. The results for  $D^0$  are shown on the top and the results for  $D^+$  are shown on the bottom. Including this factor provides notably worse results than when excluding it (see Figure 5.7).

This is most clearly seen in the ratio of  $D^+ D^-$  and  $D^0 \bar{D}^0$  cross sections, shown in Figure 5.9, where the ‘No Coulomb’ method sets  $z_{D^+ D^-}$  in Equations (5.7) and (5.11) to unity, the ‘Partial Coulomb’ sets this factor to unity only for Equation (5.7), and the ‘Full Coulomb’ is the default assumption. Agreement of the measured cross section ratio with the ‘No Coulomb’ calculation is substantially better. This is also true for the high statistics points measured at the  $\psi(3770)$  peak by Derrick Toth [44] (light blue). As the data tend to follow the ‘No Coulomb’ method, we choose this as our nominal method for the results, presented in Section 5.8. However, the interpretation for this

behavior is still undetermined.




Figure 5.9: The ratio of measured  $D^+$  to  $D^0$  cross sections.  
Several levels of Coulomb interactions are examined.

## 5.7 Systematics

To assess the systematic uncertainties in our results, we look at a variety of factors. Many of these affect all BESIII analyses, such as luminosity and tracking. Others, like the modification to KKMC generation (see Section 5.3.4), are specific to this analysis. The remaining systematics are typically due to less well-known parameters, like the radii used to describe the  $\psi(2S)$  and  $\psi(3770)$  (see Section 5.2). Each of these contributions, as well as their total, can be found in Table 5.16.

### 5.7.1 $\psi(3770)$ Parameter Systematics

Each systematic is obtained by changing a specific aspect, and re-fitting to the altered values using the VDM method. The uncertainties for each parameter are obtained by

taking the difference between this result and the nominal fit (Figure 5.7). Generally, each change was done both positively and negatively, and the values used are the largest differences seen between the two changes. The systematics examined in this analysis are summarized below, where a \* denotes aspects which were deemed negligible.

### Luminosity

A 1 % change [50] was applied to  $\mathcal{L}$  in Equation (5.21). As this is an overall scale change, the only variable significantly affected is  $\Gamma_{ee}^{\psi(3770) \rightarrow D\bar{D}}$ .

### $\pi^\pm/K^\pm$ Tracking

A 1.0 % change [51] was applied for each  $\pi^\pm$  or  $K^\pm$  in a given decay mode. The summed contribution for each mode is applied to  $\epsilon_m$  in Equation (5.19). As this is an overall scale change, the only variable significantly affected is  $\Gamma_{ee}^{\psi(3770) \rightarrow D\bar{D}}$ .

### $\pi^0$ Tracking

A 2.0 % change [52] was applied for each  $\pi^0$  in a given decay mode. The summed contribution for each mode is applied to  $\epsilon_m$  in Equation (5.19). As this is an overall scale change, the only variable significantly affected is  $\Gamma_{ee}^{\psi(3770) \rightarrow D\bar{D}}$ .

### $K_S^0$ Tracking

A 1.5 % change [53] was applied for each  $K_S^0$  in a given decay mode. The summed contribution for each mode is applied to  $\epsilon_m$  in Equation (5.19). As this is an overall scale change, the only variable significantly affected is  $\Gamma_{ee}^{\psi(3770) \rightarrow D\bar{D}}$ .

### Single Tag Fitting

A mode-dependent change [44] was applied to  $N$  in Equation (5.21). Differences from fitting were obtained by Derrick Toth after examining the use of single-Gaussian convolved signal shapes, and are shown in Table 5.13. The changes applied were obtained from the sums of the mode-dependent values averaged over their efficiencies. As this is an overall scale change, the only variable significantly affected is  $\Gamma_{ee}^{\psi(3770) \rightarrow D\bar{D}}$ .

Tag Mode	Difference (%)
$D^0 \rightarrow K^- \pi^+$	0.27
$D^0 \rightarrow K^- \pi^+ \pi^0$	0.10
$D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$	0.47
$D^0$ Average:	0.25 %
$D^+ \rightarrow K^- \pi^+ \pi^+$	0.20
$D^+ \rightarrow K^- \pi^+ \pi^+ \pi^0$	0.00
$D^+ \rightarrow K_S^0 \pi^+$	0.17
$D^+ \rightarrow K_S^0 \pi^+ \pi^0$	0.29
$D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-$	0.17
$D^+ \rightarrow K^+ K^- \pi^+$	0.74
$D^+$ Average:	0.20 %

Table 5.13: Single-tag fitting differences by mode.  
The total  $D^0$  and  $D^+$  values are averaged over the efficiencies for each mode.

### PDG Branching Fractions

A mode-dependent change based on the PDG branching fraction errors (Table 5.9) was applied to  $\epsilon_m$  in Equation (5.19). The branching fraction for each mode was changed by its uncertainty. As this is an overall scale change, the only variable significantly affected is  $\Gamma_{ee}^{\psi(3770) \rightarrow D\bar{D}}$ .

### Meson Radii

The most uncertain parameters used in the analysis are the radii of the mesons  $\psi(2S)$  and  $\psi(3770)$ . We take the same values as used by KEDR, however, each of these is quoted to have an  $\sim 25\%$  uncertainty. With this, we adjust the two radii values up or down by 25% over the four possible combinations (both up, both down, and each opposite). The maximum deviations from the nominal method seen across all four cases is used as the systematic uncertainty. Due to the high level of uncertainty on these parameters, this effect provides one of the largest systematic sources in the analysis.

### MC Iteration\*

In generating MC for this analysis, the  $D\bar{D}$  samples used a modified form of KKMC which generates events based off an input Born cross-section shape for the  $\psi(3770)$ .

However, as this shape is also the final output of the procedure, only an estimation of the results can be used for generation. To assess the variation from the input shape, we compared the output fit parameters to those used in the generation process. This process used the Exponential method, and the results are shown in Table 5.14. The numbers listed are from an earlier iteration of the MC than shown in Section 5.6, but the consistency seen is representative of all iterations. Very little difference is seen in the primary fitting parameters of the  $\psi(3770)$ . These similarities show the fit result values converging, even after only a single iteration. From this, we treat variations due to MC iteration as negligible.

Parameter		KKMC Input	Fit Results	Difference
$M^{\psi(3770)}$	[GeV]	$3.7815 \pm 0.0003$	$3.7814 \pm 0.0003$	0.0001
$\Gamma^{\psi(3770)}$	[MeV]	$24.887 \pm 0.686$	$24.839 \pm 0.681$	0.048
$\Gamma_{ee}^{\psi(3770) \rightarrow D\bar{D}}$	[eV]	$217.55 \pm 11.18$	$214.65 \pm 11.10$	2.90
$\phi^{\psi(3770)}$		$3.6374 \pm 0.0513$	$3.6375 \pm 0.0518$	0.0001
$F_{\text{NR}}$		$21.394 \pm 1.866$	$20.147 \pm 1.765$	0.992
$a_{\text{NR}}$		$-1.6202 \pm 0.5271$	$-1.5265 \pm 0.5119$	0.0937

Table 5.14: Comparison of input and output fit parameters.

The MC generation is done using the Exponential form factor model as an input Born cross section to generate  $D\bar{D}$  events using KKMC.

### MC ISR Generation\*

To compare to the generation process of KKMC, we also generated alternative MC samples of  $D\bar{D}$  using the ConExc ISR generator. This process used an input Born level shape identical to a previous iteration produced with KKMC. Each of the background samples used (such as  $q\bar{q}$  and  $\tau^+\tau^-$ ) were the same as in the nominal procedure. The cross section results using the VDM model are shown in Table 5.15, and provide  $\psi(3770)$  fit parameters within the statistical errors of the nominal method. From this, we treat variations due to the MC ISR generator as negligible.

Parameter		ConExc Fit Results	KKMC Fit Results
$M^{\psi(3770)}$	[GeV]	$3.7803 \pm 0.0003$	$3.7804 \pm 0.0003$
$\Gamma^{\psi(3770)}$	[MeV]	$23.784 \pm 0.616$	$23.732 \pm 0.612$
$\Gamma_{ee}^{\psi(3770) \rightarrow D\bar{D}}$	[eV]	$204.68 \pm 10.28$	$207.35 \pm 10.02$
$\phi^{\psi(3770)}$		$3.5954 \pm 0.0559$	$3.5952 \pm 0.0525$
$\Gamma_{ee}^{\psi(2S)}$	[MeV]	$12.229 \pm 1.336$	$14.070 \pm 1.431$
$F_0$		$-2.3415 \pm 0.4898$	$-2.0768 \pm 0.4924$

Table 5.15: Comparison of output fit parameters between ISR generators.

The MC generation is done using both the ConExc and KKMC generators and the final VDM fit results are shown. The values shown are from a previous iteration than the final results, but the output from each method remains very similar.

### Intermediate Resonances\*

In looking at the mode  $D^+ \rightarrow K^- \pi^+ \pi^+$ , we also analyzed the contribution of intermediate resonances, like the  $\rho^0$ . Using the  $2.93 \text{ fb}^{-1}$  data sample of  $\psi(3770)$  events at  $E_{\text{cm}} = 3.773 \text{ GeV}$ , we split the signal region of this mode based on a  $1.0 \text{ (GeV)}^2$  cut in both the invariant masses of  $K\pi$  and  $\pi\pi$  separately. These cuts were chosen to separate the sample into distinctly different regions, as can be seen in Figure 5.10. Fitting the signal distributions for each of these subsamples, we found no statistically significant deviations in the measured yields. From this, we treat variations due to intermediate resonances as negligible.




Figure 5.10: The  $K\pi$  vs.  $\pi\pi$  invariant masses for the mode  $D^+ \rightarrow K^- \pi^+ \pi^+$ . The data used are from the on-peak  $\psi(3770)$  sample, as the statistics are significantly higher than for the scan data.

### Non-Resonant Form Factor

In addition to the systematics described above, there is a significant source of uncertainty coming from the non-resonant form factor used. Each of the models examined, Exponential and VDM, provided quality fit results for the cross section shapes. From this, we conservatively assign an uncertainty equal to the differences in fit parameters provided by these two methods. Following the example of KEDR, we treat this as separate from the other systematics, and list it as a model uncertainty.

Systematic	$M^{\psi(3770)}$ [%]	$\Gamma^{\psi(3770)}$ [%]	$\Gamma_{ee}^{\psi(3770) \rightarrow D\bar{D}}$ [%]	$\phi^{\psi(3770)}$ [%]
Luminosity	0.000	0.004	1.005	0.014
$K^\pm/\pi^\pm$ Tracking	0.000	0.008	2.646	0.033
$\pi^0$ Tracking	0.000	0.012	0.746	0.028
$K_S^0$ Tracking	0.000	0.004	0.260	0.019
Single Tag Fits	0.000	0.012	0.213	0.008
PDG Errors	0.000	0.017	2.840	0.036
Meson Radii	0.016	2.411	3.512	1.477
Form Factor <sup>†</sup>	0.034	7.909	8.016	3.613
Total [%]	0.016	2.411	5.389	1.479
Relative Stat. Error [ $\sigma$ ]	3.000	1.088	1.342	1.229

Table 5.16: Systematic uncertainties relative to the measured parameters of the  $\psi(3770)$ .

<sup>†</sup>The Form Factor is treated as an uncertainty on the model, and is not included in the total systematic error.

### 5.7.2 Cross Section Systematics

In addition to the statistical uncertainties provided in Table 5.12, we also provide systematic uncertainties on the cross section measurements. These are calculated from the systematic shifts ( $\Delta S$ ) which affect Equation (5.21): Luminosity,  $K^\pm/\pi^\pm$  Tracking,  $\pi^0$  Tracking,  $K_S^0$  Tracking, and Single Tag Fits. For the luminosity, this value is the same 1% shift used previously. The others are calculated by weighting the efficiency from each decay mode:

$$\Delta S = \frac{\sum_m \epsilon_m \Delta S_m}{\sum_m \epsilon_m} \quad (5.22)$$

The value of  $\Delta S_m$  for  $K^\pm/\pi^\pm$  Tracking is 1.0% per  $K^\pm$  or  $\pi^\pm$  in each decay mode, and defined similarly for the other systematics using the previously listed values. The calculated shifts are listed in Table 5.17, and the resulting cross section values are listed in Table 5.18.

Systematic	$\Delta S(D^0) [\%]$	$\Delta S(D^+) [\%]$
Luminosity	1.00	1.00
$K^\pm/\pi^\pm$ Tracking	2.58	2.59
$\pi^0$ Tracking	0.94	0.63
$K_S^0$ Tracking	0.00	0.42
Single Tag Fits	0.25	0.20
PDG Error	0.31	0.18
Total	2.95	2.89

Table 5.17: Systematics shifts affecting the cross section measurements.  
The total shifts are used to calculate the systematic uncertainties of the  $D^0$  and  $D^+$  cross sections.

$E_{\text{mid}}$ [GeV]	$\sigma_{D^0\bar{D}^0}^{RC}$ [nb]	$\sigma_{D^+D^-}^{RC}$ [nb]
3.7342	$0.164 \pm 0.046 \pm 0.005$	-
3.7368	$0.218 \pm 0.068 \pm 0.006$	-
3.7447	$0.765 \pm 0.070 \pm 0.023$	$0.150 \pm 0.038 \pm 0.004$
3.7483	$0.838 \pm 0.064 \pm 0.025$	$0.360 \pm 0.048 \pm 0.010$
3.7501	$0.968 \pm 0.053 \pm 0.029$	$0.439 \pm 0.040 \pm 0.013$
3.7517	$1.237 \pm 0.052 \pm 0.036$	$0.574 \pm 0.040 \pm 0.017$
3.7534	$1.377 \pm 0.051 \pm 0.041$	$0.748 \pm 0.042 \pm 0.022$
3.7556	$1.566 \pm 0.055 \pm 0.046$	$0.797 \pm 0.045 \pm 0.023$
3.7562	$1.629 \pm 0.051 \pm 0.048$	$0.914 \pm 0.042 \pm 0.026$
3.7592	$1.948 \pm 0.052 \pm 0.057$	$1.192 \pm 0.045 \pm 0.034$
3.7624	$2.385 \pm 0.058 \pm 0.070$	$1.534 \pm 0.051 \pm 0.044$
3.7650	$2.715 \pm 0.071 \pm 0.080$	$1.882 \pm 0.066 \pm 0.054$
3.7676	$3.102 \pm 0.087 \pm 0.092$	$2.200 \pm 0.081 \pm 0.064$
3.7713	$3.406 \pm 0.101 \pm 0.100$	$2.634 \pm 0.098 \pm 0.076$
3.7742	$3.714 \pm 0.111 \pm 0.110$	$3.067 \pm 0.111 \pm 0.089$
3.7775	$3.692 \pm 0.111 \pm 0.109$	$3.078 \pm 0.111 \pm 0.089$
3.7802	$3.444 \pm 0.104 \pm 0.102$	$2.599 \pm 0.100 \pm 0.075$
3.7829	$2.791 \pm 0.090 \pm 0.082$	$2.206 \pm 0.088 \pm 0.064$
3.7869	$1.995 \pm 0.072 \pm 0.059$	$1.627 \pm 0.073 \pm 0.047$
3.7891	$1.361 \pm 0.058 \pm 0.040$	$1.178 \pm 0.062 \pm 0.034$
3.7926	$0.920 \pm 0.045 \pm 0.027$	$0.680 \pm 0.045 \pm 0.020$
3.7970	$0.562 \pm 0.037 \pm 0.017$	$0.414 \pm 0.039 \pm 0.012$
3.8003	$0.424 \pm 0.035 \pm 0.013$	$0.241 \pm 0.037 \pm 0.007$
3.8024	$0.325 \pm 0.038 \pm 0.010$	$0.236 \pm 0.043 \pm 0.007$
3.8070	$0.352 \pm 0.050 \pm 0.010$	$0.289 \pm 0.051 \pm 0.008$
3.8093	$0.233 \pm 0.050 \pm 0.007$	$0.132 \pm 0.055 \pm 0.004$
3.8135	$0.060 \pm 0.049 \pm 0.002$	$0.153 \pm 0.066 \pm 0.004$
3.8153	$0.056 \pm 0.051 \pm 0.002$	$0.089 \pm 0.063 \pm 0.003$
3.8229	$0.140 \pm 0.083 \pm 0.004$	$0.197 \pm 0.107 \pm 0.006$
3.8320	$0.069 \pm 0.086 \pm 0.002$	$0.046 \pm 0.099 \pm 0.001$
3.8390	$0.237 \pm 0.105 \pm 0.007$	$0.254 \pm 0.124 \pm 0.007$
3.8494	$0.186 \pm 0.104 \pm 0.005$	$0.186 \pm 0.118 \pm 0.005$
3.8555	$0.337 \pm 0.111 \pm 0.010$	$0.273 \pm 0.104 \pm 0.008$
3.8632	$0.340 \pm 0.127 \pm 0.010$	$0.099 \pm 0.091 \pm 0.003$

Table 5.18: Measurements of the  $D^0 / D^+$  cross sections.  
The first errors are statistical and the second are systematic.

## 5.8 Results

After incorporating the systematic and model uncertainties, the total results for the main  $\psi(3770)$  parameters are shown in Table 5.19. The results shown are from the VDM model, as we treat this as the nominal results. The Exponential model is used as a measure of uncertainty, however, the quality of fits found by this approach means it cannot be excluded as a viable option.

$M^{\psi(3770)}$	$3780.8 \pm 0.2 \pm 0.6 \pm 1.3$	[MeV]
$\Gamma^{\psi(3770)}$	$24.1 \pm 0.5 \pm 0.6 \pm 1.9$	[MeV]
$\Gamma_{ee}^{\psi(3770) \rightarrow D\bar{D}}$	$216 \pm 9 \pm 11 \pm 17$	[eV]
$\phi^{\psi(3770)}$	$207 \pm 3 \pm 3 \pm 7$	[ $^\circ$ ]

Table 5.19: Final results for the  $\psi(3770)$  parameters.

The first error listed is statistical, the second is systematic, and the third is from the form factor model.

Additionally, since this analysis is based on an approach developed by the KEDR collaboration, a comparison to their results is also shown in table 5.20. For their measurement of  $\Gamma_{ee}^{\psi(3770)}$ , two solutions were found with very close  $\chi^2$  values, so both are quoted in their final results. With the larger statistics available at BESIII, no alternate solution was found during searches over the parameter space.

Method	$M^{\psi(3770)}$ [MeV]	$\Gamma^{\psi(3770)}$ [MeV]	$\Gamma_{ee}^{\psi(3770) \rightarrow D\bar{D}}$ [eV]
Exponential	$3782.1 \pm 0.3 \pm 0.6$	$26.0 \pm 0.6 \pm 0.7$	$233 \pm 10 \pm 13$
VDM	$3780.8 \pm 0.2 \pm 0.6$	$24.1 \pm 0.6 \pm 0.6$	$216 \pm 9 \pm 12$
KEDR	$3779.2^{+1.8+0.5+0.3}_{-1.7-0.7-0.3}$	$24.9^{+4.6+0.5+0.2}_{-4.0-0.6-0.9}$	$154^{+79+17+13}_{-58-9-25},$ $414^{+72+24+90}_{-80-26-10}$
PDG	$3773.15 \pm 0.33$	$27.2 \pm 0.9$	$[262 \pm 18] \times \mathcal{B}_{D\bar{D}}$

Table 5.20: Fit results compared to the KEDR results and the PDG.

The first errors listed are statistical, while the second are systematic. In the case of KEDR, the third error is from the model.

We can also compare our results to the cross section values found in the previous analysis of BESIII's on-peak  $\psi(3770)$  data sample of  $2.93 \text{ fb}^{-1}$  by Derrick Toth [44]. The values displayed on Figures 5.6 and 5.7 are his double-tag (DT) values. Each of these cross sections are shown in Table 5.21. Further analysis to better understand

the differences seen is still in progress, however both methods used in this analysis are within  $\sim 1\sigma$  of the high statistics method.

Model	$\sigma_{D^0 \overline{D}^0}$ [nb]	$\sigma_{D^+ D^-}$ [nb]
Derrick (DT)	$3.615 \pm 0.010 \pm 0.035$	$2.830 \pm 0.011 \pm 0.026$
Exponential	$3.662 \pm 0.131 \pm 0.108$	$2.947 \pm 0.118 \pm 0.085$
VDM	$3.748 \pm 0.131 \pm 0.111$	$2.951 \pm 0.118 \pm 0.085$

Table 5.21: Comparison of cross section calculations at  $E_{\text{cm}} = 3.7732 \text{ GeV}$ . The values measured previously used a  $2.93 \text{ fb}^{-1}$  data sample of  $\psi(3770)$  events to reconstruct double-tag (DT) decays. The first errors listed are statistical, while the second is systematic.

## Chapter 6

# Measurement of Hadronic Production and $\Gamma(\psi(3770) \rightarrow \text{non-}D\bar{D})$

The second half of this analysis concerns the measurement of hadronic production. Namely, we are examining hadronic events in the regions above the  $D\bar{D}$  threshold (scan data) and below the  $\psi(2S)$  peak (continuum data). After identifying the total number of hadronic events in data, we can determine  $\sigma_{\psi(3770) \rightarrow \text{non-}D\bar{D}}$  in the scan region by subtracting off all other known decay processes. Using this, along with the previous values of  $\sigma_{\psi(3770) \rightarrow D\bar{D}}$ , we can measure the non- $D\bar{D}$  branching fraction:

$$\Gamma(\psi(3770) \rightarrow \text{non-}D\bar{D}) = \frac{\sigma_{\psi(3770) \rightarrow \text{non-}D\bar{D}}}{\sigma_{\psi(3770) \rightarrow D\bar{D}} + \sigma_{\psi(3770) \rightarrow \text{non-}D\bar{D}}}. \quad (6.1)$$

### 6.1 Data and Monte Carlo Samples

#### 6.1.1 Data Samples

The data used for this analysis was also produced by BEPCII and collected by BESIII. The samples used include continuum data taken at 3.650 GeV in 2009 (old continuum), as well as multiple other continuum points taken around this energy in 2013 (new continuum). We also use Round 1 (R1) and Round 2 (R2) of the high-statistics  $\psi(3770)$

data taken in 2010 and 2011, respectively. Each of these samples, and their measured luminosities, can be seen in Table 6.1. The values of luminosity were measured during a previous version of this analysis using the procedure described in Section 5.3.3. The labels given to each continuum point are the intended energy targets, but generally differ from their true values. In addition to these datasets, the scan data described previously (Section 5.3.1) is also used.

### 6.1.2 Center-of-Mass Energy Measurement

As before, a precise measurement of each energy point is vital to the accuracy of the final results. Most notably, due to the rapidly increasing shape of the  $\psi(2S)$  cross section near the high end of the continuum points, the value of the 3671 (New) point is highly susceptible to the assumed cross section of the  $\psi(2S)$ . Following the procedure of Section 5.3.2, we measured the  $E_{\text{cm}}$  value of each continuum point. This resulted in a 4 MeV to 6 MeV shift downwards for each of the new continuum points, but virtually no shift for the old continuum point. The measured energies and luminosities of each sample are shown in Table 6.1.

Sample Name	$E_{\text{cm}}$ [GeV]	Luminosity pb $^{-1}$
3500 (New)	3.496	$3.680 \pm 0.009$
3542 (New)	3.538	$3.481 \pm 0.009$
3600 (New)	3.596	$0.395 \pm 0.019$
3650 (New)	3.644	$5.420 \pm 0.009$
3671 (New)	3.665	$4.669 \pm 0.009$
3650 (Old)	3.650	$44.334 \pm 0.009$
$\psi(3770)$ (R1)	3.773	$926.922 \pm 0.092$
$\psi(3770)$ (R2)	3.773	$1978.920 \pm 0.091$

Table 6.1: Data samples used for the inclusive measurement.

While the 3600 (New) sample was intended to be similar in luminosity to the other continuum points, accelerator issues inhibited the data collection procedure. The new continuum points also have a much smaller luminosity compared to the other datasets used in this analysis.

## 6.2 Event Selection

In order to select the number of hadronic events in each sample, we apply a variety of cuts. For charged tracks in the MDC, these include the cuts shown in Table 6.2.

Vertex ( $xy$ )	$ V_{xy}  < 1 \text{ cm}$
Vertex ( $z$ )	$ Vz  < 10 \text{ cm}$
MDC Angle	$ \cos \theta  < 0.93$

Table 6.2: Selection cuts on charged tracks used to count hadronic events.

For neutral tracks in the EMC, these include the cuts shown in Table 6.3.

Minimum Energy (Barrel)	$E_{\text{EMC}} > 25 \text{ MeV}$	$( \cos \theta  < 0.80)$
Minimum Energy (Endcap)	$E_{\text{EMC}} > 50 \text{ MeV}$	$(0.86 <  \cos \theta  < 0.92)$
TDC Timing	$(0 \leq t \leq 14) \times 50 \text{ ns}$	

Table 6.3: Selection cuts on neutral tracks used to count hadronic events.

To reject background events of  $e^+e^- \rightarrow e^+e^-$  or  $e^+e^- \rightarrow \gamma\gamma$ , we also employ cuts related to the highest energy and highest momentum tracks in the event. These are listed in Table 6.4.

Highest Energy	$\cos \theta_+^{\max} < 0.8$	$(N_{\text{tracks}} = 2)$
	$\cos \theta_-^{\max} > -0.8$	
Highest Momentum	$\cos \theta_+^{\max} < 0.8 \text{ or } (p/E_{\text{cm}})_+^{\max} \leq 0.3$	$(N_{\text{tracks}} = 3, 4)$
	$\cos \theta_-^{\max} > -0.8 \text{ or } (p/E_{\text{cm}})_-^{\max} \leq 0.3$	
Highest Momentum	$0.8 \leq (E_{\text{EMC}}/p)_+^{\max} \leq 1.1$	
	$0.8 \leq (E_{\text{EMC}}/p)_-^{\max} \leq 1.1$	

Table 6.4: Selection cuts to remove Bhabha and two-photon backgrounds.

The  $+$  and  $-$  denote positively and negatively charged tracks, respectively. The  $^{\max}$  notation indicates the highest energy or momenta track for the corresponding charge. The energy cuts depend on the total number of charged tracks in the event,  $N_{\text{tracks}}$ .

After applying these sets of cuts, there are three groups of cuts which are considered: Standard (SHAD), Loose (LHAD), and Tight (THAD). For the nominal procedure, SHAD is used, while LHAD and THAD are for future systematic considerations. The cuts included in each of these sets are shown in Tables 6.5 to 6.7. These apply to the number of charged tracks ( $N_{\text{tracks}}$ ), the visible energy ( $E_{\text{vis}}$ ), the total visible momentum

in the  $z$ -direction ( $p_{z\text{vis}}$ ), the maximum shower energy ( $E_{\text{EMC}}^{\text{max}}$ ), and the total shower energy ( $E_{\text{EMC}}^{\text{tot}}$ ). Here, ‘visible’ refers to the sum over charged and neutral tracks.

Number of Tracks	$N_{\text{tracks}} > 2$	
Visible Energy	$(E_{\text{vis}}/E_{\text{cm}}) > 0.3$	
Visible Momentum	$(p_{z\text{vis}}/E_{\text{vis}}) < 0.6$	$(N_{\text{tracks}} = 3, 4)$
Maximum Shower Energy	$(E_{\text{EMC}}^{\text{max}}/E_{\text{beam}}) < 0.75$	$(N_{\text{tracks}} = 3, 4)$
Total Shower Energy	$0.25 < (E_{\text{EMC}}^{\text{tot}}/E_{\text{cm}}) < 0.75$	$(N_{\text{tracks}} = 3)$
	$0.15 < (E_{\text{EMC}}^{\text{tot}}/E_{\text{cm}}) < 0.75$	$(N_{\text{tracks}} = 4)$

Table 6.5: Standard selection cuts (SHAD) for counting hadronic events.

Number of Tracks	$N_{\text{tracks}} > 1$	
Visible Energy	$(E_{\text{vis}}/E_{\text{cm}}) > 0.4$	$(N_{\text{tracks}} = 2)$
	$(E_{\text{vis}}/E_{\text{cm}}) > 0.3$	$(N_{\text{tracks}} \geq 3)$
Visible Momentum	$(p_{z\text{vis}}/E_{\text{vis}}) < 0.3$	$(N_{\text{tracks}} = 2)$
	$(p_{z\text{vis}}/E_{\text{vis}}) < 0.6$	$(N_{\text{tracks}} = 3, 4)$
Maximum Shower Energy	$(E_{\text{EMC}}^{\text{max}}/E_{\text{beam}}) < 0.50$	$(N_{\text{tracks}} = 2)$
	$(E_{\text{EMC}}^{\text{max}}/E_{\text{beam}}) < 0.75$	$(N_{\text{tracks}} = 3, 4)$
Total Shower Energy	$0.25 < (E_{\text{EMC}}^{\text{tot}}/E_{\text{cm}}) < 0.75$	$(N_{\text{tracks}} = 2, 3)$
	$0.15 < (E_{\text{EMC}}^{\text{tot}}/E_{\text{cm}}) < 0.75$	$(N_{\text{tracks}} = 4)$

Table 6.6: Loose selection cuts (LHAD) for counting hadronic events.

Number of Tracks	$N_{\text{tracks}} > 3$	
Visible Energy	$(E_{\text{vis}}/E_{\text{cm}}) > 0.4$	
Visible Momentum	$(p_{z\text{vis}}/E_{\text{vis}}) < 0.6$	$(N_{\text{tracks}} = 4)$
Maximum Shower Energy	$(E_{\text{EMC}}^{\text{max}}/E_{\text{beam}}) < 0.75$	$(N_{\text{tracks}} = 4, 5)$
Total Shower Energy	$0.15 < (E_{\text{EMC}}^{\text{tot}}/E_{\text{cm}}) < 0.75$	$(N_{\text{tracks}} = 4)$
	$0.00 < (E_{\text{EMC}}^{\text{tot}}/E_{\text{cm}}) < 0.75$	$(N_{\text{tracks}} = 5)$

Table 6.7: Tight selection cuts (THAD) for counting hadronic events.

### 6.3 Hadron Counting

In order to determine the total number of hadronic events in each data sample, we use the average velocity in the  $z$ -direction. Signal events should have tracks which average to zero around the collision point, while background events will have randomly oriented

momenta. The fits are done using a double Gaussian shape for the signal and a 2<sup>nd</sup> order polynomial for the background.




Figure 6.1: The number of hadrons found in the 3650 (Old) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).

## 6.4 Background Subtraction

To precisely determine the number of hadronic events in the old continuum sample, we must subtract off a variety of backgrounds. The samples considered for this measurement include two-track QED processes ( $e^+e^-$ ,  $\mu^+\mu^-$ ,  $\tau^+\tau^-$ ,  $\gamma\gamma$ ), radiative  $J/\psi$  ( $\gamma J/\psi$ ), two photon fusion ( $2\gamma$ ), and events coming from  $\psi(2S)$ . Initially, we assume the  $\psi(2S)$  has a standard Breit-Wigner shape.

Each background contributes to the total number of reconstructed events based on their cross section ( $\sigma$ ) and reconstruction efficiency ( $\epsilon_{\text{MC}}$ ):

$$N_{\text{had}} = \mathcal{L} \times \sigma \times \epsilon_{\text{MC}}. \quad (6.2)$$

The efficiency is simply the fraction of reconstructed tracks compared to the total generated in a given MC sample:

$$\epsilon_{\text{MC}} = \left( \frac{N_{\text{rec}}}{N_{\text{gen}}} \right) \quad (6.3)$$

The MC samples were generated with  $2.5 \times 10^3$  events for each of the 79 runs in the old continuum data for each of the included backgrounds. Each sample was analyzed with all three cut selection groups (see Section 6.2). The reconstruction efficiencies for each, along with their cross sections at 3.650 GeV are shown in Table 6.8.

3650 (Old) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$e^+e^-$	554.562	$0.0006 \pm 0.0002$	$0.0008 \pm 0.0002$	$0.0001 \pm 0.0001$
$\mu^+\mu^-$	5.560	$0.0033 \pm 0.0004$	$0.0044 \pm 0.0005$	$0.0029 \pm 0.0004$
$\tau^+\tau^-$	1.844	$12.8351 \pm 0.0255$	$28.7692 \pm 0.0382$	$9.9371 \pm 0.0224$
$\gamma J/\psi$	1.260	$45.9222 \pm 0.0482$	$55.1722 \pm 0.0529$	$34.1250 \pm 0.0416$
$\gamma\gamma$	21.530	$0.0009 \pm 0.0002$	$0.0010 \pm 0.0002$	$0.0005 \pm 0.0002$
$2\gamma$	1.257	$2.4109 \pm 0.0110$	$4.6297 \pm 0.0153$	$1.6468 \pm 0.0091$
$\psi(2S)^{\dagger}$	0.150	$62.9891 \pm 0.0078$	$69.2882 \pm 0.0082$	$51.6942 \pm 0.0071$

Table 6.8: Reconstruction of background samples for the old continuum data. These include standard QED two-track processes ( $e^+e^-$ ,  $\mu^+\mu^-$ ,  $\tau^+\tau^-$ ,  $\gamma\gamma$ ), radiative  $J/\psi$  ( $\gamma J/\psi$ ) and two photon fusion ( $2\gamma$ ) events, and a contribution coming from  $\psi(2S)$ .

$\dagger$  The  $\psi(2S)$  is assumed to have a standard Breit-Wigner shape.

Using each of these values, we can determine the total number of hadronic events in the data. This is done by subtracting the expected amount of background from the measured number of events passing each selection method in data. The results for the old continuum data are shown in Table 6.9. Given that the  $e^+e^-$ ,  $\mu^+\mu^-$ , and  $\gamma\gamma$  samples have contributions which are much smaller than the uncertainty on the total result, we exclude these from the final calculation.

3650 (Old) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$477001 \pm 691$	$546546 \pm 739$	$375380 \pm 613$
$e^+e^-*$	$149 \pm 43$	$187 \pm 48$	$12 \pm 12$
$\mu^+\mu^-*$	$8 \pm 1$	$11 \pm 1$	$7 \pm 1$
$\tau^+\tau^-$	$10490 \pm 30$	$23514 \pm 59$	$8122 \pm 25$
$\gamma J/\psi$	$25658 \pm 60$	$30826 \pm 71$	$19067 \pm 46$
$\gamma\gamma^*$	$9 \pm 2$	$10 \pm 2$	$4 \pm 1$
$2\gamma$	$1443 \pm 7$	$2771 \pm 11$	$986 \pm 6$
$\psi(2S)^{\dagger}$	$4175 \pm 9$	$4593 \pm 10$	$3427 \pm 7$
Hadrons	$435234 \pm 694$	$484842 \pm 745$	$343779 \pm 615$

Table 6.9: Hadronic events selected in the old continuum data. As expected, SHAD finds less events than LHAD and more than THAD.

\* The contribution is neglected for the total results.

$\dagger$  The  $\psi(2S)$  is assumed to have a standard Breit-Wigner shape.

## 6.5 Efficiency Extrapolation

Due to complicated interactions above the  $D\bar{D}$  threshold,  $q\bar{q}$  events not coming from  $\psi(3770)$  are currently not well modeled by our MC generators. In order to accurately count these events, we utilize the continuum data, as below the  $\psi(2S)$  the process is more reliable. Assuming the production of these events remains consistent over energy, we can use the measurements at lower energies to extrapolate to higher energies:

Measuring the hadronic events for the new continuum data follows exactly as for the old continuum data, but with the negligible backgrounds excluded. The number of hadrons found in each data sample can be seen in Figures 6.2 to 6.6. Reconstruction efficiencies are shown in Table 6.10 with the total results shown in Table 6.11.




Figure 6.2: The number of hadrons found in the 3500 (New) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure 6.3: The number of hadrons found in the 3542 (New) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure 6.4: The number of hadrons found in the 3600 (New) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure 6.5: The number of hadrons found in the 3650 (New) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure 6.6: The number of hadrons found in the 3671 (New) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).

3500 (New) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$\tau^+\tau^-$	0.000	-	-	-
$\gamma J/\psi$	1.831	$47.079 \pm 0.077$	$56.117 \pm 0.084$	$35.320 \pm 0.066$
$2\gamma$	1.240	$2.380 \pm 0.017$	$4.924 \pm 0.025$	$1.644 \pm 0.014$
$\psi(2S)^{\dagger}$	0.006	$62.989 \pm 0.008$	$69.288 \pm 0.008$	$51.694 \pm 0.007$

3542 (New) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$\tau^+\tau^-$	0.000	-	-	-
$\gamma J/\psi$	1.632	$47.188 \pm 0.072$	$56.430 \pm 0.079$	$35.355 \pm 0.063$
$2\gamma$	1.270	$2.386 \pm 0.016$	$5.046 \pm 0.024$	$1.633 \pm 0.013$
$\psi(2S)^{\dagger}$	0.009	$62.989 \pm 0.008$	$69.288 \pm 0.008$	$51.694 \pm 0.007$

3600 (New) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$\tau^+\tau^-$	1.262	$12.851 \pm 0.080$	$29.096 \pm 0.121$	$10.040 \pm 0.071$
$\gamma J/\psi$	1.412	$47.524 \pm 0.154$	$56.902 \pm 0.169$	$35.703 \pm 0.134$
$2\gamma$	1.311	$2.651 \pm 0.036$	$5.089 \pm 0.050$	$1.897 \pm 0.031$
$\psi(2S)^{\dagger}$	0.024	$62.989 \pm 0.008$	$69.288 \pm 0.008$	$51.694 \pm 0.007$

3650 (New) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$\tau^+\tau^-$	1.844	$12.964 \pm 0.033$	$28.939 \pm 0.049$	$10.154 \pm 0.029$
$\gamma J/\psi$	1.260	$47.414 \pm 0.063$	$57.043 \pm 0.069$	$35.701 \pm 0.055$
$2\gamma$	1.346	$2.410 \pm 0.014$	$4.675 \pm 0.020$	$1.682 \pm 0.012$
$\psi(2S)^{\dagger}$	0.110	$62.989 \pm 0.008$	$69.288 \pm 0.008$	$51.694 \pm 0.007$

3671 (New) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$\tau^+\tau^-$	2.026	$12.997 \pm 0.047$	$28.851 \pm 0.069$	$10.169 \pm 0.041$
$\gamma J/\psi$	1.205	$47.496 \pm 0.089$	$57.237 \pm 0.098$	$35.745 \pm 0.077$
$2\gamma$	1.361	$2.473 \pm 0.020$	$4.787 \pm 0.028$	$1.698 \pm 0.017$
$\psi(2S)^{\dagger}$	0.436	$62.989 \pm 0.008$	$69.288 \pm 0.008$	$51.694 \pm 0.007$

Table 6.10: Reconstruction of background samples for the new continuum data.

<sup>†</sup> The  $\psi(2S)$  is assumed to have a standard Breit-Wigner shape.

3500 (New) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$42106 \pm 205$	$47942 \pm 219$	$32999 \pm 182$
$\gamma J/\psi$	$3173 \pm 10$	$3782 \pm 11$	$2380 \pm 8$
$2\gamma$	$109 \pm 1$	$225 \pm 1$	$75 \pm 1$
$\psi(2S)^{\dagger}$	$13 \pm 1$	$14 \pm 1$	$14 \pm 1$
Hadrons	$38812 \pm 205$	$43921 \pm 219$	$30533 \pm 182$
3542 (New) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$50253 \pm 224$	$56812 \pm 238$	$39448 \pm 199$
$\gamma J/\psi$	$3450 \pm 9$	$4126 \pm 10$	$2585 \pm 7$
$2\gamma$	$136 \pm 1$	$287 \pm 1$	$93 \pm 1$
$\psi(2S)^{\dagger}$	$26 \pm 1$	$28 \pm 1$	$21 \pm 1$
Hadrons	$46641 \pm 224$	$52371 \pm 239$	$36749 \pm 199$
3600 (New) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$4293 \pm 66$	$4912 \pm 70$	$3285 \pm 57$
$\tau^+\tau^-$	$64 \pm 3$	$145 \pm 7$	$50 \pm 2$
$\gamma J/\psi$	$265 \pm 13$	$317 \pm 16$	$199 \pm 10$
$2\gamma$	$14 \pm 1$	$26 \pm 1$	$10 \pm 1$
$\psi(2S)^{\dagger}$	$6 \pm 1$	$7 \pm 1$	$5 \pm 1$
Hadrons	$3944 \pm 67$	$4417 \pm 72$	$3023 \pm 58$
3650 (New) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$58733 \pm 242$	$67000 \pm 259$	$46536 \pm 216$
$\tau^+\tau^-$	$1295 \pm 4$	$2892 \pm 7$	$1015 \pm 3$
$\gamma J/\psi$	$3239 \pm 7$	$3896 \pm 8$	$2439 \pm 6$
$2\gamma$	$176 \pm 1$	$341 \pm 2$	$123 \pm 1$
$\psi(2S)^{\dagger}$	$376 \pm 1$	$414 \pm 1$	$309 \pm 1$
Hadrons	$53647 \pm 242$	$59458 \pm 259$	$42652 \pm 216$
3671 (New) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$49868 \pm 223$	$56804 \pm 238$	$39537 \pm 199$
$\tau^+\tau^-$	$1229 \pm 5$	$2729 \pm 8$	$962 \pm 4$
$\gamma J/\psi$	$2671 \pm 7$	$3219 \pm 8$	$2010 \pm 6$
$2\gamma$	$157 \pm 1$	$304 \pm 2$	$108 \pm 1$
$\psi(2S)^{\dagger}$	$1282 \pm 3$	$1410 \pm 3$	$1052 \pm 2$
Hadrons	$44528 \pm 224$	$49141 \pm 239$	$35405 \pm 199$

Table 6.11: Hadronic events selected in the new continuum data.

<sup>†</sup> The  $\psi(2S)$  is assumed to have a standard Breit-Wigner shape.

Since the rate of  $q\bar{q}$  production varies smoothly with energy, the reconstruction efficiency (relative to the old continuum data) for a given  $E_{\text{cm}}$  point (in MeV) can be calculated as follows:

$$\frac{\epsilon(E_{\text{cm}})}{\epsilon(3650)} = \left[ \frac{N_{\text{had}}(E_{\text{cm}})}{N_{\text{had}}(3650)} \right] \left[ \frac{\mathcal{L}(3650)}{\mathcal{L}(E_{\text{cm}})} \right] \left[ \frac{E_{\text{cm}}}{3650} \right]^2. \quad (6.4)$$

This efficiency ratio is calculated for each of the new continuum points, and a linear fit is performed for each of the selection cut methods. From this slope, we can extrapolate to find the expected number of hadronic events for the scan data. As the old continuum data was taken at a relatively similar time as the scan data, we use it as a normalization point for the efficiency extrapolation. The results for each cut are shown in Figures 6.7 to 6.9, where the dashed black line is the fit to the new continuum, and the solid green line is the extrapolation to higher energies.

From these extrapolations, it is evident the highest energy point (3.665 GeV) is generally lower than the other new continuum points. This is most likely due to the assumption of a standard Breit-Wigner shape for the  $\psi(2S)$ . Recent experimental evidence indicates this resonance may be susceptible to interference effects which distort the shape away from its peak. If the actual shape is lower than expected for the higher energy continuum points, it would decrease the background contribution thereby raising the efficiency ratio. This behavior is explored in Section 6.7, as BESIII plans to take considerably more data across this region in the near future. For now, we continue with the default Breit-Wigner assumption, and apply the procedure for the  $\psi(3770)$  data. While the final results may not be indicative of the true values, it will provide limits for measuring various important other parameters of the  $\psi(3770)$ .

### Efficiency Extrapolation - SHAD




Figure 6.7: The extrapolation for SHAD events using the new continuum data. The new continuum points (blue) are fit using a linear slope (dashed black), then extrapolated to higher energies (solid green) based on the old continuum energy point at 3.650 GeV (cyan). The energy point for the  $\psi(3770)$  samples at 3.773 GeV is also shown (solid red).

### Efficiency Extrapolation - LHAD




Figure 6.8: The extrapolation for LHAD events using the new continuum data. The new continuum points (blue) are fit using a linear slope (dashed black), then extrapolated to higher energies (solid green) based on the old continuum energy point at 3.650 GeV (cyan). The energy point for the  $\psi(3770)$  samples at 3.773 GeV is also shown (solid red).

### Efficiency Extrapolation - THAD




Figure 6.9: The extrapolation for THAD events using the new continuum data. The new continuum points (blue) are fit using a linear slope (dashed black), then extrapolated to higher energies (solid green) based on the old continuum energy point at 3.650 GeV (cyan). The energy point for the  $\psi(3770)$  samples at 3.773 GeV is also shown (solid red).

## 6.6 Results for $\psi(3770)$ Data

The procedure for determining the hadronic events in the  $\psi(3770)$  data is similar to the continuum region, but with certain modifications introduced due to the different relevant backgrounds. Namely, above the  $D\bar{D}$  peak, contributions from  $D^0\bar{D}^0$  and  $D^+D^-$  events must be accounted for. Also, instead of the  $\psi(2S)$  component, there is now a background from radiative  $\psi(2S)$  production ( $\gamma\psi(2S)$ ). Lastly, due to the minimal contribution of two photon fusion events ( $2\gamma$ ) in this region, this component is neglected for the  $\psi(3770)$  samples. The counting of total hadronic events, however, functions identically to the continuum data, and the results are shown in Figures 6.10 and 6.11.




Figure 6.10: The number of hadrons found in the  $\psi(3770)$  (R1) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure 6.11: The number of hadrons found in the  $\psi(3770)$  (R2) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).

### 6.6.1 $D\bar{D}$ Correction

Since the  $D\bar{D}$  cross section is not a smooth scaling of energy (like  $q\bar{q}$ ), we must precisely determine its reconstruction efficiency using data. Instead of using the standard definition of  $\epsilon_{MC}$  for  $D\bar{D}$ , we apply a correction based on the multiplicity of  $D\bar{D}$  events in both data and MC. This is done by finding single-tagged  $D$  candidates and counting the number of tracks not used for reconstruction. By randomly sampling pairs of points from this distribution, and assuming the decays are uncorrelated, we can produce an average representation of multiplicity in  $D\bar{D}$  events. From this new distribution, selections methods for SHAD, LHAD, and THAD are performed based off number of tracks selected by the simplified cut criteria shown in Table 6.12 relative to the total. The corrections applied to each efficiency are the ratio of these selections in data and MC. As the  $\psi(3770)$  samples for R1 and R2 were taken at different times, they are treated separately for this process. The results for each are shown in Table 6.13 with the corresponding distributions shown in Figures 6.12 to 6.15.

Selection Method	Number of Tracks
SHAD	$N_{\text{tracks}} > 2$
LHAD	$N_{\text{tracks}} > 1$
THAD	$N_{\text{tracks}} > 3$

Table 6.12: Selection methods for the  $D\bar{D}$  efficiency correction.

Selection	$\psi(3770)$ R1		$\psi(3770)$ R2	
	$(\epsilon_{\text{Data}}/\epsilon_{\text{MC}})(D^0)$	$(\epsilon_{\text{Data}}/\epsilon_{\text{MC}})(D^+)$	$(\epsilon_{\text{Data}}/\epsilon_{\text{MC}})(D^0)$	$(\epsilon_{\text{Data}}/\epsilon_{\text{MC}})(D^+)$
SHAD	0.9751	0.9992	0.9759	0.9999
LHAD	0.9930	1.0018	0.9935	1.0024
THAD	0.9662	1.0064	0.9684	1.0108

Table 6.13: Efficiency corrections for the  $\psi(3770)$  samples.

The corrections are impactful for  $D^0$ , but minimal for  $D^+$ . This is due to the differences in their low-side other  $D$  multiplicities, as seen in Figures 6.12 to 6.15.




Figure 6.12: The other-side  $D^0$  tracks and corresponding  $D^0\overline{D}^0$  multiplicities for R1. The distribution of good tracks not used for reconstruction of single-tagged  $D^0$  particles (left) is randomly sampled for pairs of points which comprise the total multiplicity distribution (right). The tracks in the  $D^0\overline{D}^0$  multiplicity distribution are used to determine the efficiency correction based off the cuts in Table 6.12.




Figure 6.13: The other-side  $D^+$  tracks and corresponding  $D^+D^-$  multiplicities for R1. The distribution of good tracks not used for reconstruction of single-tagged  $D^+$  particles (left) is randomly sampled for pairs of points which comprise the total multiplicity distribution (right). The tracks in the  $D^+D^-$  multiplicity distribution are used to determine the efficiency correction based off the cuts in Table 6.12.




Figure 6.14: The other-side  $D^0$  tracks and corresponding  $D^0\bar{D}^0$  multiplicities for R2. The distribution of good tracks not used for reconstruction of single-tagged  $D^0$  particles (left) is randomly sampled for pairs of points which comprise the total multiplicity distribution (right). The tracks in the  $D^0\bar{D}^0$  multiplicity distribution are used to determine the efficiency correction based off the cuts in Table 6.12.




Figure 6.15: The other-side  $D^+$  tracks and corresponding  $D^+D^-$  multiplicities for R2. The distribution of good tracks not used for reconstruction of single-tagged  $D^+$  particles (left) is randomly sampled for pairs of points which comprise the convolved distribution (right). The number of tracks in the  $D^+D^-$  multiplicity distribution is used to determine the efficiency correction based off the cuts in Table 6.12.

### 6.6.2 Reconstruction Efficiencies

MC samples for each of the new backgrounds introduced above the  $D\bar{D}$  threshold were generated in the same way as for the continuum data. The reconstruction efficiencies for each are shown in Table 6.14. Values for the  $D^0\bar{D}^0$  and  $D^+D^-$  samples have already been multiplied by the correction factors discussed in Section 6.6.1.

For the cross section of  $\gamma\psi(2S)$ , rather than the radiative return formula (as used with  $\gamma J/\psi$ ), we use results from CLEO-c and BESIII to determine the value. These values are  $\sigma(e^+e^- \rightarrow \gamma\psi(2S), \psi(2S) \rightarrow \pi^+\pi^-J/\psi) = (1036 \pm 13 \pm 23) \text{ pb}$  from the CLEO-c measurement, and  $\Gamma(\psi(2S) \rightarrow \pi^+\pi^-J/\psi) = (34.43 \pm 0.30) \%$  after averaging the BESIII measurement with the value from the PDG. From this, we determine

$$\sigma(e^+e^- \rightarrow \gamma\psi') = \frac{\sigma(e^+e^- \rightarrow \gamma\psi', \psi' \rightarrow \pi^+\pi^-J/\psi)}{\Gamma(\psi' \rightarrow \pi^+\pi^-J/\psi)} = (3009 \pm 81) \text{ pb}. \quad (6.5)$$

Additionally, the  $D^0\bar{D}^0$  and  $D^+D^-$  cross sections are taken from the on-peak  $\psi(3770)$  measurement performed by Derrick Toth [44].

$\psi(3770)$ (R1) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	3.615	$73.9324 \pm 0.0142$	$79.8496 \pm 0.0147$	$60.3601 \pm 0.0128$
$D^+D^-$	2.830	$61.4048 \pm 0.0146$	$68.8212 \pm 0.0154$	$49.4007 \pm 0.0131$
$\tau^+\tau^-$	2.652	$12.7566 \pm 0.0253$	$28.0142 \pm 0.0374$	$9.8776 \pm 0.0222$
$\gamma J/\psi$	0.986	$46.6185 \pm 0.0206$	$56.2494 \pm 0.0227$	$34.7544 \pm 0.0178$
$\gamma\psi(2S)$	3.009	$63.2551 \pm 0.0137$	$69.9696 \pm 0.0144$	$51.5643 \pm 0.0123$

$\psi(3770)$ (R2) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	3.615	$74.5111 \pm 0.0097$	$80.3399 \pm 0.0101$	$61.0386 \pm 0.0088$
$D^+D^-$	2.830	$61.8444 \pm 0.0100$	$69.1974 \pm 0.0106$	$49.9163 \pm 0.0090$
$\tau^+\tau^-$	2.652	$12.8646 \pm 0.0254$	$28.2140 \pm 0.0376$	$10.0198 \pm 0.0224$
$\gamma J/\psi$	0.986	$47.0066 \pm 0.0146$	$56.6679 \pm 0.0161$	$35.1951 \pm 0.0127$
$\gamma\psi(2S)$	3.009	$63.7345 \pm 0.0097$	$70.4050 \pm 0.0102$	$52.1189 \pm 0.0088$

Table 6.14: Reconstruction of background samples for the  $\psi(3770)$  data.

### 6.6.3 Signal Amounts

Using the reconstruction efficiencies from Section 6.6.2, we can compute the contribution of background events for each sample. For the  $q\bar{q}$  component, we scale the number of hadronic events found in the 3650 (Old) sample based on equation (6.4) for both R1 and R2 separately. Due to the uncertainty on the extrapolation procedure, this becomes the dominant source of error on the resulting hadronic events. This is also highly susceptible to the assumption of the  $\psi(2S)$  shape for calculating the continuum points in the extrapolation fit. The resulting number of hadronic events is shown in Table 6.15.

$\psi(3770)$ (R1) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$15694505 \pm 3962$	$17722728 \pm 4210$	$12580701 \pm 3547$
$q\bar{q}^\dagger$	$8522688 \pm 71353$	$9330411 \pm 76320$	$6789405 \pm 61599$
$D^0\bar{D}^0$	$2477345 \pm 534$	$2675620 \pm 560$	$2022561 \pm 473$
$D^+D^-$	$1610764 \pm 414$	$1805311 \pm 442$	$1295875 \pm 366$
$\tau^+\tau^-$	$313542 \pm 622$	$688559 \pm 922$	$242781 \pm 547$
$\gamma J/\psi$	$425891 \pm 193$	$513875 \pm 213$	$317504 \pm 166$
$\gamma\psi(2S)$	$1764254 \pm 419$	$1951528 \pm 445$	$1438185 \pm 372$
Hadrons	$490569 \pm 71795$	$658730 \pm 76807$	$401064 \pm 61995$

$\psi(3770)$ (R2) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$33867464 \pm 5820$	$38217129 \pm 6182$	$27271529 \pm 5222$
$q\bar{q}^\dagger$	$18314683 \pm 154300$	$20015495 \pm 164688$	$14644571 \pm 133785$
$D^0\bar{D}^0$	$5330375 \pm 738$	$5747352 \pm 770$	$4366577 \pm 662$
$D^+D^-$	$3463499 \pm 583$	$3875291 \pm 620$	$2795485 \pm 520$
$\tau^+\tau^-$	$675063 \pm 1331$	$1480514 \pm 1972$	$525781 \pm 1175$
$\gamma J/\psi$	$916819 \pm 288$	$1105253 \pm 317$	$686446 \pm 249$
$\gamma\psi(2S)$	$3795113 \pm 603$	$4192317 \pm 636$	$3103459 \pm 541$
Hadrons	$1179686 \pm 155135$	$1589187 \pm 165610$	$991049 \pm 134532$

Table 6.15: Hadronic events selected in the  $\psi(3770)$  data.

<sup>†</sup> The  $q\bar{q}$  contribution is obtained using an extrapolation from the continuum region in which the  $\psi(2S)$  is assumed to have a standard Breit-Wigner shape.

### 6.6.4 Non- $D\bar{D}$ Branching Fraction (Breit-Wigner)

The resulting hadronic production is assumed to result from non- $D\bar{D}$  decays of the  $\psi(3770)$ . We can convert this into a cross section as follows:

$$\sigma(\psi(3770) \rightarrow \text{non-}D\bar{D}) = \frac{N_{\text{non-}D\bar{D}}}{\epsilon_{\text{non-}D\bar{D}} \times \mathcal{L}}. \quad (6.6)$$

For the efficiency of non- $D\bar{D}$  events, the value of  $\epsilon_{\text{MC}}$  for  $\gamma\psi(2S)$  is used due to the assumption of similar behavior in their decays. Using the standard Breit-Wigner assumption for the  $\psi(2S)$ , the results for the cross section and branching fraction of non- $D\bar{D}$  decays from  $\psi(3770)$  are shown in Tables 6.16 and 6.17.

Sample	$\sigma_{\text{non-}D\bar{D}}$ [nb] (SHAD)	$\sigma_{\text{non-}D\bar{D}}$ [nb] (LHAD)	$\sigma_{\text{non-}D\bar{D}}$ [nb] (THAD)
$\psi(3770)$ (R1)	$0.9892 \pm 0.1219$	$1.1679 \pm 0.1179$	$0.9925 \pm 0.1291$
$\psi(3770)$ (R2)	$1.0877 \pm 0.1224$	$1.2926 \pm 0.1183$	$1.1142 \pm 0.1298$
Lum. Weighted	$1.0563 \pm 0.1223$	$1.2528 \pm 0.1182$	$1.0754 \pm 0.1296$

Table 6.16: Cross sections for  $\psi(3770) \rightarrow \text{non-}D\bar{D}$  found using the  $\psi(3770)$  data. The  $q\bar{q}$  contributions were calculated using the Breit-Wigner formulation for the  $\psi(2S)$  component in the continuum extrapolation.

Sample	$\Gamma_{\text{non-}D\bar{D}}$ (SHAD)	$\Gamma_{\text{non-}D\bar{D}}$ (LHAD)	$\Gamma_{\text{non-}D\bar{D}}$ (THAD)
$\psi(3770)$ (R1)	$0.1331 \pm 0.0183$	$0.1534 \pm 0.0185$	$0.1334 \pm 0.0190$
$\psi(3770)$ (R2)	$0.1444 \pm 0.0186$	$0.1671 \pm 0.0189$	$0.1474 \pm 0.0193$
Lum. Weighted	$0.1408 \pm 0.0185$	$0.1627 \pm 0.0187$	$0.1430 \pm 0.0192$

Table 6.17: Branching fractions for  $\psi(3770) \rightarrow \text{non-}D\bar{D}$  found using the  $\psi(3770)$  data. The  $q\bar{q}$  contributions were calculated using the Breit-Wigner formulation for the  $\psi(2S)$  component in the continuum extrapolation.

## 6.7 $\psi(2S)$ Background Investigation

The most impactful assumption for the measurement of the branching fraction is the cross section shape of  $\psi(2S)$  in the continuum region. For the procedure thus far, we have assumed this shape to be a standard Breit-Wigner. Given the drop in relative efficiency for the 3671 (New) data point (see Figure 6.7), it is likely this overestimates the branching fraction (see Table 6.17) by subtracting off too large of a background component for the  $\psi(2S)$ .

As an alternative comparison, we use the ratio of cross section productions for the resonance peak and continuum values,

$$\frac{\sigma_{\text{res}}}{\sigma_{\text{cont}}(E_{\text{cm}})} = \frac{\sqrt{2\pi} (M_{\text{res}} - E_{\text{cm}})^2}{\Gamma_{\text{res}} \times \sigma_{E_{\text{cm}}}}, \quad (6.7)$$

where  $M_{\text{res}}$  and  $\Gamma_{\text{res}}$  are the mass and total width of the resonance, respectively, and  $\sigma_{E_{\text{cm}}}$  is the center-of-mass energy spread during collection of the data. For the  $\psi(2S)$ , with  $M_{\psi(2S)} = 3686.1 \text{ MeV}$  and  $\Gamma_{\psi(2S)} = 0.299 \text{ MeV}$ , assuming an energy spread of  $\sigma_{E_{\text{cm}}} = 1.5 \text{ MeV}$  gives  $\sigma_{\psi(2S)} \sim 2500 \times \sigma_{\text{cont}}(E_{\text{cm}})$  at  $E_{\text{cm}} = 3.665 \text{ GeV}$ .

In 2009, BESIII collected  $166.25 \text{ pb}^{-1}$  of data at  $E_{\text{cm}} = 3.686 \text{ GeV}$  and found  $(106.41 \pm 0.86) \times 10^6$  events of  $\psi(2S)$  decays [54], corresponding to  $\sigma_{\psi(2S)} \sim 640 \text{ nb}$ . Using Equation (6.7) for each of the continuum points, we obtain the cross section values shown in Table 6.18, where each generally is notably smaller than from the Breit-Wigner assumption.

Sample	$E_{\text{cm}}$ [GeV]	$\sigma_{\psi(2S)}$ [nb] (Ratio)	$\sigma_{\psi(2S)}$ [nb] (BW)
3500 (New)	3.496	0.0032	0.0056
3542 (New)	3.538	0.0052	0.0092
3600 (New)	3.596	0.0141	0.0244
3650 (New)	3.644	0.0646	0.1101
3671 (New)	3.665	0.2572	0.4359
3650 (Old)	3.650	0.0879	0.1495

Table 6.18: Cross sections of  $\psi(2S)$  calculated using two different methods.

### 6.7.1 Non- $D\bar{D}$ Branching Fraction ( $\psi(2S)$ from Data)

We repeat the procedure using the cross sections calculated based off Equation (6.7) in place of the Breit-Wigner values (see Table 6.18). This leads to the continuum extrapolations shown in Figures 6.16 to 6.18. The non- $D\bar{D}$  results are shown in Tables 6.19 and 6.20, where the branching fractions are lower on average by around 1.6% compared to the original method.

Sample	$\sigma_{\text{non-}D\bar{D}}$ [nb] (SHAD)	$\sigma_{\text{non-}D\bar{D}}$ [nb] (LHAD)	$\sigma_{\text{non-}D\bar{D}}$ [nb] (THAD)
$\psi(3770)$ (R1)	$0.8367 \pm 0.1224$	$1.0157 \pm 0.1184$	$0.8391 \pm 0.1297$
$\psi(3770)$ (R2)	$0.9353 \pm 0.1230$	$1.1406 \pm 0.1189$	$0.9609 \pm 0.1304$
Lum. Weighted	$0.9039 \pm 0.1228$	$1.1008 \pm 0.1187$	$0.9220 \pm 0.1302$

Table 6.19: Cross sections for  $\psi(3770) \rightarrow \text{non-}D\bar{D}$  found using the  $\psi(3770)$  data. The  $q\bar{q}$  contributions were calculated using  $\psi(2S)$  data for the  $\psi(2S)$  components in the continuum extrapolation.

Sample	$\Gamma_{\text{non-}D\bar{D}}$ (SHAD)	$\Gamma_{\text{non-}D\bar{D}}$ (LHAD)	$\Gamma_{\text{non-}D\bar{D}}$ (THAD)
$\psi(3770)$ (R1)	$0.1149 \pm 0.0180$	$0.1361 \pm 0.0181$	$0.1152 \pm 0.0188$
$\psi(3770)$ (R2)	$0.1267 \pm 0.0183$	$0.1504 \pm 0.0185$	$0.1297 \pm 0.0190$
Lum. Weighted	$0.1230 \pm 0.0182$	$0.1458 \pm 0.0183$	$0.1251 \pm 0.0190$

Table 6.20: Branching fractions for  $\psi(3770) \rightarrow \text{non-}D\bar{D}$  found using the  $\psi(3770)$  data. The  $q\bar{q}$  contributions were calculated using  $\psi(2S)$  data for the  $\psi(2S)$  components in the continuum extrapolation.

### Efficiency Extrapolation - SHAD




Figure 6.16: The extrapolation for SHAD events using  $\psi(2S)$  data.

The new continuum points (blue) are fit using a linear slope (dashed black), then extrapolated to higher energies (solid green) based on the old continuum energy point at 3.650 GeV (cyan). The energy point for the  $\psi(3770)$  samples at 3.773 GeV is also shown (solid red).

### Efficiency Extrapolation - LHAD




Figure 6.17: The extrapolation for LHAD events using  $\psi(2S)$  data.

The new continuum points (blue) are fit using a linear slope (dashed black), then extrapolated to higher energies (solid green) based on the old continuum energy point at 3.650 GeV (cyan). The energy point for the  $\psi(3770)$  samples at 3.773 GeV is also shown (solid red).

### Efficiency Extrapolation - THAD




Figure 6.18: The extrapolation for THAD events using  $\psi(2S)$  data.

The new continuum points (blue) are fit using a linear slope (dashed black), then extrapolated to higher energies (solid green) based on the old continuum energy point at 3.650 GeV (cyan). The energy point for the  $\psi(3770)$  samples at 3.773 GeV is also shown (solid red).

### 6.7.2 Non- $D\bar{D}$ Branching Fraction ( $\psi(2S)$ Excluded)

While the Breit-Wigner formulation likely provides an upper bound to the non- $D\bar{D}$  branching fractions, it is unclear how much lower the true values are. The modification detailed in Section 6.7 provides a likely estimation of the actual value, but previous experiments have measured wildly fluctuating values. In fact, some results have shown no evidence of a non- $D\bar{D}$  branching fraction at all. To obtain a potential lower bound from our procedure, we can repeat the measurement without including any contribution from  $\psi(2S)$  events. This will increase the extrapolated reconstruction efficiency fits, which subtracts off a larger amount of hadrons from the  $\psi(3770)$  data, and corresponds to a lower branching fraction. The results for this procedure are shown in Tables 6.21 and 6.22.

Sample	$\sigma_{\text{non-}D\bar{D}}$ [nb] (SHAD)	$\sigma_{\text{non-}D\bar{D}}$ [nb] (LHAD)	$\sigma_{\text{non-}D\bar{D}}$ [nb] (THAD)
$\psi(3770)$ (R1)	$0.6190 \pm 0.1232$	$0.7986 \pm 0.1192$	$0.6203 \pm 0.1305$
$\psi(3770)$ (R2)	$0.7179 \pm 0.1238$	$0.9239 \pm 0.1197$	$0.7421 \pm 0.1313$
Lum. Weighted	$0.6864 \pm 0.1236$	$0.8839 \pm 0.1195$	$0.7033 \pm 0.1311$

Table 6.21: Cross sections for  $\psi(3770) \rightarrow \text{non-}D\bar{D}$  found using the  $\psi(3770)$  data. The  $q\bar{q}$  contributions were calculated after excluding the  $\psi(2S)$  components in the continuum extrapolation.

Sample	$\Gamma_{\text{non-}D\bar{D}}$ (SHAD)	$\Gamma_{\text{non-}D\bar{D}}$ (LHAD)	$\Gamma_{\text{non-}D\bar{D}}$ (THAD)
$\psi(3770)$ (R1)	$0.0876 \pm 0.0178$	$0.1102 \pm 0.0176$	$0.0878 \pm 0.0187$
$\psi(3770)$ (R2)	$0.1002 \pm 0.0180$	$0.1254 \pm 0.0179$	$0.1033 \pm 0.0188$
Lum. Weighted	$0.0962 \pm 0.0179$	$0.1205 \pm 0.0178$	$0.0983 \pm 0.0188$

Table 6.22: Branching fractions for  $\psi(3770) \rightarrow \text{non-}D\bar{D}$  found using the  $\psi(3770)$  data. The  $q\bar{q}$  contributions were calculated after excluding the  $\psi(2S)$  components in the continuum extrapolation.

### Efficiency Extrapolation - SHAD




Figure 6.19: The extrapolation for SHAD events without including  $\psi(2S)$  events. The new continuum points (blue) are fit using a linear slope (dashed black), then extrapolated to higher energies (solid green) based on the old continuum energy point at 3.650 GeV (cyan). The energy point for the  $\psi(3770)$  samples at 3.773 GeV is also shown (solid red).

### Efficiency Extrapolation - LHAD




Figure 6.20: The extrapolation for LHAD events without including  $\psi(2S)$  events. The new continuum points (blue) are fit using a linear slope (dashed black), then extrapolated to higher energies (solid green) based on the old continuum energy point at 3.650 GeV (cyan). The energy point for the  $\psi(3770)$  samples at 3.773 GeV is also shown (solid red).

### Efficiency Extrapolation - THAD




Figure 6.21: The extrapolation for THAD events without including  $\psi(2S)$  events. The new continuum points (blue) are fit using a linear slope (dashed black), then extrapolated to higher energies (solid green) based on the old continuum energy point at 3.650 GeV (cyan). The energy point for the  $\psi(3770)$  samples at 3.773 GeV is also shown (solid red).

## 6.8 Results for Scan Data

In addition to the  $\psi(3770)$  data, we also examine the energy points of the scan data. While the statistics for these points are far lower, it will provide insight for the behavior of hadronic production with changing center-of-mass energy. We use  $\psi(2S)$  data in order to extrapolate the  $q\bar{q}$  events throughout this region (see Section 6.7.1).

Each scan data point used will be referred to by its name listed in Table 6.23. The cross sections for each background sample are also listed, where the  $D\bar{D}$  values are taken from the results in Table 5.18. To determine the cross sections of  $\gamma\psi(2S)$  at each point, the values were calculated using the standard radiative formula (as for  $\gamma J/\psi$ ), then scaled by the ratio of the data-driven value from Equation (6.5) to the value of this formula at 3.773 GeV.

The total hadronic counting fits for the scan data can be found in Appendix D. As the  $\psi(3770)$  R1 data was taken at a similar time as the scan data, we use the same  $D\bar{D}$  corrections from R1 for all of the scan data sample points. The reconstruction efficiencies and signal amounts can be found in Appendices E and F, respectively.

In addition to the non- $D\bar{D}$  branching fraction, we can also examine the inclusive hadronic cross section as a function of center-of-mass energy: Dividing the fit counts for each selection method from Appendix D by the luminosities from Table 5.3, we find the inclusive cross sections shown in Figure 6.22. As these values are independent of the extrapolation procedure, they are immediately ready for use in future analyses.

The non- $D\bar{D}$  cross sections and branching ratios calculated for each scan point are shown in Figures 6.23 and 6.24, respectively. For comparison, the same calculation for the  $\psi(3770)$  data is also shown on each of these plots. Due to the uncertain cross section shape of the  $\psi(2S)$  required to properly extrapolate to the scan data region, these values are not representative of a true measurement for the branching fraction. For now, we have done our best to ensure the accuracy of all other background components analyzed. This means, with an improved understanding of the  $\psi(2S)$  cross section in the continuum region (likely to be analyzed in the very near future, as BESIII has data taking planned for this region), the branching fraction value can easily be updated for a final result. From our analysis of the  $\psi(2S)$  background (see Section 6.7), we expect this value to be in the range 9.6 % to 14.1 % with our current estimate being 12.3 %.

Sample	$E_{\text{cm}}$ [GeV]	$D^0\bar{D}^0$ [nb]	$D^+D^-$ [nb]	$\tau^+\tau^-$ [nb]	$\gamma J/\psi$ [nb]	$\gamma\psi(2S)$ [nb]
3734 (Scan)	3.7342	0.164	0.000	2.467	1.060	5.359
3736 (Scan)	3.7368	0.218	0.000	2.475	1.055	5.094
3744 (Scan)	3.7447	0.765	0.150	2.514	1.039	4.428
3748 (Scan)	3.7483	0.838	0.360	2.522	1.032	4.180
3750 (Scan)	3.7501	0.968	0.439	2.533	1.029	4.065
3751 (Scan)	3.7517	1.237	0.574	2.542	1.025	3.969
3753 (Scan)	3.7534	1.377	0.748	2.553	1.022	3.871
3755 (Scan)	3.7556	1.566	0.797	2.562	1.018	3.752
3756 (Scan)	3.7562	1.629	0.914	2.573	1.017	3.721
3759 (Scan)	3.7592	1.948	1.192	2.584	1.011	3.572
3762 (Scan)	3.7624	2.385	1.534	2.598	1.005	3.426
3765 (Scan)	3.7650	2.715	1.882	2.613	1.000	3.315
3767 (Scan)	3.7676	3.102	2.200	2.627	0.995	3.212
3771 (Scan)	3.7713	3.406	2.634	2.641	0.989	3.075
3774 (Scan)	3.7742	3.714	3.067	2.656	0.983	2.975
3777 (Scan)	3.7775	3.692	3.078	2.669	0.978	2.870
3780 (Scan)	3.7802	3.444	2.599	2.683	0.973	2.788
3782 (Scan)	3.7829	2.791	2.206	2.696	0.968	2.712
3786 (Scan)	3.7869	1.995	1.627	2.709	0.961	2.605
3789 (Scan)	3.7891	1.361	1.178	2.722	0.957	2.550
3792 (Scan)	3.7926	0.920	0.680	2.739	0.951	2.467
3797 (Scan)	3.7970	0.562	0.414	2.755	0.944	2.370
3800 (Scan)	3.8003	0.424	0.241	2.771	0.938	2.301
3802 (Scan)	3.8024	0.325	0.236	2.784	0.935	2.260
3807 (Scan)	3.8070	0.352	0.289	2.795	0.927	2.174
3809 (Scan)	3.8093	0.233	0.132	2.807	0.923	2.133
3813 (Scan)	3.8135	0.060	0.153	2.819	0.917	2.063
3815 (Scan)	3.8153	0.056	0.089	2.829	0.914	2.034
3822 (Scan)	3.8229	0.140	0.197	2.859	0.902	1.920
3832 (Scan)	3.8320	0.069	0.046	2.887	0.888	1.799
3839 (Scan)	3.8390	0.237	0.254	2.914	0.877	1.716
3849 (Scan)	3.8494	0.186	0.186	2.939	0.862	1.604
3855 (Scan)	3.8555	0.337	0.273	2.967	0.853	1.545
3863 (Scan)	3.8632	0.340	0.099	2.988	0.843	1.476

Table 6.23: Energy values and background cross sections for the scan data.




Figure 6.22: The inclusive cross sections measured for the scan data region. The cyan, purple, and yellow points correspond to the luminosity-averaged inclusive cross sections measured for the  $\psi(3770)$  data.




Figure 6.23: The non- $D\bar{D}$  cross sections measured for the scan data region. The cyan, purple, and yellow points correspond to the luminosity-averaged non- $D\bar{D}$  cross sections measured for the  $\psi(3770)$  data.




Figure 6.24: The non- $D\bar{D}$  branching fractions measured for the scan data region. The cyan, purple, and yellow points correspond to the luminosity-averaged non- $D\bar{D}$  branching fractions measured for the  $\psi(3770)$  data.

## **Chapter 7**

# **Conclusion**

This is where the Conclusions go!

# References

- [1] M. Ablikim et al. [BES Collaboration], eConf C070805 (2007) 02 [Phys. Lett. B 660 (2008) 315].
- [2] J. Brodzicka et al. [Belle Collaboration], Phys. Rev. Lett. 100 (2008) 092001.
- [3] B. Aubert et al. [BABAR Collaboration], Phys. Rev. D 77 (2008) 011102.
- [4] B. Aubert et al., Phys. Rev. D 76 (2007) 111105(R).
- [5] V. V. Anashin *et al.*, Phys. Lett. B **711**, 292 (2012) [arXiv:1109.4205 [hep-ex]].
- [6] K.A. Olive et al. (Particle Data Group), Chin. Phys. C, **38**, 090001 (2014).
- [7] Wikimedia Commons. Standard model of elementary particles. 2006.
- [8] C. S. Wu, E. Ambler, R. W. Hayward, D. D. Hoppes, and R. P. Hudson Phys. Rev. **105**, 1413 (1957).
- [9] J. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay Phys. Rev. Lett. **13**, 138 (1964).
- [10] M. Ablikim et al. (BESIII Collaboration) Phys. Rev. Lett. **110**, 252001 (2013).
- [11] Z. Q. Liu et al. (Belle Collaboration) Phys. Rev. Lett. **110**, 252002 (2013).
- [12] R. Aaij et al. (LHCb Collaboration) Phys. Rev. Lett. **115**, 072001 (2015).
- [13] M. Kobayashi and T. Maskawa, “CP-Violation in the Renormalizable Theory of Weak Interaction”. Prog. Theor. Phys. 1973; 49 (2): 652-657. [doi: 10.1143/PTP.49.652]

- [14] S. L. Glashow, J. Iliopoulos, and L. Maiani Phys. Rev. D **2**, 1285 (1970).
- [15] B. Pontecorvo. (1957). “Inverse beta processes and nonconservation of lepton charge”. Zhurnal eksperimentalno i Teoreticheskogo Fiziki. **34**: 247. reproduced and translated in Soviet Physics JETP. 7: 172. 1958.
- [16] Z. Maki, M. Nakagawa, S. Sakata. (1962). “Remarks on the Unified Model of Elementary Particles”. Progress of Theoretical Physics. **28**: 870.
- [17] J. L. Rosner, Phys. Rev. D **64**, 094002 (2001) [hep-ph/0105327].
- [18] J. L. Rosner, Annals Phys. **319**, 1 (2005) [hep-ph/0411003].
- [19] S. Okubo, Phys. Lett. **5**, 165 (1963).
- [20] G. Zweig, Developments in the Quark Theory of Hadrons, Volume 1. Edited by D. Lichtenberg and S. Rosen. pp. 22-101
- [21] J. Iizuka, K. Okada and O. Shito, Prog. Theor. Phys. **35**, 1061 (1966).
- [22] E. A. Kuraev and V. S. Fadin, Sov. J. Nucl. Phys. **41**, 466 (1985) [Yad. Fiz. **41**, 733 (1985)].
- [23] J. M. Blatt and V. F. Weisskopf, Theoretical Nuclear Physics, Wiley, New York, (1952).
- [24] M. Ablikim *et al.* [BESIII Collaboration], Nucl. Instrum. Meth. A **614**, 345 (2010) [arXiv:0911.4960 [physics.ins-det]].
- [25] J. D. Jackson. *Classical Electrodynamics (3rd ed.)*. Wiley. (Section 13.2)
- [26] W. Li, *et al.*, Proc. Int. Conf. Comput. High Energy and Nucl. Phys. 225 (2006).
- [27] See <http://linux.web.cern.ch/linux/scientific5>.
- [28] G. Barrand, I. Belyaev, P. Binko, M. Cattaneo, R. Chytracek, G. Corti, M. Frank and G. Gracia *et al.*, Comput. Phys. Commun. **140**, 45 (2001).
- [29] C. Arnault, “CMT: A software configuration management tool,” (2000).

- [30] See <http://root.cern.ch>.
- [31] S. Agostinelli, *et al.*, Nucl. Instr. and Meth. **506**, (3), 250 (2003); J. Allison, *et al.*, IEEE Trans. Nucl. Sci. NS **53** (1), 270 (2006); See <http://www.geant4.org/geant4>.
- [32] S. Jadach , B. F. L. Ward and Z. Was, Comp. Phys. Commun. **130**, 260 (2000); S. Jadach, B. F. L. Ward and Z. Was, Phys. Rev. D **63**, 113009 (2001).
- [33] S. Jadach, Z. Was, R. Decker and J. H. Kuhn, Comput. Phys. Commun. **76**, 361 (1993).
- [34] See <http://home.thep.lu.se/~torbjorn/Pythia.html>.
- [35] D.J. Lange, Nucl. Instrum. Meth. A **462**, 152 (2001).
- [36] R. G. Ping, Chinese Physics C **32**, 8 (2008).
- [37] E. Barberio, B. van Eijk and Z. Was, Comput. Phys. Commun. **66**, 115 (1991).
- [38] J. C. Chen, G. S. Huang, X. R. Qi, D. H. Zhang and Y. S. Zhu, Phys. Rev. D **62**, 034003 (2000).
- [39] G. Bonneau and F. Martin, Nucl. Phys. B **27**, 381 (1971).
- [40] C.M. Carloni Calame, G. Montagna, O. Nicrosini, F. Piccinini, Nucl. Phys. Proc. Suppl. **131** 48-55 (2004).
- [41] See <http://www.mysql.com/about>.
- [42] R. M. Baltrusaitis *et al.* [MARK-III Collaboration], Phys. Rev. Lett. **56**, 2140 (1986).
- [43] J. Adler *et al.* [MARK-III Collaboration], Phys. Rev. Lett. **60**, 89 (1988).
- [44] D. Toth. (2014). “Measurement of Non- $D\bar{D}$  Decays of the  $\psi(3770)$  Resonance at BESIII”.
- [45] L. Dong, “Center-of-mass beam energy calibration for  $\psi(3770)$  data with BOSS 6.6.4,” BESIII-doc-284-v6 (2014).

- [46] A. Hafner, “Luminosity Measurement for the  $\psi(3770)$  data at BESIII,” BESIII-doc-406-v2 (2015).
- [47] D. M. Asner and W. M. Sun, Phys. Rev. D, **77**, 019901 (2008).
- [48] HFAG 2015 (*CPV*-Allowed) [[http://www.slac.stanford.edu/xorg/hfag/charm/CHARM15/results\\_minimal.pdf](http://www.slac.stanford.edu/xorg/hfag/charm/CHARM15/results_minimal.pdf)]
- [49] T. Evans et al., Physics Letters B 757 (2016) 520527.
- [50] M. Ablikim et al. [BESIII Collaboration], Chin. Phys. C, **37**, 12 (2013):12300.
- [51] G. Rong et. al, “Measurements of Branching Fractions and Form Factors for  $D^0 \rightarrow K^- e^+ \nu_e, \pi^- e^+ \nu_e$  Decays”, BESIII-doc-125-v3 (2015).
- [52] B. C. Ke et al. [Carnegie Mellon University], “ $\pi^0$  Reconstruction Efficiency”, BESIII-doc-446-v4 (2015).
- [53] T. Ma , “Determination of  $K_S^0$  Systematics”, BESIII-doc-125-v3 (2014).
- [54] M. Ablikim, [BESIII Collaboration], “Determination of the number of  $\psi'$  events at BESIII”, Chin. Phys. C, **37**, 6 (2013).

# Appendix A

## Glossary and Acronyms

Care has been taken in this thesis to minimize the use of jargon, but this cannot always be achieved. This appendix defines certain terms used in a glossary, and contains a table of acronyms and initialisms used along with their meanings.

### A.1 Glossary

### A.2 Acronyms / Initialisms

Table A.1: Acronyms and Initialisms

Name	Meaning
BESIII	The third Beijing Spectrometer
IHEP	Institute of High Energy Physics
BEPCII	The second Beijing Electron-Position Collider
MDC	Multi-Layer Drift Chamber
ToF	Time-of-Flight System
EMC	Electromagnetic Calorimeter
MUC	Muon Identifier
PMTs	Photomultipler Tubes

*Continued on next page*

*Table A.1 – Continued from previous page*

Acronym	Meaning
RPC	Resistive Plate Counter
ISR	Initial State Radiation
FSR	Final State Radiation
BOSS	BESIII Offline Software System
MC	Monte Carlo
ADC	Analog-to-Digital Conversion
TDC	Time-to-Digital Conversion

## Appendix B

# $D^0$ Signal Fits




Figure B.1: The 2D signal fit for  $D^0$  in Bin 0.



















Figure B.2: The 2D signal fit for  $D^0$  in Bin 1.




Figure B.3: The 2D signal fit for  $D^0$  in Bin 2.Figure B.4: The 2D signal fit for  $D^0$  in Bin 3.Figure B.5: The 2D signal fit for  $D^0$  in Bin 4.




Figure B.6: The 2D signal fit for  $D^0$  in Bin 5.Figure B.7: The 2D signal fit for  $D^0$  in Bin 6.Figure B.8: The 2D signal fit for  $D^0$  in Bin 7.




Figure B.9: The 2D signal fit for  $D^0$  in Bin 8.Figure B.10: The 2D signal fit for  $D^0$  in Bin 9.Figure B.11: The 2D signal fit for  $D^0$  in Bin 10.




Figure B.12: The 2D signal fit for  $D^0$  in Bin 11.Figure B.13: The 2D signal fit for  $D^0$  in Bin 12.Figure B.14: The 2D signal fit for  $D^0$  in Bin 13.




Figure B.15: The 2D signal fit for  $D^0$  in Bin 14.Figure B.16: The 2D signal fit for  $D^0$  in Bin 15.Figure B.17: The 2D signal fit for  $D^0$  in Bin 16.



Figure B.18: The 2D signal fit for  $D^0$  in Bin 17.Figure B.19: The 2D signal fit for  $D^0$  in Bin 18.Figure B.20: The 2D signal fit for  $D^0$  in Bin 19.

Figure B.21: The 2D signal fit for  $D^0$  in Bin 20.Figure B.22: The 2D signal fit for  $D^0$  in Bin 21.Figure B.23: The 2D signal fit for  $D^0$  in Bin 22.

Figure B.24: The 2D signal fit for  $D^0$  in Bin 23.Figure B.25: The 2D signal fit for  $D^0$  in Bin 24.Figure B.26: The 2D signal fit for  $D^0$  in Bin 25.

Figure B.27: The 2D signal fit for  $D^0$  in Bin 26.Figure B.28: The 2D signal fit for  $D^0$  in Bin 27.Figure B.29: The 2D signal fit for  $D^0$  in Bin 28.

Figure B.30: The 2D signal fit for  $D^0$  in Bin 29.Figure B.31: The 2D signal fit for  $D^0$  in Bin 30.Figure B.32: The 2D signal fit for  $D^0$  in Bin 31.

Figure B.33: The 2D signal fit for  $D^0$  in Bin 32.Figure B.34: The 2D signal fit for  $D^0$  in Bin 33.

## Appendix C

### $D^+$ Signal Fits




Figure C.1: The 2D signal fit for  $D^+$  in Bin 2.



















Figure C.2: The 2D signal fit for  $D^+$  in Bin 3.




Figure C.3: The 2D signal fit for  $D^+$  in Bin 4.Figure C.4: The 2D signal fit for  $D^+$  in Bin 5.Figure C.5: The 2D signal fit for  $D^+$  in Bin 6.




Figure C.6: The 2D signal fit for  $D^+$  in Bin 7.Figure C.7: The 2D signal fit for  $D^+$  in Bin 8.Figure C.8: The 2D signal fit for  $D^+$  in Bin 9.




Figure C.9: The 2D signal fit for  $D^+$  in Bin 10.Figure C.10: The 2D signal fit for  $D^+$  in Bin 11.Figure C.11: The 2D signal fit for  $D^+$  in Bin 12.




Figure C.12: The 2D signal fit for  $D^+$  in Bin 13.Figure C.13: The 2D signal fit for  $D^+$  in Bin 14.Figure C.14: The 2D signal fit for  $D^+$  in Bin 15.




Figure C.15: The 2D signal fit for  $D^+$  in Bin 16.Figure C.16: The 2D signal fit for  $D^+$  in Bin 17.Figure C.17: The 2D signal fit for  $D^+$  in Bin 18.

Figure C.18: The 2D signal fit for  $D^+$  in Bin 19.Figure C.19: The 2D signal fit for  $D^+$  in Bin 20.Figure C.20: The 2D signal fit for  $D^+$  in Bin 21.

Figure C.21: The 2D signal fit for  $D^+$  in Bin 22.Figure C.22: The 2D signal fit for  $D^+$  in Bin 23.Figure C.23: The 2D signal fit for  $D^+$  in Bin 24.

Figure C.24: The 2D signal fit for  $D^+$  in Bin 25.Figure C.25: The 2D signal fit for  $D^+$  in Bin 26.Figure C.26: The 2D signal fit for  $D^+$  in Bin 27.

Figure C.27: The 2D signal fit for  $D^+$  in Bin 28.Figure C.28: The 2D signal fit for  $D^+$  in Bin 29.Figure C.29: The 2D signal fit for  $D^+$  in Bin 30.

Figure C.30: The 2D signal fit for  $D^+$  in Bin 31.Figure C.31: The 2D signal fit for  $D^+$  in Bin 32.Figure C.32: The 2D signal fit for  $D^+$  in Bin 33.

## Appendix D

# Scan Data Hadronic Counting Fits




Figure D.1: The number of hadrons found in the 3734 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.2: The number of hadrons found in the 3736 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.3: The number of hadrons found in the 3744 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.4: The number of hadrons found in the 3748 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.5: The number of hadrons found in the 3750 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.6: The number of hadrons found in the 3751 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.7: The number of hadrons found in the 3753 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.8: The number of hadrons found in the 3755 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.9: The number of hadrons found in the 3756 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.10: The number of hadrons found in the 3759 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.11: The number of hadrons found in the 3762 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.12: The number of hadrons found in the 3765 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.13: The number of hadrons found in the 3767 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.14: The number of hadrons found in the 3771 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.15: The number of hadrons found in the 3774 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.16: The number of hadrons found in the 3777 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.17: The number of hadrons found in the 3780 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.18: The number of hadrons found in the 3782 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.19: The number of hadrons found in the 3786 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.20: The number of hadrons found in the 3789 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.21: The number of hadrons found in the 3792 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.22: The number of hadrons found in the 3797 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.23: The number of hadrons found in the 3800 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.24: The number of hadrons found in the 3802 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.25: The number of hadrons found in the 3807 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.26: The number of hadrons found in the 3809 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.27: The number of hadrons found in the 3813 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.28: The number of hadrons found in the 3815 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.29: The number of hadrons found in the 3822 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.30: The number of hadrons found in the 3832 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.31: The number of hadrons found in the 3839 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.32: The number of hadrons found in the 3849 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.33: The number of hadrons found in the 3855 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).




Figure D.34: The number of hadrons found in the 3863 (Scan) data sample. This includes results for SHAD (left), LHAD (middle), and THAD (right).

## Appendix E

# Scan Data Reconstruction Efficiency

3734 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.164	$0.7360 \pm 0.0019$	$0.7956 \pm 0.0020$	$0.6015 \pm 0.0017$
$D^+D^-$	0.000	-	-	-
$\tau^+\tau^-$	2.467	$0.1284 \pm 0.0008$	$0.2834 \pm 0.0012$	$0.0987 \pm 0.0007$
$\gamma J/\psi$	1.060	$0.4662 \pm 0.0015$	$0.5625 \pm 0.0017$	$0.3475 \pm 0.0013$
$\gamma\psi(2S)$	5.359	$0.6305 \pm 0.0018$	$0.6967 \pm 0.0019$	$0.5142 \pm 0.0016$

Table E.1: Reconstruction of background samples for the 3734 (Scan) data.

3736 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.218	$0.7348 \pm 0.0027$	$0.7951 \pm 0.0028$	$0.5995 \pm 0.0024$
$D^+D^-$	0.000	-	-	-
$\tau^+\tau^-$	2.475	$0.1281 \pm 0.0011$	$0.2815 \pm 0.0017$	$0.0984 \pm 0.0010$
$\gamma J/\psi$	1.055	$0.4640 \pm 0.0022$	$0.5597 \pm 0.0024$	$0.3429 \pm 0.0019$
$\gamma\psi(2S)$	5.094	$0.6334 \pm 0.0025$	$0.6995 \pm 0.0026$	$0.5148 \pm 0.0023$

Table E.2: Reconstruction of background samples for the 3736 (Scan) data.

3744 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.765	$0.7304 \pm 0.0019$	$0.7932 \pm 0.0020$	$0.5959 \pm 0.0017$
$D^+D^-$	0.150	$0.6071 \pm 0.0017$	$0.6838 \pm 0.0018$	$0.4871 \pm 0.0016$
$\tau^+\tau^-$	2.514	$0.1270 \pm 0.0008$	$0.2812 \pm 0.0012$	$0.0983 \pm 0.0007$
$\gamma J/\psi$	1.039	$0.4626 \pm 0.0015$	$0.5613 \pm 0.0017$	$0.3431 \pm 0.0013$
$\gamma\psi(2S)$	4.428	$0.6259 \pm 0.0018$	$0.6938 \pm 0.0019$	$0.5101 \pm 0.0016$

Table E.3: Reconstruction of background samples for the 3744 (Scan) data.

3748 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.838	$0.7276 \pm 0.0016$	$0.7906 \pm 0.0016$	$0.5907 \pm 0.0014$
$D^+D^-$	0.360	$0.6037 \pm 0.0014$	$0.6800 \pm 0.0015$	$0.4820 \pm 0.0013$
$\tau^+\tau^-$	2.522	$0.1262 \pm 0.0006$	$0.2781 \pm 0.0010$	$0.0970 \pm 0.0006$
$\gamma J/\psi$	1.032	$0.4609 \pm 0.0012$	$0.5588 \pm 0.0014$	$0.3427 \pm 0.0011$
$\gamma\psi(2S)$	4.180	$0.6252 \pm 0.0014$	$0.6928 \pm 0.0015$	$0.5080 \pm 0.0013$

Table E.4: Reconstruction of background samples for the 3748 (Scan) data.

3750 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.968	$0.7351 \pm 0.0012$	$0.7960 \pm 0.0013$	$0.5996 \pm 0.0011$
$D^+D^-$	0.439	$0.6130 \pm 0.0011$	$0.6879 \pm 0.0012$	$0.4921 \pm 0.0010$
$\tau^+\tau^-$	2.533	$0.1283 \pm 0.0005$	$0.2812 \pm 0.0007$	$0.0993 \pm 0.0004$
$\gamma J/\psi$	1.029	$0.4661 \pm 0.0010$	$0.5633 \pm 0.0011$	$0.3470 \pm 0.0008$
$\gamma\psi(2S)$	4.065	$0.6322 \pm 0.0011$	$0.6986 \pm 0.0012$	$0.5162 \pm 0.0010$

Table E.5: Reconstruction of background samples for the 3750 (Scan) data.

3751 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	1.237	$0.7316 \pm 0.0009$	$0.7935 \pm 0.0009$	$0.5950 \pm 0.0008$
$D^+D^-$	0.574	$0.6084 \pm 0.0008$	$0.6841 \pm 0.0009$	$0.4872 \pm 0.0007$
$\tau^+\tau^-$	2.542	$0.1274 \pm 0.0004$	$0.2805 \pm 0.0006$	$0.0983 \pm 0.0003$
$\gamma J/\psi$	1.025	$0.4629 \pm 0.0007$	$0.5606 \pm 0.0008$	$0.3439 \pm 0.0006$
$\gamma\psi(2S)$	3.969	$0.6279 \pm 0.0008$	$0.6959 \pm 0.0009$	$0.5098 \pm 0.0008$

Table E.6: Reconstruction of background samples for the 3751 (Scan) data.

3753 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	1.377	$0.7342 \pm 0.0010$	$0.7955 \pm 0.0010$	$0.5973 \pm 0.0009$
$D^+D^-$	0.748	$0.6103 \pm 0.0009$	$0.6855 \pm 0.0009$	$0.4888 \pm 0.0008$
$\tau^+\tau^-$	2.553	$0.1267 \pm 0.0004$	$0.2798 \pm 0.0006$	$0.0976 \pm 0.0003$
$\gamma J/\psi$	1.022	$0.4636 \pm 0.0008$	$0.5606 \pm 0.0008$	$0.3443 \pm 0.0007$
$\gamma\psi(2S)$	3.871	$0.6290 \pm 0.0009$	$0.6964 \pm 0.0009$	$0.5112 \pm 0.0008$

Table E.7: Reconstruction of background samples for the 3753 (Scan) data.

3755 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	1.566	$0.7322 \pm 0.0009$	$0.7931 \pm 0.0009$	$0.5961 \pm 0.0008$
$D^+D^-$	0.797	$0.6080 \pm 0.0008$	$0.6832 \pm 0.0009$	$0.4883 \pm 0.0007$
$\tau^+\tau^-$	2.562	$0.1276 \pm 0.0004$	$0.2808 \pm 0.0006$	$0.0986 \pm 0.0003$
$\gamma J/\psi$	1.018	$0.4632 \pm 0.0007$	$0.5603 \pm 0.0008$	$0.3450 \pm 0.0006$
$\gamma\psi(2S)$	3.752	$0.6272 \pm 0.0008$	$0.6946 \pm 0.0009$	$0.5104 \pm 0.0008$

Table E.8: Reconstruction of background samples for the 3755 (Scan) data.

3756 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	1.629	$0.7410 \pm 0.0010$	$0.7995 \pm 0.0011$	$0.6067 \pm 0.0009$
$D^+D^-$	0.914	$0.6150 \pm 0.0009$	$0.6889 \pm 0.0010$	$0.4960 \pm 0.0008$
$\tau^+\tau^-$	2.573	$0.1293 \pm 0.0004$	$0.2821 \pm 0.0006$	$0.1002 \pm 0.0004$
$\gamma J/\psi$	1.017	$0.4686 \pm 0.0008$	$0.5645 \pm 0.0009$	$0.3507 \pm 0.0007$
$\gamma\psi(2S)$	3.721	$0.6347 \pm 0.0010$	$0.7011 \pm 0.0010$	$0.5184 \pm 0.0009$

Table E.9: Reconstruction of background samples for the 3756 (Scan) data.

3759 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	1.948	$0.7415 \pm 0.0009$	$0.8000 \pm 0.0009$	$0.6076 \pm 0.0008$
$D^+D^-$	1.192	$0.6162 \pm 0.0008$	$0.6894 \pm 0.0008$	$0.4967 \pm 0.0007$
$\tau^+\tau^-$	2.584	$0.1286 \pm 0.0004$	$0.2826 \pm 0.0005$	$0.0999 \pm 0.0003$
$\gamma J/\psi$	1.011	$0.4688 \pm 0.0007$	$0.5652 \pm 0.0008$	$0.3507 \pm 0.0006$
$\gamma\psi(2S)$	3.572	$0.6351 \pm 0.0008$	$0.7010 \pm 0.0008$	$0.5194 \pm 0.0007$

Table E.10: Reconstruction of background samples for the 3759 (Scan) data.

3762 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	2.385	$0.7420 \pm 0.0008$	$0.8004 \pm 0.0009$	$0.6078 \pm 0.0007$
$D^+D^-$	1.534	$0.6154 \pm 0.0007$	$0.6891 \pm 0.0008$	$0.4964 \pm 0.0007$
$\tau^+\tau^-$	2.598	$0.1285 \pm 0.0003$	$0.2819 \pm 0.0005$	$0.0998 \pm 0.0003$
$\gamma J/\psi$	1.005	$0.4680 \pm 0.0007$	$0.5642 \pm 0.0007$	$0.3496 \pm 0.0006$
$\gamma\psi(2S)$	3.426	$0.6359 \pm 0.0008$	$0.7017 \pm 0.0008$	$0.5199 \pm 0.0007$

Table E.11: Reconstruction of background samples for the 3762 (Scan) data.

3765 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	2.715	$0.7425 \pm 0.0010$	$0.8006 \pm 0.0011$	$0.6085 \pm 0.0009$
$D^+D^-$	1.882	$0.6168 \pm 0.0009$	$0.6905 \pm 0.0010$	$0.4978 \pm 0.0008$
$\tau^+\tau^-$	2.613	$0.1281 \pm 0.0004$	$0.2819 \pm 0.0006$	$0.0997 \pm 0.0004$
$\gamma J/\psi$	1.000	$0.4693 \pm 0.0008$	$0.5649 \pm 0.0009$	$0.3519 \pm 0.0007$
$\gamma\psi(2S)$	3.315	$0.6376 \pm 0.0010$	$0.7035 \pm 0.0010$	$0.5219 \pm 0.0009$

Table E.12: Reconstruction of background samples for the 3765 (Scan) data.

3767 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	3.102	$0.7432 \pm 0.0011$	$0.8008 \pm 0.0012$	$0.6090 \pm 0.0010$
$D^+D^-$	2.200	$0.6171 \pm 0.0010$	$0.6908 \pm 0.0011$	$0.4978 \pm 0.0009$
$\tau^+\tau^-$	2.627	$0.1286 \pm 0.0005$	$0.2818 \pm 0.0007$	$0.0999 \pm 0.0004$
$\gamma J/\psi$	0.995	$0.4693 \pm 0.0009$	$0.5653 \pm 0.0010$	$0.3511 \pm 0.0008$
$\gamma\psi(2S)$	3.212	$0.6369 \pm 0.0010$	$0.7025 \pm 0.0011$	$0.5214 \pm 0.0009$

Table E.13: Reconstruction of background samples for the 3767 (Scan) data.

3771 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	3.406	$0.7440 \pm 0.0014$	$0.8021 \pm 0.0014$	$0.6098 \pm 0.0012$
$D^+D^-$	2.634	$0.6179 \pm 0.0012$	$0.6913 \pm 0.0013$	$0.4987 \pm 0.0011$
$\tau^+\tau^-$	2.641	$0.1284 \pm 0.0006$	$0.2821 \pm 0.0008$	$0.1003 \pm 0.0005$
$\gamma J/\psi$	0.989	$0.4700 \pm 0.0011$	$0.5661 \pm 0.0012$	$0.3528 \pm 0.0009$
$\gamma\psi(2S)$	3.075	$0.6356 \pm 0.0013$	$0.7020 \pm 0.0013$	$0.5206 \pm 0.0011$

Table E.14: Reconstruction of background samples for the 3771 (Scan) data.

3774 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	3.714	$0.7429 \pm 0.0014$	$0.8009 \pm 0.0014$	$0.6084 \pm 0.0012$
$D^+D^-$	3.067	$0.6172 \pm 0.0012$	$0.6903 \pm 0.0013$	$0.4975 \pm 0.0011$
$\tau^+\tau^-$	2.656	$0.1286 \pm 0.0006$	$0.2819 \pm 0.0008$	$0.0996 \pm 0.0005$
$\gamma J/\psi$	0.983	$0.4695 \pm 0.0011$	$0.5657 \pm 0.0012$	$0.3509 \pm 0.0009$
$\gamma\psi(2S)$	2.975	$0.6365 \pm 0.0013$	$0.7028 \pm 0.0013$	$0.5194 \pm 0.0011$

Table E.15: Reconstruction of background samples for the 3774 (Scan) data.

3777 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	3.692	$0.7436 \pm 0.0016$	$0.8016 \pm 0.0016$	$0.6090 \pm 0.0014$
$D^+D^-$	3.078	$0.6171 \pm 0.0014$	$0.6904 \pm 0.0015$	$0.4978 \pm 0.0013$
$\tau^+\tau^-$	2.669	$0.1299 \pm 0.0007$	$0.2837 \pm 0.0010$	$0.1012 \pm 0.0006$
$\gamma J/\psi$	0.978	$0.4702 \pm 0.0013$	$0.5662 \pm 0.0014$	$0.3537 \pm 0.0011$
$\gamma\psi(2S)$	2.870	$0.6362 \pm 0.0015$	$0.7024 \pm 0.0015$	$0.5204 \pm 0.0013$

Table E.16: Reconstruction of background samples for the 3777 (Scan) data.

3780 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	3.444	$0.7460 \pm 0.0014$	$0.8036 \pm 0.0014$	$0.6126 \pm 0.0012$
$D^+D^-$	2.599	$0.6194 \pm 0.0012$	$0.6925 \pm 0.0013$	$0.5003 \pm 0.0011$
$\tau^+\tau^-$	2.683	$0.1285 \pm 0.0006$	$0.2824 \pm 0.0008$	$0.1000 \pm 0.0005$
$\gamma J/\psi$	0.973	$0.4703 \pm 0.0011$	$0.5665 \pm 0.0012$	$0.3530 \pm 0.0009$
$\gamma\psi(2S)$	2.788	$0.6380 \pm 0.0013$	$0.7043 \pm 0.0013$	$0.5228 \pm 0.0011$

Table E.17: Reconstruction of background samples for the 3780 (Scan) data.

3782 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	2.791	$0.7441 \pm 0.0012$	$0.8019 \pm 0.0013$	$0.6101 \pm 0.0011$
$D^+D^-$	2.206	$0.6195 \pm 0.0011$	$0.6923 \pm 0.0012$	$0.5004 \pm 0.0010$
$\tau^+\tau^-$	2.696	$0.1281 \pm 0.0005$	$0.2814 \pm 0.0008$	$0.0995 \pm 0.0004$
$\gamma J/\psi$	0.968	$0.4706 \pm 0.0010$	$0.5663 \pm 0.0011$	$0.3531 \pm 0.0008$
$\gamma\psi(2S)$	2.712	$0.6377 \pm 0.0011$	$0.7046 \pm 0.0012$	$0.5210 \pm 0.0010$

Table E.18: Reconstruction of background samples for the 3782 (Scan) data.

3786 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	1.995	$0.7460 \pm 0.0012$	$0.8037 \pm 0.0013$	$0.6122 \pm 0.0011$
$D^+D^-$	1.627	$0.6194 \pm 0.0011$	$0.6923 \pm 0.0012$	$0.5013 \pm 0.0010$
$\tau^+\tau^-$	2.709	$0.1289 \pm 0.0005$	$0.2810 \pm 0.0007$	$0.1007 \pm 0.0004$
$\gamma J/\psi$	0.961	$0.4698 \pm 0.0010$	$0.5652 \pm 0.0011$	$0.3524 \pm 0.0008$
$\gamma\psi(2S)$	2.605	$0.6380 \pm 0.0011$	$0.7044 \pm 0.0012$	$0.5223 \pm 0.0010$

Table E.19: Reconstruction of background samples for the 3786 (Scan) data.

3789 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	1.361	$0.7457 \pm 0.0010$	$0.8035 \pm 0.0011$	$0.6118 \pm 0.0009$
$D^+D^-$	1.178	$0.6188 \pm 0.0009$	$0.6918 \pm 0.0010$	$0.5002 \pm 0.0008$
$\tau^+\tau^-$	2.722	$0.1284 \pm 0.0004$	$0.2805 \pm 0.0006$	$0.1003 \pm 0.0004$
$\gamma J/\psi$	0.957	$0.4700 \pm 0.0008$	$0.5655 \pm 0.0009$	$0.3526 \pm 0.0007$
$\gamma\psi(2S)$	2.550	$0.6379 \pm 0.0010$	$0.7040 \pm 0.0010$	$0.5225 \pm 0.0009$

Table E.20: Reconstruction of background samples for the 3789 (Scan) data.

3792 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.920	$0.7440 \pm 0.0010$	$0.8019 \pm 0.0011$	$0.6094 \pm 0.0009$
$D^+D^-$	0.680	$0.6176 \pm 0.0009$	$0.6905 \pm 0.0010$	$0.4989 \pm 0.0008$
$\tau^+\tau^-$	2.739	$0.1286 \pm 0.0004$	$0.2800 \pm 0.0006$	$0.0998 \pm 0.0004$
$\gamma J/\psi$	0.951	$0.4677 \pm 0.0008$	$0.5637 \pm 0.0009$	$0.3498 \pm 0.0007$
$\gamma\psi(2S)$	2.467	$0.6364 \pm 0.0010$	$0.7027 \pm 0.0010$	$0.5196 \pm 0.0009$

Table E.21: Reconstruction of background samples for the 3792 (Scan) data.

3797 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.562	$0.7444 \pm 0.0010$	$0.8025 \pm 0.0010$	$0.6102 \pm 0.0009$
$D^+D^-$	0.414	$0.6183 \pm 0.0009$	$0.6916 \pm 0.0009$	$0.4998 \pm 0.0008$
$\tau^+\tau^-$	2.755	$0.1280 \pm 0.0004$	$0.2796 \pm 0.0006$	$0.0998 \pm 0.0004$
$\gamma J/\psi$	0.944	$0.4688 \pm 0.0008$	$0.5646 \pm 0.0008$	$0.3512 \pm 0.0007$
$\gamma\psi(2S)$	2.370	$0.6380 \pm 0.0009$	$0.7042 \pm 0.0009$	$0.5218 \pm 0.0008$

Table E.22: Reconstruction of background samples for the 3797 (Scan) data.

3800 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.424	$0.7417 \pm 0.0009$	$0.8004 \pm 0.0009$	$0.6068 \pm 0.0008$
$D^+D^-$	0.241	$0.6161 \pm 0.0008$	$0.6894 \pm 0.0009$	$0.4965 \pm 0.0007$
$\tau^+\tau^-$	2.771	$0.1281 \pm 0.0004$	$0.2803 \pm 0.0006$	$0.0994 \pm 0.0003$
$\gamma J/\psi$	0.938	$0.4671 \pm 0.0007$	$0.5633 \pm 0.0008$	$0.3492 \pm 0.0006$
$\gamma\psi(2S)$	2.301	$0.6358 \pm 0.0008$	$0.7029 \pm 0.0009$	$0.5183 \pm 0.0008$

Table E.23: Reconstruction of background samples for the 3800 (Scan) data.

3802 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.325	$0.7421 \pm 0.0011$	$0.8008 \pm 0.0012$	$0.6075 \pm 0.0010$
$D^+D^-$	0.236	$0.6157 \pm 0.0010$	$0.6892 \pm 0.0011$	$0.4959 \pm 0.0009$
$\tau^+\tau^-$	2.784	$0.1285 \pm 0.0005$	$0.2806 \pm 0.0007$	$0.0995 \pm 0.0004$
$\gamma J/\psi$	0.935	$0.4676 \pm 0.0009$	$0.5635 \pm 0.0010$	$0.3496 \pm 0.0008$
$\gamma\psi(2S)$	2.260	$0.6362 \pm 0.0010$	$0.7034 \pm 0.0011$	$0.5195 \pm 0.0009$

Table E.24: Reconstruction of background samples for the 3802 (Scan) data.

3807 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.352	$0.7407 \pm 0.0012$	$0.8000 \pm 0.0013$	$0.6049 \pm 0.0011$
$D^+D^-$	0.289	$0.6132 \pm 0.0011$	$0.6876 \pm 0.0012$	$0.4941 \pm 0.0010$
$\tau^+\tau^-$	2.795	$0.1286 \pm 0.0005$	$0.2800 \pm 0.0007$	$0.0995 \pm 0.0004$
$\gamma J/\psi$	0.927	$0.4666 \pm 0.0010$	$0.5628 \pm 0.0011$	$0.3480 \pm 0.0008$
$\gamma\psi(2S)$	2.174	$0.6344 \pm 0.0011$	$0.7024 \pm 0.0012$	$0.5175 \pm 0.0010$

Table E.25: Reconstruction of background samples for the 3807 (Scan) data.

3809 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.233	$0.7430 \pm 0.0016$	$0.8019 \pm 0.0016$	$0.6092 \pm 0.0014$
$D^+D^-$	0.132	$0.6170 \pm 0.0014$	$0.6907 \pm 0.0015$	$0.4977 \pm 0.0013$
$\tau^+\tau^-$	2.807	$0.1281 \pm 0.0007$	$0.2803 \pm 0.0010$	$0.0998 \pm 0.0006$
$\gamma J/\psi$	0.923	$0.4676 \pm 0.0012$	$0.5632 \pm 0.0014$	$0.3511 \pm 0.0011$
$\gamma\psi(2S)$	2.133	$0.6375 \pm 0.0015$	$0.7046 \pm 0.0015$	$0.5216 \pm 0.0013$

Table E.26: Reconstruction of background samples for the 3809 (Scan) data.

3813 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.060	$0.7393 \pm 0.0019$	$0.7994 \pm 0.0020$	$0.6045 \pm 0.0017$
$D^+D^-$	0.153	$0.6128 \pm 0.0018$	$0.6879 \pm 0.0019$	$0.4923 \pm 0.0016$
$\tau^+\tau^-$	2.819	$0.1272 \pm 0.0008$	$0.2783 \pm 0.0012$	$0.0980 \pm 0.0007$
$\gamma J/\psi$	0.917	$0.4679 \pm 0.0015$	$0.5634 \pm 0.0017$	$0.3488 \pm 0.0013$
$\gamma\psi(2S)$	2.063	$0.6341 \pm 0.0018$	$0.7020 \pm 0.0019$	$0.5160 \pm 0.0016$

Table E.27: Reconstruction of background samples for the 3813 (Scan) data.

3815 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.056	$0.7424 \pm 0.0019$	$0.8022 \pm 0.0020$	$0.6068 \pm 0.0017$
$D^+D^-$	0.089	$0.6157 \pm 0.0018$	$0.6887 \pm 0.0019$	$0.4969 \pm 0.0016$
$\tau^+\tau^-$	2.829	$0.1287 \pm 0.0008$	$0.2809 \pm 0.0012$	$0.0999 \pm 0.0007$
$\gamma J/\psi$	0.914	$0.4675 \pm 0.0015$	$0.5635 \pm 0.0017$	$0.3494 \pm 0.0013$
$\gamma\psi(2S)$	2.034	$0.6381 \pm 0.0018$	$0.7048 \pm 0.0019$	$0.5202 \pm 0.0016$

Table E.28: Reconstruction of background samples for the 3815 (Scan) data.

3822 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.140	$0.7442 \pm 0.0019$	$0.8038 \pm 0.0020$	$0.6097 \pm 0.0017$
$D^+D^-$	0.197	$0.6176 \pm 0.0018$	$0.6919 \pm 0.0019$	$0.4992 \pm 0.0016$
$\tau^+\tau^-$	2.859	$0.1295 \pm 0.0008$	$0.2813 \pm 0.0012$	$0.1003 \pm 0.0007$
$\gamma J/\psi$	0.902	$0.4676 \pm 0.0015$	$0.5651 \pm 0.0017$	$0.3497 \pm 0.0013$
$\gamma\psi(2S)$	1.920	$0.6401 \pm 0.0018$	$0.7064 \pm 0.0019$	$0.5234 \pm 0.0016$

Table E.29: Reconstruction of background samples for the 3822 (Scan) data.

3832 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.069	$0.7469 \pm 0.0027$	$0.8037 \pm 0.0028$	$0.6122 \pm 0.0025$
$D^+D^-$	0.046	$0.6209 \pm 0.0025$	$0.6924 \pm 0.0026$	$0.5009 \pm 0.0022$
$\tau^+\tau^-$	2.887	$0.1290 \pm 0.0011$	$0.2796 \pm 0.0017$	$0.1003 \pm 0.0010$
$\gamma J/\psi$	0.888	$0.4678 \pm 0.0022$	$0.5634 \pm 0.0024$	$0.3500 \pm 0.0019$
$\gamma\psi(2S)$	1.799	$0.6366 \pm 0.0025$	$0.7046 \pm 0.0027$	$0.5206 \pm 0.0023$

Table E.30: Reconstruction of background samples for the 3832 (Scan) data.

3839 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.237	$0.7435 \pm 0.0027$	$0.8017 \pm 0.0028$	$0.6102 \pm 0.0025$
$D^+D^-$	0.254	$0.6168 \pm 0.0025$	$0.6902 \pm 0.0026$	$0.4991 \pm 0.0022$
$\tau^+\tau^-$	2.914	$0.1270 \pm 0.0011$	$0.2759 \pm 0.0017$	$0.0983 \pm 0.0010$
$\gamma J/\psi$	0.877	$0.4676 \pm 0.0022$	$0.5631 \pm 0.0024$	$0.3502 \pm 0.0019$
$\gamma\psi(2S)$	1.716	$0.6374 \pm 0.0025$	$0.7059 \pm 0.0027$	$0.5231 \pm 0.0023$

Table E.31: Reconstruction of background samples for the 3839 (Scan) data.

3849 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.186	$0.7440 \pm 0.0027$	$0.8028 \pm 0.0028$	$0.6087 \pm 0.0025$
$D^+D^-$	0.186	$0.6195 \pm 0.0025$	$0.6921 \pm 0.0026$	$0.5007 \pm 0.0022$
$\tau^+\tau^-$	2.939	$0.1289 \pm 0.0011$	$0.2773 \pm 0.0017$	$0.1001 \pm 0.0010$
$\gamma J/\psi$	0.862	$0.4679 \pm 0.0022$	$0.5612 \pm 0.0024$	$0.3496 \pm 0.0019$
$\gamma\psi(2S)$	1.604	$0.6357 \pm 0.0025$	$0.7041 \pm 0.0027$	$0.5206 \pm 0.0023$

Table E.32: Reconstruction of background samples for the 3849 (Scan) data.

3855 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.337	$0.7467 \pm 0.0019$	$0.8046 \pm 0.0020$	$0.6116 \pm 0.0017$
$D^+D^-$	0.273	$0.6188 \pm 0.0018$	$0.6917 \pm 0.0019$	$0.5006 \pm 0.0016$
$\tau^+\tau^-$	2.967	$0.1290 \pm 0.0008$	$0.2785 \pm 0.0012$	$0.1000 \pm 0.0007$
$\gamma J/\psi$	0.853	$0.4676 \pm 0.0015$	$0.5621 \pm 0.0017$	$0.3498 \pm 0.0013$
$\gamma\psi(2S)$	1.545	$0.6374 \pm 0.0018$	$0.7049 \pm 0.0019$	$0.5207 \pm 0.0016$

Table E.33: Reconstruction of background samples for the 3855 (Scan) data.

3863 (Scan) Reconstruction				
Sample	$\sigma$ [nb]	$\epsilon_{\text{MC}}$ (SHAD) [%]	$\epsilon_{\text{MC}}$ (LHAD) [%]	$\epsilon_{\text{MC}}$ (THAD) [%]
$D^0\bar{D}^0$	0.340	$0.7476 \pm 0.0019$	$0.8040 \pm 0.0020$	$0.6123 \pm 0.0017$
$D^+D^-$	0.099	$0.6191 \pm 0.0018$	$0.6916 \pm 0.0019$	$0.5006 \pm 0.0016$
$\tau^+\tau^-$	2.988	$0.1280 \pm 0.0008$	$0.2763 \pm 0.0012$	$0.1001 \pm 0.0007$
$\gamma J/\psi$	0.843	$0.4667 \pm 0.0015$	$0.5620 \pm 0.0017$	$0.3499 \pm 0.0013$
$\gamma\psi(2S)$	1.476	$0.6370 \pm 0.0018$	$0.7054 \pm 0.0019$	$0.5210 \pm 0.0016$

Table E.34: Reconstruction of background samples for the 3863 (Scan) data.

,

## Appendix F

# Scan Data Signal Counts

3734 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$12047 \pm 109$	$13458 \pm 116$	$9605 \pm 98$
$q\bar{q}$	$7942 \pm 62$	$8743 \pm 66$	$6314 \pm 52$
$D^0\overline{D^0}$	$100 \pm 1$	$108 \pm 1$	$81 \pm 1$
$D^+D^-$	$0 \pm 0$	$0 \pm 0$	$0 \pm 0$
$\tau^+\tau^-$	$262 \pm 1$	$579 \pm 3$	$201 \pm 1$
$\gamma J/\psi$	$409 \pm 2$	$494 \pm 2$	$305 \pm 1$
$\gamma\psi(2S)$	$2802 \pm 12$	$3096 \pm 13$	$2285 \pm 10$
Hadrons	$529 \pm 126$	$435 \pm 134$	$416 \pm 111$

Table F.1: Hadronic events selected in the 3734 (Scan) data.

3736 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$4786 \pm 69$	$5413 \pm 73$	$3767 \pm 61$
$q\bar{q}$	$3144 \pm 28$	$3460 \pm 30$	$2500 \pm 23$
$D^0\overline{D^0}$	$52 \pm 1$	$56 \pm 1$	$42 \pm 1$
$D^+D^-$	$0 \pm 0$	$0 \pm 0$	$0 \pm 0$
$\tau^+\tau^-$	$104 \pm 1$	$229 \pm 1$	$80 \pm 1$
$\gamma J/\psi$	$160 \pm 1$	$194 \pm 1$	$118 \pm 1$
$\gamma\psi(2S)$	$1060 \pm 7$	$1171 \pm 8$	$861 \pm 6$
Hadrons	$263 \pm 75$	$301 \pm 80$	$162 \pm 66$

Table F.2: Hadronic events selected in the 3736 (Scan) data.

3744 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$13426 \pm 115$	$15255 \pm 123$	$10823 \pm 104$
$q\bar{q}$	$9082 \pm 72$	$9984 \pm 78$	$7226 \pm 61$
$D^0\bar{D}^0$	$532 \pm 2$	$577 \pm 2$	$434 \pm 1$
$D^+D^-$	$86 \pm 1$	$97 \pm 1$	$69 \pm 1$
$\tau^+\tau^-$	$304 \pm 2$	$673 \pm 3$	$235 \pm 1$
$\gamma J/\psi$	$457 \pm 2$	$555 \pm 2$	$339 \pm 1$
$\gamma\psi(2S)$	$2639 \pm 11$	$2925 \pm 12$	$2151 \pm 9$
Hadrons	$322 \pm 137$	$440 \pm 146$	$366 \pm 121$

Table F.3: Hadronic events selected in the 3744 (Scan) data.

3748 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$20024 \pm 141$	$22788 \pm 150$	$15815 \pm 125$
$q\bar{q}$	$13384 \pm 105$	$14706 \pm 113$	$10651 \pm 90$
$D^0\bar{D}^0$	$857 \pm 3$	$931 \pm 3$	$695 \pm 2$
$D^+D^-$	$305 \pm 1$	$344 \pm 1$	$243 \pm 1$
$\tau^+\tau^-$	$447 \pm 2$	$985 \pm 4$	$343 \pm 2$
$\gamma J/\psi$	$668 \pm 2$	$810 \pm 2$	$497 \pm 2$
$\gamma\psi(2S)$	$3672 \pm 13$	$4069 \pm 14$	$2984 \pm 11$
Hadrons	$688 \pm 177$	$940 \pm 189$	$398 \pm 155$

Table F.4: Hadronic events selected in the 3748 (Scan) data.

3750 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$32171 \pm 179$	$36551 \pm 191$	$25686 \pm 160$
$q\bar{q}$	$21617 \pm 168$	$23746 \pm 180$	$17205 \pm 144$
$D^0\bar{D}^0$	$1616 \pm 4$	$1750 \pm 4$	$1318 \pm 3$
$D^+D^-$	$611 \pm 1$	$686 \pm 1$	$490 \pm 1$
$\tau^+\tau^-$	$738 \pm 3$	$1618 \pm 5$	$571 \pm 2$
$\gamma J/\psi$	$1089 \pm 3$	$1316 \pm 3$	$810 \pm 2$
$\gamma\psi(2S)$	$5838 \pm 16$	$6451 \pm 17$	$4767 \pm 14$
Hadrons	$659 \pm 246$	$982 \pm 263$	$522 \pm 216$

Table F.5: Hadronic events selected in the 3750 (Scan) data.

3751 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$42694 \pm 206$	$48594 \pm 220$	$34040 \pm 184$
$q\bar{q}$	$28246 \pm 219$	$31021 \pm 234$	$22483 \pm 188$
$D^0\bar{D}^0$	$2687 \pm 6$	$2915 \pm 6$	$2186 \pm 5$
$D^+D^-$	$1037 \pm 2$	$1166 \pm 2$	$830 \pm 2$
$\tau^+\tau^-$	$961 \pm 3$	$2117 \pm 5$	$742 \pm 2$
$\gamma J/\psi$	$1409 \pm 3$	$1707 \pm 4$	$1047 \pm 2$
$\gamma\psi(2S)$	$7401 \pm 17$	$8204 \pm 18$	$6010 \pm 14$
Hadrons	$948 \pm 301$	$1461 \pm 322$	$739 \pm 264$

Table F.6: Hadronic events selected in the 3751 (Scan) data.

3753 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$48035 \pm 219$	$54668 \pm 233$	$38460 \pm 196$
$q\bar{q}$	$31437 \pm 244$	$34517 \pm 262$	$25027 \pm 210$
$D^0\bar{D}^0$	$3344 \pm 7$	$3623 \pm 7$	$2720 \pm 6$
$D^+D^-$	$1510 \pm 3$	$1696 \pm 3$	$1209 \pm 2$
$\tau^+\tau^-$	$1069 \pm 3$	$2363 \pm 6$	$824 \pm 3$
$\gamma J/\psi$	$1567 \pm 3$	$1895 \pm 4$	$1164 \pm 3$
$\gamma\psi(2S)$	$8055 \pm 18$	$8918 \pm 20$	$6547 \pm 15$
Hadrons	$1049 \pm 329$	$1652 \pm 352$	$967 \pm 288$

Table F.7: Hadronic events selected in the 3753 (Scan) data.

3755 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$49692 \pm 222$	$56270 \pm 237$	$39753 \pm 199$
$q\bar{q}$	$32437 \pm 254$	$35604 \pm 272$	$25826 \pm 219$
$D^0\bar{D}^0$	$3917 \pm 8$	$4242 \pm 9$	$3188 \pm 7$
$D^+D^-$	$1655 \pm 3$	$1860 \pm 4$	$1329 \pm 3$
$\tau^+\tau^-$	$1116 \pm 3$	$2457 \pm 6$	$862 \pm 3$
$\gamma J/\psi$	$1610 \pm 3$	$1948 \pm 4$	$1199 \pm 3$
$\gamma\psi(2S)$	$8039 \pm 17$	$8903 \pm 19$	$6541 \pm 15$
Hadrons	$915 \pm 339$	$1253 \pm 362$	$803 \pm 296$

Table F.8: Hadronic events selected in the 3755 (Scan) data.

3756 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$57313 \pm 239$	$64807 \pm 254$	$46172 \pm 214$
$q\bar{q}$	$36748 \pm 288$	$40333 \pm 309$	$29261 \pm 248$
$D^0\bar{D}^0$	$4672 \pm 10$	$5042 \pm 10$	$3825 \pm 8$
$D^+D^-$	$2176 \pm 4$	$2437 \pm 5$	$1754 \pm 4$
$\tau^+\tau^-$	$1287 \pm 4$	$2809 \pm 7$	$997 \pm 4$
$\gamma J/\psi$	$1844 \pm 4$	$2222 \pm 5$	$1380 \pm 3$
$\gamma\psi(2S)$	$9141 \pm 20$	$10098 \pm 22$	$7466 \pm 17$
Hadrons	$1441 \pm 375$	$1863 \pm 401$	$1485 \pm 328$

Table F.9: Hadronic events selected in the 3756 (Scan) data.

3759 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$66837 \pm 258$	$75711 \pm 275$	$53739 \pm 231$
$q\bar{q}$	$42080 \pm 333$	$46166 \pm 357$	$33513 \pm 287$
$D^0\bar{D}^0$	$6410 \pm 12$	$6916 \pm 13$	$5253 \pm 10$
$D^+D^-$	$3260 \pm 6$	$3647 \pm 7$	$2627 \pm 5$
$\tau^+\tau^-$	$1474 \pm 4$	$3241 \pm 7$	$1145 \pm 4$
$\gamma J/\psi$	$2103 \pm 4$	$2536 \pm 5$	$1573 \pm 3$
$\gamma\psi(2S)$	$10067 \pm 20$	$11112 \pm 21$	$8234 \pm 17$
Hadrons	$1439 \pm 422$	$2090 \pm 451$	$1391 \pm 369$

Table F.10: Hadronic events selected in the 3759 (Scan) data.

3762 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$68538 \pm 261$	$77492 \pm 278$	$55377 \pm 235$
$q\bar{q}$	$42513 \pm 341$	$46621 \pm 365$	$33866 \pm 294$
$D^0\bar{D}^0$	$7945 \pm 15$	$8570 \pm 16$	$6508 \pm 12$
$D^+D^-$	$4238 \pm 8$	$4746 \pm 9$	$3418 \pm 7$
$\tau^+\tau^-$	$1499 \pm 4$	$3288 \pm 7$	$1164 \pm 3$
$\gamma J/\psi$	$2111 \pm 4$	$2545 \pm 5$	$1577 \pm 3$
$\gamma\psi(2S)$	$9779 \pm 19$	$10791 \pm 20$	$7995 \pm 16$
Hadrons	$449 \pm 431$	$927 \pm 460$	$846 \pm 377$

Table F.11: Hadronic events selected in the 3762 (Scan) data.

3765 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$51719 \pm 227$	$58419 \pm 241$	$41736 \pm 204$
$q\bar{q}$	$31053 \pm 253$	$34042 \pm 272$	$24741 \pm 218$
$D^0\bar{D}^0$	$6617 \pm 15$	$7135 \pm 16$	$5423 \pm 12$
$D^+D^-$	$3810 \pm 9$	$4265 \pm 9$	$3075 \pm 7$
$\tau^+\tau^-$	$1098 \pm 4$	$2417 \pm 7$	$854 \pm 3$
$\gamma J/\psi$	$1540 \pm 3$	$1854 \pm 4$	$1155 \pm 3$
$\gamma\psi(2S)$	$6939 \pm 16$	$7656 \pm 17$	$5679 \pm 14$
Hadrons	$657 \pm 341$	$1046 \pm 364$	$804 \pm 299$

Table F.12: Hadronic events selected in the 3765 (Scan) data.

3767 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$40038 \pm 200$	$45288 \pm 212$	$32311 \pm 179$
$q\bar{q}$	$23074 \pm 192$	$25285 \pm 206$	$18387 \pm 165$
$D^0\bar{D}^0$	$5629 \pm 14$	$6065 \pm 15$	$4613 \pm 12$
$D^+D^-$	$3314 \pm 8$	$3711 \pm 9$	$2673 \pm 7$
$\tau^+\tau^-$	$825 \pm 3$	$1808 \pm 5$	$640 \pm 2$
$\gamma J/\psi$	$1140 \pm 3$	$1374 \pm 3$	$853 \pm 2$
$\gamma\psi(2S)$	$4994 \pm 13$	$5509 \pm 14$	$4088 \pm 11$
Hadrons	$1058 \pm 278$	$1534 \pm 297$	$1053 \pm 244$

Table F.13: Hadronic events selected in the 3767 (Scan) data.

3771 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$34012 \pm 184$	$38173 \pm 195$	$27356 \pm 165$
$q\bar{q}$	$19014 \pm 161$	$20825 \pm 173$	$15156 \pm 138$
$D^0\bar{D}^0$	$5106 \pm 15$	$5504 \pm 16$	$4185 \pm 12$
$D^+D^-$	$3279 \pm 10$	$3668 \pm 11$	$2647 \pm 8$
$\tau^+\tau^-$	$683 \pm 3$	$1501 \pm 5$	$533 \pm 2$
$\gamma J/\psi$	$936 \pm 3$	$1127 \pm 3$	$702 \pm 2$
$\gamma\psi(2S)$	$3938 \pm 12$	$4349 \pm 13$	$3225 \pm 10$
Hadrons	$1053 \pm 246$	$1194 \pm 262$	$904 \pm 216$

Table F.14: Hadronic events selected in the 3771 (Scan) data.

3774 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$31457 \pm 177$	$35438 \pm 188$	$25241 \pm 158$
$q\bar{q}$	$17211 \pm 148$	$18843 \pm 159$	$13721 \pm 127$
$D^0\bar{D}^0$	$5038 \pm 15$	$5431 \pm 16$	$4126 \pm 13$
$D^+D^-$	$3456 \pm 10$	$3865 \pm 12$	$2786 \pm 9$
$\tau^+\tau^-$	$623 \pm 3$	$1367 \pm 5$	$483 \pm 2$
$\gamma J/\psi$	$843 \pm 2$	$1015 \pm 3$	$630 \pm 2$
$\gamma\psi(2S)$	$3458 \pm 10$	$3818 \pm 11$	$2822 \pm 9$
Hadrons	$825 \pm 232$	$1095 \pm 247$	$671 \pm 204$

Table F.15: Hadronic events selected in the 3774 (Scan) data.

3777 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$31195 \pm 176$	$35150 \pm 187$	$25095 \pm 158$
$q\bar{q}$	$17165 \pm 150$	$18785 \pm 161$	$13688 \pm 129$
$D^0\bar{D}^0$	$5006 \pm 16$	$5397 \pm 17$	$4100 \pm 13$
$D^+D^-$	$3464 \pm 11$	$3875 \pm 12$	$2794 \pm 9$
$\tau^+\tau^-$	$632 \pm 3$	$1380 \pm 5$	$492 \pm 3$
$\gamma J/\psi$	$838 \pm 3$	$1009 \pm 3$	$630 \pm 2$
$\gamma\psi(2S)$	$3329 \pm 11$	$3675 \pm 12$	$2723 \pm 9$
Hadrons	$758 \pm 233$	$1026 \pm 248$	$665 \pm 205$

Table F.16: Hadronic events selected in the 3777 (Scan) data.

3780 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$32426 \pm 180$	$36485 \pm 191$	$26084 \pm 161$
$q\bar{q}$	$18339 \pm 161$	$20062 \pm 173$	$14627 \pm 138$
$D^0\bar{D}^0$	$5011 \pm 14$	$5398 \pm 15$	$4114 \pm 12$
$D^+D^-$	$3139 \pm 9$	$3510 \pm 10$	$2536 \pm 8$
$\tau^+\tau^-$	$672 \pm 3$	$1477 \pm 5$	$523 \pm 2$
$\gamma J/\psi$	$892 \pm 2$	$1074 \pm 3$	$669 \pm 2$
$\gamma\psi(2S)$	$3469 \pm 10$	$3830 \pm 11$	$2843 \pm 9$
Hadrons	$900 \pm 242$	$1130 \pm 258$	$768 \pm 213$

Table F.17: Hadronic events selected in the 3780 (Scan) data.

3782 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$34028 \pm 184$	$38558 \pm 196$	$27372 \pm 165$
$q\bar{q}$	$20194 \pm 179$	$22082 \pm 192$	$16109 \pm 154$
$D^0\bar{D}^0$	$4464 \pm 12$	$4812 \pm 13$	$3660 \pm 10$
$D^+D^-$	$2938 \pm 8$	$3283 \pm 9$	$2373 \pm 7$
$\tau^+\tau^-$	$742 \pm 3$	$1631 \pm 5$	$576 \pm 2$
$\gamma J/\psi$	$979 \pm 3$	$1178 \pm 3$	$734 \pm 2$
$\gamma\psi(2S)$	$3717 \pm 10$	$4107 \pm 11$	$3037 \pm 9$
Hadrons	$991 \pm 258$	$1462 \pm 275$	$878 \pm 226$

Table F.18: Hadronic events selected in the 3782 (Scan) data.

3786 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$37498 \pm 193$	$42325 \pm 205$	$30190 \pm 173$
$q\bar{q}$	$23901 \pm 214$	$26122 \pm 229$	$19072 \pm 184$
$D^0\bar{D}^0$	$3793 \pm 10$	$4086 \pm 10$	$3112 \pm 8$
$D^+D^-$	$2568 \pm 7$	$2870 \pm 7$	$2078 \pm 5$
$\tau^+\tau^-$	$890 \pm 3$	$1940 \pm 6$	$695 \pm 3$
$\gamma J/\psi$	$1150 \pm 3$	$1384 \pm 3$	$863 \pm 2$
$\gamma\psi(2S)$	$4236 \pm 11$	$4677 \pm 12$	$3468 \pm 9$
Hadrons	$957 \pm 289$	$1241 \pm 309$	$898 \pm 253$

Table F.19: Hadronic events selected in the 3786 (Scan) data.

3789 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$39308 \pm 198$	$44415 \pm 210$	$31651 \pm 177$
$q\bar{q}$	$26534 \pm 239$	$28991 \pm 256$	$21176 \pm 206$
$D^0\bar{D}^0$	$2874 \pm 6$	$3096 \pm 7$	$2358 \pm 5$
$D^+D^-$	$2064 \pm 5$	$2308 \pm 5$	$1668 \pm 4$
$\tau^+\tau^-$	$989 \pm 3$	$2162 \pm 6$	$773 \pm 3$
$\gamma J/\psi$	$1273 \pm 3$	$1532 \pm 3$	$955 \pm 2$
$\gamma\psi(2S)$	$4607 \pm 11$	$5084 \pm 12$	$3773 \pm 9$
Hadrons	$964 \pm 311$	$1238 \pm 332$	$944 \pm 272$

Table F.20: Hadronic events selected in the 3789 (Scan) data.

3792 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$46626 \pm 215$	$53085 \pm 230$	$37384 \pm 193$
$q\bar{q}$	$33038 \pm 300$	$36079 \pm 322$	$26373 \pm 259$
$D^0\bar{D}^0$	$2416 \pm 5$	$2604 \pm 5$	$1979 \pm 4$
$D^+D^-$	$1482 \pm 3$	$1657 \pm 3$	$1197 \pm 2$
$\tau^+\tau^-$	$1243 \pm 4$	$2707 \pm 7$	$964 \pm 4$
$\gamma J/\psi$	$1570 \pm 3$	$1893 \pm 4$	$1174 \pm 3$
$\gamma\psi(2S)$	$5543 \pm 12$	$6120 \pm 13$	$4526 \pm 11$
Hadrons	$1330 \pm 370$	$2020 \pm 396$	$1166 \pm 323$

Table F.21: Hadronic events selected in the 3792 (Scan) data.

3797 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$51348 \pm 226$	$58331 \pm 241$	$41249 \pm 203$
$q\bar{q}$	$37808 \pm 349$	$41264 \pm 373$	$30191 \pm 300$
$D^0\bar{D}^0$	$1693 \pm 3$	$1825 \pm 3$	$1388 \pm 3$
$D^+D^-$	$1036 \pm 2$	$1158 \pm 2$	$837 \pm 1$
$\tau^+\tau^-$	$1427 \pm 5$	$3118 \pm 8$	$1112 \pm 4$
$\gamma J/\psi$	$1790 \pm 4$	$2156 \pm 4$	$1341 \pm 3$
$\gamma\psi(2S)$	$6119 \pm 13$	$6755 \pm 14$	$5005 \pm 11$
Hadrons	$1471 \pm 416$	$2050 \pm 445$	$1372 \pm 363$

Table F.22: Hadronic events selected in the 3797 (Scan) data.

3800 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$48859 \pm 221$	$55719 \pm 236$	$38965 \pm 197$
$q\bar{q}$	$36644 \pm 342$	$39976 \pm 367$	$29268 \pm 295$
$D^0\bar{D}^0$	$1235 \pm 2$	$1333 \pm 2$	$1010 \pm 2$
$D^+D^-$	$583 \pm 1$	$652 \pm 1$	$470 \pm 1$
$\tau^+\tau^-$	$1395 \pm 4$	$3051 \pm 7$	$1081 \pm 4$
$\gamma J/\psi$	$1721 \pm 3$	$2076 \pm 4$	$1286 \pm 3$
$\gamma\psi(2S)$	$5748 \pm 12$	$6354 \pm 13$	$4686 \pm 10$
Hadrons	$1530 \pm 408$	$2273 \pm 436$	$1160 \pm 355$

Table F.23: Hadronic events selected in the 3800 (Scan) data.

3802 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$33032 \pm 181$	$37538 \pm 193$	$26268 \pm 162$
$q\bar{q}$	$25099 \pm 238$	$27373 \pm 255$	$20050 \pm 205$
$D^0\bar{D}^0$	$649 \pm 1$	$700 \pm 1$	$531 \pm 1$
$D^+D^-$	$391 \pm 1$	$438 \pm 1$	$315 \pm 1$
$\tau^+\tau^-$	$963 \pm 3$	$2103 \pm 6$	$746 \pm 3$
$\gamma J/\psi$	$1177 \pm 3$	$1418 \pm 3$	$880 \pm 2$
$\gamma\psi(2S)$	$3871 \pm 10$	$4280 \pm 10$	$3161 \pm 8$
Hadrons	$880 \pm 300$	$1223 \pm 320$	$582 \pm 261$

Table F.24: Hadronic events selected in the 3802 (Scan) data.

3807 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$21499 \pm 146$	$24410 \pm 156$	$17234 \pm 131$
$q\bar{q}$	$16378 \pm 159$	$17850 \pm 171$	$13087 \pm 137$
$D^0\bar{D}^0$	$459 \pm 1$	$495 \pm 1$	$374 \pm 1$
$D^+D^-$	$311 \pm 1$	$349 \pm 1$	$251 \pm 1$
$\tau^+\tau^-$	$632 \pm 2$	$1377 \pm 5$	$489 \pm 2$
$\gamma J/\psi$	$761 \pm 2$	$918 \pm 2$	$567 \pm 1$
$\gamma\psi(2S)$	$2427 \pm 7$	$2688 \pm 8$	$1980 \pm 6$
Hadrons	$527 \pm 217$	$729 \pm 232$	$482 \pm 190$

Table F.25: Hadronic events selected in the 3807 (Scan) data.

3809 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$15124 \pm 122$	$17177 \pm 131$	$12089 \pm 109$
$q\bar{q}$	$11655 \pm 116$	$12699 \pm 124$	$9315 \pm 99$
$D^0\bar{D}^0$	$217 \pm 1$	$234 \pm 1$	$177 \pm 1$
$D^+D^-$	$102 \pm 1$	$114 \pm 1$	$82 \pm 1$
$\tau^+\tau^-$	$450 \pm 2$	$986 \pm 4$	$351 \pm 2$
$\gamma J/\psi$	$541 \pm 2$	$652 \pm 2$	$406 \pm 1$
$\gamma\psi(2S)$	$1705 \pm 6$	$1885 \pm 7$	$1395 \pm 5$
Hadrons	$451 \pm 169$	$605 \pm 181$	$360 \pm 148$

Table F.26: Hadronic events selected in the 3809 (Scan) data.

3813 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$10836 \pm 104$	$12308 \pm 110$	$8700 \pm 93$
$q\bar{q}$	$8323 \pm 85$	$9063 \pm 91$	$6653 \pm 73$
$D^0\overline{D}^0$	$39 \pm 1$	$43 \pm 1$	$32 \pm 1$
$D^+D^-$	$84 \pm 1$	$94 \pm 1$	$67 \pm 1$
$\tau^+\tau^-$	$321 \pm 2$	$703 \pm 3$	$247 \pm 1$
$\gamma J/\psi$	$384 \pm 1$	$463 \pm 2$	$286 \pm 1$
$\gamma\psi(2S)$	$1173 \pm 5$	$1298 \pm 5$	$954 \pm 4$
Hadrons	$509 \pm 135$	$641 \pm 144$	$456 \pm 118$

Table F.27: Hadronic events selected in the 3813 (Scan) data.

3815 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$8159 \pm 90$	$9190 \pm 95$	$6536 \pm 80$
$q\bar{q}$	$6308 \pm 66$	$6868 \pm 71$	$5044 \pm 56$
$D^0\overline{D}^0$	$28 \pm 1$	$30 \pm 1$	$23 \pm 1$
$D^+D^-$	$37 \pm 1$	$41 \pm 1$	$30 \pm 1$
$\tau^+\tau^-$	$247 \pm 1$	$540 \pm 3$	$192 \pm 1$
$\gamma J/\psi$	$290 \pm 1$	$350 \pm 1$	$217 \pm 1$
$\gamma\psi(2S)$	$883 \pm 4$	$975 \pm 4$	$719 \pm 3$
Hadrons	$363 \pm 112$	$383 \pm 119$	$309 \pm 99$

Table F.28: Hadronic events selected in the 3815 (Scan) data.

3822 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$4748 \pm 68$	$5379 \pm 73$	$3754 \pm 61$
$q\bar{q}$	$3695 \pm 41$	$4019 \pm 44$	$2956 \pm 35$
$D^0\overline{D}^0$	$41 \pm 1$	$44 \pm 1$	$34 \pm 1$
$D^+D^-$	$48 \pm 1$	$54 \pm 1$	$39 \pm 1$
$\tau^+\tau^-$	$147 \pm 1$	$321 \pm 2$	$114 \pm 1$
$\gamma J/\psi$	$168 \pm 1$	$203 \pm 1$	$126 \pm 1$
$\gamma\psi(2S)$	$491 \pm 2$	$542 \pm 3$	$401 \pm 2$
Hadrons	$154 \pm 80$	$193 \pm 86$	$81 \pm 70$

Table F.29: Hadronic events selected in the 3822 (Scan) data.

3832 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$3364 \pm 58$	$3842 \pm 61$	$2729 \pm 52$
$q\bar{q}$	$2622 \pm 31$	$2848 \pm 34$	$2098 \pm 26$
$D^0\overline{D}^0$	$14 \pm 1$	$15 \pm 1$	$12 \pm 1$
$D^+D^-$	$8 \pm 1$	$9 \pm 1$	$6 \pm 1$
$\tau^+\tau^-$	$105 \pm 1$	$229 \pm 1$	$82 \pm 1$
$\gamma J/\psi$	$118 \pm 1$	$142 \pm 1$	$88 \pm 1$
$\gamma\psi(2S)$	$326 \pm 2$	$360 \pm 2$	$266 \pm 2$
Hadrons	$168 \pm 66$	$236 \pm 70$	$174 \pm 58$

Table F.30: Hadronic events selected in the 3832 (Scan) data.

3839 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$3156 \pm 56$	$3567 \pm 59$	$2569 \pm 50$
$q\bar{q}$	$2574 \pm 31$	$2793 \pm 34$	$2061 \pm 27$
$D^0\overline{D}^0$	$49 \pm 1$	$53 \pm 1$	$40 \pm 1$
$D^+D^-$	$43 \pm 1$	$49 \pm 1$	$35 \pm 1$
$\tau^+\tau^-$	$103 \pm 1$	$225 \pm 1$	$80 \pm 1$
$\gamma J/\psi$	$114 \pm 1$	$138 \pm 1$	$86 \pm 1$
$\gamma\psi(2S)$	$306 \pm 2$	$339 \pm 2$	$251 \pm 1$
Hadrons	$-36 \pm 64$	$-32 \pm 68$	$13 \pm 57$

Table F.31: Hadronic events selected in the 3839 (Scan) data.

3849 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$3266 \pm 57$	$3738 \pm 61$	$2603 \pm 51$
$q\bar{q}$	$2529 \pm 32$	$2740 \pm 34$	$2027 \pm 27$
$D^0\overline{D}^0$	$38 \pm 1$	$41 \pm 1$	$31 \pm 1$
$D^+D^-$	$31 \pm 1$	$35 \pm 1$	$25 \pm 1$
$\tau^+\tau^-$	$104 \pm 1$	$225 \pm 1$	$81 \pm 1$
$\gamma J/\psi$	$111 \pm 1$	$133 \pm 1$	$83 \pm 1$
$\gamma\psi(2S)$	$281 \pm 2$	$312 \pm 2$	$230 \pm 1$
Hadrons	$168 \pm 65$	$249 \pm 70$	$123 \pm 57$

Table F.32: Hadronic events selected in the 3849 (Scan) data.

3855 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$3640 \pm 60$	$4097 \pm 64$	$2905 \pm 53$
$q\bar{q}$	$2910 \pm 36$	$3151 \pm 39$	$2333 \pm 31$
$D^0\overline{D^0}$	$80 \pm 1$	$86 \pm 1$	$65 \pm 1$
$D^+D^-$	$53 \pm 1$	$60 \pm 1$	$43 \pm 1$
$\tau^+\tau^-$	$122 \pm 1$	$263 \pm 1$	$94 \pm 1$
$\gamma J/\psi$	$127 \pm 1$	$152 \pm 1$	$95 \pm 1$
$\gamma\psi(2S)$	$314 \pm 2$	$347 \pm 2$	$256 \pm 1$
Hadrons	$32 \pm 70$	$35 \pm 75$	$16 \pm 62$

Table F.33: Hadronic events selected in the 3855 (Scan) data.

3863 (Scan) Results			
Sample	$N_{\text{had}}$ (SHAD)	$N_{\text{had}}$ (LHAD)	$N_{\text{had}}$ (THAD)
Data	$3472 \pm 58$	$3921 \pm 62$	$2773 \pm 52$
$q\bar{q}$	$2732 \pm 35$	$2955 \pm 38$	$2191 \pm 30$
$D^0\overline{D^0}$	$76 \pm 1$	$82 \pm 1$	$62 \pm 1$
$D^+D^-$	$18 \pm 1$	$20 \pm 1$	$14 \pm 1$
$\tau^+\tau^-$	$114 \pm 1$	$247 \pm 1$	$89 \pm 1$
$\gamma J/\psi$	$118 \pm 1$	$142 \pm 1$	$88 \pm 1$
$\gamma\psi(2S)$	$282 \pm 1$	$312 \pm 2$	$230 \pm 1$
Hadrons	$129 \pm 68$	$160 \pm 73$	$94 \pm 60$

Table F.34: Hadronic events selected in the 3863 (Scan) data.

,