

ECEN 303: Random Signals and Systems

Chapter 1: Mathematical Review

References

- Chapter 1, Class Notes

Set

- A **set** is a collection of objects, which are called the **elements** of the set.

Set

- A **set** is a collection of objects, which are called the **elements** of the set.
- **Examples:**

Set

- A **set** is a collection of objects, which are called the **elements** of the set.
- **Examples:**
 - ▶ Set of the non-negative integers less than five

Set

- A **set** is a collection of objects, which are called the **elements** of the set.
- **Examples:**
 - ▶ Set of the non-negative integers less than five
 - ▶ Set of the rivers in North America

Set

- A **set** is a collection of objects, which are called the **elements** of the set.
- **Examples:**
 - ▶ Set of the non-negative integers less than five
 - ▶ Set of the rivers in North America
 - ▶ Set of the ECEN courses offered in this semesters

Set

- A **set** is a collection of objects, which are called the **elements** of the set.
- **Examples:**
 - ▶ Set of the non-negative integers less than five
 - ▶ Set of the rivers in North America
 - ▶ Set of the ECEN courses offered in this semesters
 - ▶ Set of the positive numbers divisible by three

Set

- A **set** is a collection of objects, which are called the **elements** of the set.
- **Examples:**
 - ▶ Set of the non-negative integers less than five
 - ▶ Set of the rivers in North America
 - ▶ Set of the ECEN courses offered in this semesters
 - ▶ Set of the positive numbers divisible by three
 - ▶ Set of the real numbers between two and three

Set

- A **set** is a collection of objects, which are called the **elements** of the set.
- **Examples:**
 - ▶ Set of the non-negative integers less than five
 - ▶ Set of the rivers in North America
 - ▶ Set of the ECEN courses offered in this semesters
 - ▶ Set of the positive numbers divisible by three
 - ▶ Set of the real numbers between two and three
 - ▶ \vdots

Representing a Set

- If it contains only a few elements, one can simply list the objects in the set

Representing a Set

- If it contains only a few elements, one can simply list the objects in the set
 - ▶ Set of the non-negative integers less than five = $\{0, 1, 2, 3, 4\}$

Representing a Set

- If it contains only a few elements, one can simply list the objects in the set
 - ▶ Set of the non-negative integers less than five = $\{0, 1, 2, 3, 4\}$
 - ▶ Set of the rivers in North America = $\{ \text{Mackenzie, Colorado, Yukon, Mississippi, Brazos, } \dots, \text{Hudson} \}$

Representing a Set

- If it contains only a few elements, one can simply list the objects in the set
 - ▶ Set of the non-negative integers less than five = $\{0, 1, 2, 3, 4\}$
 - ▶ Set of the rivers in North America = $\{ \text{Mackenzie, Colorado, Yukon, Mississippi, Brazos, } \dots, \text{Hudson} \}$
- The content of a set can also be enumerated whenever it has a **countable** number of elements

Representing a Set

- If it contains only a few elements, one can simply list the objects in the set
 - ▶ Set of the non-negative integers less than five = $\{0, 1, 2, 3, 4\}$
 - ▶ Set of the rivers in North America = $\{ \text{Mackenzie, Colorado, Yukon, Mississippi, Brazos, } \dots, \text{Hudson} \}$
- The content of a set can also be enumerated whenever it has a **countable** number of elements
 - ▶ Set of the positive numbers divisible by three = $\{3, 6, 9, \dots\}$

Representing a Set

- If it contains only a few elements, one can simply list the objects in the set
 - ▶ Set of the non-negative integers less than five = $\{0, 1, 2, 3, 4\}$
 - ▶ Set of the rivers in North America = $\{ \text{Mackenzie, Colorado, Yukon, Mississippi, Brazos, } \dots, \text{Hudson} \}$
- The content of a set can also be enumerated whenever it has a **countable** number of elements
 - ▶ Set of the positive numbers divisible by three = $\{3, 6, 9, \dots\}$
- Usually, the way to specify a set is to take some collection S of objects and some property that elements of S may or may not possess, and to form the set consisting of all elements of S having that property

Representing a Set

- If it contains only a few elements, one can simply list the objects in the set
 - ▶ Set of the non-negative integers less than five = $\{0, 1, 2, 3, 4\}$
 - ▶ Set of the rivers in North America = $\{ \text{Mackenzie, Colorado, Yukon, Mississippi, Brazos, } \dots, \text{Hudson} \}$
- The content of a set can also be enumerated whenever it has a **countable** number of elements
 - ▶ Set of the positive numbers divisible by three = $\{3, 6, 9, \dots\}$
- Usually, the way to specify a set is to take some collection S of objects and some property that elements of S may or may not possess, and to form the set consisting of all elements of S having that property
 - ▶ Set of the positive numbers divisible by three = $\{x \mid x > 0 \text{ and } x \text{ is divisible by } 3\}$

Representing a Set

- If it contains only a few elements, one can simply list the objects in the set
 - ▶ Set of the non-negative integers less than five = $\{0, 1, 2, 3, 4\}$
 - ▶ Set of the rivers in North America = $\{ \text{Mackenzie, Colorado, Yukon, Mississippi, Brazos, } \dots, \text{Hudson} \}$
- The content of a set can also be enumerated whenever it has a **countable** number of elements
 - ▶ Set of the positive numbers divisible by three = $\{3, 6, 9, \dots\}$
- Usually, the way to specify a set is to take some collection S of objects and some property that elements of S may or may not possess, and to form the set consisting of all elements of S having that property
 - ▶ Set of the positive numbers divisible by three = $\{x \mid x > 0 \text{ and } x \text{ is divisible by } 3\}$
 - ▶ Set of the real numbers between two and three = $\{x \in \mathbb{R} \mid 2 \leq x \leq 3\}$

Set Equivalence

- If an element x belongs to a set S , we express this fact by writing $x \in S$.

Set Equivalence

- If an element x belongs to a set S , we express this fact by writing $x \in S$.
- If x does not belong to S , we write $x \notin S$.

Set Equivalence

- If an element x belongs to a set S , we express this fact by writing $x \in S$.
- If x does not belong to S , we write $x \notin S$.
- We use the equality symbol to denote *logical identity*.

Set Equivalence

- If an element x belongs to a set S , we express this fact by writing $x \in S$.
- If x does not belong to S , we write $x \notin S$.
- We use the equality symbol to denote *logical identity*.
 - ▶ $x = y$ means that x and y are symbols denoting the same object.

Set Equivalence

- If an element x belongs to a set S , we express this fact by writing $x \in S$.
- If x does not belong to S , we write $x \notin S$.
- We use the equality symbol to denote *logical identity*.
 - ▶ $x = y$ means that x and y are symbols denoting the same object.
- $S = T$ states that S and T are two symbols for the same set.

Set Equivalence

- If an element x belongs to a set S , we express this fact by writing $x \in S$.
- If x does not belong to S , we write $x \notin S$.
- We use the equality symbol to denote *logical identity*.
 - ▶ $x = y$ means that x and y are symbols denoting the same object.
- $S = T$ states that S and T are two symbols for the same set.
 - ▶ The sets S and T contain precisely the same elements

Set Equivalence

- If an element x belongs to a set S , we express this fact by writing $x \in S$.
- If x does not belong to S , we write $x \notin S$.
- We use the equality symbol to denote *logical identity*.
 - ▶ $x = y$ means that x and y are symbols denoting the same object.
- $S = T$ states that S and T are two symbols for the same set.
 - ▶ The sets S and T contain precisely the same elements
- $S \neq T$ expresses the fact that S and T are different

Subset, Empty Set

- A set S is a **subset** of T if every element of S is also contained in T .

Subset, Empty Set

- A set S is a **subset** of T if every element of S is also contained in T .
 - ▶ Example: $S = \{2, 4, 6, 8\}, T = \{1, 2, 3, 4, \dots, 10\}$

Subset, Empty Set

- A set S is a **subset** of T if every element of S is also contained in T .
 - ▶ Example: $S = \{2, 4, 6, 8\}, T = \{1, 2, 3, 4, \dots, 10\}$
 - ▶ We express this relation by writing $S \subset T$

Subset, Empty Set

- A set S is a **subset** of T if every element of S is also contained in T .
 - ▶ Example: $S = \{2, 4, 6, 8\}, T = \{1, 2, 3, 4, \dots, 10\}$
 - ▶ We express this relation by writing $S \subset T$
 - ▶ A set can be its own subset

Subset, Empty Set

- A set S is a **subset** of T if every element of S is also contained in T .
 - ▶ Example: $S = \{2, 4, 6, 8\}, T = \{1, 2, 3, 4, \dots, 10\}$
 - ▶ We express this relation by writing $S \subset T$
 - ▶ A set can be its own subset
- $S = T$ **if and only if** $S \subset T$ and $T \subset S$.

Subset, Empty Set

- A set S is a **subset** of T if every element of S is also contained in T .
 - ▶ Example: $S = \{2, 4, 6, 8\}$, $T = \{1, 2, 3, 4, \dots, 10\}$
 - ▶ We express this relation by writing $S \subset T$
 - ▶ A set can be its own subset
- $S = T$ **if and only if** $S \subset T$ and $T \subset S$.
- The **empty set**, denoted by \emptyset , is a set that contains no elements.

Universal Set, Complement of a Set

- The **universal set** is the collection of all objects of interest in a particular context, and it is represented by Ω .

Universal Set, Complement of a Set

- The **universal set** is the collection of all objects of interest in a particular context, and it is represented by Ω .
 - ▶ Example: If the objects of interest are positive integers, then $\Omega = \{1, 2, 3, \dots\}$

Universal Set, Complement of a Set

- The **universal set** is the collection of all objects of interest in a particular context, and it is represented by Ω .
 - ▶ Example: If the objects of interest are positive integers, then $\Omega = \{1, 2, 3, \dots\}$
 - ▶ Example: If the objects of interest are positive integers less than or equal to 10, then $\Omega = \{1, 2, 3, \dots, 10\}$

Universal Set, Complement of a Set

- The **universal set** is the collection of all objects of interest in a particular context, and it is represented by Ω .
 - ▶ Example: If the objects of interest are positive integers, then $\Omega = \{1, 2, 3, \dots\}$
 - ▶ Example: If the objects of interest are positive integers less than or equal to 10, then $\Omega = \{1, 2, 3, \dots, 10\}$
- The **complement** of a set S , denoted as S^c , is the collection of all objects in Ω that do not belong to S , $S^c = \{x \in \Omega | x \notin S\}$

Universal Set, Complement of a Set

- The **universal set** is the collection of all objects of interest in a particular context, and it is represented by Ω .
 - ▶ Example: If the objects of interest are positive integers, then $\Omega = \{1, 2, 3, \dots\}$
 - ▶ Example: If the objects of interest are positive integers less than or equal to 10, then $\Omega = \{1, 2, 3, \dots, 10\}$
- The **complement** of a set S , denoted as S^c , is the collection of all objects in Ω that do not belong to S , $S^c = \{x \in \Omega | x \notin S\}$
 - ▶ $S = \{2, 4, 6, 8\}$, $\Omega = \{1, 2, 3, \dots, 10\}$, $S^c = ?$

Union and Intersection

- The **union** of sets S and T is the collection of all elements that belong to S or T (or both), and it is denoted by $S \cup T$.

Union and Intersection



- The **union** of sets S and T is the collection of all elements that belong to S or T (or both), and it is denoted by $S \cup T$.
- Formally, $S \cup T = \{x | x \in S \text{ OR } x \in T\}$.

Union and Intersection

- The **union** of sets S and T is the collection of all elements that belong to S or T (or both), and it is denoted by $S \cup T$.
- Formally, $S \cup T = \{x | x \in S \text{ OR } x \in T\}$.

Union and Intersection

- The **union** of sets S and T is the collection of all elements that belong to S or T (or both), and it is denoted by $S \cup T$.
- Formally, $S \cup T = \{x | x \in S \text{ OR } x \in T\}$.



Union and Intersection

- The **intersection** of sets S and T is the collection of all elements that belong to both S and T , and it is denoted by $S \cap T$.

Union and Intersection



- The **intersection** of sets S and T is the collection of all elements that belong to both S and T , and it is denoted by $S \cap T$.
- Formally, $S \cap T = \{x | x \in S \text{ AND } x \in T\}$.

Union and Intersection

- The **intersection** of sets S and T is the collection of all elements that belong to both S and T , and it is denoted by $S \cap T$.
- Formally, $S \cap T = \{x | x \in S \text{ AND } x \in T\}$.

Union and Intersection

- The **intersection** of sets S and T is the collection of all elements that belong to both S and T , and it is denoted by $S \cap T$.
- Formally, $S \cap T = \{x | x \in S \text{ AND } x \in T\}$.



Disjoint Sets and Partition

- The sets S and T are **disjoint** if they have no elements in common

Disjoint Sets and Partition

- The sets S and T are **disjoint** if they have no elements in common
 - ▶ We write it as $S \cap T = \emptyset$

Disjoint Sets and Partition



- The sets S and T are **disjoint** if they have no elements in common
 - ▶ We write it as $S \cap T = \emptyset$
- A collection of sets is said to form a **partition** of S if the sets in the collection are disjoint and their union is S .

Disjoint Sets and Partition

- The sets S and T are **disjoint** if they have no elements in common
 - ▶ We write it as $S \cap T = \emptyset$
- A collection of sets is said to form a **partition** of S if the sets in the collection are disjoint and their union is S .

Disjoint Sets and Partition

- The sets S and T are **disjoint** if they have no elements in common
 - ▶ We write it as $S \cap T = \emptyset$
- A collection of sets is said to form a **partition** of S if the sets in the collection are disjoint and their union is S .



Algebra of Sets: Distributive laws

Distributive laws

$$R \cap (S \cup T) = (R \cap S) \cup (R \cap T)$$

$$R \cup (S \cap T) = (R \cup S) \cap (R \cup T).$$

- Ordering of operation is important



Algebra of Sets: Distributive laws

Distributive laws

$$R \cap (S \cup T) = (R \cap S) \cup (R \cap T)$$

$$R \cup (S \cap T) = (R \cup S) \cap (R \cup T).$$

- Ordering of operation is important



Algebra of Sets: De Morgan's laws

De Morgan's laws

① $(A \cup B)^c = A^c \cap B^c$

② $(A \cap B)^c = A^c \cup B^c$

Algebra of Sets: De Morgan's laws

De Morgan's laws

$$① \quad (A \cup B)^c = A^c \cap B^c$$

$$② \quad (A \cap B)^c = A^c \cup B^c$$

De Morgan's laws

$$① \quad (\cup_{i=1}^n A_i)^c = \cap_{i=1}^n A_i^c$$

$$② \quad (\cap_{i=1}^n A_i)^c = \cup_{i=1}^n A_i^c$$


Proof of De Morgan's laws

Cartesian Products

- Given sets S and T , the **cartesian product** $S \times T$ is the set of all **ordered pairs** (x, y) such that x is an element of S and y is an element of T ,
 $S \times T = \{(x, y) | x \in S \text{ and } y \in T\}$.

Cartesian Products

- Given sets S and T , the **cartesian product** $S \times T$ is the set of all **ordered pairs** (x, y) such that x is an element of S and y is an element of T ,
 $S \times T = \{(x, y) | x \in S \text{ and } y \in T\}$.



Function

- A **function** is a relation that assigns **exactly one** value to each possible input.

Function



- A **function** is a relation that assigns **exactly one** value to each possible input.
- A common representation is $f(x) = y$, where f denotes the rule of correspondence.

Function

- A **function** is a relation that assigns **exactly one** value to each possible input.
- A common representation is $f(x) = y$, where f denotes the rule of correspondence.


Function

- A **function** is a relation that assigns **exactly one** value to each possible input.
- A common representation is $f(x) = y$, where f denotes the rule of correspondence.




Function

- A **function** is a relation that assigns **exactly one** value to each possible input.
- A common representation is $f(x) = y$, where f denotes the rule of correspondence.



This is a function



This is NOT a function

Domain, Codomain, Image/Range

- The **domain** of a function is the set over which this function is defined; that is, the collection of arguments that can be used as input.

Domain, Codomain, Image/Range

- The **domain** of a function is the set over which this function is defined; that is, the collection of arguments that can be used as input.
- The **codomain** is the set into which all the outputs of the function are constrained to lie.


Domain, Codomain, Image/Range

- The **domain** of a function is the set over which this function is defined; that is, the collection of arguments that can be used as input.
- The **codomain** is the set into which all the outputs of the function are constrained to lie.
- To specify these sets explicitly, the notation $f : X \rightarrow Y$ is frequently used, where X indicates the domain and Y is the codomain.

Domain, Codomain, Image/Range

- The **domain** of a function is the set over which this function is defined; that is, the collection of arguments that can be used as input.
- The **codomain** is the set into which all the outputs of the function are constrained to lie.
- To specify these sets explicitly, the notation $f : X \rightarrow Y$ is frequently used, where X indicates the domain and Y is the codomain.
- The **image** of X under function $f : X \rightarrow Y$ is the set of all objects of the form $f(x)$, where x ranges over the elements of X : $\{f(x) \in Y | x \in X\}$

Domain, Codomain, Image/Range



$$f : X \rightarrow Y$$

Function: Examples

- Example: Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ where $f(x) = x^2$. In this case, the domain \mathbb{R} and the codomain \mathbb{R} are identical. The rule of correspondence for this function is $x \mapsto x^2$, which should be read “ x maps to x^2 .”

Function: Examples

- Example: Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ where $f(x) = x^2$. In this case, the domain \mathbb{R} and the codomain \mathbb{R} are identical. The rule of correspondence for this function is $x \mapsto x^2$, which should be read “ x maps to x^2 .”
- Example: Suppose S is a subset of the real numbers. We define the **indicator function** of set S , denoted $\mathbf{1}_S : \mathbb{R} \rightarrow \{0, 1\}$, by

$$\mathbf{1}_S(x) = \begin{cases} 1, & x \in S \\ 0, & x \notin S. \end{cases}$$

Preimage

- The **preimage** of a set $T \subset Y$ under $f : X \rightarrow Y$ is the subset of X defined by $f^{-1}(T) = \{x \in X \mid f(x) \in T\}$


Preimage

- The **preimage** of a set $T \subset Y$ under $f : X \rightarrow Y$ is the subset of X defined by $f^{-1}(T) = \{x \in X \mid f(x) \in T\}$


Preimage

- The **preimage** of a set $T \subset Y$ under $f : X \rightarrow Y$ is the subset of X defined by $f^{-1}(T) = \{x \in X \mid f(x) \in T\}$

Person Favorite Food



$f: \text{Person} \rightarrow \text{Favorite Food}$



$f: X \rightarrow Y$

Preimage

- The **preimage** of a set $T \subset Y$ under $f : X \rightarrow Y$ is the subset of X defined by $f^{-1}(T) = \{x \in X | f(x) \in T\}$

Preimage

- The **preimage** of a set $T \subset Y$ under $f : X \rightarrow Y$ is the subset of X defined by $f^{-1}(T) = \{x \in X \mid f(x) \in T\}$
- Example: Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ where $f(x) = x^2$. What is the preimage of $\{4\}$?

Preimage

- The **preimage** of a set $T \subset Y$ under $f : X \rightarrow Y$ is the subset of X defined by $f^{-1}(T) = \{x \in X \mid f(x) \in T\}$
- Example: Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ where $f(x) = x^2$. What is the preimage of $\{4\}$?
- Example: Consider the function $f : [0, \infty) \rightarrow [0, \infty)$ where $f(x) = x^2$. What is the preimage of $\{4\}$?

One-to-one (Injective) function

- The function $f : X \rightarrow Y$ is **injective** or **one-to-one** if $f(x_1) = f(x_2)$ implies $x_1 = x_2$ for all $x_1, x_2 \in X$.

One-to-one (Injective) function

- The function $f : X \rightarrow Y$ is **injective** or **one-to-one** if $f(x_1) = f(x_2)$ implies $x_1 = x_2$ for all $x_1, x_2 \in X$.
- Example: Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ where $f(x) = x^2$. Is this injective?

One-to-one (Injective) function

- The function $f : X \rightarrow Y$ is **injective** or **one-to-one** if $f(x_1) = f(x_2)$ implies $x_1 = x_2$ for all $x_1, x_2 \in X$.
- Example: Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ where $f(x) = x^2$. Is this injective?
- Example: Consider the function $f : [0, \infty) \rightarrow [0, \infty)$ where $f(x) = x^2$. Is this injective?

Onto (Surjective) function

- The function f is **surjective** or **onto** if its image is equal to the codomain. More specifically, a function $f : X \rightarrow Y$ is surjective if and only if, for every $y \in Y$, there exists $x \in X$ such that $f(x) = y$.

Onto (Surjective) function

- The function f is **surjective** or **onto** if its image is equal to the codomain. More specifically, a function $f : X \rightarrow Y$ is surjective if and only if, for every $y \in Y$, there exists $x \in X$ such that $f(x) = y$.
- Example: Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ where $f(x) = 2x + 1$. Is this surjective?

Onto (Surjective) function

- The function f is **surjective** or **onto** if its image is equal to the codomain. More specifically, a function $f : X \rightarrow Y$ is surjective if and only if, for every $y \in Y$, there exists $x \in X$ such that $f(x) = y$.
- Example: Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ where $f(x) = 2x + 1$. Is this surjective?
- Example: Consider the function $f : [0, \infty) \rightarrow [0, \infty)$ where $f(x) = x^2$. Is this surjective?

Onto (Surjective) function

- The function f is **surjective** or **onto** if its image is equal to the codomain. More specifically, a function $f : X \rightarrow Y$ is surjective if and only if, for every $y \in Y$, there exists $x \in X$ such that $f(x) = y$.
- Example: Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ where $f(x) = 2x + 1$. Is this surjective?
- Example: Consider the function $f : [0, \infty) \rightarrow [0, \infty)$ where $f(x) = x^2$. Is this surjective?
- Example: Consider the function $f : \mathbb{R} \rightarrow \mathbb{R}$ where $f(x) = x^2$. Is this surjective?

Bijjective function

- A function that is both one-to-one and onto is called a **bijection**

Bijjective function

- A function that is both one-to-one and onto is called a **bijection**
- A function f is bijective if and only if its inverse relation f^{-1} is itself a function.

Examples

