

# ECEN 303: Random Signals and Systems

## Chapter 3: Basic Concepts of Probability

# References

- Chapter 3, Class Notes
- D. Bertsekas and J. Tsitsiklis, *Introduction to Probability*, Chapter 1.2
- M. Harchol-Balter, *Introduction to Probability for Computing*, Chapter 2
- S. Ross, *A First Course in Probability*, Chapter 2

# Experiment

In the context of probability, an **experiment** is a random occurrence that produces one of several outcomes.

# Experiment

In the context of probability, an **experiment** is a random occurrence that produces one of several outcomes.

- Experiment of tossing a coin

# Experiment

In the context of probability, an **experiment** is a random occurrence that produces one of several outcomes.

- Experiment of tossing a coin
- Experiment of rolling a die

# Experiment

In the context of probability, an **experiment** is a random occurrence that produces one of several outcomes.

- Experiment of tossing a coin
- Experiment of rolling a die
- Experiment of rolling a die twice

# Experiment

In the context of probability, an **experiment** is a random occurrence that produces one of several outcomes.

- Experiment of tossing a coin
- Experiment of rolling a die
- Experiment of rolling a die twice
- Experiment of conducting a survey (who will be the next president?)

# Experiment

In the context of probability, an **experiment** is a random occurrence that produces one of several outcomes.

- Experiment of tossing a coin
- Experiment of rolling a die
- Experiment of rolling a die twice
- Experiment of conducting a survey (who will be the next president?)
- Experiment of measuring the temperature at 2PM in College Station in August



# Sample Space

The set of all possible outcomes of an experiment is called the **sample space** of the experiment, and it is denoted by  $\Omega$ .

# Sample Space

The set of all possible outcomes of an experiment is called the **sample space** of the experiment, and it is denoted by  $\Omega$ .

- Experiment of tossing a coin:  $\Omega = \{H, T\}$

# Sample Space

The set of all possible outcomes of an experiment is called the **sample space** of the experiment, and it is denoted by  $\Omega$ .

- Experiment of tossing a coin:  $\Omega = \{H, T\}$
- Experiment of rolling a die:  $\Omega = \{1, 2, 3, \dots, 6\}$

# Sample Space

The set of all possible outcomes of an experiment is called the **sample space** of the experiment, and it is denoted by  $\Omega$ .

- Experiment of tossing a coin:  $\Omega = \{H, T\}$
- Experiment of rolling a die:  $\Omega = \{1, 2, 3, \dots, 6\}$
- Experiment of measuring the temperature:  $\Omega = [60, 120]$

# Sample Space

The set of all possible outcomes of an experiment is called the **sample space** of the experiment, and it is denoted by  $\Omega$ .

- Experiment of tossing a coin:  $\Omega = \{H, T\}$
- Experiment of rolling a die:  $\Omega = \{1, 2, 3, \dots, 6\}$
- Experiment of measuring the temperature:  $\Omega = [60, 120]$
- **Joint experiment and sample space**: Consider two experiments with sample spaces  $\Omega_1$  and  $\Omega_2$ , respectively. Then, we can consider a joint experiment with sample space  $\Omega = \Omega_1 \times \Omega_2$

# Sample Space

The set of all possible outcomes of an experiment is called the **sample space** of the experiment, and it is denoted by  $\Omega$ .

- Experiment of tossing a coin:  $\Omega = \{H, T\}$
- Experiment of rolling a die:  $\Omega = \{1, 2, 3, \dots, 6\}$
- Experiment of measuring the temperature:  $\Omega = [60, 120]$
- **Joint experiment and sample space:** Consider two experiments with sample spaces  $\Omega_1$  and  $\Omega_2$ , respectively. Then, we can consider a joint experiment with sample space  $\Omega = \Omega_1 \times \Omega_2$
- Experiment of rolling a die twice:  
 $\Omega = \{1, 2, 3, \dots, 6\} \times \{1, 2, 3, \dots, 6\} = \{(1, 1), (1, 2), \dots, (6, 6)\}$

# Sample Space

The set of all possible outcomes of an experiment is called the **sample space** of the experiment, and it is denoted by  $\Omega$ .

- Experiment of tossing a coin:  $\Omega = \{H, T\}$
- Experiment of rolling a die:  $\Omega = \{1, 2, 3, \dots, 6\}$
- Experiment of measuring the temperature:  $\Omega = [60, 120]$
- **Joint experiment and sample space:** Consider two experiments with sample spaces  $\Omega_1$  and  $\Omega_2$ , respectively. Then, we can consider a joint experiment with sample space  $\Omega = \Omega_1 \times \Omega_2$
- Experiment of rolling a die twice:  
 $\Omega = \{1, 2, 3, \dots, 6\} \times \{1, 2, 3, \dots, 6\} = \{(1, 1), (1, 2), \dots, (6, 6)\}$
- Experiment of flipping a coin and rolling a die:  $\Omega = ?$

# Event

An **event** is any subset of the sample space.



# Event

An **event** is any subset of the sample space.

- Experiment of tossing a coin, event of getting tails:  $E = \{T\}$

# Event

An **event** is any subset of the sample space.

- Experiment of tossing a coin, event of getting tails:  $E = \{T\}$
- Experiment of rolling a die, event of getting an odd number:  $E = \{1, 3, 5\}$

# Event

An **event** is any subset of the sample space.

- Experiment of tossing a coin, event of getting tails:  $E = \{T\}$
- Experiment of rolling a die, event of getting an odd number:  $E = \{1, 3, 5\}$
- Experiment of measuring the temperature, event of temperature within 80 and 90:  
 $E = [80, 90]$

# Event

An **event** is any subset of the sample space.

- Experiment of tossing a coin, event of getting tails:  $E = \{T\}$
- Experiment of rolling a die, event of getting an odd number:  $E = \{1, 3, 5\}$
- Experiment of measuring the temperature, event of temperature within 80 and 90:  $E = [80, 90]$
- **Joint events:** Consider two experiments with sample spaces  $\Omega_1$  and  $\Omega_2$ , respectively. Also, consider two events  $E_1 \subset \Omega_1$  and  $E_2 \subset \Omega_2$ . then, we can consider a joint event  $E = E_1 \times E_2 \subset \Omega_1 \times \Omega_2$

# Event

An **event** is any subset of the sample space.

- Experiment of tossing a coin, event of getting tails:  $E = \{T\}$
- Experiment of rolling a die, event of getting an odd number:  $E = \{1, 3, 5\}$
- Experiment of measuring the temperature, event of temperature within 80 and 90:  $E = [80, 90]$
- **Joint events:** Consider two experiments with sample spaces  $\Omega_1$  and  $\Omega_2$ , respectively. Also, consider two events  $E_1 \subset \Omega_1$  and  $E_2 \subset \Omega_2$ . then, we can consider a joint event  $E = E_1 \times E_2 \subset \Omega_1 \times \Omega_2$
- Experiment of rolling a die twice, event of getting the first number odd and the second number even:  $E = \{1, 3, 5\} \times \{2, 4, 6\} = \{(1, 2), (1, 4), \dots, (5, 6)\}$

# Event

An **event** is any subset of the sample space.

- Experiment of tossing a coin, event of getting tails:  $E = \{T\}$
- Experiment of rolling a die, event of getting an odd number:  $E = \{1, 3, 5\}$
- Experiment of measuring the temperature, event of temperature within 80 and 90:  $E = [80, 90]$
- **Joint events:** Consider two experiments with sample spaces  $\Omega_1$  and  $\Omega_2$ , respectively. Also, consider two events  $E_1 \subset \Omega_1$  and  $E_2 \subset \Omega_2$ . then, we can consider a joint event  $E = E_1 \times E_2 \subset \Omega_1 \times \Omega_2$
- Experiment of rolling a die twice, event of getting the first number odd and the second number even:  $E = \{1, 3, 5\} \times \{2, 4, 6\} = \{(1, 2), (1, 4), \dots, (5, 6)\}$
- Experiment of flipping a coin and rolling a die, event of getting a tails and a prime number:  $E = ?$

## Events: Example

- Consider the experiment of rolling a die twice. What is the sample space?

## Events: Example

- Consider the experiment of rolling a die twice. What is the sample space?

	$E_1$		$E_2$		
$\Omega =$	(1,2)	(1,3)	(1,4)	(1,5)	(1,6)
	(2,2)	(2,3)	(2,4)	(2,5)	(2,6)
	(3,2)	(3,3)	(3,4)	(3,5)	(3,6)
	(4,2)	(4,3)	(4,4)	(4,5)	(4,6)
	(5,2)	(5,3)	(5,4)	(5,5)	(5,6)
	(6,2)	(6,3)	(6,4)	(6,5)	(6,6)



## Events: Example

- Consider the experiment of rolling a die twice. What is the sample space?

	$E_1$		$E_2$		
$\Omega = \left\{ \begin{array}{l} (1,1) \\ (2,1) \\ (3,1) \\ (4,1) \\ (5,1) \\ (6,1) \end{array} \right.$	(1,2)	(1,3)	(1,4)	(1,5)	(1,6)
	(2,2)	(2,3)	(2,4)	(2,5)	(2,6)
	(3,2)	(3,3)	(3,4)	(3,5)	(3,6)
	(4,2)	(4,3)	(4,4)	(4,5)	(4,6)
	(5,2)	(5,3)	(5,4)	(5,5)	(5,6)
	(6,2)	(6,3)	(6,4)	(6,5)	(6,6)

- What does event  $E_1$  represent?

## Events: Example

- Consider the experiment of rolling a die twice. What is the sample space?

	$E_1$		$E_2$		
$\Omega = \left\{ \begin{array}{l} (1,1) \\ (2,1) \\ (3,1) \\ (4,1) \\ (5,1) \\ (6,1) \end{array} \right.$	(1,2)	(1,3)	(1,4)	(1,5)	(1,6)
	(2,2)	(2,3)	(2,4)	(2,5)	(2,6)
	(3,2)	(3,3)	(3,4)	(3,5)	(3,6)
	(4,2)	(4,3)	(4,4)	(4,5)	(4,6)
	(5,2)	(5,3)	(5,4)	(5,5)	(5,6)
	(6,2)	(6,3)	(6,4)	(6,5)	(6,6)

- What does event  $E_1$  represent?
- What is  $E_1 \cup E_2$ ?

## Events: Example

- Consider the experiment of rolling a die twice. What is the sample space?

	$E_1$		$E_2$		
$\Omega = \left\{ \begin{array}{l} (1,1) \\ (2,1) \\ (3,1) \\ (4,1) \\ (5,1) \\ (6,1) \end{array} \right.$	(1,2)	(1,3)	(1,4)	(1,5)	(1,6)
	(2,2)	(2,3)	(2,4)	(2,5)	(2,6)
	(3,2)	(3,3)	(3,4)	(3,5)	(3,6)
	(4,2)	(4,3)	(4,4)	(4,5)	(4,6)
	(5,2)	(5,3)	(5,4)	(5,5)	(5,6)
	(6,2)	(6,3)	(6,4)	(6,5)	(6,6)

- What does event  $E_1$  represent?
- What is  $E_1 \cup E_2$ ?
- What is  $E_1^c$ ?

# Disjoint Events and Partition

Two events  $E_1$  and  $E_2$  are **disjoint (a.k.a. mutually exclusive)** events if  $E_1 \cap E_2 = \emptyset$ .

# Disjoint Events and Partition



Two events  $E_1$  and  $E_2$  are **disjoint (a.k.a. mutually exclusive)** events if  $E_1 \cap E_2 = \emptyset$ .

If  $E_1, E_2, \dots, E_n$  are events such that  $E_i \cap E_j = \emptyset$  for all  $i, j, i \neq j$ , and  $\cup_{i=1}^n E_i = F$ , then we say that events  $E_1, E_2, \dots, E_n$  are a **partition** of the event  $F$ .

# Disjoint Events and Partition

Two events  $E_1$  and  $E_2$  are **disjoint (a.k.a. mutually exclusive)** events if  $E_1 \cap E_2 = \emptyset$ .


If  $E_1, E_2, \dots, E_n$  are events such that  $E_i \cap E_j = \emptyset$  for all  $i, j, i \neq j$ , and  $\cup_{i=1}^n E_i = F$ , then we say that events  $E_1, E_2, \dots, E_n$  are a **partition** of the event  $F$ .



# Disjoint Events and Partition

Two events  $E_1$  and  $E_2$  are **disjoint (a.k.a. mutually exclusive)** events if  $E_1 \cap E_2 = \emptyset$ .

If  $E_1, E_2, \dots, E_n$  are events such that  $E_i \cap E_j = \emptyset$  for all  $i, j, i \neq j$ , and  $\cup_{i=1}^n E_i = F$ , then we say that events  $E_1, E_2, \dots, E_n$  are a **partition** of the event  $F$ .



# Set Difference

- For any two sets  $A$  and  $B$ ,  $A - B$  is defined as the set consisting of those elements of  $A$  that are not in  $B$ ,


$$A - B = \{x | x \in A \text{ and } x \notin B\}$$



# Set Difference

- For any two sets  $A$  and  $B$ ,  $A - B$  is defined as the set consisting of those elements of  $A$  that are not in  $B$ ,


$$A - B = \{x | x \in A \text{ and } x \notin B\}$$



# Set Difference

- For any two sets  $A$  and  $B$ ,  $A - B$  is defined as the set consisting of those elements of  $A$  that are not in  $B$ ,

$$A - B = \{x | x \in A \text{ and } x \notin B\}$$




- By definition, we can write that  $A - B = A \cap B^c$

# Set Difference

- For any two sets  $A$  and  $B$ ,  $A - B$  is defined as the set consisting of those elements of  $A$  that are not in  $B$ ,

$$A - B = \{x | x \in A \text{ and } x \notin B\}$$




- By definition, we can write that  $A - B = A \cap B^c$
- $B$  and  $A - B$  are disjoint sets

# Set Difference

- For any two sets  $A$  and  $B$ ,  $A - B$  is defined as the set consisting of those elements of  $A$  that are not in  $B$ ,

$$A - B = \{x | x \in A \text{ and } x \notin B\}$$



- By definition, we can write that  $A - B = A \cap B^c$
- $B$  and  $A - B$  are disjoint sets
- $B$  and  $A - B$  form a partition of the set  $A \cup B$

# Axioms of Probability

- A **probability law** assigns to every event  $A$  a number  $P(A)$ , called the *probability of event  $A$* , such that the following **axioms** are satisfied.

# Axioms of Probability

- A **probability law** assigns to every event  $A$  a number  $P(A)$ , called the *probability of event  $A$* , such that the following **axioms** are satisfied.

1 (Nonnegativity)  $P(A) \geq 0$ , for every event  $A$ .

# Axioms of Probability

- A **probability law** assigns to every event  $A$  a number  $P(A)$ , called the *probability of event  $A$* , such that the following **axioms** are satisfied.

① **(Nonnegativity)**  $P(A) \geq 0$ , for every event  $A$ .

② **(Normalization)** The probability of the sample space  $\Omega$  is equal to one,  
$$P(\Omega) = 1.$$

# Axioms of Probability

- A **probability law** assigns to every event  $A$  a number  $P(A)$ , called the *probability of event  $A$* , such that the following **axioms** are satisfied.

① **(Nonnegativity)**  $P(A) \geq 0$ , for every event  $A$ .

② **(Normalization)** The probability of the sample space  $\Omega$  is equal to one,  
$$P(\Omega) = 1.$$

③ **(Countable Additivity)** If  $A$  and  $B$  are disjoint events,  $A \cap B = \emptyset$ , then the probability of their union satisfies

$$P(A \cup B) = P(A) + P(B).$$




# Results from the Axioms

If  $A \subset B$ , then  $P(A) \leq P(B)$ .

# Results from the Axioms

If  $A \subset B$ , then  $P(A) \leq P(B)$ .



# Results from the Axioms

If  $A \subset B$ , then  $P(A) \leq P(B)$ .

## Results from the Axioms

If  $A \subset B$ , then  $P(A) \leq P(B)$ .

**Proof:** Since  $A \subset B$ , we have  $B = A \cup (B - A)$ . Noting that  $A$  and  $B - A$  are disjoint sets, we get

$$P(B) = P(A) + P(B - A) \geq P(A),$$


where the inequality follows from the nonnegativity of probability laws. □

## Results from the Axioms

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

## Results from the Axioms

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$



# Results from the Axioms

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

## Results from the Axioms

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

**Proof:** We can write  $A \cup B = A \cup (B - A)$ .



## Results from the Axioms

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

**Proof:** We can write  $A \cup B = A \cup (B - A)$ . Using the third axiom of probability on the disjoint sets  $A$  and  $(B - A)$ , we can write

$$P(A \cup B) = P(A) + P(B - A).$$

## Results from the Axioms

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

**Proof:** We can write  $A \cup B = A \cup (B - A)$ . Using the third axiom of probability on the disjoint sets  $A$  and  $(B - A)$ , we can write

$$P(A \cup B) = P(A) + P(B - A).$$

Similarly, we can write  $B = (A \cap B) \cup (B - A)$ .

## Results from the Axioms

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

**Proof:** We can write  $A \cup B = A \cup (B - A)$ . Using the third axiom of probability on the disjoint sets  $A$  and  $(B - A)$ , we can write

$$P(A \cup B) = P(A) + P(B - A).$$

Similarly, we can write  $B = (A \cap B) \cup (B - A)$ . Then, applying the third axiom, we obtain

$$P(B) = P(A \cap B) + P(B - A) \implies P(B - A) = P(B) - P(A \cap B).$$

## Results from the Axioms

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

**Proof:** We can write  $A \cup B = A \cup (B - A)$ . Using the third axiom of probability on the disjoint sets  $A$  and  $(B - A)$ , we can write

$$P(A \cup B) = P(A) + P(B - A).$$

Similarly, we can write  $B = (A \cap B) \cup (B - A)$ . Then, applying the third axiom, we obtain

$$P(B) = P(A \cap B) + P(B - A) \implies P(B - A) = P(B) - P(A \cap B).$$


Combining these two equations yields the desired result. □

# Results from the Axioms

$$P(A \cup B) \leq P(A) + P(B).$$

# Results from the Axioms

$$P(A \cup B) \leq P(A) + P(B).$$



## Results from the Axioms

$$P(A \cup B) \leq P(A) + P(B).$$

# Results from the Axioms

$$P(A \cup B) \leq P(A) + P(B).$$

## Union Bound

Let  $A_1, A_2, \dots, A_n$  be a collection of events, then

$$P\left(\bigcup_{k=1}^n A_k\right) \leq \sum_{k=1}^n P(A_k).$$



# Examples

- Show that  $P(A^c) = 1 - P(A)$

# Examples

- Show that  $P(A^c) = 1 - P(A)$ 
  - ▶ Note that  $\Omega = A \cup A^c$ . Using the third axiom, we get  $P(\Omega) = P(A) + P(A^c)$ . From the second axiom,  $P(\Omega) = 1$ . Combining these, we get the result.

# Examples

- Show that  $P(A^c) = 1 - P(A)$ 
  - ▶ Note that  $\Omega = A \cup A^c$ . Using the third axiom, we get  $P(\Omega) = P(A) + P(A^c)$ .  
From the second axiom,  $P(\Omega) = 1$ . Combining these, we get the result.
- Amy is taking two books along on her spring break. With probability 0.5, she will like the first book; with probability 0.4, she will like the second book; and with probability 0.3, she will like both books. What is the probability that she likes neither book?

# Examples

- Show that  $P(A^c) = 1 - P(A)$ 
  - ▶ Note that  $\Omega = A \cup A^c$ . Using the third axiom, we get  $P(\Omega) = P(A) + P(A^c)$ .  
From the second axiom,  $P(\Omega) = 1$ . Combining these, we get the result.
- Amy is taking two books along on her spring break. With probability 0.5, she will like the first book; with probability 0.4, she will like the second book; and with probability 0.3, she will like both books. What is the probability that she likes neither book?
  - ▶ Let  $B_i$  denote the event that  $K$  likes book  $i, i = 1, 2$ .

# Examples

- Show that  $P(A^c) = 1 - P(A)$ 
  - ▶ Note that  $\Omega = A \cup A^c$ . Using the third axiom, we get  $P(\Omega) = P(A) + P(A^c)$ .  
From the second axiom,  $P(\Omega) = 1$ . Combining these, we get the result.
- Amy is taking two books along on her spring break. With probability 0.5, she will like the first book; with probability 0.4, she will like the second book; and with probability 0.3, she will like both books. What is the probability that she likes neither book?
  - ▶ Let  $B_i$  denote the event that  $K$  likes book  $i, i = 1, 2$ . Then the probability that she likes at least one of the books is

# Examples

- Show that  $P(A^c) = 1 - P(A)$ 
  - ▶ Note that  $\Omega = A \cup A^c$ . Using the third axiom, we get  $P(\Omega) = P(A) + P(A^c)$ .  
From the second axiom,  $P(\Omega) = 1$ . Combining these, we get the result.
- Amy is taking two books along on her spring break. With probability 0.5, she will like the first book; with probability 0.4, she will like the second book; and with probability 0.3, she will like both books. What is the probability that she likes neither book?
  - ▶ Let  $B_i$  denote the event that  $K$  likes book  $i, i = 1, 2$ . Then the probability that she likes at least one of the books is

$$P(B_1 \cup B_2) = P(B_1) + P(B_2) - P(B_1 \cap B_2) = 0.5 + 0.4 - 0.3 = 0.6$$

# Examples

- Show that  $P(A^c) = 1 - P(A)$ 
  - ▶ Note that  $\Omega = A \cup A^c$ . Using the third axiom, we get  $P(\Omega) = P(A) + P(A^c)$ . From the second axiom,  $P(\Omega) = 1$ . Combining these, we get the result.
- Amy is taking two books along on her spring break. With probability 0.5, she will like the first book; with probability 0.4, she will like the second book; and with probability 0.3, she will like both books. What is the probability that she likes neither book?
  - ▶ Let  $B_i$  denote the event that  $K$  likes book  $i, i = 1, 2$ . Then the probability that she likes at least one of the books is

$$P(B_1 \cup B_2) = P(B_1) + P(B_2) - P(B_1 \cap B_2) = 0.5 + 0.4 - 0.3 = 0.6$$

Probability that he likes neither book is

$$P(B_1^c \cap B_2^c) = P((B_1 \cup B_2)^c) = 1 - P((B_1 \cup B_2)) = 1 - 0.6 = 0.4$$

# Examples

- Consider two events  $A, B$  that are mutually exclusive. Let  $P(A) = 0.3$  and  $P(B) = 0.5$ . What is the probability that
  - 1 Either  $A$  or  $B$  occurs?



# Examples

- Consider two events  $A, B$  that are mutually exclusive. Let  $P(A) = 0.3$  and  $P(B) = 0.5$ . What is the probability that
  - 1 Either  $A$  or  $B$  occurs?

$$P(A \cup B) = P(A) + P(B) = 0.3 + 0.5 = 0.8$$

# Examples

- Consider two events  $A, B$  that are mutually exclusive. Let  $P(A) = 0.3$  and  $P(B) = 0.5$ . What is the probability that

- 1 Either  $A$  or  $B$  occurs?

$$P(A \cup B) = P(A) + P(B) = 0.3 + 0.5 = 0.8$$

- 2  $A$  occurs but  $B$  does not?

# Examples

- Consider two events  $A, B$  that are mutually exclusive. Let  $P(A) = 0.3$  and  $P(B) = 0.5$ . What is the probability that

- 1 Either  $A$  or  $B$  occurs?

$$P(A \cup B) = P(A) + P(B) = 0.3 + 0.5 = 0.8$$

- 2  $A$  occurs but  $B$  does not?

$$P(A \cap B^c) = P(A) = 0.3$$

# Examples

- Consider two events  $A, B$  that are mutually exclusive. Let  $P(A) = 0.3$  and  $P(B) = 0.5$ . What is the probability that

- 1 Either  $A$  or  $B$  occurs?

$$P(A \cup B) = P(A) + P(B) = 0.3 + 0.5 = 0.8$$

- 2  $A$  occurs but  $B$  does not?

$$P(A \cap B^c) = P(A) = 0.3$$

- 3 Both  $A$  and  $B$  occur?

# Examples

- Consider two events  $A, B$  that are mutually exclusive. Let  $P(A) = 0.3$  and  $P(B) = 0.5$ . What is the probability that

- 1 Either  $A$  or  $B$  occurs?

$$P(A \cup B) = P(A) + P(B) = 0.3 + 0.5 = 0.8$$

- 2  $A$  occurs but  $B$  does not?

$$P(A \cap B^c) = P(A) = 0.3$$

- 3 Both  $A$  and  $B$  occur?

$$P(A \cap B) = P(\emptyset) = 0$$

# Finite Sample Spaces

$\Omega$  is **finite sample space** if it contains a finite number of elements

# Finite Sample Spaces

$\Omega$  is **finite sample space** if it contains a finite number of elements

- **(Example):** The rolling of a fair die is an experiment with a finite number of equally likely outcomes.

# Finite Sample Spaces

$\Omega$  is **finite sample space** if it contains a finite number of elements

- **(Example):** The rolling of a fair die is an experiment with a finite number of equally likely outcomes. The probability of observing any of the faces labeled one through six is therefore equal to  $1/6$ .



# Finite Sample Spaces

$\Omega$  is **finite sample space** if it contains a finite number of elements

- **(Example):** The rolling of a fair die is an experiment with a finite number of equally likely outcomes. The probability of observing any of the faces labeled one through six is therefore equal to  $1/6$ . The probability of any event can be computed by counting the number of distinct outcomes included in the event.

# Finite Sample Spaces

$\Omega$  is **finite sample space** if it contains a finite number of elements

- **(Example):** The rolling of a fair die is an experiment with a finite number of equally likely outcomes. The probability of observing any of the faces labeled one through six is therefore equal to  $1/6$ . The probability of any event can be computed by counting the number of distinct outcomes included in the event. For example, the probability of rolling a prime number is

# Finite Sample Spaces

$\Omega$  is **finite sample space** if it contains a finite number of elements

- **(Example):** The rolling of a fair die is an experiment with a finite number of equally likely outcomes. The probability of observing any of the faces labeled one through six is therefore equal to  $1/6$ . The probability of any event can be computed by counting the number of distinct outcomes included in the event. For example, the probability of rolling a prime number is

$$P(\{2, 3, 5\}) = P(2) + P(3) + P(5) = \frac{3}{6}.$$

# Finite Sample Spaces

- **(Example):** Consider the experiment of rolling a pair of 4-sided fair dice.

# Finite Sample Spaces

- **(Example):** Consider the experiment of rolling a pair of 4-sided fair dice. We interpret 'fair' to mean that each of the sixteen possible outcomes (ordered pairs  $(i, j)$ , with  $i, j = 1, 2, 3, 4$ ), has the same probability of  $1/16$ .

# Finite Sample Spaces

- **(Example):** Consider the experiment of rolling a pair of 4-sided fair dice. We interpret 'fair' to mean that each of the sixteen possible outcomes (ordered pairs  $(i, j)$ , with  $i, j = 1, 2, 3, 4$ ), has the same probability of  $1/16$ . To calculate the probability of an event, we must count the number of elements of event and divide by 16 (the total number of possible outcomes).

# Finite Sample Spaces

- **(Example):** Consider the experiment of rolling a pair of 4-sided fair dice. We interpret 'fair' to mean that each of the sixteen possible outcomes (ordered pairs  $(i, j)$ , with  $i, j = 1, 2, 3, 4$ ), has the same probability of  $1/16$ . To calculate the probability of an event, we must count the number of elements of event and divide by 16 (the total number of possible outcomes).
  - 1  $P(\text{the sum of the rolls is even})$

# Finite Sample Spaces

- **(Example):** Consider the experiment of rolling a pair of 4-sided fair dice. We interpret 'fair' to mean that each of the sixteen possible outcomes (ordered pairs  $(i, j)$ , with  $i, j = 1, 2, 3, 4$ ), has the same probability of  $1/16$ . To calculate the probability of an event, we must count the number of elements of event and divide by 16 (the total number of possible outcomes).
  - 1  $P(\text{the sum of the rolls is even})$
  - 2  $P(\text{the sum of the rolls is odd})$



# Finite Sample Spaces

- **(Example):** Consider the experiment of rolling a pair of 4-sided fair dice. We interpret 'fair' to mean that each of the sixteen possible outcomes (ordered pairs  $(i, j)$ , with  $i, j = 1, 2, 3, 4$ ), has the same probability of  $1/16$ . To calculate the probability of an event, we must count the number of elements of event and divide by 16 (the total number of possible outcomes).
  - 1  $P(\text{the sum of the rolls is even})$
  - 2  $P(\text{the sum of the rolls is odd})$
  - 3  $P(\text{the first roll is equal to the second})$

# Finite Sample Spaces

- **(Example):** Consider the experiment of rolling a pair of 4-sided fair dice. We interpret 'fair' to mean that each of the sixteen possible outcomes (ordered pairs  $(i, j)$ , with  $i, j = 1, 2, 3, 4$ ), has the same probability of  $1/16$ . To calculate the probability of an event, we must count the number of elements of event and divide by 16 (the total number of possible outcomes).
  - 1  $P(\text{the sum of the rolls is even})$
  - 2  $P(\text{the sum of the rolls is odd})$
  - 3  $P(\text{the first roll is equal to the second})$
  - 4  $P(\text{the first roll is larger than the second})$

# Finite Sample Spaces

- **(Example):** Consider the experiment of rolling a pair of 4-sided fair dice. We interpret 'fair' to mean that each of the sixteen possible outcomes (ordered pairs  $(i, j)$ , with  $i, j = 1, 2, 3, 4$ ), has the same probability of  $1/16$ . To calculate the probability of an event, we must count the number of elements of event and divide by 16 (the total number of possible outcomes).
  - 1  $P(\text{the sum of the rolls is even})$
  - 2  $P(\text{the sum of the rolls is odd})$
  - 3  $P(\text{the first roll is equal to the second})$
  - 4  $P(\text{the first roll is larger than the second})$
  - 5  $P(\text{at least one roll is equal to 4})$

# Countably Infinite Sample Spaces

$\Omega$  is **countably infinite sample space** if it contains a countably infinite number of elements

# Countably Infinite Sample Spaces

- **(Example):** Suppose that a fair coin is tossed repetitively until heads is observed. What is the probability the probability of the number of coin tosses being even?

# Countably Infinite Sample Spaces

- **(Example):** Suppose that a fair coin is tossed repetitively until heads is observed. What is the probability the probability of the number of coin tosses being even?

Sample space for this experiment is  $\Omega = \{1, 2, \dots\}$ , a countably infinite set.

# Countably Infinite Sample Spaces

- **(Example):** Suppose that a fair coin is tossed repetitively until heads is observed. What is the probability the probability of the number of coin tosses being even?

Sample space for this experiment is  $\Omega = \{1, 2, \dots\}$ , a countably infinite set.

The probability of observing heads on the first trial is  $1/2$

# Countably Infinite Sample Spaces

- **(Example):** Suppose that a fair coin is tossed repetitively until heads is observed. What is the probability the probability of the number of coin tosses being even?

Sample space for this experiment is  $\Omega = \{1, 2, \dots\}$ , a countably infinite set.

The probability of observing heads on the first trial is  $1/2$

The probability of observing heads for the first time on trial  $k$  is  $2^{-k}$ .



# Countably Infinite Sample Spaces

- **(Example):** Suppose that a fair coin is tossed repetitively until heads is observed. What is the probability the probability of the number of coin tosses being even?

Sample space for this experiment is  $\Omega = \{1, 2, \dots\}$ , a countably infinite set.

The probability of observing heads on the first trial is  $1/2$

The probability of observing heads for the first time on trial  $k$  is  $2^{-k}$ .

The probability of the entire sample space is therefore equal to

$$P(\Omega) = \sum_{k=1}^{\infty} P(k) = \sum_{k=1}^{\infty} \frac{1}{2^k} = 1, \text{ as expected.}$$

# Countably Infinite Sample Spaces

- **(Example):** Suppose that a fair coin is tossed repetitively until heads is observed. What is the probability the probability of the number of coin tosses being even?

Sample space for this experiment is  $\Omega = \{1, 2, \dots\}$ , a countably infinite set.

The probability of observing heads on the first trial is  $1/2$

The probability of observing heads for the first time on trial  $k$  is  $2^{-k}$ .

The probability of the entire sample space is therefore equal to

$$P(\Omega) = \sum_{k=1}^{\infty} P(k) = \sum_{k=1}^{\infty} \frac{1}{2^k} = 1, \text{ as expected.}$$

The probability of the number of coin tosses being even can be computed as

$$P(\{2, 4, 6, \dots\}) = \sum_{k=1}^{\infty} P(2k) = \sum_{k=1}^{\infty} \frac{1}{2^{2k}} = \frac{1}{4} \frac{1}{(1 - \frac{1}{4})} = \frac{1}{3}.$$

# Uncountably Infinite Sample Spaces

$\Omega$  is **uncountably infinite sample space** if it contains uncountably infinite number of elements

- Consider  $\Omega = [0, 1]$ . Choose an element from this interval, uniformly at random. What is the probability that:

# Uncountably Infinite Sample Spaces

$\Omega$  is **uncountably infinite sample space** if it contains uncountably infinite number of elements

- Consider  $\Omega = [0, 1]$ . Choose an element from this interval, uniformly at random. What is the probability that:
  - ▶ the chosen element lies in the interval  $(0, 0.25)$ ?

# Uncountably Infinite Sample Spaces

$\Omega$  is **uncountably infinite sample space** if it contains uncountably infinite number of elements

- Consider  $\Omega = [0, 1]$ . Choose an element from this interval, uniformly at random. What is the probability that:
  - ▶ the chosen element lies in the interval  $(0, 0.25)$ ?
  - ▶ the chosen element lies in the interval  $(0.75, 1)$ ?

# Uncountably Infinite Sample Spaces

$\Omega$  is **uncountably infinite sample space** if it contains uncountably infinite number of elements

- Consider  $\Omega = [0, 1]$ . Choose an element from this interval, uniformly at random. What is the probability that:
  - ▶ the chosen element lies in the interval  $(0, 0.25)$ ?
  - ▶ the chosen element lies in the interval  $(0.75, 1)$ ?
  - ▶ the chosen element lies in the interval  $(a, b)$ , for  $0 \leq a < b \leq 1$ ?

# Uncountably Infinite Sample Spaces

$\Omega$  is **uncountably infinite sample space** if it contains uncountably infinite number of elements

- Consider  $\Omega = [0, 1]$ . Choose an element from this interval, uniformly at random. What is the probability that:
  - ▶ the chosen element lies in the interval  $(0, 0.25)$ ?
  - ▶ the chosen element lies in the interval  $(0.75, 1)$ ?
  - ▶ the chosen element lies in the interval  $(a, b)$ , for  $0 \leq a < b \leq 1$ ?
  - ▶ the chosen element is 0.55?

# Uncountably Infinite Sample Spaces

$\Omega$  is **uncountably infinite sample space** if it contains uncountably infinite number of elements

- Consider  $\Omega = [0, 2]$ . Choose an element from this interval, uniformly at random. What is the probability that:



# Uncountably Infinite Sample Spaces

$\Omega$  is **uncountably infinite sample space** if it contains uncountably infinite number of elements

- Consider  $\Omega = [0, 2]$ . Choose an element from this interval, uniformly at random. What is the probability that:
  - ▶ the chosen element lies in the interval  $(0, 0.25)$ ?

# Uncountably Infinite Sample Spaces

$\Omega$  is **uncountably infinite sample space** if it contains uncountably infinite number of elements

- Consider  $\Omega = [0, 2]$ . Choose an element from this interval, uniformly at random. What is the probability that:
  - ▶ the chosen element lies in the interval  $(0, 0.25)$ ?
  - ▶ the chosen element lies in the interval  $(0.75, 1)$ ?

# Uncountably Infinite Sample Spaces

$\Omega$  is **uncountably infinite sample space** if it contains uncountably infinite number of elements

- Consider  $\Omega = [0, 2]$ . Choose an element from this interval, uniformly at random. What is the probability that:
  - ▶ the chosen element lies in the interval  $(0, 0.25)$ ?
  - ▶ the chosen element lies in the interval  $(0.75, 1)$ ?
  - ▶ the chosen element lies in the interval  $(a, b)$ , for  $0 \leq a < b \leq 1$ ?

# Uncountably Infinite Sample Spaces

$\Omega$  is **uncountably infinite sample space** if it contains uncountably infinite number of elements

- Consider  $\Omega = [0, 2]$ . Choose an element from this interval, uniformly at random. What is the probability that:
  - ▶ the chosen element lies in the interval  $(0, 0.25)$ ?
  - ▶ the chosen element lies in the interval  $(0.75, 1)$ ?
  - ▶ the chosen element lies in the interval  $(a, b)$ , for  $0 \leq a < b \leq 1$ ?
  - ▶ the chosen element is 0.55?

# Uncountably Infinite Sample Spaces

- R and J plans to meet a coffee shop at a given time. Each will arrive at the meeting place with a delay between 0 and 1 hour, with all pairs of delays being equally likely. The first to arrive will wait for 15 minutes and will leave if the other has not yet arrived. What is the probability that they will meet?

# Uncountably Infinite Sample Spaces

- R and J plans to meet a coffee shop at a given time. Each will arrive at the meeting place with a delay between 0 and 1 hour, with all pairs of delays being equally likely. The first to arrive will wait for 15 minutes and will leave if the other has not yet arrived. What is the probability that they will meet?

Let us use as sample space the square  $\Omega = [0, 1] \times [0, 1]$ , whose elements are the possible pairs of delays for the two of them.

# Uncountably Infinite Sample Spaces

- R and J plans to meet a coffee shop at a given time. Each will arrive at the meeting place with a delay between 0 and 1 hour, with all pairs of delays being equally likely. The first to arrive will wait for 15 minutes and will leave if the other has not yet arrived. What is the probability that they will meet?

Let us use as sample space the square  $\Omega = [0, 1] \times [0, 1]$ , whose elements are the possible pairs of delays for the two of them.

Our interpretation of “equally likely” pairs of delays is to let the probability of a subset of  $\Omega$  be equal to its area divided by total area.

# Uncountably Infinite Sample Spaces

- R and J plans to meet a coffee shop at a given time. Each will arrive at the meeting place with a delay between 0 and 1 hour, with all pairs of delays being equally likely. The first to arrive will wait for 15 minutes and will leave if the other has not yet arrived. What is the probability that they will meet?

Let us use as sample space the square  $\Omega = [0, 1] \times [0, 1]$ , whose elements are the possible pairs of delays for the two of them.

Our interpretation of “equally likely” pairs of delays is to let the probability of a subset of  $\Omega$  be equal to its area divided by total area.

This probability law satisfies the three probability axioms.



# Uncountably Infinite Sample Spaces

- R and J plans to meet a coffee shop at a given time. Each will arrive at the meeting place with a delay between 0 and 1 hour, with all pairs of delays being equally likely. The first to arrive will wait for 15 minutes and will leave if the other has not yet arrived. What is the probability that they will meet?

# Uncountably Infinite Sample Spaces

- R and J plans to meet a coffee shop at a given time. Each will arrive at the meeting place with a delay between 0 and 1 hour, with all pairs of delays being equally likely. The first to arrive will wait for 15 minutes and will leave if the other has not yet arrived. What is the probability that they will meet?

