

UNIVERSITY OF LJUBLJANA  
FACULTY OF MATHEMATICS AND PHYSICS

Financial mathematics – 1st cycle

Anej Rozman, Tanja Luštrek  
**Rich-Neighbor Edge Colorings**

Term Paper in Finance Lab  
Short Presentation

Advisers: Assistant Professor Janoš Vidali, Professor Riste  
Škrekovski

Ljubljana, 2023

## 1. INTRODUCTION


In this paper we set out to analyse an open conjecture in a modern graph theory problem known as rich-neighbor edge coloring.

**Definition 1.1.** In an edge coloring, an edge  $e$  is called *rich* if all edges adjacent to  $e$  have different colors. An edge coloring is called a *rich-neighbor edge coloring* if every edge is adjacent to some rich edge.

**Definition 1.2.**  $X'_{rn}(G)$  denotes the smallest number of colors for which there exists a rich-neighbor edge coloring.

**Conjecture 1.3.** For every graph  $G$  of maximum degree  $\Delta$ ,  $X'_{rn}(G) \leq 2\Delta - 1$  holds.

**Example 1.4.** Let's take a look at the Petersen graph and an example of a rich-neighbor edge coloring.



We can see that for the Petersen graph (which is 3-regular)  $X'_{rn} = 5 \leq 5$ . ◇

## 2. PLAN

Our assignment is to create an algorithm that proves the conjecture for graphs of maximum degree 4 (So it finds a rich-neighbor edge coloring for every graph, for example for all graphs with 5 vertices and a maximum degree of 5), and to make a random search algorithm for graphs of maximum degree  $\geq 5$ .