2026 3

> | v1.7 | 2026-03-18 | wenping.wang@remegen.com
>
> - *v1.0 2026-03-08 — Q1-Q3 C016 OS*
> - *v1.1 2026-03-09 — Q4 DECIDING / vs HOPE-03*
> - *v1.2 2026-03-09 — Q4 6+45=51*
> - *v1.3 2026-03-11 — Q2/Q3 KM*
> - *v1.4 2026-03-13 — " " " " " DV "*
> - *v1.5 2026-03-17 — Q5 G001 C Q6*
> - *v1.6 2026-03-18 — Q6 G001 ESMO poster Q7 " "*
> - *v1.7 2026-03-18 — Q6 C016 K*

- Q1 DV 2.0 mg/kg G001 1.5 mg/kg
- Q2 RC48-C016 UBC OS UTUC OS UTUC
- Q3 UBC OS UTUC
- Q4 DECIDING-I T HOPE-03
- Q5 G001 C
- Q6
- Q7 C016 " " " " CrCl 50

Q1 DV 2.0 mg/kg G001 1.5 mg/kg

RC48-C005 C009 C016 C017 DV 2.0 mg/kg III DV-001 G001 1.5 mg/kg

Extinction Coefficient EC

EC	BSA-based EC	DV-based EC
	2.0 mg/kg	1.5 mg/kg
	BSA	

DV-based dose = BSA-based dose × (1.07 ÷ 1.41) = 2.0 × 0.759 ≈ 1.52 ≈ 1.5 mg/kg

- 1.07 = BSA-based EC
- 1.41 = DV-based EC

G001 DV-based EC

Q2 RC48-C016 UBC OS UTUC OS UTUC

RC48-C016 NCT05302284 DV+ OS

			OS

UTUC	111	37 33%	31.5
UBC	130	44 34%	23.2
	243	81 33%	31.5

31.5 23.2 OS 31.5

OS KM 50%
OS

OS 33%

- OS 24 34 2 50%
OS 31.5
- **UTUC** 31.5 OS UTUC UTUC 50%
UTUC OS OS 24 UTUC OS

KM OS
OS

Q3 UBC OS UTUC

Q2 DV+T UTUC OS 31.5 vs UBC OS 23.2 8.3

	OS
UTUC	16.9
UBC	16.5

OS 16.9 vs 16.5 DV+T HR 0.53 vs 0.55
DV+T OS Q2

1. 33% KM OS
HR UTUC 0.53 vs UBC 0.55 OS 8.3

2. UTUC UBC ECOG
HR OS
HR 0.53 vs 0.55 DV+T UTUC UBC OS

Q4 DECIDING-I T HOPE-03

DECIDING NCT05979740 HER2
MIBC *Neoplasia* Phase 1 6 "
6 "

1.

DECIDING **6 + 45 = 51**

Phase 1	N=6	DLT	28 ≤2 DLT
Phase 2	N=45	3 CR	—

6 Phase 1 Phase 2 45

2. T Phase 1 n=6

Supplementary Table 1

cT2N0M0	5	83.3%
cT3N0M0	1	16.7%

cT2-4a T2 T3 HER2 3+ CR

3. DECIDING

2-3

Phase 1	±	46 Gy	23 2 Gy/ 5
Phase 2 boost		50-60 Gy	14 Gy/7f 2f

>50 Gy TMT 60-66 Gy

4. HOPE-03

	DECIDING-I	HOPE-03
	concurrent 2-3	sequential
	50-60 Gy	

HOPE-03 V3.0 " "

6 Phase 1 51 6+45 Phase 1 T2 83.3%
50-60 Gy HOPE-03
DECIDING " + " HOPE " "

Q5 G001 C

G001 NCT04879329 II C disitamab vedotin /RC48
pembrolizumab /Keytruda HER2

1.

ClinicalTrials.gov C

	1.5 mg/kg DV-based EC	Q2W 2	
	400 mg	Q6W 6	

400 mg Q6W FDA 2020 4 200 mg Q3W 1.5 mg/kg DV-based EC 2.0 mg/kg BSA-based EC Q1

6 1 15 1 29 3 1

2.

G001 C016 RC48-C016 Protocol v1.0 " RC48 30 "

3. PD-1

	G001 C	**C016**
	400 mg Q6W	3 mg/kg Q2W
	3 DV 1	DV
	6	2

G001 C 1.5 mg/kg Q2W 400 mg Q6W C016 ADC C016

Q6

" "

G001 C016 C016

1.

C016 10 Inclusion #10

> *Blood creatinine ≤ 1.5 × ULN or creatinine clearance (CrCl) ≥ 50 mL/min as calculated by the Cockcroft-Gault equation*

" "

134 µmol/L WS/T 404.5-2015 ULN 97 µmol/L 20-59 111 µmol/L 60-79 ULN 97 1.5 × 97 = 145.5 µmol/L 134 < 145.5 " ≤ 1.5 × ULN"

2.

	C016	
	9	

	6 1	1 LVEF ≥ 50% NYHA < 3
	8 "stably controlled"	
		≥ 100 × 10^9/L
		/CrCl

3. G001

G001 ESMO 2024 1967MO Slide 4 C b

> *ECOG PS of 2 was allowed if hemoglobin ≥10 g/dL and CrCl ≥50 mL/min.*

CrCl ≥50 mL/min G001 ECOG PS 2 ECOG PS 0-1 protocol C016

C016 " ≤ 1.5 × ULN CrCl ≥ 50 mL/min" " " G001 ESMO 2024 CrCl ≥50 ECOG PS 2 >1 LVEF

Q7 C016 " " " " CrCl 50

C016 10 " ≤ 1.5 × ULN CrCl ≥ 50 mL/min" CrCl 40

1. " CrCl "

CrCl Cockcroft-Gault CrCl

> *80 55 kg 90 μmol/L 1.02 mg/dL*
>
> - *90 < 111 × 1.5 = 166.5 μmol/L →*
> - *CrCl = (140−80) × 55 ÷ (72 × 1.02) ≈ 45 mL/min →*

protocol " "

2. " "

CrCl CrCl " " " " C016

3. Protocol

C016 v4.0 10

> *Blood creatinine ≤ 1.5 × ULN or creatinine clearance ≥ 50 mL/min as calculated by the Cockcroft-Gault equation*

Appendix 4 Cockcroft-Gault " "

C016 " ≤ 1.5 × ULN CrCl ≥ 50"
CrCl

---

1. Sheng X, Yan X, Wang L, et al. Disitamab Vedotin plus Toripalimab versus Chemotherapy in HER2-Expressing Advanced Urothelial Cancer: A Randomized Phase III Trial (RC48-C016). *N Engl J Med*. 2025. NCT05302284
2. Powles T, et al. RC48G001: Disitamab Vedotin in Previously Treated HER2-Expressing Advanced Urothelial Cancer. LBA631, 2026 ASCO GU.
3. Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. *J Am Stat Assoc*. 1958;53(282):457-481.
4. Zhang L, Jin D, Zang J, et al. Disitamab vedotin combined with toripalimab and radiotherapy for multimodal organ-sparing treatment of muscle invasive bladder cancer: a proof-of-concept study. *Neoplasia*. 2025;68:101216.
5. Wen F, Lin T, Zhang P, Shen Y. RC48-ADC combined with tislelizumab as neoadjuvant treatment in patients with HER2-positive locally advanced muscle-invasive urothelial bladder cancer: a multi-center phase Ib/II study (HOPE-03). *Front Oncol*. 2024;13:1233196.
6. RC48G001 (NCT04879329). ClinicalTrials.gov.
7. FDA approves new dosing regimen for pembrolizumab. FDA, April 2020.
8. WS/T 404.5-2015. 5 .
9. RC48-C016 Protocol v4.0. RemeGen Co., Ltd. Apr 2024.
10. Sheng X, Powles T, Gupta S, et al. Disitamab vedotin combined with pembrolizumab in treatment-naive patients with HER2-expressing locally advanced or metastatic urothelial cancer: Cohort C of the RC48G001 study. ESMO 2024, Abstract 1967MO.