

TJ Age (or Weight)	Uncuffed ETT	Cuffed ETT	Laryngoscope Blade	DL ETT	CVL	A-line	LMA	Foley	Chest Tube	Salem Sump
1kg	2.5		Miller00		5Fr	24G	1	5Fr	8Fr	
2 kg	3		Miller00		5Fr	24G	1	5Fr	10Fr	10Fr
3 kg	3		Miller0		5Fr	24G	1	5Fr	12Fr	10Fr
Term Newborn	3.5		Miller1		5Fr	24G	1	5Fr	12Fr	10Fr
5 kg	3.5		Miller1		5Fr	24G	1.5	5Fr	12Fr	10Fr
1year	4		Miller1		5Fr	24G	1.5	5Fr	16Fr	10Fr
2 years	4.5		Miller/Mac2		5Fr	24G	2	8Fr	16Fr	14Fr
3 years	4.5		Miller/Mac2		5Fr	22G	2	8Fr	20Fr	14Fr
4 years	5		Miller/Mac2		5Fr	22G	2	8Fr	20Fr	14Fr
5 years		5	Miller/Mac2		5Fr	22G	2	8Fr	20Fr	14Fr
6 years		5	Miller/Mac2		7Fr	22G	2.5	8Fr	20Fr	14Fr
7 years		5	Miller/Mac2		7Fr	22G	2.5	8Fr	20Fr	14Fr
8 years		5.5	Miller/Mac2	26Fr	7Fr	22G	2.5	8Fr	20Fr	14Fr
9 years		5.5	Miller/Mac2	26Fr	7Fr	22G	2.5	10Fr	20Fr	14Fr
10 years		6	Miller/Mac3	28 Fr	7Fr	20G	2.5	10Fr	20Fr	14Fr
11years		6	Miller/Mac3	28 Fr	7Fr	20G	3	10Fr	24 Fr	14Fr
12years		6	Miller/Mac3	28 Fr	7Fr	20G	3	10Fr	24 Fr	14Fr
13years		6	Miller/Mac3	32 Fr	9Fr	20G	3	12 Fr	24 Fr	16 Fr
14 years		7	Miller/Mac3	32 Fr	9Fr	20G	3	12 Fr	32 Fr	16 Fr
Age (or Weight)	Uncuffed ETT	Cuffed ETT	Laryngoscope Blade	DL ETT	CVL	A-line	LMA	Foley	Chest Tube	Salem Sump
15 years		7	Miller/Mac3	35Fr	9Fr	20G	4	12 Fr	32 Fr	16 Fr
16 years		7	Miller/Mac3	35Fr	9Fr	20G	4	12 Fr	32 Fr	16 Fr
17 years		7	Miller/Mac3	35Fr	9Fr	20G	4	12 Fr	36 Fr	16 Fr
18 years		7	Miller/Mac3	35Fr	9Fr	20G	4	12 Fr	36 Fr	16 Fr

出處:Handbook of Pediatric Anesthesia

General Guidelines for Pediatric Patient

	早 產 兒	出 生	6 個月	1 歲	2 歲	3 歲	4 歲	5 歲	6 歲	7 歲	8 歲	10 歲	12 歲	> 15歲
Kg	1.5	3.5	7	10	12	14	17	18	20	22	25	33	40	50
SBP	40	50			65	75								
	~	~	60~115		~	~	80~120	90~120	95~135	100~140				
	60	70			115	125								
HR	120~180	100~180		100	~	90~120	~	~	80	65	70~115	60~110	55~105	
				130			110	135						
RR	55	30												
	~	~	25~40		20~30		14~25	14~22	13~20	10~16				
	60	50												
ETT	2.5	~	3.5~4.0	4.0~5.0	5.0~5.5		5.5~6.0				6.0~6.5			
		3.5					(氣囊)							
Blade		0~1 Miller		1 Miller			2 Miller 1~3 MAC				>8歲 2Miller或3MAC			
Vent Bag			1L				2L				3L			
Oral airway	0	0	1	1	2	2	3	3	3	4	4	4	5	5

Endo大小

4+(年齡÷4)

Endo深度

BW<0.5kg 深度小於/等於5 cm

BW<1.0kg 深度小於/等於6 cm (不適用體重+6公式)

<2歲 內徑mm ×3

>2歲 12+(年齡÷2)

鼻管深度

15+(年齡÷2)

LMA大小選擇

號碼(#) 體重(kg) Cuff(ml)

1	<5	<4
1.5	5~10	<7
2	10~20	<10
2.5	20~30	<14

General Guidelines for Pediatric Patient

Age	BW	Pulse (bpm)	RR (bpm)	SBP (mmHg)	DBP (mmHg)	Minute ventilation (L/min;mL/min/kg)	VC(mL;mL/kg)	Hct(%)
Premature	1	130-150	40-60	42-52	21±8			35-65
Premature	1-2	125-150	40-60	50-60	28±8			35-65
Newborn	2-4	120-150	40-60	60-70	37±8	1.05;200-260	120;40	45-65

出 處 :Pediatric

Anesthesia

Normal vital Signs By Age

Age group HR(beat/min) SBP(mmHg) RR(breaths/min)

Newborn	94-145	60-90	30-60
Infant	125-170	75-100	30-60
Toddler	100-160	80-110	24-40
Preschool	70-110	80-110	22-34
School age	70-110	85-120	18-30
Adolescent	55-100	95-120	12-16

出 處 :Pediatric

Anesthesia

