

Neostigmine

【INEO】Vagostin® 0.5mg/1mL/Amp

ATC Code : N07AA01

中文名：安可健注射液 «信東»

適應症：腸管麻痺、筋無力症、血管痙攣性疾患、眼筋無力症。

藥理分類：Acetylcholinesterase Inhibitor.

用法用量：Administration:

1. **IM, SC or slow IV**(at least one minute, concomitantly with or just after IV atropine sulfate).
2. Acute colonic pseudo-obstruction (off-label use):
Note: Continuous cardiac monitoring with clinical assessment for 30 minutes after administration. Atropine should be available at the bedside if bradyarrhythmia occur.
 - **Bolus: IV:** Administer over 3 to 5 minutes.
 - **Intermittent infusion: IV:** Administration over 60 minutes.
 - **Continuous infusion: IV:** 0.4 mg per hour for up to 8 hours; if no response increase to 0.8 mg per hour for up to 24 hours.

Indications and dosage regimens:

Diagnosis of myasthenia gravis: IM.

- Adults: 0.022 mg/kg preceded by atropine sulfate 0.011 mg/kg IM.
- Children: 0.025-0.04 mg/kg preceded by atropine sulfate 0.011 mg/kg SC.

Myasthenia gravis: IM, SC, IV (IV route is hazardous)

- Adults: 0.5-2.5 mg
- Children: 0.01-0.04 mg/kg Q2-4h as needed.

Neonatal myasthenia gravis: IM

0.03 mg/kg Q2-4h; SC, 0.1-0.2 mg.

Neuromuscular blockade reversal: Slow IV

- Adults:
0.5-2.5 mg; repeated if necessary, totally not exceed 5 mg. (0.6-1.2 mg atropine sulfate should be given to counteract the adverse muscarinic effects.)
- Neonates & infants:
0.04 mg/kg with 0.02 mg/kg atropine sulfate.

Postoperative distention and urinary retention:

- Prophylaxis: Adults, IM or SC, 0.25 mg Q4-6h for 2-3 days.
- Treatment: Adults, IM or SC, 0.5-1 mg, repeated Q3-5h for 5 doses after the bladder has been emptied.

Acute colonic pseudo-obstruction (Ogilvie syndrome) (off-label use):

Note: Continuous cardiac monitoring with clinical assessment for 30 minutes after administration. Atropine should be available at the bedside if bradyarrhythmia occurs.

Bolus: IV: 2 mg over 3 to 5 minutes.

Intermittent infusion: IV: 2.5 mg over 60 minutes.

Continuous infusion: IV: 0.4 mg per hour for up to 8 hours; if no response increase to 0.8 mg per hour for up to 24 hours.

不良反應：唾液或汗水增多、胃腸不適、抽筋等。

交互作用：● Corticosteroids 對重症肌無力患者產生嚴重肌肉強度衰竭的現象，作用從服

藥後第 2 天開始在第 7 天達到最強，並且對 anticholinesterase 的治療不反應，嚴重的甚至需要靠呼吸器才能維持呼吸，併用時仍須嚴密監控病人反應並隨時提供生命維持系統待命。

● Donepezil, bupropion: reduced seizure threshold, ↑ risk of seizures.

● Hydrocortisone, prednisolone: ↓ neostigmine effectiveness.

注意事項：

1. 仿單:可肌肉注射或皮下注射(用於眼科) 。仿單未提到 IV 給藥。

2. 腸或尿道阻塞之患者禁用本劑。

3. (仿單)皮下及肌肉注射之，每日 1~3 次，每次 0.25~0.5mg。必要時(Antidote to curare Principle)以 0.5mg 之本劑與 0.6mg 之 Atropine Sulfate 合併靜脈注射。

懷 孕 期：

1. 無相關資料，宜審慎評估。

2. Anticholinesterases have caused uterine irritability and induced **premature labor** with IV use in near-term pregnant women. Neostigmine may be used to treat myasthenia gravis in pregnant women; however, if an acetylcholinesterase inhibitor is needed during pregnancy, another agent may be preferred.

授 乳 期：

1. 未有授乳婦使用 neostigmine 產生問題之文獻，故宜慎用。

2. It is not known if neostigmine is present in breast milk.

3. The decision to breastfeed during therapy should consider the risk of infant exposure, the benefits of breastfeeding to the infant, and the benefits of treatment to the mother.

貯 存：

避光貯存。