



Metamodeling under Inequality Constraints

Andrés F. López-Lopera¹, François Bachoc², Nicolas Durrande^{1,3}, and Olivier Roustant¹

¹École des Mines de Saint-Étienne (EMSE), France.

²Institut de Mathématiques de Toulouse (IMT), France.

³PROWLER.io, Cambridge, UK.

Nov 22, 2017

Table of Contents

1 Motivation

- Gaussian CDF example
- Constrained Kriging: Maatouk and Bay (2016)

2 Contributions

- Old contributions (Nice, 2017)
- Constrained maximum likelihood estimation (CMLE)
- Finite-dimensional approximation for 2D input spaces
- 2D example (IRSN)

3 Conclusions and Future Works

4 Questions?

5 References

Table of Contents

1 Motivation

- Gaussian CDF example
- Constrained Kriging: Maatouk and Bay (2016)

2 Contributions

- Old contributions (Nice, 2017)
- Constrained maximum likelihood estimation (CMLE)
- Finite-dimensional approximation for 2D input spaces
- 2D example (IRSN)

3 Conclusions and Future Works

4 Questions?

5 References

Table of Contents

1 Motivation

- Gaussian CDF example
- Constrained Kriging: Maatouk and Bay (2016)

2 Contributions

- Old contributions (Nice, 2017)
- Constrained maximum likelihood estimation (CMLE)
- Finite-dimensional approximation for 2D input spaces
- 2D example (IRSN)


3 Conclusions and Future Works

4 Questions?

5 References

Motivation


Toy example. $y(x) = \frac{1}{2} \left[1 + \operatorname{erf} \left\{ \frac{x-0.5}{0.2\sqrt{2}} \right\} \right]$ (Gaussian CDF).



⇒ Data are bounded and monotonic!!

Motivation


Toy example. $y(x) = \frac{1}{2} \left[1 + \operatorname{erf} \left\{ \frac{x-0.5}{0.2\sqrt{2}} \right\} \right]$ (Gaussian CDF).



Unconstrained Kriging

Motivation

Toy example. $y(x) = \frac{1}{2} \left[1 + \operatorname{erf} \left\{ \frac{x-0.5}{0.2\sqrt{2}} \right\} \right]$ (Gaussian CDF).



Constrained Kriging

Table of Contents

1 Motivation

- Gaussian CDF example
- Constrained Kriging: Maatouk and Bay (2016)

2 Contributions

- Old contributions (Nice, 2017)
- Constrained maximum likelihood estimation (CMLE)
- Finite-dimensional approximation for 2D input spaces
- 2D example (IRSN)


3 Conclusions and Future Works

4 Questions?

5 References

Constrained Kriging: finite-dimensional approximation

Figure: toy example $y(x) = \Phi(\frac{x-0.5}{0.2})$ (Gaussian CDF).




⇒ The finite approx. is also bounded and monotonic!!

Constrained Kriging: Maatouk and Bay (2016)

Let the finite-one-dimensional GP-based approximation be defined as


$$Y_m(x) = \sum_{j=1}^m \xi_j \phi_j(x), \quad \text{s.t.} \quad \begin{cases} Y_m(x_i) = y_i & (\text{interpolation conditions}), \\ Y_m \in \mathcal{E} & (\text{inequality conditions}), \end{cases}$$

where $\boldsymbol{\xi} = [\xi_1, \dots, \xi_m]^\top \sim \mathcal{N}(\mathbf{0}, \boldsymbol{\Gamma})$, and $\phi_j : [0, 1] \rightarrow \mathbb{R}$ are hat functions.



Constrained Kriging: finite-dimensional approximation


Figure: toy example $y(x) = \Phi(\frac{x-0.5}{0.2})$ (Gaussian CDF).



Finite-dimensional approximation.

Constrained Kriging: finite-dimensional approximation


Figure: toy example $y(x) = \Phi(\frac{x-0.5}{0.2})$ (Gaussian CDF).



Unconstrained GP.

Constrained Kriging: finite-dimensional approximation


Figure: toy example $y(x) = \Phi(\frac{x-0.5}{0.2})$ (Gaussian CDF).



Constrained GP with boundedness constraint.

Constrained Kriging: finite-dimensional approximation

Figure: toy example $y(x) = \Phi(\frac{x-0.5}{0.2})$ (Gaussian CDF).



Constrained GP with monotonicity constraint.

Table of Contents

1 Motivation

- Gaussian CDF example
- Constrained Kriging: Maatouk and Bay (2016)

2 Contributions

- Old contributions (Nice, 2017)
- Constrained maximum likelihood estimation (CMLE)
- Finite-dimensional approximation for 2D input spaces
- 2D example (IRSN)

3 Conclusions and Future Works

4 Questions?

5 References

Table of Contents

1 Motivation

- Gaussian CDF example
- Constrained Kriging: Maatouk and Bay (2016)

2 Contributions

- Old contributions (Nice, 2017)
- Constrained maximum likelihood estimation (CMLE)
- Finite-dimensional approximation for 2D input spaces
- 2D example (IRSN)

3 Conclusions and Future Works

4 Questions?

5 References

Old contributions (Nice, 2017)

1. First, since $Y_m \in \mathcal{E} \Leftrightarrow \boldsymbol{\xi} \in \mathcal{C}$, and assuming that \mathcal{C} is composed by a set of q linear inequalities of the form

$$\mathcal{C} = \left\{ \mathbf{c} \in \mathbb{R}^m; \forall j = 1, \dots, m, \forall k = 1, \dots, q, \lambda_{k,j} \in \mathbb{R} : \ell_k \leq \sum_{j=1}^m \lambda_{k,j} c_j \leq u_k \right\},$$

the posterior distribution is given by a truncated multinormal

$$\Lambda \boldsymbol{\xi} | \{ \Phi \boldsymbol{\xi} = \mathbf{y}, \mathbf{l} \leq \Lambda \boldsymbol{\xi} \leq \mathbf{u} \} \sim \mathcal{TN}(\Lambda \boldsymbol{\mu}, \Lambda \Sigma \Lambda^\top, \mathbf{l}, \mathbf{u}), \quad (1)$$

where $\Lambda = (\lambda_{k,j})_{1 \leq k \leq q, 1 \leq j \leq m}$, $\mathbf{l} = (\ell_k)_{1 \leq k \leq q}$, $\mathbf{u} = (u_k)_{1 \leq k \leq q}$, and

$$\boldsymbol{\mu} = \Gamma \Phi^\top [\Phi \Gamma \Phi^\top]^{-1} \mathbf{y}, \quad \text{and} \quad \Sigma = \Gamma - \Gamma \Phi^\top [\Phi \Gamma \Phi^\top]^{-1} \Phi \Gamma. \quad (2)$$

Old contributions (Nice, 2017)

1. First, since $Y_m \in \mathcal{E} \Leftrightarrow \boldsymbol{\xi} \in \mathcal{C}$, and assuming that \mathcal{C} is composed by a set of q linear inequalities of the form

$$\mathcal{C} = \left\{ \mathbf{c} \in \mathbb{R}^m; \forall j = 1, \dots, m, \forall k = 1, \dots, q, \lambda_{k,j} \in \mathbb{R} : \ell_k \leq \sum_{j=1}^m \lambda_{k,j} c_j \leq u_k \right\},$$

the posterior distribution is given by a truncated multinormal

$$\Lambda \boldsymbol{\xi} | \{ \Phi \boldsymbol{\xi} = \mathbf{y}, \mathbf{l} \leq \Lambda \boldsymbol{\xi} \leq \mathbf{u} \} \sim \mathcal{TN}(\Lambda \boldsymbol{\mu}, \Lambda \Sigma \Lambda^\top, \mathbf{l}, \mathbf{u}), \quad (1)$$

where $\Lambda = (\lambda_{k,j})_{1 \leq k \leq q, 1 \leq j \leq m}$, $\mathbf{l} = (\ell_k)_{1 \leq k \leq q}$, $\mathbf{u} = (u_k)_{1 \leq k \leq q}$, and

$$\boldsymbol{\mu} = \Gamma \Phi^\top [\Phi \Gamma \Phi^\top]^{-1} \mathbf{y}, \quad \text{and} \quad \Sigma = \Gamma - \Gamma \Phi^\top [\Phi \Gamma \Phi^\top]^{-1} \Phi \Gamma. \quad (2)$$

⇒ The distribution of Equation (1) can be approximated using MCMC.

Old contributions (Nice, 2017)

1. First, since $Y_m \in \mathcal{E} \Leftrightarrow \boldsymbol{\xi} \in \mathcal{C}$, and assuming that \mathcal{C} is composed by a set of q linear inequalities of the form

$$\mathcal{C} = \left\{ \mathbf{c} \in \mathbb{R}^m; \forall j = 1, \dots, m, \forall k = 1, \dots, q, \lambda_{k,j} \in \mathbb{R} : \ell_k \leq \sum_{j=1}^m \lambda_{k,j} c_j \leq u_k \right\},$$

the posterior distribution is given by a truncated multinormal

$$\boldsymbol{\Lambda} \boldsymbol{\xi} | \{ \boldsymbol{\Phi} \boldsymbol{\xi} = \mathbf{y}, \mathbf{l} \leq \boldsymbol{\Lambda} \boldsymbol{\xi} \leq \mathbf{u} \} \sim \mathcal{TN}(\boldsymbol{\Lambda} \boldsymbol{\mu}, \boldsymbol{\Lambda} \boldsymbol{\Sigma} \boldsymbol{\Lambda}^\top, \mathbf{l}, \mathbf{u}), \quad (1)$$

where $\boldsymbol{\Lambda} = (\lambda_{k,j})_{1 \leq k \leq q, 1 \leq j \leq m}$, $\mathbf{l} = (\ell_k)_{1 \leq k \leq q}$, $\mathbf{u} = (u_k)_{1 \leq k \leq q}$, and


$$\boldsymbol{\mu} = \boldsymbol{\Gamma} \boldsymbol{\Phi}^\top [\boldsymbol{\Phi} \boldsymbol{\Gamma} \boldsymbol{\Phi}^\top]^{-1} \mathbf{y}, \quad \text{and} \quad \boldsymbol{\Sigma} = \boldsymbol{\Gamma} - \boldsymbol{\Gamma} \boldsymbol{\Phi}^\top [\boldsymbol{\Phi} \boldsymbol{\Gamma} \boldsymbol{\Phi}^\top]^{-1} \boldsymbol{\Phi} \boldsymbol{\Gamma}. \quad (2)$$

⇒ The distribution of Equation (1) can be approximated using MCMC.

⇒ What about $\boldsymbol{\Lambda}$, \mathbf{l} , \mathbf{u} ?

Old contributions (Nice, 2017)


Boundedness constraint: $y(x) = \Phi\left(\frac{x-0.5}{0.2}\right)$.



$$\underbrace{\begin{bmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{bmatrix}}_{l_b} \preceq \underbrace{\begin{bmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & \cdots & 0 & 1 \end{bmatrix}}_{\Lambda_b} \underbrace{\begin{bmatrix} \xi_1 \\ \xi_2 \\ \xi_3 \\ \vdots \\ \xi_{m-1} \\ \xi_m \end{bmatrix}}_{\xi} \preceq \underbrace{\begin{bmatrix} 1 \\ 1 \\ 1 \\ \vdots \\ 1 \\ 1 \end{bmatrix}}_{u_b}$$

Old contributions (Nice, 2017)


Monotonicity constraint: $y(x) = \Phi\left(\frac{x-0.5}{0.2}\right)$.



$$\underbrace{\begin{bmatrix} -\infty \\ 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{bmatrix}}_{l_m} \preceq \underbrace{\begin{bmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ -1 & 1 & 0 & \cdots & 0 & 0 \\ 0 & -1 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & \cdots & -1 & 1 \end{bmatrix}}_{\Lambda_m} \underbrace{\begin{bmatrix} \xi_1 \\ \xi_2 \\ \xi_3 \\ \vdots \\ \xi_{m-1} \\ \xi_m \end{bmatrix}}_{\xi} \preceq \underbrace{\begin{bmatrix} \infty \\ \infty \\ \infty \\ \vdots \\ \infty \\ \infty \end{bmatrix}}_{u_m}$$

Old contributions (Nice, 2017)


Boundedness and monotonicity constraints: $y(x) = \Phi\left(\frac{x-0.5}{0.2}\right)$.



$$\underbrace{\begin{bmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 0 \\ 0 \\ 0 \end{bmatrix}}_l \preceq \underbrace{\begin{bmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ -1 & 1 & 0 & \cdots & 0 & 0 \\ 0 & -1 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & \cdots & -1 & 1 \\ 0 & 0 & 0 & \cdots & 0 & 1 \end{bmatrix}}_{\Lambda} \underbrace{\begin{bmatrix} \xi_1 \\ \xi_2 \\ \xi_3 \\ \vdots \\ \xi_{m-1} \\ \xi_m \end{bmatrix}}_{\xi} \preceq \underbrace{\begin{bmatrix} 1 \\ \infty \\ \infty \\ \vdots \\ \infty \\ \infty \\ 1 \end{bmatrix}}_u.$$

Old contributions (Nice, 2017)

Boundedness and monotonicity constraints: $y(x) = \Phi\left(\frac{x-0.5}{0.2}\right)$.



or simply,

$$\underbrace{\begin{bmatrix} l_b \\ l_m \end{bmatrix}}_l \leq \underbrace{\begin{bmatrix} \Lambda_b \\ \Lambda_m \end{bmatrix}}_{\Lambda} \xi \leq \underbrace{\begin{bmatrix} u_b \\ u_m \end{bmatrix}}_u.$$

Old contributions (Nice, 2017)

2. Second, the truncated multinormal $\Lambda\xi | \{\Phi\xi = \mathbf{y}, \mathbf{l} \leq \Lambda\xi \leq \mathbf{u}\}$ can be sampled efficiently via Hamiltonian Monte-Carlo (HMC) (Pakman and Paninski, 2014).

Old contributions (Nice, 2017)

2. Second, the truncated multinormal $\Lambda\xi | \{\Phi\xi = \mathbf{y}, \mathbf{l} \leq \Lambda\xi \leq \mathbf{u}\}$ can be sampled efficiently via Hamiltonian Monte-Carlo (HMC) (Pakman and Paninski, 2014).

⇒ Denoting $\eta = \Lambda\xi$ with Λ injective, notice that the samples of ξ can be obtained using the samples of η by solving the linear system.

Old contributions (Nice, 2017)

2. Second, the truncated multinormal $\Lambda\xi | \{\Phi\xi = \mathbf{y}, \mathbf{l} \leq \Lambda\xi \leq \mathbf{u}\}$ can be sampled efficiently via Hamiltonian Monte-Carlo (HMC) (Pakman and Paninski, 2014).

\Rightarrow Denoting $\eta = \Lambda\xi$ with Λ injective, notice that the samples of ξ can be obtained using the samples of η by solving the linear system.

3. Third, a constrained likelihood was suggested empirically

$$\begin{aligned}\mathcal{L}_{\mathcal{C},m}(\theta) &= \log p_{\theta}(\mathbf{Y}_m | \xi \in \mathcal{C}) \\ &= \log \frac{p_{\theta}(\mathbf{Y}_m) P_{\theta}(\xi \in \mathcal{C} | \Phi\xi = \mathbf{Y}_m)}{P_{\theta}(\xi \in \mathcal{C})} \\ &= \log p_{\theta}(\mathbf{Y}_m) + \log P_{\theta}(\xi \in \mathcal{C} | \Phi\xi = \mathbf{Y}_m) - \log P_{\theta}(\xi \in \mathcal{C}).\end{aligned}$$

Old contributions (Nice, 2017)

2. Second, the truncated multinormal $\Lambda\xi | \{\Phi\xi = \mathbf{y}, \mathbf{l} \leq \Lambda\xi \leq \mathbf{u}\}$ can be sampled efficiently via Hamiltonian Monte-Carlo (HMC) (Pakman and Paninski, 2014).

\Rightarrow Denoting $\eta = \Lambda\xi$ with Λ injective, notice that the samples of ξ can be obtained using the samples of η by solving the linear system.

3. Third, a constrained likelihood was suggested empirically

$$\begin{aligned}\mathcal{L}_{\mathcal{C},m}(\theta) &= \log p_{\theta}(\mathbf{Y}_m | \xi \in \mathcal{C}) \\ &= \log \frac{p_{\theta}(\mathbf{Y}_m) P_{\theta}(\xi \in \mathcal{C} | \Phi\xi = \mathbf{Y}_m)}{P_{\theta}(\xi \in \mathcal{C})} \\ &= \log p_{\theta}(\mathbf{Y}_m) + \log P_{\theta}(\xi \in \mathcal{C} | \Phi\xi = \mathbf{Y}_m) - \log P_{\theta}(\xi \in \mathcal{C}).\end{aligned}$$

\Rightarrow unconstrained log-likelihood.

\Rightarrow truncated multinormals to be estimated!!

Old contributions (Nice, 2017)

2. Second, the truncated multinormal $\Lambda \xi | \{\Phi \xi = \mathbf{y}, \mathbf{l} \leq \Lambda \xi \leq \mathbf{u}\}$ can be sampled efficiently via Hamiltonian Monte-Carlo (HMC) (Pakman and Paninski, 2014).

\Rightarrow Denoting $\eta = \Lambda \xi$ with Λ injective, notice that the samples of ξ can be obtained using the samples of η by solving the linear system.

3. Third, a constrained likelihood was suggested empirically

$$\begin{aligned} \mathcal{L}_{\mathcal{C},m}(\theta) &= \log p_{\theta}(\mathbf{Y}_m | \xi \in \mathcal{C}) \\ &= \log \frac{p_{\theta}(\mathbf{Y}_m) P_{\theta}(\xi \in \mathcal{C} | \Phi \xi = \mathbf{Y}_m)}{P_{\theta}(\xi \in \mathcal{C})} \\ &= \log p_{\theta}(\mathbf{Y}_m) + \log P_{\theta}(\xi \in \mathcal{C} | \Phi \xi = \mathbf{Y}_m) - \log P_{\theta}(\xi \in \mathcal{C}). \end{aligned}$$

\Rightarrow unconstrained log-likelihood.

\Rightarrow truncated multinormals to be estimated!!

4. Finally, the full framework was assessed under different inequality constraints in synthetic examples.

Old contributions (Nice, 2017)

In this sense, we proposed at (Nice, 2017):

- to implement a gradient-based method to estimate automatically the covariance parameters of the model;
- to investigate theoretical properties of the proposed constrained likelihood;
- to evaluate the proposed approach with real-world datasets;
- to build an R package;
- and to extend this approach when the input space is multi-dimensional, i.e. $\mathbf{x} \in [0 \ 1]^d$.

Old contributions (Nice, 2017)

In this sense, we proposed at (Nice, 2017):

- to implement a gradient-based method to estimate automatically the covariance parameters of the model; ✓
- to investigate theoretical properties of the proposed constrained likelihood; ✓
- to evaluate the proposed approach with real-world datasets; ✓
- to build an R package; ✓
- and to extend this approach when the input space is multi-dimensional, i.e. $\mathbf{x} \in [0 \ 1]^d$. **2D ✓ More than 2D ✗**

Table of Contents

1 Motivation

- Gaussian CDF example
- Constrained Kriging: Maatouk and Bay (2016)

2 Contributions

- Old contributions (Nice, 2017)
- **Constrained maximum likelihood estimation (CMLE)**
- Finite-dimensional approximation for 2D input spaces
- 2D example (IRSN)

3 Conclusions and Future Works

4 Questions?

5 References

Maximum likelihood (ML): asymptotic properties.

Let \mathcal{E}_κ be one of the following convex set of functions

$$\mathcal{E}_\kappa = \begin{cases} f : \mathbb{X} \rightarrow \mathbb{R}, f \text{ is } C^0 \text{ and } \forall \mathbf{x} \in \mathbb{X}, \ell \leq f(\mathbf{x}) \leq u & \text{if } \kappa = 0, \\ f : \mathbb{X} \rightarrow \mathbb{R}, f \text{ is } C^1 \text{ and } \forall \mathbf{x} \in \mathbb{X}, \forall i = 1, \dots, d, \frac{\partial}{\partial x_i} f(\mathbf{x}) \geq 0 & \text{if } \kappa = 1, \\ f : \mathbb{X} \rightarrow \mathbb{R}, f \text{ is } C^2 \text{ and } \forall \mathbf{x} \in \mathbb{X}, \frac{\partial^2}{\partial \mathbf{x}^2} f(\mathbf{x}) \text{ is a non-negative} & \text{if } \kappa = 2, \\ & \text{definite matrix} \end{cases}$$

which corresponds to boundedness, monotonicity, and convexity constraints. We will focus on the GP Y and the observation vector

$$\mathbf{Y}_n = [Y(x_1), \dots, Y(x_n)]^\top.$$

Maximum likelihood (ML)

Proposition 1: asymptotic consistency of ML

Let Y be a centred GP on $\mathbb{X} \subset \mathbb{R}^d$ with covariance k satisfying **Condition A.1 from (López-Lopera et al., 2017)**. Let $\mathbf{Y}_n = [Y(x_1), \dots, Y(x_n)]^\top$. Let

$$\mathcal{L}_n(\boldsymbol{\theta}) = -\frac{1}{2} \log(\det(\mathbf{R}_{\boldsymbol{\theta}})) - \frac{1}{2} \mathbf{Y}_n^\top \mathbf{R}_{\boldsymbol{\theta}}^{-1} \mathbf{Y}_n - \frac{n}{2} \log 2\pi, \quad (\text{Unconstrained likelihood})$$

with $\mathbf{R}_{\boldsymbol{\theta}} = (k_{\boldsymbol{\theta}}(x_i, x_j))_{1 \leq i, j \leq n}$. Let $\hat{\boldsymbol{\theta}} \in \arg \max_{\boldsymbol{\theta} \in \Theta} \mathcal{L}_n(\boldsymbol{\theta})$. Assume $\forall \varepsilon > 0$,

$$P(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\| \geq \varepsilon) \xrightarrow{n \rightarrow \infty} 0. \quad (\text{Consistency of the unconditional ML})$$

Let $\kappa \in \{0, 1, 2\}$. Let \mathcal{E}_κ . Then, we have $P(Y \in \mathcal{E}_\kappa) > 0$ from Lemmas A.3, A.4 and A.5 of (López-Lopera et al., 2017), and thus

$$P(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\| \geq \varepsilon \mid Y \in \mathcal{E}_\kappa) \xrightarrow{n \rightarrow \infty} 0. \quad (\text{Consistency of the conditional ML})$$

Maximum likelihood (ML)

Proof 1: asymptotic consistency of ML

We have

$$P(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\| \geq \varepsilon \mid Y \in \mathcal{E}_\kappa) = \frac{P(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\| \geq \varepsilon, Y \in \mathcal{E}_\kappa)}{P(Y \in \mathcal{E}_\kappa)} \leq \frac{P(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\| \geq \varepsilon)}{P(Y \in \mathcal{E}_\kappa)}.$$

Since $P(Y \in \mathcal{E}_\kappa) > 0$ is fixed, and $P(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\| \geq \varepsilon) \xrightarrow[n \rightarrow \infty]{} 0$, the result follows.

Constrained maximum likelihood (CML)

Proposition 2: asymptotic consistency of CML

We use the same notations and assumptions as in Proposition 1. Let P_{θ} be the distribution of Y with covariance function k_{θ} . Let

$$\mathcal{L}_{C,n}(\theta) = \mathcal{L}_n(\theta) + \log P_{\theta}(Y \in \mathcal{E}_{\kappa} | \mathbf{Y}_n) - \log P_{\theta}(Y \in \mathcal{E}_{\kappa}). \quad (\text{Constrained ML})$$

Assume that $\forall \varepsilon > 0$ and $\forall M < \infty$, (Consistency of the unconditional ML)

$$P\left(\sup_{\|\theta - \theta^*\| \geq \varepsilon} (\mathcal{L}_n(\theta) - \mathcal{L}_n(\theta^*)) \geq -M\right) \xrightarrow{n \rightarrow \infty} 0.$$

Then, (Consistency of the conditional CML)

$$P\left(\sup_{\|\theta - \theta^*\| \geq \varepsilon} (\mathcal{L}_{C,n}(\theta) - \mathcal{L}_{C,n}(\theta^*)) \geq -M \mid Y \in \mathcal{E}_{\kappa}\right) \xrightarrow{n \rightarrow \infty} 0.$$

Consequently (Consistency of ML and CML estimators)

$$\operatorname{argmax}_{\theta \in \Theta} \mathcal{L}_n(\theta) \xrightarrow[n \rightarrow \infty]{P} \theta^*, \quad \text{and} \quad \operatorname{argmax}_{\theta \in \Theta} \mathcal{L}_{C,n}(\theta) \xrightarrow[n \rightarrow \infty]{P|Y \in \mathcal{E}_{\kappa}} \theta^*.$$

Table of Contents

- 1 Motivation
 - Gaussian CDF example
 - Constrained Kriging: Maatouk and Bay (2016)
- 2 Contributions
 - Old contributions (Nice, 2017)
 - Constrained maximum likelihood estimation (CMLE)
 - **Finite-dimensional approximation for 2D input spaces**
 - 2D example (IRSN)
- 3 Conclusions and Future Works
- 4 Questions?
- 5 References

Finite-dimensional approximation for 2D input spaces

The approximation can be extended to two dimensional input spaces by tensorisation:


$$Y_{m_1, m_2}(x_1, x_2) := \sum_{j_1=1}^{m_1} \sum_{j_2=1}^{m_2} \xi_{j_2, j_1} \phi_{j_1}^1(x_1) \phi_{j_2}^2(x_2), \text{ s.t. } \begin{cases} Y_{m_1, m_2}(x_1^i, x_2^i) = y_i, \\ \xi_{j_2, j_1} \in \mathcal{C}, \end{cases}$$

where $\xi_{j_2, j_1} = Y(t_{j_1}, t_{j_2})$ and $(x_1^1, x_2^1), \dots, (x_1^n, x_2^n)$ constitute a DoE, and $\phi_{j_1}^1, \phi_{j_2}^2 : [0, 1] \rightarrow \mathbb{R}$ are hat functions.

\Rightarrow We can also assume that $\boldsymbol{\xi} = [\xi_{1,1}, \dots, \xi_{1,m_1}, \dots, \xi_{m_2,1}, \dots, \xi_{m_2,m_1}]^\top$ is a zero-mean Gaussian vector with covariance matrix $\boldsymbol{\Gamma}$.

Finite-dimensional approximation for 2D input spaces


Boundedness in 2D.



Examples of 2D Gaussian models with different types of constraints.

Finite-dimensional approximation for 2D input spaces


Monotonicity in 2D.



Examples of 2D Gaussian models with different types of constraints.

Finite-dimensional approximation for 2D input spaces


Convexity in 2D (a weak version).



Examples of 2D Gaussian models with different types of constraints.

Finite-dimensional approximation for 2D input spaces

Boundedness and monotonicity in 2D.



Examples of 2D Gaussian models with different types of constraints.

Table of Contents

1 Motivation

- Gaussian CDF example
- Constrained Kriging: Maatouk and Bay (2016)

2 Contributions


- Old contributions (Nice, 2017)
- Constrained maximum likelihood estimation (CMLE)
- Finite-dimensional approximation for 2D input spaces
- 2D example (IRSN)

3 Conclusions and Future Works

4 Questions?

5 References

2D example (IRSN)



Nuclear criticality safety assessments: IRSN's dataset.

$\Rightarrow k_{eff}$ is positive and non-decreasing.


2D example (IRSN)

Procedure:


- 1 We used a Latin hypercube design (LHD) with different number of training points n .
- 2 We trained unconstrained and constrained models using either MLE or CMLE.
- 3 For the constrained models, we imposed both positivity and monotonicity constraints.
- 4 We evaluated their performances over the test points (i.e. $121 - n$).

2D example (IRSN)

Unconstrained model + MLE



(a)




(b)


2D Gaussian models for interpolating the IRSN's dataset.

2D example (IRSN)

Constrained model + MLE



(c)




(d)


2D Gaussian models for interpolating the IRSN's dataset.

2D example (IRSN)

Constrained model + CMLE



(e)



(f)

2D Gaussian models for interpolating the IRSN's dataset.

2D example (IRSN)

Now, we repeat the procedure for 20 random LHDs, and we compute the Q^2 and predictive variance adequation (PVA) criteria...

2D example (IRSN)

Let n_t be the number of test points, z_1, \dots, z_{n_t} and $\hat{z}_1, \dots, \hat{z}_{n_t}$ the sets of test and predicted observations (respectively), then...

Q^2 criterion:

$$Q^2 = 1 - \frac{\sum_{i=1}^{n_t} (\hat{z}_i - z_i)^2}{\sum_{i=1}^{n_t} (\bar{z} - z_i)^2}, \quad (3)$$


where \bar{z} is the mean of the test data. $\Rightarrow Q^2 \rightarrow 1 \checkmark$

Predictive variance adequation (PVA) criterion:


$$\text{PVA} = \left| \log \left(\frac{1}{n_t} \sum_{i=1}^{n_t} \frac{(\hat{z}_i - z_i)^2}{\hat{\sigma}_i^2} \right) \right|, \quad (4)$$

where $\hat{\sigma}_i^2$ are the predictive variances. $\Rightarrow \text{PVA} \rightarrow 0 \checkmark$

2D example (IRSN)





Model
 1. UM-MLE
 2. CM-MLE
 3. CM-CMLE



Model
 1. UM-MLE
 2. CM-MLE
 3. CM-CMLE

Assessment of the models for interpolating the IRSN's dataset using different number of training points n and using twenty different Latin hypercube designs.



2D example (IRSN)



Assessment of the models for interpolating the IRSN's dataset using different number of training points n and using twenty different Latin hypercube designs.

⇒ Unconstrained model was often outperformed by constrained ones.

2D example (IRSN)





Assessment of the models for interpolating the IRSN's dataset using different number of training points n and using twenty different Latin hypercube designs.

⇒ Unconstrained model was often outperformed by constrained ones.

⇒ MLE achieves a good tradeoff between prediction accuracy and computational cost.

2D example (IRSN)



Assessment of the models for interpolating the IRSN's dataset using different number of training points n and using twenty different Latin hypercube designs.

⇒ Unconstrained model was often outperformed by constrained ones.

⇒ MLE achieves a good tradeoff between prediction accuracy and computational cost.

⇒ CMLE is unstable due to numerical approximations.

Table of Contents

1 Motivation

- Gaussian CDF example
- Constrained Kriging: Maatouk and Bay (2016)

2 Contributions

- Old contributions (Nice, 2017)
- Constrained maximum likelihood estimation (CMLE)
- Finite-dimensional approximation for 2D input spaces
- 2D example (IRSN)

3 Conclusions and Future Works

4 Questions?

5 References

Conclusions and Future Works

Conclusions

- We further investigated the approach proposed in (Maatouk and Bay, 2017): now it works for any linear set of inequality constraints in 1D or 2D.
- We proved the consistency of the constrained likelihood for covariance parameter estimation.
- We implemented the R codes: `lineqGP` package (second round!!).

Conclusions and Future Works

Conclusions

- We further investigated the approach proposed in (Maatouk and Bay, 2017): now it works for any linear set of inequality constraints in 1D or 2D.
- We proved the consistency of the constrained likelihood for covariance parameter estimation.
- We implemented the R codes: `lineqGP` package (second round!!).

♦ Working paper:

López-Lopera, A.F., Bachoc, F., Durrande, N., and Roustant, O. (2017). Finite-dimensional Gaussian approximation with linear inequality constraints. ArXiv e-prints. (Bon courage!!)

Conclusions and Future Works

Future works

- To find an efficient and more reliable estimator of orthant multinormal distributions

$$\mathcal{L}_{\mathcal{C},m}(\boldsymbol{\theta}) = \log p_{\boldsymbol{\theta}}(\mathbf{Y}_m) + \log P_{\boldsymbol{\theta}}(\boldsymbol{\xi} \in \mathcal{C} | \Phi \boldsymbol{\xi} = \mathbf{Y}_m) - \log P_{\boldsymbol{\theta}}(\boldsymbol{\xi} \in \mathcal{C}).$$


Conclusions and Future Works

Future works

- To find an efficient and more reliable estimator of orthant multinormal distributions

$$\mathcal{L}_{\mathcal{C},m}(\theta) = \log p_{\theta}(\mathbf{Y}_m) + \log P_{\theta}(\xi \in \mathcal{C} | \Phi \xi = \mathbf{Y}_m) - \log P_{\theta}(\xi \in \mathcal{C}).$$

- To work in the full framework for higher dimensions...



For example, in multidimensional problems with specific constrained dimensions.


Conclusions and Future Works

Future works

- To find an efficient and more reliable estimator of orthant multinormal distributions

$$\mathcal{L}_{C,m}(\theta) = \log p_{\theta}(\mathbf{Y}_m) + \log P_{\theta}(\xi \in \mathcal{C} | \Phi \xi = \mathbf{Y}_m) - \log P_{\theta}(\xi \in \mathcal{C}).$$

- To work in the full framework for higher dimensions...



For example, in multidimensional problems with specific constrained dimensions.

- To study more asymptotic properties of the proposed framework.

Acknowledgement

We thank Yann Richet (IRSN) for providing the nuclear criticality safety data.

Table of Contents

1 Motivation

- Gaussian CDF example
- Constrained Kriging: Maatouk and Bay (2016)

2 Contributions

- Old contributions (Nice, 2017)
- Constrained maximum likelihood estimation (CMLE)
- Finite-dimensional approximation for 2D input spaces
- 2D example (IRSN)

3 Conclusions and Future Works

4 Questions?

5 References

Table of Contents

1 Motivation

- Gaussian CDF example
- Constrained Kriging: Maatouk and Bay (2016)

2 Contributions

- Old contributions (Nice, 2017)
- Constrained maximum likelihood estimation (CMLE)
- Finite-dimensional approximation for 2D input spaces
- 2D example (IRSN)

3 Conclusions and Future Works

4 Questions?

5 References

References I

- Altmann, Y., McLaughlin, S., and Dobigeon, N. (2014). Sampling from a multivariate Gaussian distribution truncated on a simplex: A review. In *2014 IEEE Workshop on Statistical Signal Processing (SSP)*, pages 113–116.
- Anderes, E. (2010). On the consistent separation of scale and variance for Gaussian random fields. *Ann. Statist.*, 38(2):870–893.
- Azaïs, J.-M. and Wschebor, M. (2009). *Level sets and extrema of random processes and fields*. John Wiley & Sons.
- Azzimonti, D. and Ginsbourger, D. (2017). Estimating orthant probabilities of high dimensional Gaussian vectors with an application to set estimation (to appear). *Journal of Computational and Graphical Statistics*.
- Bachoc, F. (2013). Cross validation and maximum likelihood estimations of hyperparameters of Gaussian processes with model misspecification. *Computational Statistics & Data Analysis*, 66:55 – 69.
- Bachoc, F. (2014). Asymptotic analysis of the role of spatial sampling for covariance parameter estimation of gaussian processes. *Journal of Multivariate Analysis*, 125(Supplement C):1 – 35.
- Bay, X., Grammont, L., and Maatouk, H. (2016). Generalization of the Kimeldorf-Wahba correspondence for constrained interpolation. *Electronic journal of statistics*, 10(1):1580–1595.
- Bect, J., Bachoc, F., and Ginsbourger, D. (2016). A supermartingale approach to Gaussian process based sequential design of experiments. *ArXiv e-prints*.
- Betancourt, M. (2017). A Conceptual Introduction to Hamiltonian Monte Carlo. *ArXiv e-prints*.

References II

- Bishop, C. M. (2007). *tern Recognition And Machine Learning (Information Science And Statistics)*. Springer.
- Botev, Z. I. (2017). The normal law under linear restrictions: simulation and estimation via minimax tilting. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 79(1):125–148.
- Brogan, W. L. (1991). *Modern Control Theory (3rd Ed.)*. Prentice-Hall, Inc., Upper Saddle River, NJ, USA.
- Brooks, S., Gelman, A., Jones, G., and Meng, X. (2011). *Handbook of Markov Chain Monte Carlo*. Chapman & Hall/CRC Handbooks of Modern Statistical Methods. CRC Press.
- Cousin, A., Maatouk, H., and Rullière, D. (2016). Kriging of financial term-structures. *European Journal of Operational Research*, 255(2):631–648.
- Cramér, H. and Leadbetter, M. (1967). *Stationary and related stochastic processes: sample function properties and their applications*. Wiley series in probability and mathematical statistics. Tracts on probability and statistics. Wiley.
- Da Veiga, S. and Marrel, A. (2012). Gaussian process modeling with inequality constraints. *Annales de la faculté des sciences de Toulouse Mathématiques*, 21(3):529–555.
- Du, J., Zhang, H., and Mandrekar, V. S. (2009). Fixed-domain asymptotic properties of tapered maximum likelihood estimators. *Ann. Statist.*, 37(6A):3330–3361.
- Duane, S., Kennedy, A. D., Pendleton, B. J., and Roweth, D. (1987). Hybrid Monte Carlo. *Physics letters B*, 195(2):216–222.

References III

- Fernex, F., Heulers, L., Jacquet, O., Miss, J., and Richet, Y. (2005). The MORET 4B Monte Carlo code - New features to treat complex criticality systems. In *M&C international conference on mathematics and computation supercomputing, reactor physics and nuclear and biological application, Avignon, France*, volume 59.
- Genton, M. G. (2001). Classes of kernels for machine learning: A statistics perspective. *Journal of Machine Learning Research*, 2:299–312.
- Genz, A. (1992). Numerical computation of multivariate normal probabilities. *Journal of Computational and Graphical Statistics*, 1:141–150.
- Genz, A., Bretz, F., Miwa, T., and Mi, X. (2017). mvtnorm: Multivariate normal and t distributions. <https://cran.r-project.org/web/packages/mvtnorm/mvtnorm.pdf>. [Online; accessed 04-April-2017].
- Golchi, S., Bingham, D. R., Chipman, H., and Campbell, D. A. (2015). Monotone emulation of computer experiments. *SIAM/ASA Journal on Uncertainty Quantification*, 3(1):370–392.
- Gong, L. and Flegal, J. M. (2016). A practical sequential stopping rule for high-dimensional Markov Chain Monte Carlo. *Journal of Computational and Graphical Statistics*, 25(3):684–700.
- Gramacy, R. B. and Apley, D. W. (2015). Local Gaussian process approximation for large computer experiments. *Journal of Computational and Graphical Statistics*, 24(2):561–578.
- Hensman, J., Matthews, A. G. d. G., Filippone, M., and Ghahramani, Z. (2015). MCMC for variationally sparse Gaussian processes. In *NIPS 2015, 29th Annual Conference on Neural Information Processing Systems, December 7-12, 2015, Montreal, Quebec, Canada / Also on ArXiv, Montreal, CANADA*.

References IV

- Holsclaw, T., Sansó, B., Lee, H. K. H., Heitmann, K., Habib, S., Higdon, D., and Alam, U. (2013). Gaussian process modeling of derivative curves. *Technometrics*, 55(1):57–67.
- IAEA (2014). *Criticality Safety in the Handling of Fissile Material: Specific Safety Guide*. International Atomic Energy Agency (IAEA) Safety Standards Series No. SSG-27.
- Kallenberg, O. (2002). *Foundations of Modern Probability*. Probability and its Applications. Springer New York.
- Kim, Y., Kim, B., and Jang, W. (2010). Asymptotic properties of the maximum likelihood estimator for the proportional hazards model with doubly censored data. *Journal of Multivariate Analysis*, 101(6):1339 – 1351.
- Kocijan, J. (2016). *Modelling and Control of Dynamic Systems Using Gaussian Process Models*. Advances in Industrial Control. Springer International Publishing, 1 edition.
- Lan, S. and Shahbaba, B. (2016). *Sampling Constrained Probability Distributions Using Spherical Augmentation*, pages 25–71. Springer International Publishing, Cham.
- L'Ecuyer, P., Blanchet, J. H., Tuffin, B., and Glynn, P. W. (2010). Asymptotic robustness of estimators in rare-event simulation. *ACM Trans. Model. Comput. Simul.*, 20(1):6:1–6:41.
- Li, W. V. and Linde, W. (1999). Approximation, metric entropy and small ball estimates for Gaussian measures. *Ann. Probab.*, 27(3):1556–1578.
- Liu, J. S. (2008). *Monte Carlo Strategies in Scientific Computing*. Springer Publishing Company, Incorporated.
- Loh, W. L. (2005). Fixed-domain asymptotics for a subclass of Matérn-type Gaussian random fields. *Ann. Statist.*, 33(5):2344–2394.

References V

- Loh, W. L. and Lam, T. K. (2000). Estimating structured correlation matrices in smooth Gaussian random field models. *Ann. Statist.*, 28(3):880–904.
- López-Lopera, A. F., Bachoc, F., Durrande, N., and Roustant, O. (2017). Finite-dimensional Gaussian approximation with linear inequality constraints. *ArXiv e-prints*.
- Maatouk, H. and Bay, X. (2016). *A New Rejection Sampling Method for Truncated Multivariate Gaussian Random Variables Restricted to Convex Sets*, pages 521–530. Springer International Publishing, Cham.
- Maatouk, H. and Bay, X. (2017). Gaussian process emulators for computer experiments with inequality constraints. *Mathematical Geosciences*, 49(5):557–582.
- Maatouk, H., Bay, X., and Grammont, L. (2014). Simulation of Gaussian processes with interpolation and inequality constraints - A correspondence with optimal smoothing splines. In *GRF - Sim workshop : Simulation of Gaussian and related Random Fields*, Bern, Switzerland.
- Maatouk, H., Roustant, O., and Richet, Y. (2015). Cross-validation estimations of hyper-parameters of Gaussian processes with inequality constraints. *Procedia Environmental Sciences*, 27:38–44.
- Meyer, C. D. (2000). *Matrix Analysis and Applied Linear Algebra*. Society for Industrial and Applied Mathematics, Philadelphia, PA, USA.
- Murphy, K. P. (2012). *Machine Learning: A Probabilistic Perspective (Adaptive Computation And Machine Learning Series)*. The MIT Press.
- Neal, R. M. (1996). *Bayesian Learning for Neural Networks*. Springer-Verlag New York, Inc., Secaucus, NJ, USA.
- Neal, R. M. (2012). MCMC using Hamiltonian dynamics. *ArXiv e-prints*.

References VI

- Nickisch, H. and Rasmussen, C. (2008). Approximations for binary Gaussian process classification. *Journal of Machine Learning Research*, 9:2035–2078.
- Paciorek, C. J. and Schervish, M. J. (2004). Nonstationary covariance functions for Gaussian process regression. In *In Proc. of the Conf. on Neural Information Processing Systems*. MIT Press.
- Pakman, A. and Paninski, L. (2014). Exact Hamiltonian Monte Carlo for truncated multivariate Gaussians. *Journal of Computational and Graphical Statistics*, 23(2):518–542.
- Rasmussen, C. E. and Williams, C. K. I. (2005). *Gaussian Processes for Machine Learning (Adaptive Computation and Machine Learning)*. The MIT Press.
- Riihimäki, J. and Vehtari, A. (2010). Gaussian processes with monotonicity information. In *Journal of Machine Learning Research: Workshop and Conference Proceedings*, volume 9, pages 645–652.
- Ripley, B. D. (1987). *Stochastic Simulation*. John Wiley & Sons, Inc., New York, NY, USA.
- Robbins, H. and Monro, S. (1951). A stochastic approximation method. *Ann. Math. Statist.*, 22(3):400–407.
- Rogozhnikov, A. (2016). Brilliantly wrong thoughts on science and programming: Hamiltonian Monte Carlo explained. http://arogozhnikov.github.io/2016/12/19/markov_chain_monte_carlo.html. [Online; accessed 04-April-2017].
- Snelson, E. and Ghahramani, Z. (2006). Sparse Gaussian processes using pseudo-inputs. In Weiss, Y., Schölkopf, P. B., and Platt, J. C., editors, *Advances in Neural Information Processing Systems 18*, pages 1257–1264. MIT Press.

References VII

- Stein, M. (1999). *Interpolation of Spatial Data: Some Theory for Kriging*. Springer Series in Statistics. Springer New York.
- Taylor, J. and Benjamini, Y. (2017). RestrictedMVN: multivariate normal restricted by affine constraints. <https://cran.r-project.org/web/packages/restrictedMVN/index.html>. [Online; 02-Feb-2017].
- Tolvanen, V. (2014). Gaussian processes with monotonicity constraint for Big Data. Master's thesis, School of Electrical Engineering, Aalto University.
- Vats, D., Flegal, J. M., and Jones, G. L. (2017). Multivariate output analysis for Markov Chain Monte Carlo. *ArXiv e-prints*.
- Yadrenko, M. I. (1983). *Spectral theory of random fields*. Translation series in mathematics and engineering. Optimization Software, New York, NY. Transl. from the Russian.
- Ying, Z. (1993). Maximum likelihood estimation of parameters under a spatial sampling scheme. *Ann. Statist.*, 21(3):1567–1590.
- Yoo, E.-H. and Kyriakidis, P. C. (2006). Area-to-point kriging with inequality-type data. *Journal of Geographical Systems*, 8(4):357–390.
- Zhang, H. (2004). Inconsistent estimation and asymptotically equal interpolations in model-based geostatistics. *Journal of the American Statistical Association*, 99(465):250–261.