

# Finite-Dimensional Gaussian Approximation with Linear Inequality Constraints

Andrés F. López-Lopera<sup>1</sup>, François Bachoc<sup>2</sup>, Nicolas Durrande<sup>1,3</sup>, and  
Olivier Roustant<sup>1</sup>

<sup>1</sup>École des Mines de Saint-Étienne (EMSE), France.

<sup>2</sup>Institut de Mathématiques de Toulouse (IMT), France.

<sup>3</sup>PROWLER.io, Cambridge, UK.


This work is funded by the chair of applied mathematics OQUAIDO.

SIAM UQ 18

April 17, 2018

# Gaussian process models: motivation

**Target function:** is bounded and monotonic.




Unconstrained GP.


- true function
- predictive mean
- training points
- confidence intervals

# Gaussian process models: motivation

**Target function:** is bounded and monotonic.



Unconstrained GP.



Constrained GP.

- true function
- predictive mean
- training points
- confidence intervals

# Table of Contents

- 1 Constrained GPs under inequality conditions
  - Finite-dimensional Gaussian approximation
  - Constrained maximum likelihood estimation (cMLE)
- 2 Nuclear criticality dataset (IRSN)
- 3 Conclusions and Future Works
- 4 References

# Table of Contents


- 1 Constrained GPs under inequality conditions
  - Finite-dimensional Gaussian approximation
  - Constrained maximum likelihood estimation (cMLE)
- 2 Nuclear criticality dataset (IRSN)
- 3 Conclusions and Future Works
- 4 References

# Table of Contents

- 1 Constrained GPs under inequality conditions
  - Finite-dimensional Gaussian approximation
  - Constrained maximum likelihood estimation (cMLE)
- 2 Nuclear criticality dataset (IRSN)
- 3 Conclusions and Future Works
- 4 References

# Finite-dimensional Gaussian approximation

Finite representation: is also bounded and monotonic.




Imposing both inequality constraints on the knots is enough.

# Finite-dimensional Gaussian approximation

Let the finite-dimensional GP approximation be defined as

$$Y_m(x) = \sum_{j=1}^m \xi_j \phi_j(x), \quad \text{s.t.} \quad \begin{cases} Y_m(x_i) = y_i & \text{(interpolation conditions),} \\ Y_m \in \mathcal{E} & \text{(inequality conditions),} \end{cases}$$

where  $\boldsymbol{\xi} = [\xi_1, \dots, \xi_m]^\top \sim \mathcal{N}(\mathbf{0}, \boldsymbol{\Gamma})$ , with covariance matrix  $\boldsymbol{\Gamma}$  and  $\phi_j : [0, 1] \rightarrow \mathbb{R}$  are hat functions (Maatouk and Bay, 2017):





# Finite-dimensional Gaussian approximation

## Satisfying linear inequality constraints:

Since  $Y_m \in \mathcal{E} \Leftrightarrow \boldsymbol{\xi} \in \mathcal{C}$ , we consider  $\mathcal{C}$  is composed by

$$\mathcal{C} = \left\{ \mathbf{c} \in \mathbb{R}^m; \forall k = 1, \dots, q : \ell_k \leq \sum_{j=1}^m \lambda_{k,j} c_j \leq u_k \right\}.$$

Therefore, we have  $\boldsymbol{\Lambda}\boldsymbol{\xi} | \{\boldsymbol{\Phi}\boldsymbol{\xi} = \mathbf{y}\} \sim \mathcal{N}(\boldsymbol{\Lambda}\boldsymbol{\mu}, \boldsymbol{\Lambda}\boldsymbol{\Sigma}\boldsymbol{\Lambda}^\top)$  where

$$\boldsymbol{\mu} = \boldsymbol{\Gamma}\boldsymbol{\Phi}^\top[\boldsymbol{\Phi}\boldsymbol{\Gamma}\boldsymbol{\Phi}^\top]^{-1}\mathbf{y}, \quad \text{and} \quad \boldsymbol{\Sigma} = \boldsymbol{\Gamma} - \boldsymbol{\Gamma}\boldsymbol{\Phi}^\top[\boldsymbol{\Phi}\boldsymbol{\Gamma}\boldsymbol{\Phi}^\top]^{-1}\boldsymbol{\Phi}\boldsymbol{\Gamma}.$$

# Finite-dimensional Gaussian approximation

## Satisfying linear inequality constraints:

Since  $Y_m \in \mathcal{E} \Leftrightarrow \boldsymbol{\xi} \in \mathcal{C}$ , we consider  $\mathcal{C}$  is composed by

$$\mathcal{C} = \left\{ \mathbf{c} \in \mathbb{R}^m; \forall k = 1, \dots, q : \ell_k \leq \sum_{j=1}^m \lambda_{k,j} c_j \leq u_k \right\}.$$

Therefore, we have  $\boldsymbol{\Lambda}\boldsymbol{\xi} | \{\boldsymbol{\Phi}\boldsymbol{\xi} = \mathbf{y}\} \sim \mathcal{N}(\boldsymbol{\Lambda}\boldsymbol{\mu}, \boldsymbol{\Lambda}\boldsymbol{\Sigma}\boldsymbol{\Lambda}^\top)$  where

$$\boldsymbol{\mu} = \boldsymbol{\Gamma}\boldsymbol{\Phi}^\top[\boldsymbol{\Phi}\boldsymbol{\Gamma}\boldsymbol{\Phi}^\top]^{-1}\mathbf{y}, \quad \text{and} \quad \boldsymbol{\Sigma} = \boldsymbol{\Gamma} - \boldsymbol{\Gamma}\boldsymbol{\Phi}^\top[\boldsymbol{\Phi}\boldsymbol{\Gamma}\boldsymbol{\Phi}^\top]^{-1}\boldsymbol{\Phi}\boldsymbol{\Gamma}.$$

Posterior distribution (López-Lopera et al., 2017)

Then, the **posterior** follows a **truncated multinormal distribution**

$$\boldsymbol{\Lambda}\boldsymbol{\xi} | \{\boldsymbol{\Phi}\boldsymbol{\xi} = \mathbf{y}, l \leq \boldsymbol{\Lambda}\boldsymbol{\xi} \leq \mathbf{u}\} \sim \mathcal{TN}(\boldsymbol{\Lambda}\boldsymbol{\mu}, \boldsymbol{\Lambda}\boldsymbol{\Sigma}\boldsymbol{\Lambda}^\top, l, \mathbf{u}). \quad (1)$$

# Finite-dimensional Gaussian approximation

## Satisfying linear inequality constraints:

Since  $Y_m \in \mathcal{E} \Leftrightarrow \xi \in \mathcal{C}$ , we consider  $\mathcal{C}$  is composed by

$$\mathcal{C} = \left\{ \mathbf{c} \in \mathbb{R}^m; \forall k = 1, \dots, q : \ell_k \leq \sum_{j=1}^m \lambda_{k,j} c_j \leq u_k \right\}.$$

Therefore, we have  $\Lambda \xi | \{ \Phi \xi = \mathbf{y} \} \sim \mathcal{N}(\Lambda \mu, \Lambda \Sigma \Lambda^\top)$  where

$$\mu = \Gamma \Phi^\top [\Phi \Gamma \Phi^\top]^{-1} \mathbf{y}, \quad \text{and} \quad \Sigma = \Gamma - \Gamma \Phi^\top [\Phi \Gamma \Phi^\top]^{-1} \Phi \Gamma.$$

Posterior distribution (López-Lopera et al., 2017)

Then, the posterior follows a truncated multinormal distribution


$$\Lambda \xi | \{ \Phi \xi = \mathbf{y}, l \leq \Lambda \xi \leq \mathbf{u} \} \sim \mathcal{TN}(\Lambda \mu, \Lambda \Sigma \Lambda^\top, l, \mathbf{u}). \quad (1)$$

$\Rightarrow$  (1) can be approximated via Hamiltonian MC (Pakman and Paninski, 2014).

$\Rightarrow$  What about  $\Lambda, l, \mathbf{u}$  ?

# Finite-dimensional Gaussian approximation


## Defining boundedness constraints



$$\underbrace{\begin{bmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{bmatrix}}_{l_b} \preceq \underbrace{\begin{bmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & \cdots & 0 & 1 \end{bmatrix}}_{\Lambda_b} \underbrace{\begin{bmatrix} \xi_1 \\ \xi_2 \\ \xi_3 \\ \vdots \\ \xi_{m-1} \\ \xi_m \end{bmatrix}}_{\xi} \preceq \underbrace{\begin{bmatrix} 1 \\ 1 \\ 1 \\ \vdots \\ 1 \\ 1 \end{bmatrix}}_{u_b}$$

# Finite-dimensional Gaussian approximation


## Defining boundedness and monotonicity constraints



$$\underbrace{\begin{bmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{bmatrix}}_l \leq \underbrace{\begin{bmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ -1 & 1 & 0 & \cdots & 0 & 0 \\ 0 & -1 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & \cdots & -1 & 1 \\ 0 & 0 & 0 & \cdots & 0 & 1 \end{bmatrix}}_{\Lambda} \underbrace{\begin{bmatrix} \xi_1 \\ \xi_2 \\ \xi_3 \\ \vdots \\ \xi_{m-1} \\ \xi_m \end{bmatrix}}_{\xi} \leq \underbrace{\begin{bmatrix} 1 \\ \infty \\ \infty \\ \vdots \\ \infty \\ \infty \\ 1 \end{bmatrix}}_u$$

# Finite-dimensional Gaussian approximation


## Defining boundedness and monotonicity constraints



or simply,

$$\underbrace{\begin{bmatrix} l_b \\ l_m \end{bmatrix}}_l \leq \underbrace{\begin{bmatrix} \Lambda_b \\ \Lambda_m \end{bmatrix}}_{\Lambda} \boldsymbol{\xi} \leq \underbrace{\begin{bmatrix} u_b \\ u_m \end{bmatrix}}_u$$

# Examples on 2D input spaces



**Figure:** Examples of 2D Gaussian models with different types of constraints.

# Table of Contents

- 1 Constrained GPs under inequality conditions
  - Finite-dimensional Gaussian approximation
  - Constrained maximum likelihood estimation (cMLE)
- 2 Nuclear criticality dataset (IRSN)
- 3 Conclusions and Future Works
- 4 References



# Maximum likelihood (ML): asymptotic properties.

Let  $\mathcal{E}_\kappa$  be one of the following convex set of functions

$$\mathcal{E}_\kappa = \begin{cases} f : \mathbb{X} \rightarrow \mathbb{R}, f \text{ is } C^0 \text{ and } \forall \mathbf{x} \in \mathbb{X}, \ell \leq f(\mathbf{x}) \leq u & \text{if } \kappa = 0, \\ f : \mathbb{X} \rightarrow \mathbb{R}, f \text{ is } C^1 \text{ and } \forall \mathbf{x} \in \mathbb{X}, \forall i = 1, \dots, d, \frac{\partial}{\partial x_i} f(\mathbf{x}) \geq 0 & \text{if } \kappa = 1, \\ f : \mathbb{X} \rightarrow \mathbb{R}, f \text{ is } C^2 \text{ and } \forall \mathbf{x} \in \mathbb{X}, \frac{\partial^2}{\partial \mathbf{x}^2} f(\mathbf{x}) \text{ is a non-negative} & \text{if } \kappa = 2, \\ & \text{definite matrix} \end{cases}$$

which corresponds to boundedness, monotonicity, and convexity constraints. We will focus on the GP  $Y$  and the observation vector

$$\mathbf{Y}_n = [Y(x_1), \dots, Y(x_n)]^\top.$$

Let the unconstrained likelihood

$$\mathcal{L}_n(\boldsymbol{\theta}) = -\frac{1}{2} \log(\det(\mathbf{R}_{\boldsymbol{\theta}})) - \frac{1}{2} \mathbf{Y}_n^\top \mathbf{R}_{\boldsymbol{\theta}}^{-1} \mathbf{Y}_n - \frac{n}{2} \log 2\pi,$$

with  $\mathbf{R}_{\boldsymbol{\theta}} = (k_{\boldsymbol{\theta}}(x_i, x_j))_{1 \leq i, j \leq n}$ .

# Constrained maximum likelihood (CML)

## Proposition: asymptotic consistency of CML

Let  $P_{\theta}$  be the distribution of  $Y$  with covariance function  $k_{\theta}$ . Let

$$\mathcal{L}_{C,n}(\theta) = \mathcal{L}_n(\theta) + \log P_{\theta}(Y \in \mathcal{E}_{\kappa} | \mathbf{Y}_n) - \log P_{\theta}(Y \in \mathcal{E}_{\kappa}). \quad (\text{Constrained ML})$$

Assume that  $\forall \varepsilon > 0$  and  $\forall M < \infty$ , (Consistency of the unconditional ML)

$$P\left(\sup_{\|\theta - \theta^*\| \geq \varepsilon} (\mathcal{L}_n(\theta) - \mathcal{L}_n(\theta^*)) \geq -M\right) \xrightarrow{n \rightarrow \infty} 0.$$

Then, (Consistency of the conditional CML)

$$P\left(\sup_{\|\theta - \theta^*\| \geq \varepsilon} (\mathcal{L}_{C,n}(\theta) - \mathcal{L}_{C,n}(\theta^*)) \geq -M \mid Y \in \mathcal{E}_{\kappa}\right) \xrightarrow{n \rightarrow \infty} 0.$$


Consequently (Consistency of ML and CML estimators)

$$\operatorname{argmax}_{\theta \in \Theta} \mathcal{L}_n(\theta) \xrightarrow[n \rightarrow \infty]{P} \theta^*, \quad \text{and} \quad \operatorname{argmax}_{\theta \in \Theta} \mathcal{L}_{C,n}(\theta) \xrightarrow[n \rightarrow \infty]{P|Y \in \mathcal{E}_{\kappa}} \theta^*.$$

# Table of Contents

- 1 Constrained GPs under inequality conditions
  - Finite-dimensional Gaussian approximation
  - Constrained maximum likelihood estimation (cMLE)
- 2 Nuclear criticality dataset (IRSN)
- 3 Conclusions and Future Works
- 4 References

# Nuclear criticality dataset (IRSN)




Nuclear criticality safety assessments: IRSN's dataset.

$\Rightarrow k_{eff}$  is positive and non-decreasing.

# Nuclear criticality dataset (IRSN)

2D Gaussian models for interpolating the IRSN's dataset using  $n = 4$ .




(a) Standard GP+MLE (b) Constrained GP+MLE (c) Constrained GP+cMLE

× test data      • training points

# Nuclear criticality dataset (IRSN)

2D Gaussian models for interpolating the IRSN's dataset using  $n = 8$ .



(d) Standard GP+MLE    (e) Constrained GP+MLE    (f) Constrained GP+cMLE

× test data    • training points

Now, we repeat the procedure for 20 random LHDs, and we compute the  $Q^2$  and coverage accuracy (CA) criteria...

### Prediction accuracy

Let  $n_t$  be the number of test points,  $z_1, \dots, z_{n_t}$  and  $\hat{z}_1, \dots, \hat{z}_{n_t}$  the sets of test and predicted observations (respectively), then:

**$Q^2$  criterion:**



$$Q^2 = 1 - \frac{\sum_{i=1}^{n_t} (\hat{z}_i - z_i)^2}{\sum_{i=1}^{n_t} (\bar{z} - z_i)^2}, \quad (2)$$

where  $\bar{z}$  is the mean of the test data.  $\Rightarrow Q^2 \rightarrow 1 \checkmark$

**Coverage accuracy (CA) criterion:**

$$\text{CA}_{\pm\sigma} = \frac{1}{n_t} \sum_{i=1}^{n_t} \mathbb{1}_{z_i \in [\hat{z}_i - \hat{\sigma}_i, \hat{z}_i + \hat{\sigma}_i]} \quad (3)$$

where  $\hat{\sigma}_i$  are the predictive standard deviations.  $\Rightarrow \text{CA}_{\pm\sigma} \rightarrow 0.68 \checkmark$



Assessment of the models for interpolating the IRSN's dataset using different number of training points  $n$  and using twenty different Latin hypercube designs.

⇒ Unconstrained model was often outperformed by constrained ones.

⇒ MLE yields good tradeoff between prediction accuracy and computational cost.

⇒ cMLE provides more accurate confidence intervals.



# Table of Contents

- 1 Constrained GPs under inequality conditions
  - Finite-dimensional Gaussian approximation
  - Constrained maximum likelihood estimation (cMLE)
- 2 Nuclear criticality dataset (IRSN)
- 3 Conclusions and Future Works
- 4 References

# Conclusions and Future Works

## Conclusions

- We extended the approach from (Maatouk and Bay, 2017): [now it works for any linear inequality constraint in 1D or 2D](#).
  - We suggested [Hamiltonian MC](#) to approximate the posterior.
  - We proved the [consistency of the constrained maximum likelihood](#).
  - We implemented the R package: [lineqGPR](#) ([available in June!!](#)).
- ◆ A.F. López-Lopera, F. Bachoc, N. Durrande, and O. Roustant (2017). Finite-dimensional Gaussian approximation with linear inequality constraints. ArXiv e-prints.
- ◆ F. Bachoc, A. Lagnoux, and A.F. López-Lopera (2018). Maximum likelihood estimation for Gaussian processes under inequality constraints. ArXiv e-prints.

# Table of Contents

- 1 Constrained GPs under inequality conditions
  - Finite-dimensional Gaussian approximation
  - Constrained maximum likelihood estimation (cMLE)
- 2 Nuclear criticality dataset (IRSN)
- 3 Conclusions and Future Works
- 4 References

# References I

- Bachoc, F. (2014). Asymptotic analysis of the role of spatial sampling for covariance parameter estimation of Gaussian processes. *Journal of Multivariate Analysis*, 125(Supplement C):1 – 35.
- Bachoc, F., Lagnoux, A., and López-Lopera, A. F. (2018). Maximum likelihood estimation for Gaussian processes under inequality constraints. *ArXiv e-prints*.
- Bay, X., Grammont, L., and Maatouk, H. (2016). Generalization of the Kimeldorf-Wahba correspondence for constrained interpolation. *Electronic journal of statistics*, 10(1):1580–1595.
- Botev, Z. I. (2017). The normal law under linear restrictions: simulation and estimation via minimax tilting. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 79(1):125–148.
- Genz, A. (1992). Numerical computation of multivariate normal probabilities. *Journal of Computational and Graphical Statistics*, 1:141–150.
- López-Lopera, A. F., Bachoc, F., Durrande, N., and Roustant, O. (2017). Finite-dimensional Gaussian approximation with linear inequality constraints. *ArXiv e-prints*.
- Maatouk, H. and Bay, X. (2017). Gaussian process emulators for computer experiments with inequality constraints. *Mathematical Geosciences*, 49(5):557–582.
- Pakman, A. and Paninski, L. (2014). Exact Hamiltonian Monte Carlo for truncated multivariate Gaussians. *Journal of Computational and Graphical Statistics*, 23(2):518–542.
- Rasmussen, C. E. and Williams, C. K. I. (2005). *Gaussian Processes for Machine Learning (Adaptive Computation and Machine Learning)*. The MIT Press.
- Stein, M. (1999). *Interpolation of Spatial Data: Some Theory for Kriging*. Springer Series in Statistics. Springer New York.