

# Vietoris-Rips Complexes of Regular Polygons

Samir Chowdhury  
The Ohio State University

Adam Jaffe  
Stanford University

Joint work with  
Henry Adams (Colorado State University)  
Bonginkosi Sibanda (Brown University)

January 13, 2018

AMS Special Session on Topological Data Analysis  
JMM 2018

# Setup and overview

## Definition

For metric space  $(X, d)$  and scale  $r \geq 0$ , the *Vietoris–Rips simplicial complex*  $\mathbf{VR}_{<}(X; r)$  is the set of all finite  $\sigma \subseteq X$  with  $\text{diam}(\sigma) < r$ .

## Definition

For metric space  $(X, d)$  and scale  $r \geq 0$ , the *Vietoris–Rips simplicial complex*  $\mathbf{VR}_{\leq}(X; r)$  is the set of all finite  $\sigma \subseteq X$  with  $\text{diam}(\sigma) \leq r$ .

## Remark

The *Vietoris–Rips simplicial complex* can be fully determined by the underlying graph of its one skeleton, i.e the graph made by the zero and one dimensional simplices.





# Setup and overview

Theorem (Chazal, de Silva, Oudot, 2013)

*Suppose  $X, M$  are totally bounded metric spaces. Then for any  $k \geq 0$ ,*

$$d_B(dgm_k^{VR}(X), dgm_k^{VR}(M)) \leq 2d_{GH}(X, M)$$




One application:  $X$  a uniform sample from a manifold  $M$ .

- Problem:  $dgm_k^{VR}(M)$  is known for very few manifolds!
- Past work: circle (Adamaszek, Adams 2017), ellipses of small eccentricity (Adamaszek, Adams, Reddy 2018).

# Regular Polygons

## Definition

Given an integer  $n \geq 3$ , let the *regular  $n$ -gon*  $P_n \subseteq \mathbb{R}^2$  be a set of  $n$  points equally spaced on  $S^1$ , with line segments connecting adjacent points together. We endow  $P_n$  with the Euclidean metric of  $\mathbb{R}^2$ .



# Problem statement and strategy

- We want to describe the homotopy types and persistent homology of  $\text{VR}(P_n; r)$ .
- Method of cyclic graphs (Adamaszek et al 2016) has been successful in the circle and ellipse case.
- First we quantify scales parameters for which  $\text{VR}(P_n; r)$  supports cyclic graphs.
- Theorem of Adamaszek, Adams, and Reddy asserts that  $\text{VR}(P_n; r) \simeq S^{2\ell+1}$  or  $\text{VR}(P_n; r) \simeq \bigvee^{P+F-1} S^{2\ell}$ , depending on an invariant called the *winding fraction* of cyclic graphs supported on  $\text{VR}(P_n; r)$ . Here  $P, F$  are integers depending on the geometry of  $P_n$  that we explain later.
- **Main result:** We characterize the scale parameters  $r$  at which  $\text{VR}(P_n; r)$  is homotopy equivalent to an odd sphere or a wedge of  $P + F - 1$  even spheres. In the latter case, we precisely quantify  $P$  and  $F$ .

# Main Result


## Theorem

*For fixed  $n$ , we have sequences of reals  $\{s_{n,\ell}\}$  and  $\{t_{n,\ell}\}$  that correspond to the first and last scale parameters for which an equilateral  $(2\ell + 1)$ -star can be inscribed within  $P_n$ . Then:*


$$\begin{aligned}\mathbf{VR}_{<}(P_n; r) &\simeq \begin{cases} V^{q-1} S^{2\ell} & \text{when } s_{n,\ell} < r \leq t_{n,\ell} \\ S^{2\ell+1} & \text{when } t_{n,\ell} < r \leq s_{n,\ell+1} \end{cases} \\ \mathbf{VR}_{\leq}(P_n; r) &\simeq \begin{cases} V^{3q-1} S^{2\ell} & \text{when } s_{n,\ell} < r < t_{n,\ell} \\ S^{2\ell+1} & \text{when } t_{n,\ell} < r < s_{n,\ell+1}, \end{cases}\end{aligned}$$

*where  $q = n/\gcd(n, 2\ell + 1)$ . Furthermore, all of the above homological features are persistent, except for  $2q$  copies of  $S^{2\ell}$  during the even sphere regimes for  $\leq$ .*

# Main Result: Example




$$\mathbf{VR}_{<}(P_{15}; r)$$




$$\mathbf{VR}_{\leq}(P_{15}; r)$$



# Main Result: Example



$$\mathbf{VR}_{<}(P_{15}; r)$$



$$\mathbf{VR}_{\leq}(P_{15}; r)$$

Why do we get homology above dimension 1?








Figure:  $\text{VR}_{\leq}(6 \text{ points}; \frac{1}{3}) \simeq S^2$

# Intuition














Figure:  $\text{VR}_{\leq}(9 \text{ points}; \frac{1}{3}) \simeq V^2 S^2$

# Intuition








# Intuition





# Intuition











# Cyclic Graphs

## Definition

A directed graph  $G$  is *cyclic* if its vertices can be placed in a cyclic order such that, whenever there is an edge  $v \rightarrow u$ , then there are also edges  $v \rightarrow w \rightarrow u$  for all  $v \prec w \prec u$ .

## Definition

For a cyclic graph  $G$  and a vertex  $v$ , define  $f(v)$  to be the clockwise-most point  $u$  such that there exists an edge  $v \rightarrow u$ .

## Definition

The *winding fraction* of a cyclic graph  $G$  is

$$\text{wf}(G) = \sup \left\{ \frac{\omega}{k} \mid \begin{array}{l} G \text{ contains an } f\text{-periodic orbit of} \\ \text{length } k \text{ which "winds" } \omega \text{ times} \\ \text{around the center of } G. \end{array} \right\}.$$

# Cyclic Graphs




Figure: cyclic graphs of winding fraction  $\frac{1}{4}$  (left) and  $\frac{2}{5}$  (right).



Every vertex in a cyclic graph can be classified as exactly one of *fast*, *slow*, or *periodic* (to be defined later).

## Theorem (Adamaszek, Adams, Reddy 2018)

Let  $G$  be a cyclic graph with  $P$  periodic orbits and  $F$  invariant sets of fast points. Then:

- If  $\frac{\ell}{2\ell+1} < \text{wf}(G) \leq \frac{\ell+1}{2\ell+3}$  for some integer  $\ell \geq 0$ , then  $\text{Cl}(G) \simeq S^{2\ell+1}$ .
- If  $\text{wf}(G) = \frac{\ell}{2\ell+1}$ , then  $\text{Cl}(G) \simeq V^{P+F-1} S^{2\ell}$ .

# Geometric Lemmas for Regular Polygons

## Question

For which scale parameters  $r > 0$  does  $\mathbf{VR}(P_n; r)$  form a cyclic graph?

## Answer

The graph  $\mathbf{VR}(P_n; r)$  is cyclic up to the scale parameter shown.





## Remark

Since  $r_3 = 0$ , we conclude that  $\mathbf{VR}(P_3; r)$  is not a cyclic graph for any  $r > 0$ .

# Geometric Lemmas for Regular Polygons

## Definition

In a cyclic graph  $G$ , an  $f$ -periodic orbit which has length  $2\ell + 1$  and which “winds”  $\ell$  times around the center of  $G$  is called an *inscribed equilateral  $(2\ell + 1)$ -pointed star*, or simply a  $(2\ell + 1)$ -star.



# Geometric Lemmas for Regular Polygons

## Definition

Let  $s_{n,\ell}$  and  $t_{n,\ell}$  be the smallest and largest scale parameters  $r > 0$  for which a  $(2\ell + 1)$ -star can be inscribed into  $P_n$ .

## Remark

The winding fraction of  $\mathbf{VR}(P_n; r)$  equals  $\frac{\ell}{2\ell+1}$  for all scales  $r \in (s_{n,\ell}, t_{n,\ell})$ .

# Geometric Lemmas for Regular Polygons

## Lemma

*For any integers  $\ell \geq 1$  and  $n \geq 3$ , there exists a unique  $(2\ell + 1)$ -star inscribed in  $P_n$  containing any given basepoint if and only if  $n \geq 4\ell + 2$ .*

## Definition

For  $\ell \geq 1$ ,  $n \geq 4\ell + 2$ , and  $x \in P_n$ , denote the unique inscribed  $(2\ell + 1)$ -star containing  $x$  by  $S_{2\ell+1}(x)$ , and its side length by  $s_{2\ell+1}(x)$ .

## Lemma

*The function  $s_{2\ell+1} : P_n \rightarrow \mathbb{R}$  is continuous.*

# Geometric Lemmas for Regular Polygons




Figure:  $n = 6$  and  $2\ell + 1 = 3$





Figure:  $n = 11$  and  $2\ell + 1 = 5$

# Geometric Lemmas for Regular Polygons

## Definition

Given an integer  $n \geq 3$  and a real  $r > 0$ , let  $\ell \geq 0$  be the largest integer satisfying  $n \geq 4\ell + 2$ . Then any point  $x \in P_n$  can be classified as one of:

- *fast*, if  $s_{2\ell+1}(x) < r$
- *slow*, if  $s_{2\ell+1}(x) > r$
- *periodic*, if  $s_{2\ell+1}(x) = r$



## Definition

The integer  $F$ , the number of invariant sets of fast points in  $\mathbf{VR}(P_n; r)$ , is equal to the number of connected components in  $s_{2\ell+1}^{-1}((-\infty, r))$ , divided by  $2\ell + 1$ .

## Definition

The integer  $P$ , the number of periodic orbits in  $\mathbf{VR}(P_n; r)$ , is equal to the cardinality of  $s_{2\ell+1}^{-1}(\{r\})$ , divided by  $2\ell + 1$ .

# Geometric Lemmas for Regular Polygons




Figure:  $n = 6$  and  $2\ell + 1 = 3$



# Geometric Lemmas for Regular Polygons




Figure:  $n = 7$  and  $2\ell + 1 = 3$

# Geometric Lemmas for Regular Polygons




Figure:  $n = 10$  and  $2\ell + 1 = 5$

# Geometric Lemmas for Regular Polygons




Figure:  $n = 11$  and  $2\ell + 1 = 5$

# Geometric Lemmas for Regular Polygons

## Question

How many distinct equilateral  $(2\ell + 1)$ -stars of side length  $r$  can be inscribed into  $P_n$ ?

## Answer

The number of equilateral  $(2\ell + 1)$ -stars of side length  $r$  that can be inscribed into  $P_n$  is equal to:

$$\begin{cases} n/\gcd(n, 2\ell + 1) & \text{if } r = s_{n,\ell} \text{ or } t_{n,\ell} \\ 2n/\gcd(n, 2\ell + 1) & \text{if } s_{n,\ell} < r < t_{n,\ell} \\ 0 & \text{otherwise} \end{cases}$$

# Main Result

## Theorem


For  $r \in (0, r_n)$  we have:

$$\mathbf{VR}_{<}(P_n; r) \simeq \begin{cases} V^{q-1} S^{2\ell} & \text{when } s_{n,\ell} < r \leq t_{n,\ell} \\ S^{2\ell+1} & \text{when } t_{n,\ell} < r \leq s_{n,\ell+1} \end{cases}$$
$$\mathbf{VR}_{\leq}(P_n; r) \simeq \begin{cases} V^{3q-1} S^{2\ell} & \text{when } s_{n,\ell} < r < t_{n,\ell} \\ S^{2\ell+1} & \text{when } t_{n,\ell} < r < s_{n,\ell+1}, \end{cases}$$


where  $q = n/\gcd(n, 2\ell + 1)$ . Furthermore,

- For  $s_{n,\ell} < r < \tilde{r} \leq t_{n,\ell}$  or  $t_{n,\ell} < r < \tilde{r} \leq s_{n,\ell+1}$ , inclusion  $\mathbf{VR}_{<}(P_n; r) \hookrightarrow \mathbf{VR}_{<}(P_n; \tilde{r})$  is a homotopy equivalence.
- For  $t_{n,\ell} < r < \tilde{r} < s_{n,\ell+1}$ , inclusion  $\mathbf{VR}_{\leq}(P_n; r) \hookrightarrow \mathbf{VR}_{\leq}(P_n; \tilde{r})$  is a homotopy equivalence.
- For  $s_{n,\ell} \leq r < \tilde{r} \leq t_{n,\ell}$ , inclusion  $\mathbf{VR}_{\leq}(P_n; r) \hookrightarrow \mathbf{VR}_{\leq}(P_n; \tilde{r})$  induces a rank  $q - 1$  map on  $2\ell$ -dimensional homology  $H_{2\ell}(-; \mathbb{F})$  for any field  $\mathbb{F}$ .

# Main Result: Example




$$\mathbf{VR}_{<}(P_{15}; r)$$



$$\mathbf{VR}_{\leq}(P_{15}; r)$$

# Future Work

- Finish paper and post to arXiv



Thank you!