

First Order Jet Bundles.

Ángel Blasco Muñoz

Escuela Internacional de Doctorado, Universidad Nacional de Educación a Distancia
(UNED), España.

16 de febrero de 2026



- [1] D. J. Saunders. *The Geometry of Jet Bundles*. London Mathematical Society Lecture Note Series. Cambridge: Cambridge University Press, 1989. DOI: [10.1017/CB09780511526411](https://doi.org/10.1017/CB09780511526411).

Contents.

1 First Order Jets.

2 Prolongations of Morphisms.

Contents.

1 First Order Jets.

2 Prolongations of Morphisms.

First Order Jets.

For this section and the following, please refer to [1] for further details.

First Order Jets.

Let (E, π, M) be a bundle, and let $p \in M$.

Definition

Define the local sections $\phi, \psi \in \Gamma_p(\pi)$ to be 1-equivalent at p if $\phi(p) = \psi(p)$ and if, in some adapted coordinate systems (x^i, u^α) around $\phi(p)$,

$$\frac{\partial \phi^\alpha}{\partial x^i} \Big|_p = \frac{\partial \psi^\alpha}{\partial x^i} \Big|_p$$

First Order Jets.

Let (E, π, M) be a bundle, and let $p \in M$.

Definition

Define the local sections $\phi, \psi \in \Gamma_p(\pi)$ to be 1-equivalent at p if $\phi(p) = \psi(p)$ and if, in some adapted coordinate systems (x^i, u^α) around $\phi(p)$,

$$\frac{\partial \phi^\alpha}{\partial x^i} \Big|_p = \frac{\partial \psi^\alpha}{\partial x^i} \Big|_p$$

Definition

The equivalence class containing ϕ is called *1-jet of ϕ at p* and is denoted $j_p^1 \phi$.

Definition

The *first jet manifold* of π is the set:

$$\{ j_p^1 \phi : p \in M, \phi \in \Gamma_p \pi \}$$

and is denoted $J^1 \pi$

Definition

The *first jet manifold* of π is the set:

$$\{ j_p^1 \phi : p \in M, \phi \in \Gamma_p \pi \}$$

and is denoted $J^1 \pi$

$$\pi_1 : J^1 \pi \longrightarrow M$$

$$j_p^1 \phi \longmapsto p$$

$$\pi_{1,0} : J^1 \pi \longrightarrow E$$

$$j_p^1 \phi \longmapsto \phi(p)$$

First Order Jets.

Let (U, u) be an adapted coordinate system on E , where $u = (x^i, u^\alpha)$.

Definition

The *induced coordinate system* (U^1, u^1) on $J^1\pi$ is defined by

$$U^1 = \{j_p^1\phi : \phi(p) \in U\}$$

$$u^1 = (x^i, u^\alpha, u_i^\alpha)$$

First Order Jets.

Let (U, u) be an adapted coordinate system on E , where $u = (x^i, u^\alpha)$.

Definition

The *induced coordinate system* (U^1, u^1) on $J^1\pi$ is defined by

$$U^1 = \{j_p^1\phi : \phi(p) \in U\}$$

$$u^1 = (x^i, u^\alpha, u_i^\alpha)$$

$$x^i(j_p^1\phi) = x^i(p)$$

$$u^\alpha(j_p^1\phi) = u^\alpha(\phi(p))$$

$$u_i^\alpha(j_p^1\phi) = \frac{\partial \phi^\alpha}{\partial x^i} \Big|_p$$

Contents.

1 First Order Jets.

2 Prolongations of Morphisms.

Prolongations of Morphisms.

Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

END.