

Abstract Algebra

Notes - Year 1, Semester 2

Angel Cervera Roldan
21319203

Contents

Fields	3
Vector Spaces	4
Subspaces	4

Fields

A field is a set F containing at least two elements, along with two operations:

- $+: F \times F \rightarrow F$

- $\cdot: F \times F \rightarrow F$

that satisfies the following axioms:

1. $a + b = b + a$, and $a \cdot b = b \cdot a \quad \forall a, b \in F$
2. $(a + b) + c = a + (b + c)$, and $(a \cdot b) \cdot c = a \cdot (b \cdot c) \quad \forall a, b \in F$
3. $\exists 0 \in F$ such that $a + 0 = a \quad \forall a \in F$
4. $\exists 1 \in F$ such that $a \cdot 1 = a \quad \forall a \in F$, where $0 \neq 1$
5. $\forall a \in F$, $\exists(-a)$ such that $a + (-a) = 0$
6. $\forall a \in F - \{0\}$, $\exists(a^{-1})$ such that $a \cdot (-a) = 1$
7. $a \cdot (b + c) = a \cdot b + a \cdot c \quad \forall a, b, c \in F$

Theorem

\mathbb{Z}_p is the set of integers mod p , and it is a field iff p is a prime.

From the axioms, $\forall a, b, c \in F$ the following can be proven

1. $a + b = a + c \implies b = c$
2. $a \neq 0, ab = ac \implies b = c$
3. $-(-a) = a$
4. $a \cdot 0 = 0$

Vector Spaces

Let F be a field, a vector space over F is a non-empty set V along with two operations:

- $+: V \times V \rightarrow V$
- $\cdot: F \times V \rightarrow V$

that satisfy the following axioms:

1. $u + (v + w) = (u + v) + w \quad \forall u, v, w \in V$
2. $u + v = v + u \quad \forall u, v \in V$
3. $\exists 0 \in V$ such that $0 + v = v \quad \forall v \in V$
4. $\forall u \in V, \exists -u \in V$ such that $u + (-u) = 0$
5. $\alpha(\beta u) = (\alpha\beta)u \quad \forall \alpha, \beta \in F, u \in V$
6. $\alpha(u + v) = \alpha u + \alpha v \quad \forall \alpha \in F, u, v \in V$
7. $\exists 1 \in F$ such that $\forall u \in V \quad 1u = u$

Subspaces

If V is a vector space, and $B \subset V$ is also a vector space, then we say that B is a subspace of V .

Lemma

A non-empty subset $S \subseteq V$ is a vector space iff

- $u, s \in S \implies u + v \in S$
- $u \in S, \lambda \in F \implies \lambda u \in S$

To prove the last result, the key steps are to show that the zero vector $0 \in S$ and that for any $u \in S, (-1)u = -u \in S$. When S is a subset of a vector space, to verify that it is a subspace, we need to check that it is closed under the two operations on V .