CPSC-354 Report

Angelina House
Chapman University

November 16, 2025

Abstract

This report documents my learning and progress throughout the Fall 2025 semester in CPSC-354: Programming Languages.

Contents

1 Introduction

This report chronicles my work in CPSC-354: Programming Languages during Fall 2025. Each weekly subsection contains restated homework prompts, fully typeset solutions, and a brief reflection or question that ties the technical material to broader themes.

2 Week by Week

2.1 Week 1

Homework

Discuss the MU puzzle from *Gödel, Escher, Bach* (Chapter 1; see course PDF pp. 41–42). Does the string `MU` have a derivation in the MIU formal system starting from `MI`? Why or why not?

The MU (MIU) System Recap

We begin with the single axiom string `MI`. The system has four production rules:

1. If a string ends in `I`, append a `U` at the end.
2. If the string has the form `Mx`, duplicate everything after `M`: `Mx` $\rightarrow$ `Mxx`.
3. Replace any occurrence of `III` by `U`.
4. Delete any occurrence of `UU`.

From `MI`, we can derive many strings by applying the rules:

`MI`(2)`MII`(2)`MIIII`(3)`MUI`.

This illustrates how the rules expand and compress substrings, but it remains unclear how to end with exactly `MU`.

Let $I(s)$ be the number of `I`'s in a string s. Initially, $I(\texttt{MI}) = 1$, i.e., 1 (mod 3). Examine how each rule affects I modulo 3:

- Rule 1 adds a `U`; I is unchanged.
- Rule 2 doubles the count of `I`'s: $I \mapsto 2I$. Thus if $I \equiv 1 \pmod 3$ then $2I \equiv 2 \pmod 3$.
- Rule 3 subtracts three `I`'s (`III` $\to$ `U`); the residue modulo 3 is unchanged.
- Rule 4 deletes `U`'s only; I is unchanged.

Therefore, starting from $I \equiv 1 \pmod 3$, any derivable string keeps $I \equiv 1$ or $I \equiv 2 \pmod 3$, but never $0 \pmod 3$. The target `MU` has $I(\texttt{MU}) = 0$ ($0 \pmod 3$), so it is *not* derivable. Hence, the MU puzzle has no solution.

The puzzle shows how stepping outside a formal system and using an invariant (here, the count of `I`'s modulo 3) can decisively answer a reachability question that seems open-ended from within the rules.

Reading Summary & Question

Summary. Hofstadter introduces formal systems via the MIU puzzle and contrasts *inside-the-system* rule manipulation with *meta-level* reasoning using invariants. This frames later topics on meaning, proof, and limitations of formal systems.

Question. Are there any situations where the MU puzzle could be true, and how much would we have to change to get to that point?

2.2 Week 2

Homework

Using the course notes on abstract reduction systems (ARS), determine for each of the seven given ARSs whether it is *terminating*, *confluent*, and whether it *has unique normal forms (UN)*. Then, fill the grid of the eight possible (Confluent, Terminating, Has UN) combinations with examples or explain why some combinations are impossible.

Quick Definitions

- **Terminating:** there is no infinite chain $x_0 \to x_1 \to x_2 \to \cdots$.
- **Confluent:** if $x \to^* y$ and $x \to^* z$, then y and z are joinable.
- **Has unique normal forms (UN):** every element has a (and in fact the) unique normal form.

Seven ARSs: Diagrams & Properties

Below, each ARS is shown as a tiny diagram (compiled with `tikz-cd`) followed by its properties and a one–line justification.

1) $A = \varnothing$**.**

(empty set, no nodes or arrows)

Props: Terminating ✓, Confluent ✓, UN ✓. *Reason:* All properties hold vacuously.

2) $A = \{a\},\ R = \varnothing$**.**

a


Props: Terminating ✓, Confluent ✓, UN ✓. *Reason:* No arrows; a is a (unique) normal form.

3) $A = \{a\},\ R = \{(a, a)\}$**.**



Props: Terminating ×, Confluent ✓, UN ×. *Reason:* Self–loop gives infinite chain; only one object so forks (if any) are trivially joinable; no normal form exists.

4) $A = \{a, b, c\},\ R = \{(a, b), (a, c)\}$**.**




Props: Terminating ✓, Confluent ×, UN ×. *Reason:* From a reach distinct NFs b and c that are not joinable.

5) $A = \{a, b\},\ R = \{(a, a), (a, b)\}$**.**




Props: Terminating ×, Confluent ✓, UN ✓. *Reason:* There exists an infinite chain via the loop on a, but all reductions from a can join at b, and every element's normal form is uniquely b.

6) $A = \{a, b, c\},\ R = \{(a, b), (b, b), (a, c)\}$**.**



Props: Terminating ×, Confluent ×, UN ×. *Reason:* Loop at b breaks termination; from a you can reach b (non–NF forever) or c (an NF), so confluence and UN fail.

7) $A = \{a, b, c\},\ R = \{(a, b), (b, b), (a, c), (c, c)\}$**.**



Props: Terminating ×, Confluent ×, UN ×. *Reason:* Loops at b and c; from a diverge to non–joinable loops; no normal forms.

Summary Table (Seven ARSs)

#	Terminating	Confluent	Has UN	Quick reason
1	✓	✓	✓	Empty set: vacuous
2	✓	✓	✓	Single node, no edges
3	×	✓	×	Self–loop; no NF
4	✓	×	×	$a \to b$ or $a \to c$ (two NFs)
5	×	✓	✓	Loop at a but unique NF b
6	×	×	×	Loop at b; $a \to b$ vs $a \to c$
7	×	×	×	Loops at b, c; $a \to b$ vs $a \to c$

Eight Combinations Grid

Confluent	Terminating	Has UN	Example or reason
T	T	T	Example #2 (or #1)
T	T	F	**Impossible:** Confluence+termination ⇒ unique NF
T	F	T	Example #5
T	F	F	Example #3
F	T	T	**Impossible:** UN in a terminating system ⇒ confluent
F	T	F	Example #4
F	F	T	**Impossible:** UN (for all elements) ⇒ confluent
F	F	F	Example #6 or #7

Reflection & Question

Question. What is a real life example of something that has two possible end states like in ARS #4 where a can go to either b or c?

2.3 Week 3

Homework

Exercise 5 — Analysis

Rules:

$$\mathtt{ab} \to \mathtt{ba}, \quad \mathtt{ba} \to \mathtt{ab}, \quad \mathtt{aa} \to \varepsilon.$$

Sample reductions.

$$\mathtt{abba} \;\Rightarrow\; \mathtt{baba\,ba} \to \mathtt{ab\,bbaa\,aa} \to \varepsilon\,\mathtt{bb}.$$

$$\mathtt{bababa} \;\Rightarrow\; \mathtt{aaabbb\,aa} \to \varepsilon\,\mathtt{abbb}.$$

Non-termination. Swaps are bidirectional, so we can permute letters indefinitely; the ARS is not terminating.

Equivalence classes & invariant. Swaps preserve counts of letters. The rule $\mathtt{aa} \to \varepsilon$ preserves $\#a$ (mod 2). Thus the complete invariant is

$$\big(\#\mathtt{b},\ \#\mathtt{a} \bmod 2\big).$$

Normal forms via terminating variant. If we orient swaps one way (e.g. $\mathtt{ba} \to \mathtt{ab}$) and keep $\mathtt{aa} \to \varepsilon$, then every word terminates in a unique normal form:

$$\mathtt{b}^k \quad (\#a \text{ even}), \qquad \mathtt{a\,b}^k \quad (\#a \text{ odd}),$$

where $k = \#\mathtt{b}$.

Exercise 5b — Analysis

Rules (variant):

$$\mathtt{ab} \to \mathtt{ba}, \quad \mathtt{ba} \to \mathtt{ab}, \quad \mathtt{aa} \to \mathtt{a}.$$

Non-termination. Still non-terminating because of two-way swaps.

Equivalence classes & invariant. Swaps preserve #b. The rule aa→ a collapses any number of a's to exactly one (and in the equivalence closure we can also expand). Thus classes are determined by

$$\big(\#\mathtt{b},\ \mathbf{1}_{\{\#a\geq 1\}}\big),$$

i.e. the number of bs and whether there is at least one a.

Normal forms via terminating variant. If we orient ba→ab and keep aa→ a, $then the unique normal forms are$: $\mathtt{b}^k \quad (\#a = 0), \qquad \mathtt{a}\,\mathtt{b}^k \quad (\#a \geq 1)$, with $k = \#\mathtt{b}$.

Reflection & Question

Question. In example 5b every word with at least one a collapses to the same class. Does this mean aaa and a are always equivalent in both directions or only in the equivalence closure?

2.4 Week 4

Homework

Using a measure function (from the termination notes), prove termination of (1) Euclid's algorithm for $\gcd(a, b)$ and (2) a standard top-down merge sort.

HW 4.1 — Euclid's Algorithm Terminates

Algorithm.

```
while b != 0:
    temp = b
    b = a mod b
    a = temp
return a
```

Assumptions. $a, b \in \mathbb{N}$ and the loop guard is first checked with $b \geq 0$.

Measure. $\phi(a, b) = b \in \mathbb{N}$.

Proof. One loop step maps (a, b) to $(b,\ a \bmod b)$. If $b \neq 0$, then $0 \leq a \bmod b < b$, hence

$$\phi(b,\ a \bmod b) = a \bmod b \ <\ b \ =\ \phi(a, b).$$

Thus ϕ strictly decreases at each taken iteration and always stays in $\mathbb{N}$. Since $>$ on $\mathbb{N}$ is well-founded, there is no infinite strictly decreasing chain; the loop terminates when $b = 0$.

HW 4.2 — Merge Sort Terminates

Code (top-down).

```
function merge_sort(arr, left, right):
    if left >= right: return
    mid = (left + right) // 2   // integer division
    merge_sort(arr, left, mid)
    merge_sort(arr, mid+1, right)
    merge(arr, left, mid, right)
```

Measure. For a call on interval $[left, right]$, let

$$\phi(left, right) \ =\ right - left + 1 \in \mathbb{N}.$$

Proof. If $left \geq right$, the function returns immediately. Otherwise $left < right$ and with integer division $left \leq mid < right$. The recursive calls are on $[left, mid]$ and $[mid{+}1, right]$ with measures

$$\phi(left, mid) = mid - left + 1 \;<\; right - left + 1 = \phi(left, right),$$

$$\phi(mid{+}1, right) = right - mid \;<\; right - left + 1 = \phi(left, right).$$

Hence each recursive call strictly decreases ϕ. Because ϕ ranges over $\mathbb{N}$ and decreases along recursion, there is no infinite descent; all recursion bottoms out and the algorithm terminates. (No measure is needed for `merge`, which is non-recursive.)

Reflection & Question

Question. What's an example of a "bad" measure function that wouldn't actually prove termination?

2.5 Week 5

Homework

Evaluate

$$(\lambda f.\, \lambda x.\, f(f(x)))\ (\lambda f.\, \lambda x.\, f(f(f\, x))).$$

Solution

Let $A \equiv \lambda f.\lambda x.\, f(f(x))$ and $B \equiv \lambda f.\lambda x.\, f(f(f\, x))$.

Step 1. Apply A to B:

$$A\, B \;\to_\beta\; \lambda x.\, B\,(B\, x).$$

Step 2. α-rename inside B as $B \equiv \lambda f.\lambda y.\, f(f(f\, y))$ and compute:

$$B\, x \;=\; (\lambda f.\lambda y.\, f(f(f\, y)))\, x \;\to_\beta\; \lambda y.\, x(x(x\, y)).$$

Write $G \equiv \lambda y.\, x(x(x\, y))$. Then

$$\lambda x.\, B\,(B\, x) \;=\; \lambda x.\, B\, G \;\to_\beta\; \lambda x.\, \lambda y.\, G\big(G(G\, y)\big).$$

Step 3. Unroll G:

$$G\, y = x^3(y), \qquad G(G\, y) = x^6(y), \qquad G\big(G(G\, y)\big) = x^9(y).$$

Answer

$$\boxed{\lambda x.\, \lambda y.\, x(x(x(x(x(x(x(x(x\, y)))))))))} \;=\; \lambda x.\lambda y.\, x^9(y).$$

(Interpreting the terms as Church numerals, this is $3^2 = 9$.)

Remark. The left term applies a function twice, the right term thrice; composing them yields ninefold application.

2.6 Week 6

Homework

Compute `fact 3` using the computation rules for `fix`, `let`, and `let rec`, and label each step with the rule used.

Solution

We use the standard sugars/rules:

$$\texttt{fix}\ F\ \to\ (F\,(\texttt{fix}\ F)) \qquad\qquad \texttt{let}\ x = e_1\ \text{in}\ e_2\ \to\ (\lambda x.\, e_2)\, e_1$$
$$\texttt{let rec}\ f = e_1\ \text{in}\ e_2\ \to\ \texttt{let}\ f = (\texttt{fix}\ (\lambda f.\, e_1))\ \text{in}\ e_2$$

and basic computation rules for conditionals and arithmetic:

$$0 = 0 \to \texttt{True}, \quad n{>}0 \Rightarrow (n = 0) \to \texttt{False}, \quad \texttt{if True then}\ A\ \texttt{else}\ B \to A, \quad \texttt{if False then}\ A\ \texttt{else}\ B \to B.$$

For brevity, write

$$F \stackrel{\text{def}}{=} \lambda f.\, \lambda n.\, \texttt{if}\ (n = 0)\ \texttt{then}\ 1\ \texttt{else}\ n * f(n-1).$$

Derivation (each step labeled).

$\texttt{let rec fact} = \lambda n.\, \texttt{if}\ (n = 0)\ \texttt{then}\ 1\ \texttt{else}\ n * \texttt{fact}\ (n-1)\ \ \texttt{in}\ \ \texttt{fact}\ 3$

$\xrightarrow{\text{def of } \texttt{let rec}} \texttt{let fact}\ = \texttt{fix}\ F\ \ \texttt{in}\ \ \texttt{fact}\ 3$

$\xrightarrow{\text{def of } \texttt{let}} (\lambda \texttt{fact}.\, \texttt{fact}\ 3)\ (\texttt{fix}\ F)$

$\xrightarrow{\beta} (\texttt{fix}\ F)\ 3$

$\xrightarrow{\text{def of } \texttt{fix}} (F\ (\texttt{fix}\ F))\ 3$

$\xrightarrow{\beta} (\lambda n.\, \texttt{if}\ (n = 0)\ \texttt{then}\ 1\ \texttt{else}\ n * (\texttt{fix}\ F)(n-1))\ 3$

$\xrightarrow{\beta} \texttt{if}\ (3 = 0)\ \texttt{then}\ 1\ \texttt{else}\ 3 * (\texttt{fix}\ F)(2)$

$\xrightarrow{\text{arith/if}} 3 * (\texttt{fix}\ F)(2)$

Evaluate $(\texttt{fix}\ F)(2)$:

$(\texttt{fix}\ F)\ 2 \xrightarrow{\text{def of } \texttt{fix}} (F\ (\texttt{fix}\ F))\ 2 \xrightarrow{\beta} (\lambda n.\, \texttt{if}\ (n = 0)\ \texttt{then}\ 1\ \texttt{else}\ n * (\texttt{fix}\ F)(n-1))\ 2$

$\xrightarrow{\beta} \texttt{if}\ (2 = 0)\ \texttt{then}\ 1\ \texttt{else}\ 2 * (\texttt{fix}\ F)(1) \xrightarrow{\text{arith/if}} 2 * (\texttt{fix}\ F)(1)$

Evaluate $(\texttt{fix}\ F)(1)$:

$(\texttt{fix}\ F)\ 1 \xrightarrow{\text{def of } \texttt{fix}} (F\ (\texttt{fix}\ F))\ 1 \xrightarrow{\beta} (\lambda n.\, \texttt{if}\ (n = 0)\ \texttt{then}\ 1\ \texttt{else}\ n * (\texttt{fix}\ F)(n-1))\ 1$

$\xrightarrow{\beta} \texttt{if}\ (1 = 0)\ \texttt{then}\ 1\ \texttt{else}\ 1 * (\texttt{fix}\ F)(0) \xrightarrow{\text{arith/if}} 1 * (\texttt{fix}\ F)(0)$

Evaluate $(\texttt{fix}\ F)(0)$:

$(\texttt{fix}\ F)\ 0 \xrightarrow{\text{def of } \texttt{fix}} (F\ (\texttt{fix}\ F))\ 0 \xrightarrow{\beta} (\lambda n.\, \texttt{if}\ (n = 0)\ \texttt{then}\ 1\ \texttt{else}\ n * (\texttt{fix}\ F)(n-1))\ 0$

$\xrightarrow{\beta} \texttt{if}\ (0 = 0)\ \texttt{then}\ 1\ \texttt{else}\ 0 * (\texttt{fix}\ F)(-1) \xrightarrow{\text{arith/if}} 1$

Back-substitute: $1 * (\texttt{fix}\ F)(0) = 1 * 1 = 1, \quad 2 * (\texttt{fix}\ F)(1) = 2 * 1 = 2, \quad 3 * (\texttt{fix}\ F)(2) = 3 * 2 = 6.$

Answer

$$\boxed{\texttt{fact 3}\ \to\ 6}$$

Reflection & Question

Question. Could fix ever terminate incorrectly (or not at all) if the function has no base case?

2.7 Week 7

Homework

Given Grammar

$$\begin{aligned}
&\text{Exp} \;\to\; \text{Exp} + \text{Exp1} \;\mid\; \text{Exp1}\\
&\text{Exp1} \;\to\; \text{Exp1} * \text{Exp2} \;\mid\; \text{Exp2}\\
&\text{Exp2} \;\to\; \text{Integer} \;\mid\; (\text{Exp}) \qquad (\text{with Integer} \;\to \texttt{n})
\end{aligned}$$

Leftmost Derivations (quick checks)

2+1: $\text{Exp} \Rightarrow \text{Exp} + \text{Exp1} \Rightarrow \text{Exp1} + \text{Exp1} \Rightarrow \text{Exp2} + \text{Exp1}$
$\Rightarrow \text{Integer} + \text{Exp1} \Rightarrow \texttt{2} + \text{Exp1} \Rightarrow \texttt{2} + \text{Exp2} \Rightarrow \texttt{2} + \text{Integer} \Rightarrow \texttt{2+1}.$


1+2*3: $\text{Exp} \Rightarrow \text{Exp} + \text{Exp1} \Rightarrow \text{Exp1} + \text{Exp1} \Rightarrow \text{Exp2} + \text{Exp1}$
$\Rightarrow \text{Integer} + (\text{Exp1} * \text{Exp2}) \Rightarrow \texttt{1} + (\text{Exp2} * \text{Exp2}) \Rightarrow \texttt{1} + (\text{Integer} * \text{Integer}) \Rightarrow \texttt{1+2*3}.$


1+(2*3): $\text{Exp} \Rightarrow \text{Exp} + \text{Exp1} \Rightarrow \text{Exp1} + \text{Exp1} \Rightarrow \text{Exp2} + \text{Exp1} \Rightarrow \text{Integer} + \text{Exp1}$
$\Rightarrow \texttt{1} + \text{Exp2} \Rightarrow \texttt{1} + (\text{Exp}) \Rightarrow \texttt{1} + (\text{Exp1} * \text{Exp2}) \Rightarrow \texttt{1} + (\text{Integer} * \text{Integer}) \Rightarrow \texttt{1+(2*3)}.$


(1+2)*3: $\text{Exp} \Rightarrow \text{Exp1} \Rightarrow \text{Exp1} * \text{Exp2} \Rightarrow \text{Exp2} * \text{Exp2} \Rightarrow (\text{Exp}) * \text{Integer}$
$\Rightarrow (\text{Exp} + \text{Exp1}) * \texttt{3} \Rightarrow (\text{Exp1} + \text{Exp1}) * \texttt{3} \Rightarrow (\text{Integer} + \text{Integer}) * \texttt{3} \Rightarrow \texttt{(1+2)*3}.$


1+2*3+4*5+6: $\text{Exp} \Rightarrow \text{Exp} + \text{Exp1} \Rightarrow (\text{Exp} + \text{Exp1}) + \text{Exp1} \Rightarrow ((\text{Exp} + \text{Exp1}) + \text{Exp1}) + \text{Exp1}$
and expand each Exp1 with $\text{Exp1} \to \text{Exp1} * \text{Exp2}$ as needed.


Parse Trees




Exp
Exp1
+
Exp1
Exp2
*
Exp1
Exp2
Integer
Exp2
Integer
3
Integer
1
2


Exp
Exp1
+
Exp1
Exp2
Exp2
)
Exp
(
Integer
Exp1
1
Exp2
*
Exp1
Integer
Exp2
3
Integer
2


Exp
Exp1
Exp2 * Exp1
Integer Exp2
3) Exp (
Exp1 + Exp
Exp2 Exp1
Integer Exp2
2 Integer
1


Exp
Exp1 + Exp
Exp2 Exp1 + Exp
Integer Exp2 * Exp1 Exp1 + Exp1
6 Integer Exp2 Exp2 * Exp1 Exp2
5 Integer Integer Exp2 Integer
4 3 Integer 1
2

2.8 Week 8

Homework

Finished *Tutorial World* (NNG4). Below are my tactic scripts for Levels 5–8 and a matching English proof for Level 8.

NNG Tutorial World — Solutions (Levels 5–8)

```
-- Level 5: Adding zero
rw [add_zero b]
rw [add_zero c]
rfl

-- Level 6: Precision rewriting (same goal shape but focused rewrites)
rw [add_zero b]
rw [add_zero c]
rfl

-- Level 7: add_succ
rw [one_eq_succ_zero]
rw [add_succ]
rw [add_zero]
rfl

-- Level 8: 2 + 2 = 4
nth_rewrite 2 [two_eq_succ_one]
rw [add_succ 2]
nth_rewrite 1 [one_eq_succ_zero]
rw [add_succ 2]
rw [add_zero 2]
rw [   three_eq_succ_two]
rw [   four_eq_succ_three]
rfl
```

Natural-language proof for Level 8 ($2 + 2 = 4$)

We use Peano naturals where $\mathrm{succ}(n)$ is the successor, and

$$0, \quad 1 = \mathrm{succ}(0), \quad 2 = \mathrm{succ}(1), \quad 3 = \mathrm{succ}(2), \quad 4 = \mathrm{succ}(3).$$

Addition is defined by $a + 0 = a$ and $a + \mathrm{succ}(b) = \mathrm{succ}(a + b)$. Then:

$$\begin{aligned}
2 + 2 &= 2 + \mathrm{succ}(1) && (\text{since } 2 = \mathrm{succ}(1)) \\
&= \mathrm{succ}(2 + 1) && (\text{def. of addition}) \\
&= \mathrm{succ}\big(2 + \mathrm{succ}(0)\big) && (\text{since } 1 = \mathrm{succ}(0)) \\
&= \mathrm{succ}\big(\mathrm{succ}(2 + 0)\big) && (\text{def. of addition}) \\
&= \mathrm{succ}\big(\mathrm{succ}(2)\big) && (\text{since } 2 + 0 = 2) \\
&= \mathrm{succ}(3) = 4.
\end{aligned}$$

2.9 Week 9

Homework

From *NNG4 Addition World*, complete Level 5 and provide two Lean solutions for

$$\texttt{add_right_comm}: \ \forall abc : \mathbb{N},\ a + b + c = a + c + b,$$

one *without induction* and one *with induction*. For each, include a corresponding pen-and-paper proof.

Level 5 — Statement

Theorem 2.1 (Right-commutativity of addition)**.** *For all* $a, b, c \in \mathbb{N}$, $\quad a + b + c = a + c + b$.

Lean solution (without induction)

```
-- Using only associativity and commutativity of +
theorem add_right_comm (a b c : ) : a + b + c = a + c + b := by
  -- a + b + c = a + (b + c)
  rw [add_assoc]
  -- a + (b + c) = a + (c + b)
  rw [add_comm b c]
  -- a + (c + b) = a + c + b
  rw [add_assoc]
```

Pen-and-paper proof (corresponding to the non-inductive Lean script)

We use associativity and commutativity of $+$ on $\mathbb{N}$:

$$\begin{aligned} a + b + c &= a + (b + c) && \text{(associativity)} \\ &= a + (c + b) && \text{(commutativity)} \\ &= a + c + b && \text{(associativity).} \end{aligned}$$

Thus $a + b + c = a + c + b$.

Lean solution (with induction on b)

```
theorem add_right_comm (a b c : ) : a + b + c = a + c + b := by
  induction b with
  | zero =>
      -- goal: a + 0 + c = a + c + 0
      rw [add_zero, add_zero]
      rfl
  | succ d ih =>
      -- goal: a + succ d + c = a + c + succ d
      -- expose the IH by rewriting both sides
      rw [add_succ]      -- a + succ d + c = succ (a + d) + c
      rw [succ_add]      -- succ (a + d) + c = succ ((a + d) + c)
      rw [ih]            -- succ ((a + c) + d)
      rw [add_succ]      -- a + c + succ d = succ ((a + c) + d)
      rfl
```

Pen-and-paper proof (corresponding to the inductive Lean script)

Claim. For all $a, c \in \mathbb{N}$ and all $b \in \mathbb{N}$, we have $a + b + c = a + c + b$.

Base case $b = 0$. Using $x + 0 = x$,

$$a + 0 + c = a + c = a + c + 0.$$

Inductive step. Suppose $a + d + c = a + c + d$ for some d (induction hypothesis). Using the successor laws

$$x + (y + 1) = (x + y) + 1 \quad \text{and} \quad (x + 1) + y = (x + y) + 1,$$

we compute

$$\begin{aligned} a + (d + 1) + c &= (a + d + 1) + c = (a + d + c) + 1 \\ &= (a + c + d) + 1 \qquad \text{(by IH)} \\ &= (a + c) + (d + 1) = a + c + (d + 1). \end{aligned}$$

Hence the property holds for $d + 1$, and by induction it holds for all $b \in \mathbb{N}$. □

Reflection & Question

Question. Could we use pattern matching on b instead of induction to do the same proof?

2.10 Week 10

Homework

Work through the “Party Snacks" implication tutorial in the Lean Logic Game and provide single-line solutions for Levels 6–9.

Lean Solutions (Levels 6–9)

```
-- Level 6
fun h c d => h  c ,  d

-- Level 7
fun h cd => h cd.left cd.right

-- Level 8
fun h s =>   (h s).left, (h s).right

-- Level 9
fun r =>   (fun _ : S => r), (fun _ :  S => r)
```

Brief Explanation

Each solution follows directly from the Curry–Howard correspondence:

- Level 6 converts a function on pairs into a curried function.
- Level 7 does the reverse: it consumes a pair and applies each projection.
- Level 8 takes an implication into a conjunction and splits it.
- Level 9 builds both branches of a conjunction from a single proof of R.

Reflection & Question

Reflection. These levels reinforced that implication is just function type and conjunction is a product type. Once I rewrote each goal as a function type, the correct lambda term became much easier to see.

Question. In the Party Snacks tutorial, is there a systematic way (using Curry–Howard) to derive lambda terms directly from the goal type, or does this mostly come from practice and pattern recognition?

2.11 Week 12

Homework

Solve the first three items of the Towers of Hanoi activity:

1. Complete the execution trace of `hanoi 5 0 2` (fill in the dots).
2. Extract from the execution the moves that solve the puzzle (in the right order).
3. Verify the execution by checking it against an online Towers of Hanoi game.

Completed Execution Trace for `hanoi 5 0 2`

Below is the full stack-style execution trace, written line by line as in the note (each line shows the current top of the stack):

```
hanoi 5 0 2
hanoi 4 0 1
hanoi 3 0 2
hanoi 2 0 1
hanoi 1 0 2
move 0 2
move 0 1
hanoi 1 2 1
move 2 1
move 0 2
hanoi 2 1 2
hanoi 1 1 0
move 1 0
move 1 2
hanoi 1 0 2
move 0 2
move 0 1
hanoi 3 2 1
hanoi 2 2 0
hanoi 1 2 1
move 2 1
move 2 0
hanoi 1 1 0
move 1 0
move 2 1
hanoi 2 0 1
hanoi 1 0 2
move 0 2
move 0 1
hanoi 1 2 1
move 2 1
move 0 2
```

```
hanoi 4 1 2
hanoi 3 1 0
hanoi 2 1 2
hanoi 1 1 0
move 1 0
move 1 2
hanoi 1 0 2
move 0 2
move 1 0
hanoi 2 2 0
hanoi 1 2 1
move 2 1
move 2 0
hanoi 1 1 0
move 1 0
move 1 2
hanoi 3 0 2
hanoi 2 0 1
hanoi 1 0 2
move 0 2
move 0 1
hanoi 1 2 1
move 2 1
move 0 2
hanoi 2 1 2
hanoi 1 1 0
move 1 0
move 1 2
hanoi 1 0 2
move 0 2
```

Moves Extracted from the Trace

From this trace, the actual moves of the puzzle are the lines beginning with `move`. There are $2^5 - 1 = 31$ moves:

```
move 0 2
move 0 1
move 2 1
move 0 2
move 1 0
move 1 2
move 0 2
move 0 1
move 2 1
move 2 0
move 1 0
move 2 1
move 0 2
move 0 1
move 2 1
move 0 2
move 1 0
move 1 2
move 0 2
```

```
move 1 0
move 2 1
move 2 0
move 1 0
move 1 2
move 0 2
move 0 1
move 2 1
move 0 2
move 1 0
move 1 2
move 0 2
```

Interpreting `move x y` as "move the top disk from peg x to peg y," following these 31 moves on the online Towers of Hanoi game for $n = 5$ correctly transfers the whole tower from peg 0 to peg 2 without ever putting a larger disk on a smaller one.

Discord Question. From this execution trace it looks like both the number of moves and the number of `hanoi` calls for n disks are $2^n - 1$. Is there a clean structural argument (using the recursion rules for `hanoi`) that shows these counts are always $2^n - 1$, without having to write out the whole trace?

3 Synthesis

4 Evidence of Participation

5 Conclusion

References