

Light Exposure Behavior Assessment (LEBA): Development of a novel instrument to capture light exposure-related behaviours

Mushfiqul Anwar Siraji^{1, *}, Rafael Robert Lazar^{2, 3, *}, Juliëtte van Duijnhoven⁴, Luc Schlangen⁵, Shamsul Haque¹, Vineetha Kalavally⁶, Céline Vetter^{7, 8}, Gena Glickman⁹, Karin Smolders¹⁰, & Manuel Spitschan^{11, 2, 3}

¹ Monash University, Department of Psychology, Jeffrey Cheah School of Medicine and Health Sciences, Malaysia

² Psychiatric Hospital of the University of Basel (UPK), Centre for Chronobiology, Basel, Switzerland

³ University of Basel, Transfaculty Research Platform Molecular and Cognitive Neurosciences, Basel, Switzerland

⁴ Eindhoven University of Technology, Department of the Built Environment, Building Lighting, Eindhoven, Netherlands

⁵ Eindhoven University of Technology, Department of Industrial Engineering and Innovation Sciences, Intelligent Lighting Institute, Eindhoven, Netherlands

⁶ Monash University, Department of Electrical and Computer Systems Engineering, Malaysia, Selangor, Malaysia

⁷ University of Colorado Boulder, Department of Integrative Physiology, Boulder, USA

⁸ Ximes GmbH, Frankfurt, Germany

⁹ Uniformed Services University of the Health Sciences, Department of Psychiatry,

Bethesda, USA

¹⁰ Eindhoven University of Technology, Human-Technology Interaction Group,
Eindhoven, Netherlands

¹¹ University of Oxford, Department of Experimental Psychology, Oxford, UK

* Joint first author

Author Note

Add complete departmental affiliations for each author here. Each new line herein must be indented, like this line.

Enter author note here.

The authors made the following contributions. Mushfiqul Anwar Siraji: Formal Analysis, Visualization, Writing – original draft, Writing – review & editing;; Rafael Robert Lazar: Data curation, Investigation, Project administration, Visualization, Writing – original draft, Writing – review & editing;; Juliëtte van Duijnhoven: Conceptualization, Methodology, Investigation, Writing – review & editing; Luc Schlangen: Conceptualization, Methodology, Investigation, Writing – review & editing; Shamsul Haque: Conceptualization, Supervision, Writing – review & editing; Vineetha Kalavally: Supervision, Writing – review & editing; Céline Vetter: Conceptualization, Writing – review & editing; Gena Glickman: Conceptualization, Methodology, Writing – review & editing; Karin Smolders: Conceptualization, Methodology, Writing – review & editing; Manuel Spitschan: Conceptualization, Data curation, Investigation, Project administration, Visualization, Methodology, Writing – original draft, Writing – review & editing.

Abstract

One or two sentences providing a **basic introduction** to the field, comprehensible to a scientist in any discipline.

Two to three sentences of **more detailed background**, comprehensible to scientists in related disciplines.

One sentence clearly stating the **general problem** being addressed by this particular study.

One sentence summarizing the main result (with the words “**here we show**” or their equivalent).

Two or three sentences explaining what the **main result** reveals in direct comparison to what was thought to be the case previously, or how the main result adds to previous knowledge.

One or two sentences to put the results into a more **general context**.

Two or three sentences to provide a **broader perspective**, readily comprehensible to a scientist in any discipline.

Keywords: keywords

Word count: X

Light Exposure Behavior Assessment (LEBA): Development of a novel instrument to capture light exposure-related behaviours

Introduction

- Light exposure is important
- Light exposure Behavior is important
- Table: Overview Existing Related Scales: items in total / items on light exposure (behaviour)
- Existing Scales: Review them in text
- None of these do light exposure behavior.

Methods

Ethical approval

The cantonal ethics commission (Ethikkommission Nordwest- und Zentralschweiz, project ID Req-2021-00488) reviewed this project and issued an official clarification of responsibility (full document see Suppl. Fig X in appendix) stating: “The research project does not fall under the scope of the Human Research Act, because your project is using only anonymised data. An authorisation from the ethics committee is therefore not required and the EKNZ is not responsible for its review.”

Data Availability

Survey characteristics

Data was collected in a quantitative cross-sectional approach via a fully anonymous online survey hosted on REDCap (Harris et al., 2019, 2009) by way of the University of

Basel sciCORE. Participants were recruited via the website of a Comic co-released with the survey (Weinzaepflen & Spitschan, 2021), social media (i.e., LinkedIn, Twitter, Facebook), mailing lists, word of mouth, the investigators' personal contacts, and supported by distribution of the survey link via f.lux software (F.lux Software LLC, 2021).

Completing the online survey took approx. 15 to 20 minutes and was not compensated. The first page of the survey comprised a participant information sheet, where participants' informed consent to participate was obtained before any of the questions were displayed. Underaged participants (<18 years) were urged to obtain assent from their parents/legal guardians, before filling in the survey. Information on the first page included the objectives of the study, inclusion criteria, estimated duration, the use, storage and sharing of the data, compensation (none), and information about the type of questions in the survey. Moreover, participants needed to confirm that they were participating the survey for the first time. To ensure high data quality, five attention check items were included in the survey (e.g., "We want to make sure you are paying attention. What is 4+5?"). The data analysed in this study was collected between 17.05.2021 and 03.09.2021. Questions incorporating retrospective recall were all aligned to the period of "past four weeks," matching the presented LEBA instrument.

In addition to the LEBA questionnaire, which is subject of the current study, the following variables and items were assessed but not included in the analysis:

- Sleep disturbance and sleep-related impairment (adult and pediatric versions) (Bevans et al., 2019; Daniel J. Buysse et al., 2010; Forrest et al., 2018; Harb, Hidalgo, & Martau, 2015; L. Yu et al., 2011)
- Sleep duration, timing, and latency, chronotype, social jetlag, time in bed, work/sleep schedule and outdoor light exposure duration (version for adults and adolescents) (Roenneberg, Wirz-Justice, & Mellow, 2003)
- Sleep environment (Olivier et al., 2016)

- Meal timing & caffeine consumption [custom items]
- Light sensitivity (photophobia vs. photophilia) (Wu & Hallett, 2017)
- Self-reported pubertal stage (only if younger than 18 years old) (Petersen, Crockett, Richards, & Boxer, 1988)

Furthermore, the following 1-item demographic variables were assessed:

- Age
- Sex
- Gender identity
- Occupational Status
- COVID-19 related Occupational setting during the past four weeks
- Time zone & country of residence
- English as native language

Participants

Table 1 summarizes the survey participants' demographic characteristics. Only participants completing the full LEBA questionnaire were included, thus there are no missing values in the item analyses. XX participants were excluded from analysis due to not passing at least one of the "attention check" items. For exploring initial factor structure (EFA), a sample of 250-300 is recommended (Comrey & Lee, 1992; Schönbrodt & Perugini, 2013). For estimating the sample size for the confirmatory factor analysis (CFA) we followed the N:q rule (Bentler & Chou, 1987; Jackson, 2003; Kline, 2015; Worthington & Whittaker, 2006), where ten participants per parameter is required to earn trustworthiness of the result. Our sample size exceeds these requirements: Anonymous responses from a total of $n = 690$ participants were included in the analysis of the current study, split into samples for exploratory (EFA: $n = 428$) and confirmatory factor analysis (CFA: $n = 262$). The EFA sample included participants filling out the

questionnaire from 17.05.2021 to XX.XX.XXXX , whereas participants who filled out the questionnaire from YY.YY.YYYY to 03.09.2021 were included in the CFA analysis. Participants indicated filling out the online survey from a diverse range of geographic locations. The four most common geographic locations included:

	x
United States - America/New_York (UTC -04:00)	63
United Kingdom - Europe/London (UTC)	57
Germany - Europe/Berlin (UTC +01:00)	53
India - Asia/Kolkata (UTC +05:30)	38

For a full list of geographic locations, see Suppl. Table X in the appendix.

Age among all participants ranged from 11 years to 84 years [EFA: *min* = 11, *max* = 84; CFA: *min* = 12, *max* = 74], with an overall mean of ~ 33 years of age [Overall: *M* = 32.95, *SD* = 14.57; EFA: *M* = 32.99, *SD* = 15.11; CFA: *M* = 32.89, *SD* = 13.66]. In total 325 (47%) of the participants indicated female sex [EFA: 189 (44%); CFA: 136 (52%)], 351 (51%) indicated male [EFA: 230 (54%); CFA: 121 (46%)] and 14 (2.0%) indicated other sex [EFA: 9 (2.1%), CFA: 5 (1.9%)]. Overall, 49 (7.2%) [EFA: 33 (7.8%); CFA: 16 (6.2%)] participants indicated a gender-variant identity. In a “Yes/No” question regarding native language, 320 (46%) of respondents [EFA: 191 (45%); CFA: 129 (49%)] indicated to be native English speakers. For their “Occupational Status,” more than half of the overall sample reported that they currently work [Overall: 396 (57%); EFA: 235 (55%); CFA: 161 (61%)], whereas 174 (25%) [EFA: 122 (29%); CFA: 52 (20%)] reported that they go to school and 120 (17%) [EFA: 71 (17%); CFA: 49 (19%)] responded that they do “Neither.” With respect to the COVID-19 pandemic we asked participants to indicate their occupational setting during the last four weeks: In the overall sample 303 (44%) [EFA: 194 (45%); CFA: 109 (42%)] of the participants indicated that they were in a home office/home schooling setting., while 109 (16%) overall [EFA: 68 (16%) ; CFA: 41 (16%)] reported face-to-face work/schooling. Lastly, 147 (21%) overall [EFA: 94 (22%) ; CFA: 53

(20%)] reported a combination of home- and face-to-face work/schooling, whereas 131 (19%) overall [EFA: 72 (17%); CFA: 59 (23%)] filled in the “Neither (no work or school, or indication)” response option. We tested all demographic variables in Table 1 for significant group differences between the EFA and CFA sample, applying Wilcoxon rank sum test for the continuous variable “Age” and Pearson’s Chi-squared test for all other categorical variables via the gtsummary R package’s “add_p” function (Sjoberg et al., 2021a) . The p-values were corrected for multiple testing applying false discovery rate (FDR) via the “add_q” function of the same package. After p-value (FDR) correction for multiple testing, none of the demographic variables were significantly different between the EFA sample and the CFA sample (all q-values $q \geq 0.2$, indicating equivalence).

1. Describe EFA and CFA sample separately.
2. Sampling technique: Convince sampling (non-probability sample)
3. Method: cross-sectional survey
4. How many missing data?
5. How incomplete data were addressed.
6. Why such sample was chosen?

Procedure

Development of the Scale.

1. How the items were generated
2. How the literature was reviewed to identify construct adequacy of the items.
3. Discuss the expert panel review process to assess content validity

Data Collection. Timeline of data collection, mode of data collection.

Analytic Strategies

We used R (version 4.1.0), including several R packages, for our analyses. Initially, our tool have six poin Likert type response scale(0:Does not apply/I don't know; 1:Never, 2:Rarely; 3:Sometimes; 4:Often; 5: Alsways). As our purpose was to capture light exposure related behavior, "Does not apply/I don't know" and "Never" were providing similar information. As such we decided to collapse "Does not apply/I don't know" and "Never" options into one making it a 5 point Likert type response scale. Necessary assumptions of EFA, including sample adequacy, normality assumptions, quality of correlation matrix, were assessed. Our data violated both the univariate and multivariate normality assumptions. Due to these violations and the ordinal nature of our response data, we used a polychoric correlation matrix (C. Desjardins & Bulut, 2018) for the EFA. We employed principal axis (PA) as a factor extraction method with varimax rotation. PA is robust to the normality assumption violations (Watkins, 2020). The obtained latent structure was confirmed by another factor extraction method: the minimum residuals extraction method as well. We used a combination factor identification method including scree plot(Cattell, 1966), Horn's parallel analysis (Horn, 1965), minimum average partials method(Velicer, 1976), and hull method (Lorenzo-Seva, Timmerman, & Kiers, 2011) to identify factor numbers. Additionally, to determine the simple structure, we followed the following guidelines recommended by psychometricians (i) no factors with fewer than three items (ii) no factors with a factor loading <0.3 (iii) no items with cross-loading greater than .3 across factors (Bandalos & Finney, 2018) We also conducted psychometric analysis on non-merged response options data (supplementary analysis) and rejected the latent structure obtained as the factors were less interpretable.

Results

Exploratory Factor Analysis

Sampling adequacy was checked using Kaiser-Meyer-Olkin (KMO) measures of sampling adequacy (Kaiser, 1974). The overall KMO value for 48 items was 0.63 which was above the cutoff value (.50) indicating a mediocre sample (Hutcheson, 1999). Table 3 summarizes the univariate descriptive statistics for the 48 items. Some of the items were skewed with high Kurtosis values. Our data violated both univariate normality (Shapiro-Wilk statistics; (Shapiro & Wilk, 1965)) and multivariate normality assumptions (Marida's test; (Mardia, 1970)). Multivariate skew was = 583.80 ($p < 0.001$) and multivariate kurtosis was = 2,749.15 ($p < 0.001$). Due to these violations and ordinal nature of the response data polychoric correlations over Pearson's correlations was chosen (C. Desjardins & Bulut, 2018). Bartlett's test of sphericity (Bartlett, 1954), $\chi^2(1128) = 5042.86$, $p < .001$ indicated the correlations between items are adequate for the EFA. However only 4.96% of the inter-item correlation coefficients were greater than .30. The inter item correlation ranged between .44 to .91. And the corrected item-total correlations ranged between .10 to .44.

Scree plot (Figure 3) suggested a six-factor solution. Horn's parallel analysis (Horn, 1965) with 500 iterations also indicated a six-factor solution. However, the minimum average partial (MAP) method (Velicer, 1976) and Hull method (Lorenzo-Seva et al., 2011) suggested a five-factor solution. As a result, we tested both five-factor and six-factor solutions.

With initial 48 items we conducted three rounds of EFA gradually discarded problematic items. (cross-loading items and poor factor loading ($< .30$) items). Finally, a five-factor EFA solution with 25 items was accepted with low RMSR = 0.08 (Brown,

2015), all factor-loading higher than .30 and no cross-loading greater than .30. We confirmed this five-factor latent structure using varimax rotation with a minimum residual extraction method (Table??). Table4 displays the factor-loading (structural coefficients) and communality of the items. The absolute value of the factor-loading ranged from -.49 to .99 indicating strong coefficients. The commonalities ranged between .11 to .99. However, the histogram of the absolute values of non-redundant residual-correlations Fig5 showed 26% correlations greater than the absolute value of .05, indicating a possible under-factoring. (C. D. Desjardins, 2018). Subsequently, we fitted a six-factor solution. However, a factor emerged with only one salient variable loading in the six-factor solution, thus disqualifying the six-factor solution (Table??).

In the five-factor solution, the first factor contained three items and explained 10.25% of the total variance with a satisfactory internal reliability coefficient ($\alpha = .86$). All the items in this factor stemmed from the individual's preference to use blue light filters in different light environments. The second factor contained six items and explained 9.93% of the total variance with a satisfactory internal reliability coefficient ($\alpha = .71$). Items under this factor commonly investigate an individual's hours spent outdoor. The third factor contained five items and explained 8.83% of the total variance. Items under this factor dealt with the specific behaviors pertaining to sleep. The internal consistency reliability coefficient was, $\alpha = .68$. The fourth factor contained five items and explained 8.44% of the total variance with an internal consistency coefficient, $\alpha = .62$. These five items stemmed from the behavior related to an individual's cellphone usage during the sleep-wakeup time. Lastly, the fifth factor contained six items and explained 6.14% of the total variance. This factor tried to measure an individual's behavior lead by the awareness of light's influence on health. However, this factor showed unsatisfactory internal consistency reliability ($\alpha = .53$). It is essential to attain a balance between psychometric properties and interpretability of the common themes when exploring the latent structure. As all of the emerged factors are highly interpretable and relevant

towards our aim to capture light exposure related behavior, regardless of the apparent low reliability of the two factors, we retain the five-factor solution with 23 items for our confirmatory factor analysis (CFA). Two items showed negative factor-loading (items 44 and 21). Upon inspection, it was understood that these items are negatively correlated to the common theme, and thus in the CFA analysis, we reversed the response code for these two items.

Confirmatory Factor Analysis

We conducted a categorical confirmatory factor analysis with robust weighted least square (WLSMV) estimator as our response data was in ordinary nature (C. Desjardins & Bulut, 2018). Several indices are suggested to measure model fit. These indices can be categorized as absolute, comparative and parsimony fit indices (Brown, 2015). Absolute fit assess the model fit at an absolute level using indices including chi-square test statistics and the standardized root mean square (SRMR). Parsimony fit indices including the root mean square error of approximation (RMSEA) considers the number of free parameters in the model to assess the parsimony of the model. Comparative fit indices evaluate the fit of the specified model solution in relation to a more restricted baseline model restricting all covariances among the indicators as zero. Comparative fit index (CFI) and the Tucker Lewis index (TLI) are such two comparative fit indices. Commonly used Model fit guidelines (Hu & Bentler, 1999; Schumacker & Lomax, 2004) includes (i) Reporting of chi-square test statistics (A non-significant test statistics is required to reflect model fit) (i) CFI and TLI (CFI/TLI close to .95 or above/ranging between 90-95 and above) (ii) RMSEA (close to .06 or below), (iii) SRMR (close to .08 or below) to estimate the model fit. Table 5 summarizes the fit indices of our fitted model. Our fitted model failed to attain an absolute fit estimated by the chi-square test. However, the chi-square test is sensitive to sample size and not recommended to be used as the sole index of absolute model fit (Brown, 2015). Another absolute fit index we obtained in our

analysis was SRMR which does not work well with categorical data (C.-Y. Yu, 2002). Subsequently, we judged the model fit based on the comparative fit indices: CFI, TLI and parsimony fit index-RMSEA. Our fitted model attained acceptable fit (CFI =.94; TLI = .93); RMSEA = .06,[.05-.07, 90% CI]) with two imposed equity constrain on item pairs 32-33 and 19-17 . However SRMR value was higher than the guideline (SRMR = .12). Further by allowing one pair of items (30-41) to covary their error variance and discarding two item (item 37 & 26) for very low r-square value, our model attained best fit (CFI =.97; TLI = .96); RMSEA = .05[.04-.06, 90% CI]) and SRMR value (SRMR = .09) was also close to the suggestions of Hu and Bentle (1999). Since reliability coefficient Cronbach's alpha tends to mis-measure reliability for multidimensional construct (Sijtsma, 2009) we report Mcdonald's omega(total) as the internal consistency reliability coefficient. Mcdonald's omega(total) is reported to be a satisfactory lower bound reliability coefficient that works for both unidimensional and multidimensional construct (Zinbarg, Revelle, Yovel, & Li, 2005). Mcdonald's omega(total) for the five factors were .90, .80, .61, .72, .45 respectively. Mcdonald's omega(total) coefficient for the total scale was .73.

Analysing the quality of items by Item Information Theory

We sought the IRT to gether information regarding the item quality. IRT complements the conventional classical test theory-based analysis by gathering information on item discrimination and item difficulty(Baker, 2017). Here, an item's quality is judged based on item information in relation to participants' latent trait level (θ). We gathered evidence on item quality by fitting each factor of LEBA with the graded response model (7 to the combined EFA sample and CFA sample (n =690). Item discrimination indicates the pattern of variation in the categorical responses with the changes in latent trait, and item information curve (IIC) indicates the amount of information an item carries along the latent trait continuum. Here, we reported the item discrimination parameter and only discarded the items with relatively flat item information

curve (information $< .2$) to develop the short form of LEBA. Baker (2017) categorized the item discrimination in as none = 0; very low = 0.01 to 0.34; low = 0.35 to 0.64; moderate = 0.65 to 1.34 ; high = 1.35 to 1.69; very high > 1.70 . Item discrimination parameters of our scale fell in very high (10 items), high (4 items), moderate (4 items), low (5 items) indicating a good range of discrimination along the latent trait. Examination of the item information curve indicated 6 items (1, 25, 9, 38, 30, & 41) had relatively flat information curves thus discarded. We also gathered evidence of item fit and person fit to our fitted model.

Test information curve (TIC) indicate the amount of information an the full-scale carry along the latent trait continuum. As we treated each factor of LEBA as an unidimensional construct we obtain 5 TICs. These information curves indicated except blue filter factor, the other factor's TICs are roughly centered on the center of the trait continuum ((θ)). Also the amount of information changed rather steadily with the change of (θ) . Thus we conferred the LEBA scale (except blue filter) estimated the light exposure related behavior with precision near the center of trait continuum (Baker, 2017) which is sufficient to discriminate between latent trait measured by the each factor. The blue filter factor had a peak to the right side of the center of latent trait indicating its ability to providing information only for people who already have some preference towards using blue-filters.

Our result also indicated all the items fitted well to the respective models as assessed by assessed by RMSEA value obtained from Signed-X2 index implementation. All of the items had RMSEA value $< .06$ indicating adequate fit. Person fit indicates the validity and meaningfulness of the fitted model at the participants latent trait level (C. Desjardins & Bulut, 2018). We estimated the person fit statistics using standardized fit index Zh statistics (Drasgow, Levine, & Williams, 1985). Zh < -2 should be considered as a misfit. Fig indicates that Zh is larger than -2 for most participants, suggesting a good fit of the selected IRT models.

The overall we can concluded that IRT analysis indicated LEBA is a psychometrically sound measure. Item fit indexes and person fit index for all five fitted model were acceptable. Items had diverse slope parameters indicating a good range of discrimination- the ability to differentiate respondents with different levels of the light exposure related behavior. All-in-all we can recommend the LEBA to be used to capture light exposure related behavior.

Discussion

References

- Anwar Siraji, M. (2021). *Tabledown: A companion pack for the book "basic & advanced psychometrics in r"*. Retrieved from <https://github.com/masiraji/taledown>
- Aust, F., & Barth, M. (2020). *papaja: Prepare reproducible APA journal articles with R Markdown*. Retrieved from <https://github.com/crsh/papaja>
- Bajaj, A., Rosner, B., Lockley, S. W., & Schernhammer, E. S. (2011). Validation of a light questionnaire with real-life photopic illuminance measurements: The harvard light exposure assessment questionnaire. *Cancer Epidemiology and Prevention Biomarkers*, 20(7), 1341–1349.
- Baker, F. B. (2017). *The Basics of Item Response Theory Using R* (1st ed. 2017.). Springer.
- Bandalos, D. L., & Finney, S. J. (2018). Factor analysis: Exploratory and confirmatory. In *The reviewer's guide to quantitative methods in the social sciences* (pp. 98–122). Routledge.
- Barnier, J., Briatte, F., & Larmarange, J. (2020). *Questionr: Functions to make surveys processing easier*. Retrieved from <https://CRAN.R-project.org/package=questionr>
- Barth, M. (2021). *tinylabls: Lightweight variable labels*. Retrieved from <https://github.com/mariusbarth/tinylabls>
- Bartlett, M. (1954). A Note on the Multiplying Factors for Various Chi-square Approximations. *Journal of the Royal Statistical Society. Series B, Methodological*, 16(2), 296–298.
- Bentler, P. M., & Chou, C.-P. (1987). Practical Issues in Structural Modeling. *Sociological Methods & Research*, 16(1), 78–117. <https://doi.org/10.1177/0049124187016001004>
- Bevans, K. B., Meltzer, L. J., La Motte, A. de, Kratchman, A., Viél, D., & Forrest, C.

B. (2019). Qualitative development and content validation of the PROMIS pediatric sleep health items. *Behavioral Sleep Medicine*, 17(5), 657–671.
<https://doi.org/10.1080/15402002.2018.1461102>

Brown, T. A. (2015). *Confirmatory factor analysis for applied research* (2nd ed.). New York, NY, US: The Guilford Press.

Bryer, J., & Speerschneider, K. (2016). *Likert: Analysis and visualization likert items*. Retrieved from <https://CRAN.R-project.org/package=likert>

Buchanan, E. M., Gillenwaters, A., Scofield, J. E., & Valentine, K. D. (2019). *MOTE: Measure of the Effect: Package to assist in effect size calculations and their confidence intervals*. Retrieved from <http://github.com/doomlab/MOTE>

Buysse, Daniel J., Reynolds III, C. F., Monk, T. H., Berman, S. R., & Kupfer, D. J. (1989). The pittsburgh sleep quality index: A new instrument for psychiatric practice and research. *Psychiatry Research*, 28(2), 193–213.

Buysse, Daniel J., Yu, L., Moul, D. E., Germain, A., Stover, A., Dodds, N. E., ... Pilkonis, P. A. (2010). Development and validation of patient-reported outcome measures for sleep disturbance and sleep-related impairments. *Sleep*, 33(6), 781–792. <https://doi.org/10.1093/sleep/33.6.781>

Cattell, R. B. (1966). The Scree Test For The Number Of Factors. *Multivariate Behavioral Research*, 1(2), 245–276.
https://doi.org/10.1207/s15327906mbr0102_10

Chalmers, R. P. (2012). mirt: A multidimensional item response theory package for the R environment. *Journal of Statistical Software*, 48(6), 1–29.
<https://doi.org/10.18637/jss.v048.i06>

Chang, W., Cheng, J., Allaire, J., Sievert, C., Schloerke, B., Xie, Y., ... Borges, B. (2021). *Shiny: Web application framework for r*. Retrieved from <https://CRAN.R-project.org/package=shiny>

Comrey, A. L., & Lee, H. B. (1992). *A first course in factor analysis*, 2nd ed.

Hillsdale, NJ, US: Lawrence Erlbaum Associates, Inc.

Conigrave, J. (2020). *Corx: Create and format correlation matrices*. Retrieved from <https://CRAN.R-project.org/package=corx>

Dahl, D. B., Scott, D., Roosen, C., Magnusson, A., & Swinton, J. (2019). *Xtable: Export tables to LaTeX or HTML*. Retrieved from <https://CRAN.R-project.org/package=xtable>

Desjardins, C. D. (2018). *Handbook of educational measurement and psychometrics using R*. (O. Bulut & ProQuest (Firm), Eds.). Boca Raton, FL : CRC Press.

Desjardins, C., & Bulut, O. (2018). *Handbook of Educational Measurement and Psychometrics Using R*. <https://doi.org/10.1201/b20498>

Dianat, I., Sedghi, A., Bagherzade, J., Jafarabadi, M. A., & Stedmon, A. W. (2013). Objective and subjective assessments of lighting in a hospital setting: Implications for health, safety and performance. *Ergonomics*, 56(10), 1535–1545.

Dinno, A. (2018). *Paran: Horn's test of principal components/factors*. Retrieved from <https://CRAN.R-project.org/package=paran>

Dragow, F., Levine, M. V., & Williams, E. A. (1985). Appropriateness measurement with polychotomous item response models and standardized indices. *British Journal of Mathematical and Statistical Psychology*, 38(1), 67–86.

Eklund, N., & Boyce, P. (1996). The development of a reliable, valid, and simple office lighting survey. *Journal of the Illuminating Engineering Society*, 25(2), 25–40.

Epskamp, S. (2019). *semPlot: Path diagrams and visual analysis of various SEM packages' output*. Retrieved from <https://CRAN.R-project.org/package=semPlot>

Epskamp, S., Cramer, A. O. J., Waldorp, L. J., Schmittmann, V. D., & Borsboom, D. (2012). qgraph: Network visualizations of relationships in psychometric data. *Journal of Statistical Software*, 48(4), 1–18.

F.lux Software LLC. (2021). F.lux (Version 4.120). Retrieved from <https://justgetflux.com/>

Forrest, C. B., Meltzer, L. J., Marcus, C. L., La Motte, A. de, Kratchman, A., Buysse, D. J., ... Bevens, K. B. (2018). Development and validation of the PROMIS pediatric sleep disturbance and sleep-related impairment item banks. *Sleep*, 41(6). <https://doi.org/10.1093/sleep/zsy054>

Fox, J., & Weisberg, S. (2019). *An R companion to applied regression* (Third). Thousand Oaks CA: Sage. Retrieved from <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>

Fox, J., Weisberg, S., & Price, B. (2020). *carData: Companion to applied regression data sets*. Retrieved from <https://CRAN.R-project.org/package=carData>

Harb, F., Hidalgo, M. P., & Martau, B. (2015). Lack of exposure to natural light in the workspace is associated with physiological, sleep and depressive symptoms. *Chronobiology International*, 32(3), 368–375. <https://doi.org/10.3109/07420528.2014.982757>

Harrell Jr, F. E., Charles Dupont, with contributions from, & others., many. (2021). *Hmisc: Harrell miscellaneous*. Retrieved from <https://CRAN.R-project.org/package=Hmisc>

Harris, P. A., Taylor, R., Minor, B. L., Elliott, V., Fernandez, M., O'Neal, L., ... others. (2019). The REDCap consortium: Building an international community of software platform partners. *Journal of Biomedical Informatics*, 95, 103208.

Harris, P. A., Taylor, R., Thielke, R., Payne, J., Gonzalez, N., & Conde, J. G. (2009). Research electronic data capture (REDCap)—a metadata-driven

methodology and workflow process for providing translational research
informatics support. *Journal of Biomedical Informatics*, 42(2), 377–381.

Henry, L., & Wickham, H. (2020). *Purrr: Functional programming tools*. Retrieved
from <https://CRAN.R-project.org/package=purrr>

Horn, J. L. (1965). A rationale and test for the number of factors in factor analysis.
Psychometrika, 30(2), 179–185. <https://doi.org/10.1007/BF02289447>

Horne, J. A., & Östberg, O. (1976). A self-assessment questionnaire to determine
morningness-eveningness in human circadian rhythms. *International Journal
of Chronobiology*.

Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure
analysis: Conventional criteria versus new alternatives. *Structural Equation
Modeling: A Multidisciplinary Journal*, 6(1), 1–55.
<https://doi.org/10.1080/10705519909540118>

Hutcheson, G. D. (1999). *The multivariate social scientist : Introductory statistics
using generalized linear models*. London : SAGE.

Iannone, R. (2016). *DiagrammeRsvg: Export DiagrammeR graphviz graphs as
SVG*. Retrieved from <https://CRAN.R-project.org/package=DiagrammeRsvg>

Iannone, R. (2021). *DiagrammeR: Graph/network visualization*. Retrieved from
<https://github.com/rich-iannone/DiagrammeR>

Jackson, D. L. (2003). Revisiting Sample Size and Number of Parameter
Estimates: Some Support for the N:q Hypothesis. *Structural Equation
Modeling*, 10(1), 128–141. https://doi.org/10.1207/S15328007SEM1001_6

Johnson, P., & Kite, B. (2020). *semTable: Structural equation modeling tables*.
Retrieved from <https://CRAN.R-project.org/package=semTable>

Johnson, P., Kite, B., & Redmon, C. (2020). *Kutils: Project management tools*.
Retrieved from <https://CRAN.R-project.org/package=kutils>

Jorgensen, T. D., Pornprasertmanit, S., Schoemann, A. M., & Rosseel, Y. (2021).

- 473 *semTools: Useful tools for structural equation modeling*. Retrieved from
474 <https://CRAN.R-project.org/package=semTools>
- 475 Kaiser, H. F. (1974). An index of factorial simplicity. *Psychometrika*, 39(1), 31–36.
476 <https://doi.org/10.1007/bf02291575>
- 477 Kassambara, A. (2019). *Ggcorrplot: Visualization of a correlation matrix using*
478 *'ggplot2'*. Retrieved from <https://CRAN.R-project.org/package=ggcorrplot>
- 479 Kline, R. B. (2015). *Principles and practice of structural equation modeling*. The
480 Guilford Press.
- 481 Kowarik, A., & Templ, M. (2016). Imputation with the R package VIM. *Journal of*
482 *Statistical Software*, 74(7), 1–16. <https://doi.org/10.18637/jss.v074.i07>
- 483 Lishinski, A. (2021). *lavaanPlot: Path diagrams for 'lavaan' models via*
484 *'DiagrammeR'*. Retrieved from
485 <https://CRAN.R-project.org/package=lavaanPlot>
- 486 Lorenzo-Seva, U., Timmerman, M., & Kiers, H. (2011). The Hull Method for
487 Selecting the Number of Common Factors. *Multivariate Behavioral Research*,
488 46, 340–364. <https://doi.org/10.1080/00273171.2011.564527>
- 489 Makowski, D., Ben-Shachar, M. S., Patil, I., & Lüdtke, D. (2020). Methods and
490 algorithms for correlation analysis in R. *Journal of Open Source Software*,
491 5(51), 2306. <https://doi.org/10.21105/joss.02306>
- 492 Mardia, K. V. (1970). Measures of multivariate skewness and kurtosis with
493 applications. *Biometrika*, 57(3), 519–530.
494 <https://doi.org/10.1093/biomet/57.3.519>
- 495 Mock, T. (2021). *gtExtras: A collection of helper functions for the gt package*.
496 Retrieved from <https://github.com/jthomasmock/gtExtras>
- 497 Müller, K., & Wickham, H. (2021). *Tibble: Simple data frames*. Retrieved from
498 <https://CRAN.R-project.org/package=tibble>
- 499 Navarro-Gonzalez, D., & Lorenzo-Seva, U. (2021). *EFA.MRFA: Dimensionality*

- 500 *assessment using minimum rank factor analysis*. Retrieved from
501 <https://CRAN.R-project.org/package=EFA.MRFA>
- 502 Olivier, K., Gallagher, R. A., Killgore, W. D. S., Carrasco, N., Alfonso-Miller, P., ...
503 Grandner, M. A. (2016). Development and initial validation of the assessment
504 of sleep environment: A novel inventory for describing and quantifying the
505 impact of environmental factors on sleep. *Sleep*, 39(Abstract Supplement:
506 A367).
- 507 Ooms, J. (2021a). *Magick: Advanced graphics and image-processing in r*.
508 Retrieved from <https://CRAN.R-project.org/package=magick>
- 509 Ooms, J. (2021b). *Rsvg: Render SVG images into PDF, PNG, PostScript, or*
510 *bitmap arrays*. Retrieved from <https://CRAN.R-project.org/package=rsvg>
- 511 Petersen, A. C., Crockett, L., Richards, M., & Boxer, A. (1988). A self-report
512 measure of pubertal status: Reliability, validity, and initial norms. *Journal of*
513 *Youth and Adolescence*, 17(2), 117–133. <https://doi.org/10.1007/BF01537962>
- 514 Pornprasertmanit, S., Miller, P., Schoemann, A., & Jorgensen, T. D. (2021).
515 *Simsem: SIMulated structural equation modeling*. Retrieved from
516 <https://CRAN.R-project.org/package=simsem>
- 517 R Core Team. (2021). *R: A language and environment for statistical computing*.
518 Vienna, Austria: R Foundation for Statistical Computing. Retrieved from
519 <https://www.R-project.org/>
- 520 Revelle, W. (2021). *Psych: Procedures for psychological, psychometric, and*
521 *personality research*. Evanston, Illinois: Northwestern University. Retrieved
522 from <https://CRAN.R-project.org/package=psych>
- 523 Roenneberg, T., Wirz-Justice, A., & Mrosovsky, M. (2003). Life between clocks: Daily
524 temporal patterns of human chronotypes. *Journal of Biological Rhythms*,
525 18(1), 80–90.
- 526 Rosseel, Y. (2012). *lavaan: An R package for structural equation modeling*.

527 *Journal of Statistical Software*, 48(2), 1–36. Retrieved from

528 <https://www.jstatsoft.org/v48/i02/>

529 Ryu, C. (2021). *Dlookr: Tools for data diagnosis, exploration, transformation*.

530 Retrieved from <https://CRAN.R-project.org/package=dlookr>

531 Sarkar, D. (2008). *Lattice: Multivariate data visualization with r*. New York:

532 Springer. Retrieved from <http://lmdvr.r-forge.r-project.org>

533 Schönbrodt, F. D., & Perugini, M. (2013). At what sample size do correlations
534 stabilize? *Journal of Research in Personality*, 47(5), 609–612.

535 <https://doi.org/10.1016/j.jrp.2013.05.009>

536 Schumacker, R. E., & Lomax, R. G. (2004). *A beginner's guide to structural
537 equation modeling*. psychology press.

538 Shapiro, S. S., & Wilk, M. B. (1965). An analysis of variance test for normality
539 (complete samples). *Biometrika*, 52(3-4), 591–611.

540 <https://doi.org/10.1093/biomet/52.3-4.591>

541 Sijtsma, K. (2009). On the use, the misuse, and the very limited usefulness of
542 cronbach's alpha. *Psychometrika*, 74(1), 107.

543 Sjoberg, D. D., Curry, M., Hannum, M., Larmarange, J., Whiting, K., & Zabor, E. C.
544 (2021b). *Gtsummary: Presentation-ready data summary and analytic result*

545 *tables*. Retrieved from <https://CRAN.R-project.org/package=gtsummary>

546 Sjoberg, D. D., Curry, M., Hannum, M., Larmarange, J., Whiting, K., & Zabor, E. C.
547 (2021a). *Gtsummary: Presentation-ready data summary and analytic result*

548 *tables*. Retrieved from <https://CRAN.R-project.org/package=gtsummary>

549 Stauffer, R., Mayr, G. J., Dabernig, M., & Zeileis, A. (2009). Somewhere over the
550 rainbow: How to make effective use of colors in meteorological visualizations.

551 *Bulletin of the American Meteorological Society*, 96(2), 203–216.

552 <https://doi.org/10.1175/BAMS-D-13-00155.1>

553 Terry M. Therneau, & Patricia M. Grambsch. (2000). *Modeling survival data:*

- 554 *Extending the Cox model*. New York: Springer.
- 555 Ushey, K., McPherson, J., Cheng, J., Atkins, A., & Allaire, J. (2021). *Packrat: A*
556 *dependency management system for projects and their r package*
557 *dependencies*. Retrieved from <https://CRAN.R-project.org/package=packrat>
- 558 van Lissa, C. J. (2021). *tidySEM: Tidy structural equation modeling*. Retrieved
559 from <https://CRAN.R-project.org/package=tidySEM>
- 560 Velicer, W. (1976). Determining the Number of Components from the Matrix of
561 Partial Correlations. *Psychometrika*, 41, 321–327.
562 <https://doi.org/10.1007/BF02293557>
- 563 Venables, W. N., & Ripley, B. D. (2002). *Modern applied statistics with s* (Fourth).
564 New York: Springer. Retrieved from <https://www.stats.ox.ac.uk/pub/MASS4/>
- 565 Verriotto, J. D., Gonzalez, A., Aguilar, M. C., Parel, J.-M. A., Feuer, W. J., Smith,
566 A. R., & Lam, B. L. (2017). New methods for quantification of visual
567 photosensitivity threshold and symptoms. *Translational Vision Science &*
568 *Technology*, 6(4), 18–18.
- 569 Watkins, M. (2020). *A Step-by-Step Guide to Exploratory Factor Analysis with R*
570 *and RStudio*. <https://doi.org/10.4324/9781003120001>
- 571 Weinzaepflen, C., & Spitschan, M. (2021). Enlighten your clock: How your body
572 tells time. Open Science Framework. <https://doi.org/10.17605/OSF.IO/ZQXVH>
- 573 Wickham, H. (2011). The split-apply-combine strategy for data analysis. *Journal*
574 *of Statistical Software*, 40(1), 1–29. Retrieved from
575 <http://www.jstatsoft.org/v40/i01/>
- 576 Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag
577 New York. Retrieved from <https://ggplot2.tidyverse.org>
- 578 Wickham, H. (2019). *Stringr: Simple, consistent wrappers for common string*
579 *operations*. Retrieved from <https://CRAN.R-project.org/package=stringr>
- 580 Wickham, H. (2021a). *Forcats: Tools for working with categorical variables*

- (*factors*). Retrieved from <https://CRAN.R-project.org/package=forcats>
- Wickham, H. (2021b). *Tidyr: Tidy messy data*. Retrieved from <https://CRAN.R-project.org/package=tidyr>
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., ... Yutani, H. (2019). Welcome to the tidyverse. *Journal of Open Source Software*, 4(43), 1686. <https://doi.org/10.21105/joss.01686>
- Wickham, H., & Bryan, J. (2019). *Readxl: Read excel files*. Retrieved from <https://CRAN.R-project.org/package=readxl>
- Wickham, H., François, R., Henry, L., & Müller, K. (2021). *Dplyr: A grammar of data manipulation*. Retrieved from <https://CRAN.R-project.org/package=dplyr>
- Wickham, H., & Hester, J. (2021). *Readr: Read rectangular text data*. Retrieved from <https://CRAN.R-project.org/package=readr>
- Wilke, C. O. (2020). *Cowplot: Streamlined plot theme and plot annotations for 'ggplot2'*. Retrieved from <https://CRAN.R-project.org/package=cowplot>
- Winston Chang. (2014). *Extrafont: Tools for using fonts*. Retrieved from <https://CRAN.R-project.org/package=extrafont>
- Worthington, R. L., & Whittaker, T. A. (2006). Scale Development Research: A Content Analysis and Recommendations for Best Practices. *The Counseling Psychologist*, 34(6), 806–838. <https://doi.org/10.1177/0011000006288127>
- Wu, Y., & Hallett, M. (2017). Photophobia in neurologic disorders. *Translational Neurodegeneration*, 6(1), 26. <https://doi.org/10.1186/s40035-017-0095-3>
- Xie, Y., Wu, X., Tao, S., Wan, Y., & Tao, F. (2021). Development and validation of the self-rating of biological rhythm disorder for chinese adolescents. *Chronobiology International*, 1–7. <https://doi.org/10.1080/07420528.2021.1989450>
- Yu, C.-Y. (2002). *Evaluating cutoff criteria of model fit indices for latent variable models with binary and continuous outcomes* (Thesis). ProQuest

Dissertations Publishing.

Yu, L., Buysse, D. J., Germain, A., Moul, D. E., Stover, A., Dodds, N. E., ...

Pilkonis, P. A. (2011). Development of short forms from the PROMIS™ sleep disturbance and sleep-related impairment item banks. *Behavioral Sleep Medicine, 10*(1), 6–24. <https://doi.org/10.1080/15402002.2012.636266>

Zeileis, A., & Croissant, Y. (2010). Extended model formulas in R: Multiple parts and multiple responses. *Journal of Statistical Software, 34*(1), 1–13. <https://doi.org/10.18637/jss.v034.i01>

Zeileis, A., Fisher, J. C., Hornik, K., Ihaka, R., McWhite, C. D., Murrell, P., ...

Wilke, C. O. (2020). colorspace: A toolbox for manipulating and assessing colors and palettes. *Journal of Statistical Software, 96*(1), 1–49. <https://doi.org/10.18637/jss.v096.i01>

Zeileis, A., Hornik, K., & Murrell, P. (2009). Escaping RGBland: Selecting colors for statistical graphics. *Computational Statistics & Data Analysis, 53*(9), 3259–3270. <https://doi.org/10.1016/j.csda.2008.11.033>

Zhu, H. (2021). *kableExtra: Construct complex table with 'kable' and pipe syntax*.

Retrieved from <https://CRAN.R-project.org/package=kableExtra>

Zinbarg, R. E., Revelle, W., Yovel, I., & Li, W. (2005). Cronbach's α , Revelle's β , and McDonald's ω h: Their relations with each other and two alternative conceptualizations of reliability. *Psychometrika, 70*(1), 123–133.

Table 1

Releated Scales

Name	Author	Description	Relevant Items
Visual Light Sensitivity Questionnaire-8	Verriotto et al., 2017	Eight-question survey to assess the presence and severity of photosensitivity symptoms	NA
Office Light Survey	Eklundet al., 1996	A survey to assess electrical lighting environment in office	NA
Harvard Light Exposure Assessment Questionnaire	Bajaj et al., 2011	Self-administered semi-quantitative light questionnaire	NA
Hospital Lighting Survey	Dianat et el.,.2013	23 items questionnaire to assess light environment in a hospital	NA
Morningness-Eveningness Questionnaire	Horne et al.,1976	19 items questionnaire to understand your body clock	NA

Table 1

Releated Scales (continued)

Name	Author	Description	Relevant Items
Munich Chronotype Questionnaire (MCTQ)	Roenneberg et al.,2003	17 items questionnaire to understand individuals phase of entrainment	NA
Assessment of Sleep Environment	Olivier et.al.,2016]	13 items questionnaire measuring your sleep environment quality	NA
The Pittsburgh Sleep Quality Index (PSQI)	Buysse ei al.,1989	9 items inventory to measure sleep quality and sleeping pattern	NA
Self-Rating of Biological Rhythm Disorder for Adolescents (SBRDA)	Xie et al.,2021	29 Items questionnaire assessing four dimensions of biological rhythm disorder in adolescents	Item 3,22-25 and 29

Table 1

Releated Scales (continued)

Name	Author	Description	Relevant Items
Photosensitivity Assessment Questionnaire (PAQ)	Wu et al.,2017	16 dichotomous (yes/no) items questionnaire to assess "photophobia" and "photophilia"	All itms

Table 2

Dempgraphics

Variable	Overall, N = 690	1. EFA Sample, N = 428	2. CFA Sample, N = 262	p-value	q-value
Age	32.95 (14.57)	32.99 (15.11)	32.89 (13.66)	0.5	0.5
Sex				0.14	0.4
Female	325 (47%)	189 (44%)	136 (52%)		
Male	351 (51%)	230 (54%)	121 (46%)		
Other	14 (2.0%)	9 (2.1%)	5 (1.9%)		
Gender-Variant Identity	49 (7.2%)	33 (7.8%)	16 (6.2%)	0.4	0.5
Native English Speaker	320 (46%)	191 (45%)	129 (49%)	0.2	0.5
Occupational Status				0.040	0.2
Work	396 (57%)	235 (55%)	161 (61%)		
School	174 (25%)	122 (29%)	52 (20%)		
Neither	120 (17%)	71 (17%)	49 (19%)		
Occupational setting				0.3	0.5
Home office/Home schooling	303 (44%)	194 (45%)	109 (42%)		
Face-to-face work/Face-to-face schooling	109 (16%)	68 (16%)	41 (16%)		
Combination of home- and face-to-face- work/schooling	147 (21%)	94 (22%)	53 (20%)		
Neither (no work or school, or in vacation)	131 (19%)	72 (17%)	59 (23%)		

¹ Mean (SD); n (%)² Wilcoxon rank sum test; Pearson's Chi-squared test³ False discovery rate correction for multiple testing

Table 3

Descriptive Statistics

	Mean	SD	Skew	Kurtosis	Shapiro-Wilk Statistics	Item-Total Correlation
Item1	2.27	1.39	0.74	-0.81	0.81*	.25
Item2	2.87	1.59	0.08	-1.60	0.83*	.19
Item3	3.36	1.38	-0.48	-1.03	0.87*	.16
Item4	1.47	1.18	2.38	4.00	0.43*	.28
Item5	4.01	1.40	-1.22	0.07	0.70*	.13
Item6	2.79	1.55	0.19	-1.48	0.85*	.20
Item7	2.26	1.25	0.70	-0.60	0.85*	.19
Item8	2.97	1.20	-0.06	-0.94	0.91*	-.10
Item9	2.94	1.03	-0.12	-0.40	0.91*	.10
Item10	2.74	1.04	0.09	-0.74	0.91*	.28
Item11	2.18	0.90	0.60	0.12	0.86*	.26
Item12	2.36	1.22	0.59	-0.62	0.87*	.25
Item13	2.73	1.46	0.20	-1.36	0.87*	.33
Item14	2.14	1.31	0.77	-0.78	0.80*	.26
Item15	3.26	1.09	-0.26	-0.45	0.91*	.14
Item16	1.56	1.23	2.00	2.45	0.50*	.32
Item17	1.54	1.21	2.07	2.75	0.49*	.31
Item18	1.12	0.49	5.02	27.80	0.25*	.16
Item19	1.05	0.36	7.23	52.98	0.13*	.18
Item20	1.04	0.33	8.99	85.28	0.10*	.16
Item21	1.14	0.59	4.79	24.05	0.25*	.16
Item22	3.57	1.07	-0.65	-0.17	0.88*	.21
Item23	2.56	1.27	0.33	-1.00	0.89*	.11

Table 3 continued

	Mean	SD	Skew	Kurtosis	Shapiro-Wilk Statistics	Item-Total Correlation
Item24	4.14	0.99	-1.23	1.14	0.79*	.19
Item25	2.59	1.41	0.27	-1.27	0.86*	.19
Item26	2.25	1.27	0.69	-0.64	0.84*	.18
Item27	3.80	1.29	-0.87	-0.42	0.82*	.17
Item28	3.76	1.14	-0.68	-0.45	0.86*	.00
Item29	2.44	1.31	0.38	-1.14	0.86*	.11
Item30	1.48	1.11	2.18	3.35	0.48*	.24
Item31	3.00	1.62	-0.08	-1.61	0.83*	.44
Item32	3.55	1.65	-0.60	-1.34	0.76*	.43
Item33	3.62	1.64	-0.68	-1.25	0.74*	.32
Item34	3.42	1.83	-0.45	-1.69	0.69*	.33
Item35	3.86	1.67	-0.99	-0.85	0.65*	.23
Item36	1.54	1.25	2.13	2.86	0.46*	.36
Item37	1.33	0.91	3.03	8.43	0.41*	.01
Item38	4.30	1.08	-1.79	2.53	0.67*	.22
Item39	1.96	0.98	1.02	0.69	0.82*	.05
Item40	2.16	1.19	0.71	-0.54	0.84*	.14
Item41	1.31	0.81	2.75	6.92	0.43*	.21
Item42	3.93	1.48	-1.06	-0.44	0.71*	.18
Item43	1.64	1.18	1.79	2.02	0.60*	.15
Item44	3.51	1.30	-0.70	-0.59	0.85*	.39
Item45	2.22	1.48	0.71	-1.02	0.76*	.30
Item46	1.76	1.23	1.35	0.44	0.66*	.38
Item47	2.11	1.17	0.77	-0.39	0.83*	.32

Table 3 continued

	Mean	SD	Skew	Kurtosis	Shapiro-Wilk Statistics	Item-Total Correlation
Item48	2.60	1.25	0.29	-0.86	0.89*	.35

Note. * $p < .001$

Table 4

Factor loadings and communality of the retained items

item	PA1	PA2	PA3	PA4	PA5	Communality	Uniqueness
item16	0.99					0.993	0.007
item36	0.94					0.899	0.101
item17	0.8					0.658	0.342
item11		0.79				0.642	0.358
item10		0.76				0.592	0.408
item12		0.65				0.465	0.535
item7		0.5				0.267	0.733
item8		-0.49				0.252	0.748
item9		0.32				0.113	0.887
item27			0.8			0.658	0.342
item3			0.8			0.682	0.318
item40			0.65			0.464	0.536
item30			0.45			0.353	0.647
item41			0.36			0.329	0.671
item33				0.74		0.555	0.445
item32				0.73		0.624	0.376
item35				0.66		0.454	0.546
item37				-0.39		0.174	0.826
item38				0.38		0.178	0.822
item46					0.6	0.422	0.578
item45					0.59	0.374	0.626
item25					0.41	0.193	0.807
item4					0.41	0.219	0.781
item1					0.4	0.17	0.83
item26					0.35	0.165	0.835
% of Variance	0.1	0.1	0.09	0.08	0.06		

Note. Only loading higher than .30 is reported

Table 5

Fit indices of CFA

Model	Chi-Square	df	CFI	TLI	RMSEA	RMSEA 90% Lower CI	RMSEA 90% Upper CI	SRMR
Five factor model:25	448.51	222.00	.94	0.93	0.06	0.05	0.07	0.12
Five factor model:23	346.59	221.00	.97	0.96	0.05	0.04	0.06	0.09

Note. df: Degrees of Freedom; CFI: Comparative Fit Index; TLI: Tucker Lewis Index; RMSEA: Root Mean Square Error of Approximation; CI: Confidence Interval; SRMR: Standardized Root Mean Square

Table 6

Invariance Analysis

	Chi-Square	df	CFI	TLI	RMSEA	RMSEA 90% Lower CI	RMSEA 90% Upper	SRMR	Chi-Sqr comparison	df*	p
Configural	632.20	442.00	0.95	0.94	0.06	0.05	0.07	0.13	-	-	-
Metric	644.58	458.00	0.95	0.95	0.06	0.05	0.07	0.13	18.019a	16	0.323
Scalar	714.19	522.00	0.95	0.95	0.05	0.04	0.06	0.13	67.961b	64	0.344
Residual	714.19	522.00	0.95	0.95	0.05	0.04	0.06	0.13	0c	0	NA
Structural	691.49	542.00	0.96	0.96	0.05	0.04	0.06	0.13	12.617d	20	0.893

Note. a = Metric vs Configural; b = Scalar vs Metric; c = Residual vs Scalar; d = Structural vs Residual; * = df of model comparison

Table 7

IRT Item parameters for the LEBA Scale

	a	b1	b2	b3	b4
item16	28.55	0.78	0.90	1.06	1.40
item36	4.49	0.94	1.08	1.23	1.40
item17	2.81	0.97	1.11	1.38	1.62
item11	3.27	-0.79	0.65	1.54	2.31
item10	3.07	-1.27	-0.09	0.82	2.00
item12	1.72	-0.67	0.44	1.28	2.11
item7	1.09	-0.50	0.73	1.63	2.97
Ritem8	1.19	-2.26	-0.48	0.64	1.91
item9	0.91	-2.63	-0.96	1.11	3.49
item27	2.21	-1.88	-1.19	-0.73	0.30
item3	3.03	-1.24	-0.77	-0.20	0.66
item40	1.55	-0.51	0.46	1.32	2.22
item30	0.49	3.27	3.74	4.64	6.52
item41	0.51	3.87	4.78	6.39	8.91
item32	1.62	-1.03	-0.78	-0.42	0.16
item35	1.36	-1.09	-0.98	-0.75	-0.40
item38	0.40	-7.50	-5.58	-4.25	-0.91
item33	13.51	-0.66	-0.48	-0.24	0.13
item46	2.22	0.68	0.89	1.38	2.17
item45	1.51	0.30	0.55	1.17	1.91
item25	0.52	-1.37	-0.04	1.89	4.22
item4	0.84	2.44	2.80	3.18	3.67
item1	0.39	-0.91	1.52	3.25	5.53

Note. a = item discrimination parameter; b(1-4)
= response category difficulty parameter




Figure 1. Development of LEBA




Figure 2. Correlation plot of the items




Figure 3. Factor Identification (A) Parallel analysis (B) Scree Plot (C) Hull Method




Figure 4. Five Factor Solution




Figure 5. Histogram of residulas: five-factor solution

































































































Summary results of LEBA (N =690)											
Items	Summary Statistics				Graphics		Response Pattern				
LEBA Items	n	Mean	Median	SD	Histogram ¹	Density ²	Never	Rarely	Sometimes	Often	Always
EFA (n = 428)											
Item01	428	2.3	2.0	1.4			22.20% (95)	12.38% (53)	10.51% (45)	12.62% (54)	42.29% (181)
Item03	428	3.4	4.0	1.4			11.45% (49)	31.07% (133)	24.30% (104)	17.29% (74)	15.89% (68)
Item04	428	1.5	1.0	1.2			3.50% (15)	2.10% (9)	8.18% (35)	2.10% (9)	84.11% (360)
Item07	428	2.3	2.0	1.2			27.80% (119)	12.38% (53)	6.54% (28)	17.29% (74)	35.98% (154)
Item08	428	3.0	3.0	1.2			25.93% (111)	22.20% (95)	13.79% (59)	27.80% (119)	10.28% (44)
Item09	428	2.9	3.0	1.0			19.63% (84)	22.43% (96)	5.84% (25)	41.82% (179)	10.28% (44)
Item10	428	2.7	3.0	1.0			31.31% (134)	21.96% (94)	3.50% (15)	31.31% (134)	11.92% (51)
Item11	428	2.2	2.0	0.9			46.26% (198)	7.01% (30)	1.17% (5)	23.13% (99)	22.43% (96)
Item12	428	2.4	2.0	1.2			29.67% (127)	12.15% (52)	6.78% (29)	21.50% (92)	29.91% (128)
Item16	428	1.6	1.0	1.2			4.21% (18)	4.67% (20)	7.48% (32)	3.97% (17)	79.67% (341)
Item17	428	1.5	1.0	1.2			3.27% (14)	3.27% (14)	7.71% (33)	5.14% (22)	80.61% (345)
Item25	428	2.6	3.0	1.4			13.79% (59)	17.99% (77)	11.68% (50)	22.20% (95)	34.35% (147)
Item26	428	3.7	4.0	1.3			10.98% (47)	23.36% (100)	38.32% (164)	20.09% (86)	7.24% (31)
Item27	428	3.8	4.0	1.3			11.21% (48)	30.37% (130)	38.79% (166)	11.21% (48)	8.41% (36)
Item30	428	1.5	1.0	1.1			3.27% (14)	5.37% (23)	4.67% (20)	4.91% (21)	81.78% (350)
Item32	428	3.6	4.0	1.6			7.01% (30)	14.95% (64)	46.73% (200)	8.18% (35)	23.13% (99)
Item33	428	3.6	4.0	1.6			7.01% (30)	14.49% (62)	49.30% (211)	7.24% (31)	21.96% (94)
Item35	428	3.9	5.0	1.7			1.87% (8)	9.35% (40)	62.15% (266)	3.74% (16)	22.90% (98)
Item36	428	1.5	1.0	1.3			3.04% (13)	2.34% (10)	9.35% (40)	3.04% (13)	82.24% (352)
Item37	428	2.3	2.0	1.3			23.36% (100)	10.98% (47)	7.24% (31)	20.09% (86)	38.32% (164)
Item38	428	4.3	5.0	1.1			3.50% (15)	27.57% (118)	58.18% (249)	5.37% (23)	5.37% (23)
Item40	428	2.2	2.0	1.2			25.00% (107)	11.45% (49)	4.44% (19)	19.63% (84)	39.49% (169)
Item41	428	1.3	1.0	0.8			4.67% (20)	3.04% (13)	1.17% (5)	6.07% (26)	85.05% (364)
Item45	428	2.2	1.0	1.5			7.01% (30)	11.92% (51)	11.68% (50)	16.36% (70)	53.04% (227)
Item46	428	1.8	1.0	1.2			7.71% (33)	8.88% (38)	4.67% (20)	11.68% (50)	67.06% (287)
CFA (n =262)											
Item01	262	2.3	2.0	1.4			22.52% (59)	10.69% (28)	11.83% (31)	14.50% (38)	40.46% (106)
Item03	262	3.7	4.0	1.3			7.25% (19)	28.24% (74)	35.11% (92)	17.56% (46)	11.83% (31)
Item04	262	1.3	1.0	0.8			2.29% (6)	3.05% (8)	1.91% (5)	3.44% (9)	89.31% (234)
Item07	262	2.1	2.0	1.2			23.66% (62)	14.12% (37)	4.58% (12)	14.50% (38)	43.13% (113)
Item08	262	3.0	3.0	1.2			32.06% (84)	22.90% (60)	14.12% (37)	20.99% (55)	9.92% (26)
Item09	262	2.9	3.0	1.1			22.14% (58)	26.34% (69)	4.20% (11)	34.35% (90)	12.98% (34)
Item10	262	2.6	3.0	1.1			29.39% (77)	21.37% (56)	2.67% (7)	29.01% (76)	17.56% (46)
Item11	262	2.1	2.0	0.9			46.56% (122)	5.34% (14)	1.91% (5)	20.23% (53)	25.95% (68)
Item12	262	2.3	2.0	1.2			30.92% (81)	11.45% (30)	6.49% (17)	19.08% (50)	32.06% (84)
Item16	262	1.6	1.0	1.3			3.44% (9)	5.73% (15)	8.40% (22)	4.20% (11)	78.24% (205)
Item17	262	1.6	1.0	1.2			3.44% (9)	2.67% (7)	8.40% (22)	5.34% (14)	80.15% (210)
Item25	262	2.5	2.0	1.4			18.32% (48)	16.79% (44)	10.31% (27)	21.76% (57)	32.82% (86)
Item27	262	4.0	4.0	1.2			7.25% (19)	33.59% (88)	45.04% (118)	8.02% (21)	6.11% (16)
Item30	262	1.4	1.0	1.1			2.67% (7)	6.11% (16)	3.44% (9)	4.20% (11)	83.59% (219)
Item32	262	3.4	4.0	1.7			4.20% (11)	16.79% (44)	41.80% (109)	11.45% (30)	25.95% (68)
Item33	262	3.1	3.0	1.7			6.11% (16)	14.12% (37)	35.50% (93)	11.83% (31)	32.44% (85)
Item35	262	3.6	5.0	1.8			2.67% (7)	6.49% (17)	56.11% (147)	7.25% (19)	27.48% (72)
Item36	262	1.6	1.0	1.3			3.44% (9)	3.44% (9)	9.54% (25)	3.05% (8)	80.53% (211)
Item38	262	4.3	5.0	1.1			7.53% (20)	21.37% (56)	60.31% (158)	6.49% (17)	4.20% (11)
Item40	262	2.5	2.0	1.3			27.10% (71)	12.21% (32)	11.07% (29)	18.70% (49)	30.92% (81)
Item41	262	1.2	1.0	0.7			3.82% (10)	2.67% (7)	1.15% (3)	2.29% (6)	90.08% (236)
Item45	262	2.0	1.0	1.4			5.34% (14)	11.83% (31)	9.16% (24)	9.54% (25)	64.12% (168)
Item46	262	1.6	1.0	1.2			2.67% (7)	9.54% (25)	4.20% (11)	8.02% (21)	75.57% (198)
¹ Histogram											
² Density											

Figure 6




Figure 7. (A) Five Factor Model of LEBA




Figure 8. Item information curves (A) blue filter (B) natural light (C) smart device (D) sleep environment (E) electric light




Figure 9. Test information curves (A) blue filter (B) natural light (C) smart device (D) sleep environment (E) electric light

Appendix A

Table A1

Factor loadings and communality of the retained items(Minmum Residual)

item	MR1	MR2	MR3	MR4	MR5	Communality	Uniqueness
item16	1					0.996	0.004
item36	0.94					0.897	0.103
item17	0.8					0.658	0.342
item11		0.79				0.642	0.358
item10		0.76				0.592	0.408
item12		0.65				0.465	0.535
item7		0.5				0.267	0.733
item8		-0.49				0.252	0.748
item9		0.32				0.113	0.887
item27			0.8			0.659	0.341
item3			0.8			0.683	0.317
item40			0.65			0.464	0.536
item30			0.45			0.353	0.647
item41			0.36			0.329	0.671
item33				0.74		0.555	0.445
item32				0.73		0.623	0.377
item35				0.66		0.455	0.545
item37				-0.39		0.175	0.825
item38				0.38		0.178	0.822
item46					0.6	0.422	0.578
item45					0.59	0.374	0.626
item25					0.41	0.193	0.807
item4					0.41	0.219	0.781
item1					0.4	0.17	0.83
item26					0.35	0.165	0.835
% of Variance	0.1	0.1	0.09	0.08	0.06		

Note. Only loading higher than .30 is reported

Appendix B

Factor Analysis with Unmerged Response Option




Figure B1. Correlation plot of the items

Horn’s parallel analysis with 500 iterations indicated a five-factor solution. However, Scree plot and the MAP method suggested 6-factor solution. five-factor solution . As a result, we tested both five-factor and six-factor solutions.

Five Factor Solution[Unmerged Responses] (24 Items)

F1

- I use light therapy applying a blue light box.
- I use light therapy applying a light visor.
- I use light therapy applying a white light box.
- I use light therapy applying another form of light device.

Five Factor Solution[Unmerged Responses] (24 Items)

I use an alarm with a dawn simulation light.

F2

I spend more than 3 hours per day (in total) outside.

I spend between 1 and 3 hours per day (in total) outside.

I spend as much time outside as possible.

I spend 30 minutes or less per day (in total) outside.

I go for a walk or exercise outside within 2 hours after waking up.

I spend between 30 minutes and 1 hour per day (in total) outside.

F3

I look at my mobile phone screen immediately after waking up.

I use my mobile phone within 1 hour before attempting to fall asleep.

I check my phone when I wake up at night.

F4

I use a blue-filter app on my computer screen within 1 hour before attempting to fall asleep.

I seek out knowledge on how to improve my light exposure.

I dim my computer screen within 1 hour before attempting to fall asleep.

I discuss the effects of light on my body with other people.

I modify my light environment to match my current needs.

I dim my room light within 1 hour before attempting to fall asleep.

I use as little light as possible when I get up during the night.

F5

I wear blue-filtering, orange-tinted, and/or red-tinted glasses indoors during the day.

I wear blue-filtering, orange-tinted, and/or red-tinted glasses outdoors during the day.

Five Factor Solution[Unmerged Responses] (24 Items)

I wear blue-filtering, orange-tinted, and/or red-tinted glasses within 1 hour before attempting to fall asleep.




Figure B2. Factor Identification (A) Parallel analysis (B) Scree Plot

Table B1

Descriptive Statistics for Unmerged response options

	Mean	SD	Skew	Kurtosis	Shapiro-Wilk Statistics	Item-Total Correlation
Item1	2.16	1.51	0.49	-0.86	0.90*	.21
Item2	2.76	1.75	-0.10	-1.42	0.88*	.20
Item3	3.34	1.43	-0.58	-0.77	0.88*	.18
Item4	1.30	1.31	1.93	2.92	0.62*	.32
Item5	3.95	1.56	-1.42	0.75	0.70*	.19
Item6	2.70	1.66	0.02	-1.33	0.90*	.18
Item7	2.23	1.28	0.60	-0.59	0.89*	.18
Item8	2.95	1.24	-0.19	-0.70	0.93*	-.07
Item9	2.92	1.09	-0.37	0.11	0.91*	.14
Item10	2.73	1.07	-0.03	-0.52	0.92*	.27
Item11	2.17	0.93	0.44	0.20	0.89*	.25
Item12	2.34	1.26	0.46	-0.58	0.91*	.24
Item13	2.71	1.49	0.14	-1.29	0.89*	.28
Item14	2.11	1.34	0.68	-0.78	0.84*	.24
Item15	3.26	1.11	-0.34	-0.21	0.91*	.11
Item16	1.46	1.31	1.71	1.90	0.65*	.33
Item17	1.43	1.30	1.76	2.12	0.64*	.30
Item18	0.92	0.67	2.00	9.41	0.62*	.32
Item19	0.85	0.56	1.71	10.74	0.55*	.34
Item20	0.83	0.54	1.76	13.92	0.53*	.31
Item21	0.94	0.75	2.46	10.66	0.58*	.27
Item22	3.57	1.08	-0.72	0.08	0.88*	.19
Item23	2.53	1.31	0.22	-0.91	0.92*	.11
Item24	4.13	1.01	-1.39	2.01	0.78*	.19
Item25	2.57	1.43	0.22	-1.23	0.88*	.17
Item26	2.22	1.22	0.52	-0.62	0.88*	.16

Table B2

Factor loadings and communality of the retained items [Unmerged Responses]

item	PA1	PA2	PA5	PA3	PA4	Communality	Uniqueness	Complexity
item19	0.99					1.007	-0.007	1.058
item20	0.91					0.874	0.126	1.114
item18	0.82					0.711	0.289	1.123
item21	0.8					0.683	0.317	1.163
item4	0.47					0.25	0.75	1.298
item11		0.83				0.687	0.313	1.007
item10		0.81				0.67	0.33	1.031
item12		0.56				0.371	0.629	1.374
item8		-0.44				0.206	0.794	1.106
item7		0.42				0.226	0.774	1.614
item9		0.33				0.115	0.885	1.1
item16			0.95			0.946	0.054	1.097
item17			0.74			0.595	0.405	1.168
item36	0.3		0.73			0.653	0.347	1.431
item3				0.85		0.746	0.254	1.048
item27				0.78		0.624	0.376	1.028
item40				0.71		0.512	0.488	1.05
item35					0.58	0.351	0.649	1.091
item48					0.57	0.354	0.646	1.144
item33					0.55	0.32	0.68	1.085
item47					0.52	0.294	0.706	1.186
item44					0.45	0.216	0.784	1.145
item31					0.41	0.206	0.794	1.477
item38					0.33	0.129	0.871	1.317
% of Variance	0.15	0.09	0.09	0.08	0.08			

Note. Only loading higher than .30 is reported




Figure B3. Histogram of residulas: five-factor solution

Appendix C

Disclaimer: This is a non-public version of LEBA (dated November 12, 2021) and still a work in progress. Please do not distribute!

LEBA captures light exposure-related behaviours on a 5 point Likert type scale ranging from 1 to 5 (Never/Does not apply/I don't know = 1; Rarely = 2; Sometimes = 3; Often = 4; Always = 5). The score of each factor is calculated by the summation of scores of items belonging to the corresponding factor. The following instruction is given before displaying the items: "Please indicate how often you performed the following behaviours in the past 4 weeks."

Appendix D
LEBA Long Form (23 Items)

	Items	Never/Does not apply/I don't know	Rarely	Sometimes	Often	Always
1	I wear blue-filtering, orange-tinted, and/or red-tinted glasses indoors during the day.					
2	I wear blue-filtering, orange-tinted, and/or red-tinted glasses outdoors during the day.					
3	I wear blue-filtering, orange-tinted, and/or red-tinted glasses within 1 hour before attempting to fall asleep.					
4	I spend 30 minutes or less per day (in total) outside.					

	Items	Never/Does not apply/I don't know	Rarely	Sometimes	Often	Always
5	I spend between 1 and 3 hours per day (in total) outside.					
6	I spend between 30 minutes and 1 hour per day (in total) outside.					
7	I spend more than 3 hours per day (in total) outside.					
8	I spend as much time outside as possible.					
9	I go for a walk or exercise outside within 2 hours after waking up.					
10	I use my mobile phone within 1 hour before attempting to fall asleep.					

	Items	Never/Does not apply/I don't know	Rarely	Sometimes	Often	Always
11	I look at my mobile phone screen immediately after waking up.					
12	I check my phone when I wake up at night.					
13	I look at my smartwatch within 1 hour before attempting to fall asleep.					
14	I look at my smartwatch when I wake up at night.					
15	I dim my mobile phone screen within 1 hour before attempting to fall asleep.					
16	I use a blue-filter app on my computer screen within 1 hour before attempting to fall asleep.					

	Items	Never/Does not apply/I don't know	Rarely	Sometimes	Often	Always
17	I use as little light as possible when I get up during the night.					
18	I dim my computer screen within 1 hour before attempting to fall asleep.					
19	I use tunable lights to create a healthy light environment.					
20	I use LEDs to create a healthy light environment.					
21	I use a desk lamp when I do focused work.					
22	I use an alarm with a dawn simulation light.					
23	I turn on the lights immediately after waking up.					

639 Latent Structure, Reliability and Structural Validity

640 The long form of LEBA consists 23 items with five factors.

Factor names	Items	Reliability Coefficients:	Reliability Coefficients:
		McDonald's Omega	Cronbach's alpha
F1: Wearing blue light filters	1-3	.93	.90
F2: Spending time outdoors	4-9 (Item 4 is reversed)	.80	.78
F3: Using phone and smartwatch in bed	10-14	.61	.62
F4: Using light before bedtime	15-18	.72	.62
F5: Using light in the morning and during daytime	19-23	.45	.41
		.73(Total scale)	

641 LEBA -long form showed satisfactory structural validity (CFI =.97; TLI = .96; RMSEA = .05[.04-.06, 90% CI]; SRMR =
642 .09).

643 **How to cite:**

Appendix E
LEBA Short Form (17 Items)

	Short Form (17 Items)	Never/Does not apply/I don't know	Rarely	Sometimes	Often	Always
01	I wear blue-filtering, orange-tinted, and/or red-tinted glasses indoors during the day.					
02	I wear blue-filtering, orange-tinted, and/or red-tinted glasses outdoors during the day.					
03	I wear blue-filtering, orange-tinted, and/or red-tinted glasses within 1 hour before attempting to fall asleep.					
04	I spend 30 minutes or less per day (in total) outside.					

	Short Form (17 Items)	Never/Does not apply/I don't know	Rarely	Sometimes	Often	Always
05	I spend between 1 and 3 hours per day (in total) outside.					
06	I spend more than 3 hours per day (in total) outside.					
07	I spend as much time outside as possible.					
08	I go for a walk or exercise outside within 2 hours after waking up.					
09	I use my mobile phone within 1 hour before attempting to fall asleep.					
10	I look at my mobile phone screen immediately after waking up.					
11	I check my phone when I wake up at night.					

	Short Form (17 Items)	Never/Does not apply/I don't know	Rarely	Sometimes	Often	Always
12	I dim my mobile phone screen within 1 hour before attempting to fall asleep.					
13	I use a blue-filter app on my computer screen within 1 hour before attempting to fall asleep.					
14	I dim my computer screen within 1 hour before attempting to fall asleep.					
15	I use tunable lights to create a healthy light environment.					
16	I use LEDs to create a healthy light environment.					
17	I use an alarm with a dawn simulation light.					

644 Latent Structure, Reliability and Structural Validity

645 The short form of LEBA consists 23 items with five factors.

Factor names	Items
F1: Wearing blue light filters	1-3
F2: Spending time outdoors	4-8 (Item 4 is reversed)
F3: Using phone and smart-watch in bed	9-11
F4: Using light before bedtime	12-14
F5: Using light in the morning and during daytime	15-17

646 **How to cite:**