

Light Exposure Behavior Assessment (LEBA): Develop of a novel instrument to capture light exposure-related behaviours

Mushfiqul Anwar Siraji¹, Rafael Robert Lazar², Manuel Spitschan³, & Manuel Spitschan³

¹ Department of Psychology, Jeffrey Cheah School of Medicine and Health Sciences, Monash University, Malaysia

² University of Basel

Author Note

Add complete departmental affiliations for each author here. Each new line herein must be indented, like this line.

Enter author note here.

The authors made the following contributions. Mushfiqul Anwar Siraji: Data Analysis, Writing - Original Draft Preparation, Data Visualization; Rafael Robert Lazar: Data Analysis, Writing - Original Draft Preparation, Data Visualization; Manuel Spitschan: Data Analysis, Writing - Original Draft Preparation, Data Visualization; Manuel Spitschan: Data Analysis, Writing - Original Draft Preparation, Data Visualization.

Correspondence concerning this article should be addressed to Manuel Spitschan, .
E-mail:

Abstract

One or two sentences providing a **basic introduction** to the field, comprehensible to a scientist in any discipline.

Two to three sentences of **more detailed background**, comprehensible to scientists in related disciplines.

One sentence clearly stating the **general problem** being addressed by this particular study.

One sentence summarizing the main result (with the words “**here we show**” or their equivalent).

Two or three sentences explaining what the **main result** reveals in direct comparison to what was thought to be the case previously, or how the main result adds to previous knowledge.

One or two sentences to put the results into a more **general context**.

Two or three sentences to provide a **broader perspective**, readily comprehensible to a scientist in any discipline.

Keywords: keywords

Word count: X

Light Exposure Behavior Assessment (LEBA): Develop of a novel instrument to capture light exposure-related behaviours

Introduction

Methods

Participants

This line is just a test for pushing in the github repo.

Material

Procedure

Our study had four objectives. First, to develop an instrument to assess individual's light exposure behavior . Second, to conduct an exploratory factor analysis(EFA) to understand the latent structure. Third to gather structural validity evidence for the latent structure obtained in EFA (Furr, 2014). Lastly, we gathered item information using Item response theory (IRT)(Baker, 2017)

Data Collection. Timeline of data collection, ethical approval, mode of data collection, how consent was recorded.

Item generation and Content Validity: Expert Panel Review. How we developed the 48 items?

Analytic Strategies

We used R (version 4.1.0), including several R-packages for our analyses. Necessary assumptions of EFA, including sample adequacy, normality assumptions, quality of correlation matrix were assessed. Our data violated both the univariate and multivariate

normality assumptions. Due to these violations and the ordinal nature of our response data we used polychoric correlation matrix (Desjardins & Bulut, 2018) for the EFA. We employed principal axis (pa) a factor extraction method with varimax rotation. PA is apparently robust to the normality assumption violations (Watkins, 2020). The obtained latent structure was confirmed by weighted least squares (WLS) extraction method as well. We used a combination factor identification method including scree plot (Cattell, 1966), Horn's parallel analysis (Horn, 1965), minimum average partials method (Velicer, 1976), and hull method (Lorenzo-Seva, Timmerman, & Kiers, 2011) to identify factor numbers. Additionally, to identify the simple structure we followed the following guidelines recommended by psychometricians (i) no factors with fewer than three items (ii) no factors with a factor loading < 0.3 (iii) no items with cross-loading greater than .3 across factors (Bandalos & Finney, 2018; Child, 2006; Mulaik, 2009; Watkins, 2020)

Results

Sampling adequacy was investigated by Kaiser-Meyer-Olkin (KMO) measures of sampling adequacy (Kaiser, 1974). The overall KMO value for 23 items was 0.63 which was above the cutoff value of .50 indicating a mediocre sample (Hutcheson, 1999).

Table 1 summarizes the univariate descriptive statistics for the 48 items. Some of the items were skewed with high Kurtosis values. The Shapiro-Wilk test of normality (Shapiro & Wilk, 1965) indicated all the items violated normality assumptions. Multivariate normality assumptions were investigated by Mardia's test (Mardia, 1970). Multivariate skew = 583.80 ($p < 0.001$) and multivariate kurtosis = 2,749.15 ($p < 0.001$) indicated multivariate normality assumptions violation. Due to these violations and ordinal nature of the response data polychoric correlations over Pearson's correlations was chosen (Desjardins & Bulut, 2018). Bartlett's test of sphericity (Bartlett, 1954), $\chi^2 (1128) = 5042.86$, $p < .001$ indicated the correlations between items are adequate for the EFA. However only 4.96% of the inter-item correlation coefficients were greater than .30 in the obtained matrix. The inter item

correlation ranged between .44 to .91. The corrected item-total correlations ranged between .10 to .44.

Scree plot (Fig@ref(fig:fac.id)) suggested a six-factor solution.Horn's parallel analysis (Horn, 1965), like the Monte Carlo study, draws several sets of random data with the same number of participants as the original data set and compares the mean eigenvalues among the simulated and original data sets to retain optimal factors.This extraction method also supported a five-factor model. In our data set parallel analysis with 500 iterations indicated six-factor solution. However, In MAP method (Velicer, 1976) and Hull method (Lorenzo-Seva, Timmerman, & Kiers, 2011) suggested a five-factor solution Parallel analysis is also more immune to the normality assumptions violation(**RN1263?**).

Confirmatory Factor Analysis

Discussion

References

- Aust, F., & Barth, M. (2020). *papaja: Prepare reproducible APA journal articles with R Markdown*. Retrieved from <https://github.com/crsh/papaja>
- Baker, F. B. (2017). *The Basics of Item Response Theory Using R* (1st ed. 2017.). Springer.
- Bandalos, D. L., & Finney, S. J. (2018). Factor analysis: Exploratory and confirmatory. In *The reviewer's guide to quantitative methods in the social sciences* (pp. 98–122). Routledge.
- Barth, M. (2021). *tinylabels: Lightweight variable labels*. Retrieved from <https://github.com/mariusbarth/tinylabels>
- Bartlett, M. (1954). A Note on the Multiplying Factors for Various Chi-square Approximations. *Journal of the Royal Statistical Society. Series B, Methodological*, 16(2), 296–298.
- Buchanan, E. M., Gillenwaters, A., Scofield, J. E., & Valentine, K. D. (2019). *MOTE: Measure of the Effect: Package to assist in effect size calculations and their confidence intervals*. Retrieved from <http://github.com/doomlab/MOTE>
- Cattell, R. B. (1966). The Scree Test For The Number Of Factors. *Multivariate Behavioral Research*, 1(2), 245–276.
https://doi.org/10.1207/s15327906mbr0102_10
- Child, D. (2006). *Essentials of factor analysis* (3rd ed.). New York: Continuum.
- Desjardins, C., & Bulut, O. (2018). *Handbook of Educational Measurement and Psychometrics Using R*. <https://doi.org/10.1201/b20498>
- Dinno, A. (2018). *Paran: Horn's test of principal components/factors*. Retrieved from <https://CRAN.R-project.org/package=paran>

- 118 Epskamp, S. (2019). *semPlot: Path diagrams and visual analysis of various SEM*
119 *packages' output*. Retrieved from <https://CRAN.R-project.org/package=semPlot>
- 120 Epskamp, S., Cramer, A. O. J., Waldorp, L. J., Schmittmann, V. D., & Borsboom, D.
121 (2012). qgraph: Network visualizations of relationships in psychometric data.
122 *Journal of Statistical Software*, 48(4), 1–18.
- 123 Furr, R. M. (2014). *Psychometrics : An introduction* (2nd ed.). Thousand Oaks:
124 Thousand Oaks : SAGE.
- 125 Henry, L., & Wickham, H. (2020). *Purrr: Functional programming tools*. Retrieved
126 from <https://CRAN.R-project.org/package=purrr>
- 127 Horn, J. L. (1965). A rationale and test for the number of factors in factor analysis.
128 *Psychometrika*, 30(2), 179–185. <https://doi.org/10.1007/BF02289447>
- 129 Hutcheson, G. D. (1999). *The multivariate social scientist : Introductory statistics*
130 *using generalized linear models*. London : SAGE.
- 131 Iannone, R. (2016). *DiagrammeRsvg: Export DiagrammeR graphviz graphs as SVG*.
132 Retrieved from <https://CRAN.R-project.org/package=DiagrammeRsvg>
- 133 Iannone, R. (2021). *DiagrammeR: Graph/network visualization*. Retrieved from
134 <https://github.com/rich-iannone/DiagrammeR>
- 135 Jorgensen, T. D., Pornprasertmanit, S., Schoemann, A. M., & Rosseel, Y. (2021).
136 *semTools: Useful tools for structural equation modeling*. Retrieved from
137 <https://CRAN.R-project.org/package=semTools>
- 138 Kaiser, H. F. (1974). An index of factorial simplicity. *Psychometrika*, 39(1), 31–36.
139 <https://doi.org/10.1007/bf02291575>
- 140 Lorenzo-Seva, U., Timmerman, M., & Kiers, H. (2011). The Hull Method for
141 Selecting the Number of Common Factors. *Multivariate Behavioral Research*, 46,
142 340–364. <https://doi.org/10.1080/00273171.2011.564527>

- Mardia, K. V. (1970). Measures of multivariate skewness and kurtosis with applications. *Biometrika*, 57(3), 519–530.
<https://doi.org/10.1093/biomet/57.3.519>
- Mulaik, S. A. (2009). *Foundations of Factor Analysis* (Vol. 7). London: London: Chapman and Hall/CRC. <https://doi.org/10.1201/b15851>
- Müller, K., & Wickham, H. (2021). *Tibble: Simple data frames*. Retrieved from <https://CRAN.R-project.org/package=tibble>
- Navarro-Gonzalez, D., & Lorenzo-Seva, U. (2021). *EFA.MRFA: Dimensionality assessment using minimum rank factor analysis*. Retrieved from <https://CRAN.R-project.org/package=EFA.MRFA>
- Ooms, J. (2021). *Rsvg: Render SVG images into PDF, PNG, PostScript, or bitmap arrays*. Retrieved from <https://CRAN.R-project.org/package=rsvg>
- R Core Team. (2021). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>
- Revelle, W. (2021). *Psych: Procedures for psychological, psychometric, and personality research*. Evanston, Illinois: Northwestern University. Retrieved from <https://CRAN.R-project.org/package=psych>
- Rosseel, Y. (2012). lavaan: An R package for structural equation modeling. *Journal of Statistical Software*, 48(2), 1–36. Retrieved from <https://www.jstatsoft.org/v48/i02/>
- Ryu, C. (2021). *Dlookr: Tools for data diagnosis, exploration, transformation*. Retrieved from <https://CRAN.R-project.org/package=dlookr>
- Shapiro, S. S., & Wilk, M. B. (1965). An analysis of variance test for normality (complete samples). *Biometrika*, 52(3-4), 591–611.

<https://doi.org/10.1093/biomet/52.3-4.591>

Velicer, W. (1976). Determining the Number of Components from the Matrix of Partial Correlations. *Psychometrika*, 41, 321–327.

<https://doi.org/10.1007/BF02293557>

Venables, W. N., & Ripley, B. D. (2002). *Modern applied statistics with s* (Fourth). New York: Springer. Retrieved from <https://www.stats.ox.ac.uk/pub/MASS4/>

Watkins, M. (2020). *A Step-by-Step Guide to Exploratory Factor Analysis with R and RStudio*. <https://doi.org/10.4324/9781003120001>

Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York. Retrieved from <https://ggplot2.tidyverse.org>

Wickham, H. (2019). *Stringr: Simple, consistent wrappers for common string operations*. Retrieved from <https://CRAN.R-project.org/package=stringr>

Wickham, H. (2021a). *Forcats: Tools for working with categorical variables (factors)*. Retrieved from <https://CRAN.R-project.org/package=forcats>

Wickham, H. (2021b). *Tidyr: Tidy messy data*. Retrieved from <https://CRAN.R-project.org/package=tidyr>

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., . . . Yutani, H. (2019). Welcome to the tidyverse. *Journal of Open Source Software*, 4(43), 1686. <https://doi.org/10.21105/joss.01686>

Wickham, H., & Bryan, J. (2019). *Readxl: Read excel files*. Retrieved from <https://CRAN.R-project.org/package=readxl>

Wickham, H., François, R., Henry, L., & Müller, K. (2021). *Dplyr: A grammar of data manipulation*. Retrieved from <https://CRAN.R-project.org/package=dplyr>

Wickham, H., & Hester, J. (2021). *Readr: Read rectangular text data*. Retrieved from <https://CRAN.R-project.org/package=readr>

- 193 Zhu, H. (2021). *kableExtra: Construct complex table with 'kable' and pipe syntax*.
- 194 Retrieved from <https://CRAN.R-project.org/package=kableExtra>

Table 1

Descriptive Statistics

	Mean	SD	Skew	Kurtosis	Shapiro-Wilk Statistics	p	Item-Total Correlation
Item1	1.12	0.49	5.02	27.80	0.25	.00	.16
Item2	2.16	1.19	0.71	-0.54	0.84	.00	.14
Item3	4.14	0.99	-1.23	1.14	0.79	.00	.19
Item4	2.87	1.59	0.08	-1.60	0.83	.00	.19
Item5	1.76	1.23	1.35	0.44	0.66	.00	.38
Item6	2.73	1.46	0.20	-1.36	0.87	.00	.33
Item7	3.86	1.67	-0.99	-0.85	0.65	.00	.23
Item8	3.76	1.14	-0.68	-0.45	0.86	.00	.00
Item9	3.42	1.83	-0.45	-1.69	0.69	.00	.33
Item10	2.74	1.04	0.09	-0.74	0.91	.00	.28
Item11	2.60	1.25	0.29	-0.86	0.89	.00	.35
Item12	2.11	1.17	0.77	-0.39	0.83	.00	.32
Item13	2.94	1.03	-0.12	-0.40	0.91	.00	.10
Item14	3.62	1.64	-0.68	-1.25	0.74	.00	.32
Item15	1.64	1.18	1.79	2.02	0.60	.00	.15
Item16	3.51	1.30	-0.70	-0.59	0.85	.00	.39
Item17	1.96	0.98	1.02	0.69	0.82	.00	.05
Item18	2.44	1.31	0.38	-1.14	0.86	.00	.11
Item19	3.80	1.29	-0.87	-0.42	0.82	.00	.17
Item20	4.01	1.40	-1.22	0.07	0.70	.00	.13
Item21	1.33	0.91	3.03	8.43	0.41	.00	.01
Item22	2.59	1.41	0.27	-1.27	0.86	.00	.19
Item23	1.31	0.81	2.75	6.92	0.43	.00	.21
Item24	1.47	1.18	2.38	4.00	0.43	.00	.28
Item25	2.56	1.27	0.33	-1.00	0.89	.00	.11
Item26	1.54	1.25	2.12	2.86	0.46	.00	.26




Figure 1. ABC




Figure 2. Iter-correlation of the items




Figure 3. Factor Identification (A) Parallel analysis (B) Scree Plot, (C) Hull method
 (#fig:fac.id)