

A More Detailed Proof of Prop. 9.3.5

April 8, 2020

In the slides we explained our goal was to show

$$\frac{1}{\tilde{M}^N} \sum_{j=1}^{\tilde{M}_N} \tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j}) \xrightarrow{\mathbb{P}} \nu L(f)$$

for any $f \in \tilde{\mathcal{C}}$. First, let's write

$$\left\{ \frac{1}{\tilde{M}^N} \sum_{j=1}^{\tilde{M}_N} \tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j}) - \frac{1}{\tilde{M}^N} \sum_{j=1}^{\tilde{M}_N} E \left[\tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j}) \mid \mathcal{F}^N \right] \right\} + \left\{ \frac{1}{\tilde{M}^N} \sum_{j=1}^{\tilde{M}_N} E \left[\tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j}) \mid \mathcal{F}^N \right] \right\}$$

The first term converges to 0, and the second converges to $\mu(f)$.

1 The Trick

Recall in the slides we discussed that

$$\begin{aligned} L(\xi^{N,i}, f) &= \int L(\xi^{N,i}, dy) f(y) \\ &= \int R(\xi^{N,i}, dy) \frac{dL(\xi^{N,i}, \cdot)}{dR(\xi^{N,i}, \cdot)}(y) f(y) \\ &= E \left[\tilde{\omega}^{N,i} f(\tilde{\xi}^{N,i}) \mid \mathcal{F}^N \right] \end{aligned}$$

The reason that this is useful is that it is a function of ν samples only. Therefore, as long as this function is sufficiently well-behaved, we only need to worry about ν -consistency.

Another thing: the sum over j (new samples) turns into a sum over i (old

samples) because $\alpha_N M_N = \tilde{M}_N$:

$$\begin{aligned}
\tilde{M}_N^{-1} \sum_{j=1}^{\tilde{M}_N} E \left[\tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j}) \mid \mathcal{F}^N \right] &= \tilde{M}_N^{-1} \sum_{j=1}^{\tilde{M}_N} L(\xi^{N,i}, f) \\
&= \alpha_N^{-1} M_N^{-1} \sum_{j=1}^{\tilde{M}_N} L(\xi^{N,i}, f) \\
&= \alpha_N^{-1} M_N^{-1} \sum_{i=1}^{M_N} \alpha_N L(\xi^{N,i}, f) \\
&= M_N^{-1} \sum_{i=1}^{M_N} L(\xi^{N,i}, f),
\end{aligned}$$

So is $L(x, f) \in \mathbb{C}$? The answer is yes, as long as we pick $f \in \tilde{\mathbb{C}}$, and that follows just by the definition of it:

$$\tilde{\mathbb{C}} = \{f \in L^1(\mathbb{X}, \mu) : x \mapsto L(x, |f|) \in \mathbb{C}\}.$$

2 Showing Convergence to Zero using Theorem 9.5.7

We showed the second part of

$$\left\{ \frac{1}{\tilde{M}_N} \sum_{j=1}^{\tilde{M}_N} \tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j}) - \frac{1}{\tilde{M}_N} \sum_{j=1}^{\tilde{M}_N} E \left[\tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j}) \mid \mathcal{F}^N \right] \right\} + \left\{ \frac{1}{\tilde{M}_N} \sum_{j=1}^{\tilde{M}_N} E \left[\tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j}) \mid \mathcal{F}^N \right] \right\}$$

converges in probability to $\nu L(f)$ for any $f \in \tilde{\mathbb{C}}$. Now we must show the first part converges to 0. To show this, we set $V_{N,j} = \frac{1}{\tilde{M}_N} \tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j})$, and check the three conditions of 9.5.7.

2.1 First condition of 9.5.7

The first condition, translated into our notation, is: the triangular array $\{V_{N,j}\}_{1 \leq j \leq M_N}$ is conditionally independent given \mathcal{F}^N , and for any $N, j = 1, \dots, \tilde{M}_N$, we have $E[|V_{N,j}| \mid \mathcal{F}^N] < \infty$.

The first condition of 9.5.7 is true because of the description of the algorithm used for 9.3.5, and because $f \in \tilde{\mathbb{C}}$ by assumption.

2.2 Second Condition of 9.5.7

The second condition is that the sequence (in N)

$$\left\{ \sum_{j=1}^{\tilde{M}_N} E[|V_{N,j}| \mid \mathcal{F}^N] \right\}_N$$

is bounded in probability. This is true because it converges in probability. To see this more easily, use the same trick we've been using a few times:

$$\begin{aligned} \sum_{j=1}^{\tilde{M}_N} E[|V_{N,j}| \mid \mathcal{F}^N] &= \frac{1}{\tilde{M}_N} \sum_{j=1}^{\tilde{M}_N} E[|\tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j})| \mid \mathcal{F}^N] \\ &= \frac{1}{\tilde{M}_N} \sum_{j=1}^{\tilde{M}_N} E[\tilde{\omega}^{N,j} |f(\tilde{\xi}^{N,j})| \mid \mathcal{F}^N] \\ &\quad \text{(Assumption 9.3.3 positive RN deriv.)} \\ &= \frac{1}{\tilde{M}_N} \sum_{j=1}^{\tilde{M}_N} L(\xi^{N,i}, |f|) \xrightarrow{P} \nu L(|f|). \end{aligned}$$

2.3 Third Condition of 9.5.7

The third condition is that for any $\epsilon > 0$ we have

$$\sum_{j=1}^{\tilde{M}_N} E[|V_{N,j}| \mathbf{1}(|V_{N,j}| \geq \epsilon) \mid \mathcal{F}^N] \xrightarrow{P} 0.$$

Before, we show the main work, note that, after we pick $f \in \tilde{\mathcal{C}}$, then for any C ,

$$L(x, |f| \mathbf{1}_{\{h(x, x') \geq C\}}) = \int L(x, dx') |f(x')| \mathbf{1}_{\{h(x, x') \geq C\}} \leq L(x, |f|).$$

This is useful because if we assume $f \in \tilde{\mathcal{C}}$, this will imply that $L(x, |f|) \in \mathcal{C}$, and by propriety, this will also imply that $L(x, |f| \mathbf{1}_{\{h(x, x') \geq C\}}) \in \mathcal{C}$. We need this in the last line of the following work:

$$\begin{aligned}
\lim_N \sum_{j=1}^{\tilde{M}_N} E [|V_{N,j}| 1_{\{|V_{N,j}| \geq \epsilon\}} \mid \mathcal{F}^N] &= \lim_N \sum_{j=1}^{\tilde{M}_N} E \left[|\tilde{M}_N^{-1} \tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j})| 1_{\{|\tilde{M}_N^{-1} \tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j})| \geq \epsilon\}} \mid \mathcal{F}^N \right] \\
&\quad \text{(defn of V)} \\
&= \lim_N \tilde{M}_N^{-1} \sum_{j=1}^{\tilde{M}_N} E \left[\tilde{\omega}^{N,j} |f(\tilde{\xi}^{N,j})| 1_{\{|\tilde{M}_N^{-1} \tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j})| \geq \epsilon\}} \mid \mathcal{F}^N \right] \\
&\quad \text{(algebra)} \\
&\leq \lim_N \tilde{M}_N^{-1} \sum_{j=1}^{\tilde{M}_N} E \left[\tilde{\omega}^{N,j} |f(\tilde{\xi}^{N,j})| 1_{\{|\tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j})| \geq C\}} \mid \mathcal{F}^N \right] \\
&\quad \text{(if } \tilde{M}_N \epsilon \geq C) \\
&= \lim_N \tilde{M}_N^{-1} \sum_{i=1}^{\tilde{M}_N} \int L(\xi^{N,i}, d\tilde{\xi}^{N,j}) \left| f(\tilde{\xi}^{N,j}) \right| 1_{\{|\tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j})| \geq C\}} \\
&\quad \text{(defn of expc. and the trick with replacing } i \text{ and } j) \\
&= \iint \nu(dx) L(x, dx') |f(x')| 1_{\{|\tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j})| \geq C\}} \\
&\quad \text{(discussion above)}
\end{aligned}$$

The last term is bounded above by $\nu L(|f|) < \infty$, so we can use dominated convergence on the above work when we take the limit with $C \uparrow \infty$:

$$\begin{aligned}
\lim_N \sum_{j=1}^{\tilde{M}_N} E [|V_{N,j}| 1_{\{|V_{N,j}| \geq \epsilon\}} \mid \mathcal{F}^N] &= \lim_C \lim_N \sum_{j=1}^{\tilde{M}_N} E [|V_{N,j}| 1_{\{|V_{N,j}| \geq \epsilon\}} \mid \mathcal{F}^N] \\
&\leq \lim_C \iint \nu(dx) L(x, dx') |f(x')| 1_{\{|\tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j})| \geq C\}} \\
&\quad \text{(above)} \\
&= \iint \nu(dx) L(x, dx') |f(x')| \lim_C 1_{\{|\tilde{\omega}^{N,j} f(\tilde{\xi}^{N,j})| \geq C\}} \\
&\quad \text{(DCT)} \\
&= 0.
\end{aligned}$$