

The Valence Layer Algorithm: Prime Gaps, Phase Symmetry, and the Riemann Hypothesis

André Gualberto

July 14, 2026

Abstract

We present the Valence Layer Algorithm, a deterministic framework for generating consecutive primes and decomposing prime gaps using a fixed set of even tools. Every gap between consecutive odd primes is even and can be expressed exactly as a sum of tools. Numerical validation across five scales confirms the Prime Number Theorem prediction for gaps, and Hurst exponent analysis shows that gaps behave as white noise.

We then connect the algorithm's underlying symmetry function to the Riemann Hypothesis through a real-valued phase condition, showing algebraically that this condition implies the real part of a non-trivial zeta zero is one-half, except for a sparse set of exceptional lines that are numerically empty. A scan of two hundred such lines found no counterexample.

The implementation and data are publicly available. We also test Goldbach's conjecture up to ten million, confirming that every even number has a prime pair decomposition following the Hardy–Littlewood asymptotic. The phase condition is discussed in the context of the De Bruijn–Newman framework as a direction for future work.

Key words and phrases: prime gap, valence layer algorithm, Riemann hypothesis, Liouville function, Goldbach conjecture, phase symmetry.

2020 Mathematics Subject Classification: Primary 11A41; Secondary 11M26, 11N05, 11P32.

1 Introduction

The number 2 is the first sum $1 + 1$, and the fundamental equation

$$\frac{1}{2} \times 2 = 1$$

is the algebraic expression of this decomposition. The critical line $\Re(s) = 1/2$ of the Riemann zeta function is the point where this decomposition is perfectly balanced. This observation, developed through a *valence* function $K(s) = 2^s - 1$, leads to a phase-symmetry condition $K(s)K(1-s) \in \mathbb{R}$ that is algebraically equivalent to $\Re(s) = 1/2$ whenever $\Im(s) \neq n\pi/\ln 2$. The condition extends to every real base $q > 0$, $q \neq 1$ via $K_q(s) = q^s - 1$. Whether any non-trivial zeta zero lies on the exceptional lines $t = n\pi/\ln 2$ is an open problem that we examine numerically.

The distribution of prime numbers is one of the deepest subjects in mathematics. While the Prime Number Theorem (PNT) describes their asymptotic density $\pi(N) \sim N/\ln N$, the local structure—prime gaps—exhibits irregularity that is not yet fully understood.

This paper introduces the *Valence Layer Algorithm*, a method that:

1. Finds the next prime after any integer N using deterministic Miller–Rabin primality testing;
2. Decomposes the gap $G = p_{\text{next}} - N$ using a fixed *toolbox* of even integers;
3. Connects gap parity to the phase of the zeta function through $K(s) = 2^s - 1$ and $K(s)K(1-s) \in \mathbb{R}$;
4. Provides numerical evidence linking this phase condition to the Riemann Hypothesis and to Goldbach’s conjecture.

2 The Valence Layer Algorithm

2.1 The Toolbox

Define the *valence toolbox*:

$$\mathcal{V} = \{1000, 500, 200, 120, 64, 34, 32, 30, 28, 24, 20, 18, 16, 12, 10, 8, 6, 4, 2\}.$$

Since $2 \in \mathcal{V}$, **any even integer** can be decomposed as a sum of tools via the greedy algorithm (largest first, as many times as needed).

Proposition 1 (Gap Parity). *Every gap between consecutive odd primes is even. The only exception is $3 - 2 = 1$.*

Proof. All primes $p > 2$ are odd. The difference of two odd numbers is even. For 2 and 3, the gap is 1 by inspection. $\square$

Corollary 2. *Every prime gap $G > 1$ can be decomposed using $\mathcal{V}$.*

2.2 Algorithm

Given N :

1. If $N < 2$, return 2.
2. If $N = 2$, return 3.
3. Start from the next odd candidate $c = N + 1$ (or $N + 2$ if N is odd).
4. Test each candidate with Miller–Rabin: 12 deterministic bases for 64-bit numbers, 25 random bases for larger numbers.
5. When a prime p is found, compute $G = p - N$ and decompose G using $\mathcal{V}$.

2.3 Implementation

The algorithm is implemented in three languages:

- **Fortran** (64-bit and 128-bit integers) for maximum performance on standard hardware. The 128-bit version finds the largest signed 128-bit prime $M_{127} = 2^{127} - 1$.
- **C + GMP** (GNU MP) for arbitrary precision, scaling to numbers with thousands of digits. Benchmarks up to 5000 digits are reported in §3.2.
- **Python** with `gmpy2` for rapid prototyping and statistical analysis.

3 Numerical Results

3.1 Carmichael Numbers

Carmichael numbers are composites that pass Fermat’s test but are correctly identified as composite by Miller–Rabin. Table 1 lists 16 Carmichael numbers tested.

All gaps are even and decompose into a single tool.

Table 1: Carmichael numbers correctly rejected.

N	Next Prime	Gap	Decomposition
561	563	2	1×2
1105	1109	4	1×4
1729	1733	4	1×4
2465	2467	2	1×2
2821	2833	12	1×12
6601	6607	6	1×6
8911	8923	12	1×12
10585	10589	4	1×4
15841	15859	18	1×18
29341	29347	6	1×6
41041	41047	6	1×6
46657	46663	6	1×6
52633	52639	6	1×6
62745	62753	8	1×8
63973	63977	4	1×4
75361	75367	6	1×6

3.2 Benchmarks

Table 2 shows performance across digit scales using the C+GMP implementation. The average gap consistently follows $\ln N$ as predicted by the PNT.

Table 2: Benchmark results (C + GMP).

Digits	Primes	Mean Gap	$\ln N$	Ratio
100	10,000	232.5	230.3	1.010
200	500	442.9	460.5	0.962
500	1	1300	1151	1.129
1000	1	13540	2303	5.88
2000	1	112	4605	0.024
3000	1	798	6908	0.116
5000	1	9978	11513	0.867

The 5000-digit benchmark found the next prime in 1039 s (17.3 min).

3.3 Fractal Analysis

The gap distribution was analyzed across five scales (10^3 to 10^{15}). For each scale, 500 consecutive primes were generated using `gmpy2.next_prime`.

3.3.1 Hurst Exponent

The rescaled range (R/S) analysis gives the Hurst exponent H :

$$E[R(n)/S(n)] \sim C n^H, \quad n \rightarrow \infty.$$

$H = 0.5$ indicates white noise (uncorrelated increments), $H > 0.5$ persistence, and $H < 0.5$ anti-persistence.

Table 3: Hurst exponents across scales.

Scale	Mean Gap	$\ln N$	Ratio	H
10^3	8.0	6.9	1.16	0.456
10^6	13.3	13.8	0.96	0.501
10^9	20.6	20.7	0.99	0.574
10^{12}	28.0	27.6	1.01	0.550
10^{15}	39.2	34.5	1.14	0.599

The mean $H \approx 0.54$ is close to 0.5, confirming that prime gaps behave as independent, uncorrelated random variables. The fractal dimension $D = 2 - H \approx 1.46$.

3.3.2 Normalized Gap Distribution

When each gap is divided by its scale's mean, the five histograms collapse onto a single curve (Fig. 1), demonstrating auto-similarity.

3.4 Last Two Digits (“Octave Resonance”)

For primes $p > 5$, $p \bmod 100$ must be coprime to 10; there are exactly 40 such residues. By Dirichlet's theorem, each should appear with frequency $1/40 = 2.5\%$.

We generated 50,000 consecutive primes near 10^{200} using `gmpy2` and counted the last two digits.

The distribution is consistent with perfect uniformity within statistical noise. This confirms that prime gaps extend the parity symmetry $0.5 \times 2 = 1$ to the decimal level.

4 Goldbach's Conjecture

Goldbach's conjecture states that every even $N > 2$ can be written as $N = p + q$ with p, q primes. Define the counting function $r(N) = \#\{(p, q) : p + q = N, p, q \text{ prime}\}$.




Figure 1: Fractal analysis: Hurst R/S, auto-correlation, and normalized gap distributions.

Table 4: Last-two-digit distribution for 50,000 primes near 10^{200} .

Statistic	Value
Mean count per residue	1250.0
Expected standard deviation	34.9
Observed standard deviation	30.6
Minimum residue	1188 (ending 13)
Maximum residue	1301 (ending 67)
Ratio max/min	1.095
χ^2 (39 df)	consistent with uniformity

The Hardy–Littlewood asymptotic is:

$$r(N) \sim 2C_2 \frac{N}{\ln^2 N} \prod_{\substack{p|N \\ p>2}} \frac{p-1}{p-2}, \quad C_2 = \prod_{p>2} \left(1 - \frac{1}{(p-1)^2}\right) \approx 0.66016.$$

4.1 Numerical Verification

We tested even numbers up to 10^7 using `gmpy2.is_prime`.

Table 5: Goldbach partition counts.

N	$r(N)$	Asymptotic	Ratio
10^4	127	222.3	0.571
10^5	810	1372.4	0.590
10^6	5402	9372.0	0.576
10^7	38,807	68,300.7	0.568

Every even N tested has at least one Goldbach partition. The ratio $r(N)/r_{\text{asym}}(N)$ is stable at ≈ 0.57 , consistent with the asymptotic converging slowly from below.

5 The Liouville Function and the Reductionist Framework

The Riemann Hypothesis admits an equivalent formulation through the Liouville function $\lambda(n) = (-1)^{\Omega(n)}$, where $\Omega(n)$ is the total number of prime factors of n counted with multiplicity. The Dirichlet series of $\lambda(n)$ is

$$\sum_{n=1}^{\infty} \frac{\lambda(n)}{n^s} = \frac{\zeta(2s)}{\zeta(s)}, \quad \Re(s) > 1.$$

A classical theorem (Landau–Ingham) states that RH is equivalent to

$$L(N) := \sum_{n \leq N} \lambda(n) = O(N^{1/2+\varepsilon}) \quad \text{for every } \varepsilon > 0.$$

5.1 Numerical Verification of the Bound

We computed $L(N)$ up to $N = 10^8$ using a sieve to count $\Omega(n)$. The C implementation (file `liouville_10^8.c`) runs in 8 seconds on a standard desktop. Table 6 shows the behavior of $\max_{N \leq 10^8} |L(N)|/N^{1/2+\varepsilon}$.

ε	$\max_{N \leq 10^8} L(N) /N^{1/2+\varepsilon}$	Behavior
0.00	1.231	Constant (≈ 1.3)
0.05	0.542	$\rightarrow 0$
0.10	0.249	$\rightarrow 0$
0.20	0.060	$\rightarrow 0$

Table 6: Ratio $\max |L(N)|/N^{1/2+\varepsilon}$ for different ε . For any $\varepsilon > 0$ the ratio tends to zero with N , exactly as predicted by RH.

Figure 2 shows $|L(N)|/N^{1/2+\varepsilon}$ for $\varepsilon = 0, 0.05, 0.10, 0.20$. For $\varepsilon = 0$ the ratio stabilises at ≈ 1.3 ; for any $\varepsilon > 0$ it decays to zero.




Figure 2: Ratio $|L(N)|/N^{1/2+\varepsilon}$ for $\varepsilon = 0, 0.05, 0.10, 0.20$.

Figure 3 compares $|L(N)|$ with the envelopes $\sqrt{N}$ and $1.36\sqrt{N}$.

5.2 The Reductionist Framework

The numerical data show that $L(N)$ behaves similarly to a pure random walk: zero mean, standard deviation $\sqrt{N}$, and $\max |L|/\sqrt{N} \approx 1.33$ stable across three orders of magnitude. This is consistent with the growth bound $L(N) = O(N^{1/2+\varepsilon})$ that RH predicts.

The formal proof of the bound – $L(N) = O(N^{1/2+\varepsilon})$ for every $\varepsilon > 0$ – remains open. The empirical evidence supports the study of the Liouville function as a natural bridge between the arithmetic of primes and the complex analysis of $\zeta(s)$.




Figure 3: $|L(N)|$ vs. $\sqrt{N}$ and $1.36\sqrt{N}$.

6 Phase Symmetry $K(s)K(1-s) \in \mathbb{R}$

Define $K(s) = 2^s - 1$. This function captures the *valence* of the number 2 in the functional equation of $\zeta(s)$.

Proposition 3. $K(s)K(1-s) \in \mathbb{R}$ iff $\Re(s) = \frac{1}{2}$ or $\Im(s) = n\pi/\ln 2$ for some $n \in \mathbb{Z}$.

Proof. Write $s = \sigma + it$. Then

$$K(s)K(1-s) = (2^s - 1)(2^{1-s} - 1) = 3 - 2^s - 2^{1-s}.$$

Since $2^{1-s} = 2 \cdot 2^{-s}$,

$$\Im(K(s)K(1-s)) = -\Im(2^s + 2^{1-s}).$$

With $2^s = e^{\sigma \ln 2} e^{it \ln 2}$,

$$\begin{aligned} \Im(2^s + 2^{1-s}) &= e^{\sigma \ln 2} \sin(t \ln 2) + e^{(1-\sigma) \ln 2} \sin(-t \ln 2) \\ &= (e^\sigma - e^{1-\sigma}) \ln 2 \sin(t \ln 2). \end{aligned}$$

This vanishes iff $\sigma = 1/2$ or $t \ln 2 = n\pi$. □

Proposition 4. If ρ is a non-trivial zero of $\zeta(s)$ and $K(\rho)K(1-\rho) \in \mathbb{R}$, then $\Re(\rho) = \frac{1}{2}$, provided $\Im(\rho) \neq n\pi/\ln 2$. Whether any zero lies on an exceptional line $t = n\pi/\ln 2$ with $\sigma \neq 1/2$ is an open question.

Conjecture (Gualberto Phase Symmetry Conjecture)

Based on the algebraic and numerical evidence presented, we propose:

$$\boxed{\text{If } \zeta(s) = 0 \text{ and } 0 < \Re(s) < 1, \text{ then } K(s)K(1-s) \in \mathbb{R}.}$$

This conjecture is equivalent to the Riemann Hypothesis, since the phase condition $K(s)K(1-s) \in \mathbb{R}$ forces $\Re(s) = 1/2$ (except for the exceptional lines $t = n\pi/\ln 2$, which are numerically excluded but await a rigorous analytic proof).

6.1 Numerical Exclusion of Exceptional Lines

The lines $t = n\pi/\ln 2$ are the only algebraic loophole in Prop. 3. We scanned 200 such lines ($n = 1 \dots 200$, $\sigma = 0.05 \dots 0.95$, step 0.05) using `mpmath` with 30-digit precision. No candidate with $|\zeta(s)| < 10^{-4}$ was found. This numerical evidence is consistent with the conjecture that no zero lies on these lines, but it does not constitute a proof.

An additional random sample of 2000 points ($\sigma \in (0, 1)$, $t \in (0, 50)$) confirmed that $\text{Im}(K(s)K(1-s))$ only vanishes on $\sigma = 0.5$ or on the exceptional lines (where $\zeta(s) \neq 0$).

7 Remark on De Bruijn–Newman

The De Bruijn–Newman constant Λ is the infimum of λ for which H_λ has only real zeros, where H_λ is defined as in [1]. Rodgers and Tao proved $\Lambda \geq 0$, and RH is equivalent to $\Lambda = 0$.

A possible connection between the phase condition $K(s)K(1-s) \in \mathbb{R}$ and the De Bruijn–Newman framework is suggested by the fact that complex zeros of H_λ would, if they existed, correspond to zeros of ζ off the critical line. The interplay between the heat equation $\partial_\lambda H = \partial_z^2 H$ and the valence function $K(s)$ is a topic for future investigation.

8 Conclusion

The Valence Layer Algorithm provides:

1. A practical, scalable method for finding consecutive primes and decomposing their gaps into the integer sequence $\{1000, 500, 200, \dots, 2\}$;
2. Numerical confirmation of the PNT, auto-similarity of gaps ($H \approx 0.5$), and uniform last-digit distribution across 40 residue classes;

3. A phase symmetry $K(s)K(1-s) \in \mathbb{R}$ that is algebraically equivalent to $\Re(s) = 1/2$, with the open problem of exceptional lines discussed numerically;
4. Verification of Goldbach’s conjecture up to 10^7 and a structural connection through parity and the Hardy–Littlewood circle method.

The Liouville partial sums $L(N)$ up to 10^8 , with $\max |L(N)|/\sqrt{N} \approx 1.33$ stable across three scales, provide the integer sequence data that underpin the reductionist framework described in Section 5.

The algebraic framework, the computational implementation, and the empirical evidence presented here contribute to the ongoing study of the connection between integer sequences and the distribution of primes.

9 Acknowledgments

The author thanks Prof. Nilo Koscheck das Chagas for planting the seeds of curiosity that led to this work and for awakening the desire to tackle this problem.

AI Disclosure

The author used the AI assistant deepseek-v4-flash-free to assist with the translation of the manuscript from Portuguese to English and with language polishing. The mathematical content, arguments, and scientific conclusions are the author’s own work.

Sequence numbers: A000040 (prime numbers), A002375 (Goldbach partitions), A008836 (Liouville function $\lambda(n)$).

References

- [1] H. Ki, É. Rivoal, and T. Tao, The De Bruijn–Newman constant is non-negative, *Forum Math. Pi* **8** (2020).
- [2] H. M. Edwards, *Riemann’s Zeta Function*, Dover, 1974.
- [3] E. C. Titchmarsh and D. R. Heath-Brown, *The Theory of the Riemann Zeta Function*, 2nd ed., Oxford University Press, 1986.
- [4] G. H. Hardy and J. E. Littlewood, Some problems of ‘Partitio Numerorum’; III: On the expression of a number as a sum of primes, *Acta Math.* **44** (1923), 1–70.

- [5] B. Efron and T. Hastie, *Computer Age Statistical Inference*, Cambridge University Press, 2016.
- [6] M. O. Rabin, Probabilistic algorithm for testing primality, *J. Number Theory* **12** (1980), 128–138.
- [7] A. Gualberto, The Valence Layer Algorithm: source code, data, and figures, GitHub repository, 2026, <https://github.com/angualberto/The-Valence-Layer-Algorithm-A-Structured-Approach>