

NIP compressible types

Pablo Andújar Guerrero

University of Leeds

Workshop?

<https://anguempa.github.io/leeds-workshop.pdf>


Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .

Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .


$$\mathcal{F} = \{S_1, S_2, S_3\}$$



Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .


$$S_1 \setminus (S_2 \cup S_3)$$



Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .

$$S_3 \setminus (S_1 \cup S_2)$$



Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .

(In model-theoretic terminology a Boolean atom can be called a type.)


Let $BA(\mathcal{F})$ denote the number of Boolean atoms of \mathcal{F} .

Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .

Let $BA(\mathcal{F})$ denote the number of Boolean atoms of \mathcal{F} .

$$BA(\mathcal{F}) = 5$$




Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .

Let $BA(\mathcal{F})$ denote the number of Boolean atoms of \mathcal{F} .

$$BA(\mathcal{F}) = 2^3 = 8$$



Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .

(In model-theoretic terminology a Boolean atom can be called a type.)

Let $BA(\mathcal{F})$ denote the number of Boolean atoms of \mathcal{F} .

Definition

Let \mathcal{S} be a family of sets. The *dual shatter function* of \mathcal{S} is the function

$$\begin{aligned}\pi_{\mathcal{S}}^* : \omega &\rightarrow \omega \\ n &\mapsto \max\{BA(\mathcal{F}) : \mathcal{F} \subseteq \mathcal{S}, |\mathcal{F}| = n\}\end{aligned}$$

Counting Boolean atoms

Observe that $\pi_{\mathcal{S}}^*(n) \leq 2^n$ for all n .

Counting Boolean atoms

Observe that $\pi_{\mathcal{S}}^*(n) \leq 2^n$ for all n .

If $\pi_{\mathcal{S}}^*(d) < 2^d$ for some d then $\pi_{\mathcal{S}}^*(n) = O(n^{d-1})$ (Sauer's Lemma).

Counting Boolean atoms

Observe that $\pi_{\mathcal{S}}^*(n) \leq 2^n$ for all n .

If $\pi_{\mathcal{S}}^*(d) < 2^d$ for some d then $\pi_{\mathcal{S}}^*(n) = O(n^{d-1})$ (Sauer's Lemma).

In this case we call \mathcal{S} a *VC class* (VC=Vapnik–Chervonenkis).

Counting Boolean atoms

Observe that $\pi_{\mathcal{S}}^*(n) \leq 2^n$ for all n .

If $\pi_{\mathcal{S}}^*(d) < 2^d$ for some d then $\pi_{\mathcal{S}}^*(n) = O(n^{d-1})$ (Sauer's Lemma).

In this case we call \mathcal{S} a *VC class* (VC=Vapnik–Chervonenkis).

Definition

The VC-codensity of a VC class \mathcal{S} is

$$\text{vc}^*(\mathcal{S}) = \inf\{r \in \mathbb{R}^{\geq 0} : \pi_{\mathcal{S}}^*(n) = O(n^r)\}.$$

Counting Boolean atoms

Observe that $\pi_{\mathcal{S}}^*(n) \leq 2^n$ for all n .

If $\pi_{\mathcal{S}}^*(d) < 2^d$ for some d then $\pi_{\mathcal{S}}^*(n) = O(n^{d-1})$ (Sauer's Lemma).

In this case we call \mathcal{S} a *VC class* (VC=Vapnik–Chervonenkis).

Definition

The VC-codensity of a VC class \mathcal{S} is

$$\text{vc}^*(\mathcal{S}) = \inf\{r \in \mathbb{R}^{\geq 0} : \pi_{\mathcal{S}}^*(n) = O(n^r)\}.$$

E.g. If $\mathcal{S} = \{\text{intervals in } \mathbb{R}\}$ then $\pi_{\mathcal{S}}^*(n) = 2n$ and $\text{vc}^*(\mathcal{S}) = 1$.

Definition

A family of sets \mathcal{S} is *(downward) directed* if, for any **finite** subfamily $\mathcal{F} \subseteq \mathcal{S}$, there exists some $S \in \mathcal{S}$ such that $S \subseteq \bigcap \mathcal{F}$.


Equivalently, \mathcal{S} is directed if, for any $S_1, S_2 \in \mathcal{S}$ there exists $S \in \mathcal{S}$ with $S \subseteq S_1 \cap S_2$.

E.g.

- In a poset (P, \leq) the sets $\{a \in P : a \leq b\}$ for $b \in P$.
- The neighborhoods of a given point in a topological spaces.
- Any filter basis.


Definition

A family of sets \mathcal{S} has a finite transversal $T = \{a_1, \dots, a_n\}$ if every $S \in \mathcal{S}$ satisfies that $S \cap T \neq \emptyset$.



Definition


A family of sets \mathcal{S} has a finite transversal $T = \{a_1, \dots, a_n\}$ if every $S \in \mathcal{S}$ satisfies that $S \cap T \neq \emptyset$.



Definition

A family of sets \mathcal{S} has a finite transversal $T = \{a_1, \dots, a_n\}$ if every $S \in \mathcal{S}$ satisfies that $S \cap T \neq \emptyset$.


We care about the property that \mathcal{S} does **not** have a finite transversal.



Definition

A family of sets \mathcal{S} has a finite transversal $T = \{a_1, \dots, a_n\}$ if every $S \in \mathcal{S}$ satisfies that $S \cap T \neq \emptyset$.

We care about the property that \mathcal{S} does **not** have a finite transversal.



Is there a connection between directedness and the dual shatter function?


Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.

Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.


\mathcal{S} could be directed.



Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.

\mathcal{S} could be directed.




Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.

\mathcal{S} could be directed.


$$\pi_{\mathcal{S}}^*(3) \geq 5$$



Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .




Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .

$$\pi_{\mathcal{S}}^*(3) \geq 8$$




Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .


$$\pi_{\mathcal{S}}^*(3) \geq 6$$



Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.


$$\pi_{\mathcal{S}}^*(3) \geq 6$$



Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.


$$\pi_{\mathcal{S}}^*(3) \geq 8$$



Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.


$$BA(S_1, S_2) \geq 3$$



Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.


$$BA(S_1, S_2) \geq 3$$



Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.


$$BA(S_1, S_2, S_3) \geq 3 + 2 = 5$$



Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.


$$BA(S_1, S_2, S_3) \geq 3 + 1 + \textcolor{red}{1}.$$



Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.

$$F = S_1 \cap S_2 \cap S_3$$



Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does not have a finite transversal.

$$F = S_1 \cap S_2 \cap S_3$$



Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.


$$F = S_1 \cap S_2 \cap S_3$$



Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.


$F \cap \mathcal{S}$ could be directed.



Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.


$F \cap \mathcal{S}$ could be directed.



Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.

$F \cap \mathcal{S}$ could be directed.




Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.

Suppose that $F \cap \mathcal{S}$ is **not** directed, witnessed by S_4 and S_5 .

$$BA(S_1, S_2, S_3, S_4, S_5) \geq BA(S_1, S_2, S_3) + 3$$




Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.

Suppose that $F \cap \mathcal{S}$ is **not** directed, witnessed by S_4 and S_5 .

$$BA(S_1, S_2, S_3, S_4, S_5) \geq BA(S_1, S_2, S_3) + 3$$




Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.

Suppose that $F \cap \mathcal{S}$ is **not** directed, witnessed by S_4 and S_5 .

$$BA(S_1, S_2, S_3, S_4, S_5) \geq BA(S_1, S_2, S_3) + 3$$




Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.

Suppose that $F \cap \mathcal{S}$ is **not** directed, witnessed by S_4 and S_5 .

$$BA(S_1, \dots, S_6) \geq BA(S_1, S_2, S_3) + 3 + 2.$$




Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.

Suppose that $F \cap \mathcal{S}$ is **not** directed, witnessed by S_4 and S_5 .

$$BA(S_1, \dots, S_6) \geq BA(S_1, S_2, S_3) + 3 + 2 + \textcolor{red}{1}.$$



Is there a connection between directedness and the dual shatter function?

We continue this process, and one of two things can happen:

- We find a finite intersection F of sets in \mathcal{S} such that $F \cap \mathcal{S}$ is directed.
- There is a sequence $(S_i)_i$ of sets in \mathcal{S} such that

$$BA(B_i : i \leq 3n) \geq \sum_{j=1}^n j.$$

Meaning that

$$\pi_{\mathcal{S}}^*(3n) \geq \frac{n^2 + n}{2}.$$

Lemma A (A.G. '23)

Let \mathcal{S} be a family of sets with $\text{vc}^*(\mathcal{S}) < 2$. Then at least one holds:

- (a) There exists a finite intersection F of sets in \mathcal{S} such that $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ is directed.
- (b) \mathcal{S} has a finite transversal.

Lemma A (A.G. '23)

Let \mathcal{S} be a family of sets with $\text{vc}^*(\mathcal{S}) < 2$. Then at least one holds:

- (a) There exists a finite intersection F of sets in \mathcal{S} such that $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ is directed.
- (b) \mathcal{S} has a finite transversal.

In fact we can improve (b) in the above lemma.

Lemma A (A.G. '23)

Let \mathcal{S} be a family of sets with $\text{vc}^*(\mathcal{S}) < 2$. Then at least one holds:

- (a) There exists a finite intersection F of sets in \mathcal{S} such that $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ is directed.
- (b) \mathcal{S} has a finite transversal.

In fact we can improve (b) in the above lemma.

Better Lemma A (A.G. '23)

Let \mathcal{S} be a family of sets with $\text{vc}^*(\mathcal{S}) < 2$. Then at least one holds:

- (a) There exists a finite intersection F of sets in \mathcal{S} such that $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ is directed.
- (b) There exists F_1, \dots, F_m , finite non-empty Boolean combinations of sets in \mathcal{S} such that, for every $S \in \mathcal{S}$ there is some $i \leq m$ such that $F_i \subseteq S$.

Now let us talk compressibility.

Our two main references are:

- Martin Bays, Itay Kaplan and Pierre Simon. *Density of compressible types and some consequences*.
ArXiv: 2107.05197, 2022.
- Xi Chen, Yu Cheng, and Bo Tang. *On the recursive teaching dimension of VC classes*.
Advances in Neural Information Processing Systems 29:
Annual Conference on Neural Information Processing
Systems, 2016, Barcelona, Spain, p 2164–2171.

Model-theoretic framework

Let T be a theory and $\varphi(x, y)$ be a (partitioned) formula. For any $M \models T$ and $b \in M^y$ let $\varphi(M, b) = \{a \in M^x : M \models \varphi(a, b)\}$. We identify $\varphi(x, y)$ with the family of sets $\{\varphi(M, b) : b \in M^y\}$:

- $\varphi(x, y)$ may be directed.
- It has dual shatter function π_φ^* .
- It has VC-codensity ...

(All these are independent of M .)

Model-theoretic framework

Let T be a theory and $\varphi(x, y)$ be a (partitioned) formula. For any $M \models T$ and $b \in M^y$ let $\varphi(M, b) = \{a \in M^x : M \models \varphi(a, b)\}$. We identify $\varphi(x, y)$ with the family of sets $\{\varphi(M, b) : b \in M^y\}$:

- $\varphi(x, y)$ may be directed.
- It has dual shatter function π_φ^* .
- It has VC-codensity ...

(All these are independent of M .)

A formula φ is *NIP* if it is a VC class.

A theory T is *NIP* if every formula φ is NIP.

Model-theoretic framework

Let T be a theory and $\varphi(x, y)$ be a (partitioned) formula. For any $M \models T$ and $b \in M^y$ let $\varphi(M, b) = \{a \in M^x : M \models \varphi(a, b)\}$. We identify $\varphi(x, y)$ with the family of sets $\{\varphi(M, b) : b \in M^y\}$:

- $\varphi(x, y)$ may be directed.
- It has dual shatter function π_φ^* .
- It has VC-codensity ...

(All these are independent of M .)

A formula φ is *NIP* if it is a VC class.

A theory T is *NIP* if every formula φ is NIP.

Conjecture: in any dp-minimal theory (a subclass of NIP theories) any formula $\varphi(x, y)$ with $|x| = 1$ has VC-codensity at most one.

Theorem (Baldwin-Saxl '76)

Let G be a group definable in an NIP structure. Let \mathcal{H} be a (uniformly) definable family of subgroups of G . Then there is $k < \omega$ such that, for any finite family $\mathcal{F} \subseteq \mathcal{H}$, there is a subfamily $\mathcal{C} \subseteq \mathcal{F}$ with $|\mathcal{C}| \leq k$ such that $\cap \mathcal{C} = \cap \mathcal{F}$.

Definition

Let $k < \omega$. A family of sets \mathcal{S} is *k-compressible* if, for any **finite** subfamily $\mathcal{F} \subseteq \mathcal{S}$, there exists a subfamily $\mathcal{G} \subseteq \mathcal{S}$ of size **at most** k such that $\cap \mathcal{G} \subseteq \cap \mathcal{F}$.

Equivalently, \mathcal{S} is *k-compressible* if, for any subfamily $\mathcal{F} \subseteq \mathcal{S}$ of size $k + 1$, there exists a subfamily $\mathcal{G} \subseteq \mathcal{S}$ of size **at most** k such that $\cap \mathcal{G} \subseteq \cap \mathcal{F}$.

Definition

Let $k < \omega$. A family of sets \mathcal{S} is *k-compressible* if, for any **finite** subfamily $\mathcal{F} \subseteq \mathcal{S}$, there exists a subfamily $\mathcal{G} \subseteq \mathcal{S}$ of size **at most** k such that $\cap \mathcal{G} \subseteq \cap \mathcal{F}$.


Equivalently, \mathcal{S} is *k-compressible* if, for any subfamily $\mathcal{F} \subseteq \mathcal{S}$ of size $k + 1$, there exists a subfamily $\mathcal{G} \subseteq \mathcal{S}$ of size **at most** k such that $\cap \mathcal{G} \subseteq \cap \mathcal{F}$.

Note: Directed = 1-compressible.


“This terminology is inspired by the terminology around compression schemes in the statistical learning literature.”

- The set of real intervals in 0^+ is 1-compressible.
- The set of unbounded real intervals in 0^+ is 2-compressible.
- In \mathbb{R}^k , the set of “stripes” of the form $\{(x_1, \dots, x_k) : a < x_i < b\}$ for $a \leq 0 < b$ and $i \leq n$, is k -compressible.


- The set of real intervals in 0^+ is 1-compressible.
- The set of unbounded real intervals in 0^+ is 2-compressible.
- In \mathbb{R}^k , the set of “stripes” of the form $\{(x_1, \dots, x_k) : a < x_i < b\}$ for $a \leq 0 < b$ and $i \leq n$, is k -compressible.




- The set of real intervals in 0^+ is 1-compressible.
- The set of unbounded real intervals in 0^+ is 2-compressible.
- In \mathbb{R}^k , the set of “stripes” of the form $\{(x_1, \dots, x_k) : a < x_i < b\}$ for $a \leq 0 < b$ and $i \leq n$, is k -compressible.



- The set of real intervals in 0^+ is 1-compressible.
- The set of unbounded real intervals in 0^+ is 2-compressible.
- In \mathbb{R}^k , the set of “stripes” of the form $\{(x_1, \dots, x_k) : a < x_i < b\}$ for $a \leq 0 < b$ and $i \leq n$, is k -compressible.



- The set of real intervals in 0^+ is 1-compressible.
- The set of unbounded real intervals in 0^+ is 2-compressible.
- In \mathbb{R}^k , the set of “stripes” of the form $\{(x_1, \dots, x_k) : a < x_i < b\}$ for $a \leq 0 < b$ and $i \leq n$, is k -compressible.



- The set of real intervals in 0^+ is 1-compressible.
- The set of unbounded real intervals in 0^+ is 2-compressible.
- In \mathbb{R}^k , the set of “stripes” of the form $\{(x_1, \dots, x_k) : a < x_i < b\}$ for $a \leq 0 < b$ and $i \leq n$, is k -compressible.

A k -compressible family of sets is consistent iff it is k -consistent (every k sets have non-empty intersection).

So for k -compressible **definable** families of sets consistency is expressible with a single first order sentence.

Let T be a theory and $M \models T$.

For a formula $\varphi(x, y)$ let $\varphi^1(x, y) = \varphi(x, y)$ and $\varphi^0(x, y) = \neg\varphi(x, y)$.

Let T be a theory and $M \models T$.

For a formula $\varphi(x, y)$ let $\varphi^1(x, y) = \varphi(x, y)$ and $\varphi^0(x, y) = \neg\varphi(x, y)$.

By a φ -type $p(x)$ over $B \subseteq M^y$ we mean a maximal consistent collection of formulas of the form $\varphi^i(x, b)$, for $b \in B$ and $i \in \{0, 1\}$.

Let T be a theory and $M \models T$.

For a formula $\varphi(x, y)$ let $\varphi^1(x, y) = \varphi(x, y)$ and $\varphi^0(x, y) = \neg\varphi(x, y)$.

By a φ -type $p(x)$ over $B \subseteq M^y$ we mean a maximal consistent collection of formulas of the form $\varphi^i(x, b)$, for $b \in B$ and $i \in \{0, 1\}$.

We identify types $p(x)$ with the family of sets defined in M by the formulas in it.

Let T be a theory and $M \models T$.

For a formula $\varphi(x, y)$ let $\varphi^1(x, y) = \varphi(x, y)$ and $\varphi^0(x, y) = \neg\varphi(x, y)$.

By a φ -type $p(x)$ over $B \subseteq M^y$ we mean a maximal consistent collection of formulas of the form $\varphi^i(x, b)$, for $b \in B$ and $i \in \{0, 1\}$.

We identify types $p(x)$ with the family of sets defined in M by the formulas in it.

Definition

A φ -type $p(x)$ is k -compressible if for any **finite** partial type $q(x) \subseteq p(x)$ there exists $r(x) \subseteq p(x)$ with $|r(x)| \leq k$ such that

$$r(x) \vdash q(x) \quad (\text{i.e. } r(M) \subseteq q(M)).$$

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$.

There exists $k, n < \omega$ such that, for every $B \subseteq M^y$ and φ -type $p(x)$ over B , there exist n many k -compressible φ -types over B ,

$$q_1(x), \dots, q_n(x),$$

such that any formula $\varphi(x, b)$, $b \in B$, belongs in $p(x)$ if and only if it belongs in more than $n/2$ of the types $q_i(x)$:

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > n/2, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$.

There exists $k, n < \omega$ such that, for every $B \subseteq M^y$ and φ -type $p(x)$ over B , there exist n many k -compressible φ -types over B ,

$$q_1(x), \dots, q_n(x),$$

such that any formula $\varphi(x, b)$, $b \in B$, belongs in $p(x)$ if and only if it belongs in more than $n/2$ of the types $q_i(x)$:

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > n/2, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

The type $p(x)$ is the *rounded average* of the k -compressible types $q_1(x), \dots, q_n(x)$.

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$.

There exists $k, n < \omega$ such that, for every $B \subseteq M^y$ and φ -type $p(x)$ over B , there exist n many k -compressible φ -types over B ,

$$q_1(x), \dots, q_n(x),$$

such that any formula $\varphi(x, b)$, $b \in B$, belongs in $p(x)$ if and only if it belongs in more than $n/2$ of the types $q_i(x)$:

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > n/2, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

The type $p(x)$ is the *rounded average* of the k -compressible types $q_1(x), \dots, q_n(x)$.

The n and k depend only on the dual VC-dimension of φ (the maximum d such that $\pi_\varphi^*(d) = 2^d$).

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$. Fix $1/2 \leq \alpha < 1$.

There exists $k, n < \omega$ such that, for every $B \subseteq M^y$ and φ -type $p(x)$ over B , there exist n many k -compressible φ -types over B ,

$$q_1(x), \dots, q_n(x),$$

such that

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > \alpha n, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

The $n/2$ in the theorem can be improved to αn for any fixed $1/2 \leq \alpha < 1$.

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$. Fix $1/2 \leq \alpha < 1$.

There exists $k, n < \omega$ such that, for every $B \subseteq M^y$ and φ -type $p(x)$ over B , there exist n many k -compressible φ -types over B ,

$$q_1(x), \dots, q_n(x),$$

such that

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > \alpha n, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

The $n/2$ in the theorem can be improved to αn for any fixed $1/2 \leq \alpha < 1$.

Theorem B was used to prove existence of uniform honest definitions for NIP formulas.

Two considerations around Theorem B

- Does VC-codensity play a role in refining Theorem B?
- If the type $p(x)$ is definable, can the k -compressible types $q_1(x), \dots, q_n(x)$ be chosen definable too?

Does VC-codensity play a role in refining Theorem B?

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$. Then there exists $k, n < \omega$ such that every φ -type over any $B \subseteq M^y$ is the rounded average of n many k -compressible φ -types over B .

Does VC-codensity play a role in refining Theorem B?

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$. Then there exists $k, n < \omega$ such that every φ -type over any $B \subseteq M^y$ is the rounded average of n many k -compressible φ -types over B .

Proposition C (A.G. '23) - *Uniform Better Lemma A*

Let $\varphi(x, y)$ be a formula in a theory T with $\mathbf{vc}^*(\varphi) < 2$. Let $M \models T$. Then there exist $k, n < \omega$ such, for every φ -type $p(x)$ over any $B \subseteq M^y$, at least one holds:

- (a) $p(x)$ is k -compressible.
- (b) There are n many partial φ -types $q_1(x), \dots, q_n(x)$, with $\mathbf{vc}^*(q_i) < 1$ for every $i \leq n$, such that every formula in $p(x)$ belongs in at least one of the $q_i(x)$.

Does VC-codensity play a role in refining Theorem B?

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$. Then there exists $k, n < \omega$ such that every φ -type over any $B \subseteq M^y$ is the rounded average of n many k -compressible φ -types over B .

Proposition C (A.G. '23) - *Uniform Better Lemma A*

Let $\varphi(x, y)$ be a formula in a theory T with $\mathbf{vc}^*(\varphi) < 2$. Let $M \models T$. Then there exist $k, n < \omega$ such, for every φ -type $p(x)$ over any $B \subseteq M^y$, at least one holds:

- (a) $p(x)$ is k -compressible.
- (b) There are n many partial φ -types $q_1(x), \dots, q_n(x)$, with $\mathbf{vc}^*(q_i) < 1$ for every $i \leq n$, such that every formula in $p(x)$ belongs in at least one of the $q_i(x)$.

Does Proposition C have a generalization to $\mathbf{vc}^*(\varphi) < n$?

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

Why do we care?

Because we are interested in further understanding the density of k -compressible types.

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

Why do we care?

Because we are interested in further understanding the density of k -compressible types.

Corollary D (of Theorem B)

Let M be a distal structure, $\varphi(x, y)$ a formula, and $p(x)$ a φ -type. Then there exists another formula $\psi(x, z)$ and a directed (i.e. 1-compressible) ψ -type $q(x)$ such that

$$q(x) \vdash p(x).$$

($q(x)$ can be chosen over the same parameters as $p(x)$)

This corollary is a strong “uniform” density result for k -compressible types.

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

Question E

In Corollary D, if the φ -type $p(x)$ extends to a definable type over M (in the full language), can the directed ψ -type $q(x)$ be chosen definable?

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

Question E

In Corollary D, if the φ -type $p(x)$ extends to a definable type over M (in the full language), can the directed ψ -type $q(x)$ be chosen definable?

A positive answer to our question about Theorem B, would yield a positive answer to Question E.

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?



This one.

Question E

In Corollary D, if the φ -type $p(x)$ extends to a definable type over M (in the full language), can the directed ψ -type $q(x)$ be chosen definable?

A positive answer to our question about Theorem B, would yield a positive answer to Question E.

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

A theory T is *of VC-density one* if every formula $\varphi(x, y)$ with $|x| = 1$ has VC-codensity less than 2.

Using Proposition C we can show that Question E has a positive answer in distal theories of VC-density one.

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

A theory T is *of VC-density one* if every formula $\varphi(x, y)$ with $|x| = 1$ has VC-codensity less than 2.

Using Proposition C we can show that Question E has a positive answer in distal theories of VC-density one.

Theorem (A.G. '23)

Let M be a distal structure of VC-density one, $\varphi(x, y)$ be a formula, and $p(x)$ a φ -type. If $p(x)$ extends to a **definable** type in $S_x(M)$, then there exists another formula $\psi(x, z)$ and a directed **definable** ψ -type $q(x)$ such that

$$q(x) \vdash p(x).$$

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

Itay Kaplan <[REDACTED]>



Para: Pablo Andujar Guerrero

Mar 07/03/2023 18:57

I talked about it with Martin and Pierre.

It seems that there might be a counterexample but maybe it's not so easy to verify.

The idea is to take a meet-tree with a height function. So let's take M to be $(\omega^{\omega}, \omega, h)$ where h goes from $w^{<w}$ to w and gives the level. The language has the tree order, the ω order, h (and the meet). Now if you take a point s in the tree, you can look at the type saying $x \geq s$ and x is incomparable to any other point $\geq s$ (the generic type of a "closed ball") and $h(x) > M$ (height = infinity).

Then this type is the average of compressible types by taking 3 different branches $> s$, say B_1, B_2, B_3 , and looking at the types saying $x > B_i$, $h(x) > M$, but these branches won't be definable.

In order to verify that this actually works, one would need to understand what are definable types, etc.

Itay

In the 2-sorted structure $(\omega^{<\omega} \cup \omega, <, <_\omega, h)$, with $h : \omega^{<\omega} \rightarrow \omega$ the height function, consider the formulas

$$\varphi_1(x, y_1) := "(x = y_1) \vee (x < y_1) \vee (x > y_1)" \text{ for } y_1 \in \omega^{<\omega},$$

$$\varphi_2(x, y_2) := "y_2 <_\omega h(x)" \text{ for } y_2 < \omega.$$

In the 2-sorted structure $(\omega^{<\omega} \cup \omega, <, <_\omega, h)$, with $h : \omega^{<\omega} \rightarrow \omega$ the height function, consider the formulas

$$\varphi_1(x, y_1) := "(x = y_1) \vee (x < y_1) \vee (x > y_1)" \text{ for } y_1 \in \omega^{<\omega},$$

$$\varphi_2(x, y_2) := "y_2 <_\omega h(x)" \text{ for } y_2 < \omega.$$

Consider the $\{\varphi_1, \varphi_2\}$ -type $p(x)$ given by

$$\varphi_1(x, ()),$$

$$\neg \varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\},$$

$$b_2 <_\omega h(x) \text{ for every } b_2 < \omega.$$

The type $p(x)$ is clearly definable.

In the 2-sorted structure $(\omega^{<\omega} \cup \omega, <, <_\omega, h)$, with $h : \omega^{<\omega} \rightarrow \omega$ the height function, consider the formulas

$$\varphi_1(x, y_1) := "(x = y_1) \vee (x < y_1) \vee (x > y_1)" \text{ for } y_1 \in \omega^{<\omega},$$

$$\varphi_2(x, y_2) := "y_2 <_\omega h(x)" \text{ for } y_2 < \omega.$$

Consider the $\{\varphi_1, \varphi_2\}$ -type $p(x)$ given by

$$\varphi_1(x, ()),$$

$$\neg \varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\},$$

$$b_2 <_\omega h(x) \text{ for every } b_2 < \omega.$$

The type $p(x)$ is clearly definable.

(I believe that we can apply Theorem B to $p(x)$.)

Consider the $\{\varphi_1, \varphi_2\}$ -type $p(x)$ given by

$$\begin{aligned} &\varphi_1(x, ()), \\ &\neg\varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\}, \\ &b_2 <_\omega h(x) \text{ for every } b_2 < \omega. \end{aligned}$$

The type $p(x)$ is clearly definable.

(I believe that we can apply Theorem B to $p(x)$.)

Consider the $\{\varphi_1, \varphi_2\}$ -type $p(x)$ given by

$$\begin{aligned} &\varphi_1(x, ()), \\ &\neg\varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\}, \\ &b_2 <_\omega h(x) \text{ for every } b_2 < \omega. \end{aligned}$$

The type $p(x)$ is clearly definable.

(I believe that we can apply Theorem B to $p(x)$.)

Pick 3 disjoint branches B_1, B_2, B_3 . For each $i \in \{1, 2, 3\}$ let $q_i(x)$ be the $\{\varphi_1, \varphi_2\}$ -type:

$$\begin{aligned} &\varphi_1(x, b_1) \text{ iff } b_1 \in B_i, \\ &b_2 <_\omega h(x) \text{ for every } b_2 < \omega. \end{aligned}$$

Consider the $\{\varphi_1, \varphi_2\}$ -type $p(x)$ given by

$$\begin{aligned} &\varphi_1(x, ()), \\ &\neg\varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\}, \\ &b_2 <_\omega h(x) \text{ for every } b_2 < \omega. \end{aligned}$$

The type $p(x)$ is clearly definable.

(I believe that we can apply Theorem B to $p(x)$.)

Pick 3 disjoint branches B_1, B_2, B_3 . For each $i \in \{1, 2, 3\}$ let $q_i(x)$ be the $\{\varphi_1, \varphi_2\}$ -type:

$$\begin{aligned} &\varphi_1(x, b_1) \text{ iff } b_1 \in B_i, \\ &b_2 <_\omega h(x) \text{ for every } b_2 < \omega. \end{aligned}$$

The type $p(x)$ is the rounded average of the types $q_1(x), q_2(x), q_3(x)$. Moreover these are 2-compressible. However they are probably not definable.

Email (Bays-Kaplan-Simon)

The $\{\varphi_1, \varphi_2\}$ -type $p(x)$ is definable but might not be the rounded average of finitely many k -compressible (for any k) **definable** $\{\varphi_1, \varphi_2\}$ -types.

In particular $p(x)$ is the rounded average of the 2-compressible types $q_1(x)$, $q_2(x)$ and $q_3(x)$, but these are probably not definable.

The big picture: what do we know about NIP types?

1. Few types over finite sets.
2. Indiscernible sequences converge to types.
3. Equivalence of invariance, non-dividing and non-forking over models.
4. Existence of honest definitions.
5. Described by **compressible types** (Theorem B).

The big picture: what do we know about NIP types?

1. Few types over finite sets.
2. Indiscernible sequences converge to types.
3. Equivalence of invariance, non-dividing and non-forking over models.
4. Existence of honest definitions.
5. Described by **compressible types** (Theorem B).

Opinion: VC-(co)density is conspicuously absent.