

NIP compressible types

Pablo Andújar Guerrero

University of Leeds

anguempa.github.io/handout.pdf

Theorem: For any NIP formula $\varphi(x, y)$ there exists $n, k < \omega$ such any φ -type $p(x)$ is a rounded average of n many k -compressible φ -types.

Theorem: For any NIP formula $\varphi(x, y)$ there exists $n, k < \omega$ such any φ -type $p(x)$ is a rounded average of n many k -compressible φ -types.


Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .

Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .


$$\mathcal{F} = \{S_1, S_2, S_3\}$$



Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .


$$S_1 \setminus (S_2 \cup S_3)$$



Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .

$$S_3 \setminus (S_1 \cup S_2)$$



Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .

(In model-theoretic terminology a Boolean atom can be called a type.)


Let $BA(\mathcal{F})$ denote the number of Boolean atoms of \mathcal{F} .

Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .

Let $BA(\mathcal{F})$ denote the number of Boolean atoms of \mathcal{F} .

$$BA(\mathcal{F}) = 5$$




Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .

Let $BA(\mathcal{F})$ denote the number of Boolean atoms of \mathcal{F} .

$$BA(\mathcal{F}) = 2^3 = 8$$



Counting Boolean atoms

A *Boolean atom* of a finite family of sets \mathcal{F} is a maximal non-empty Boolean combination of sets in \mathcal{F} .

Let $BA(\mathcal{F})$ denote the number of Boolean atoms of \mathcal{F} .

Definition

Let \mathcal{S} be a family of sets. The *dual shatter function* of \mathcal{S} is the function

$$\begin{aligned}\pi_{\mathcal{S}}^* : \omega &\rightarrow \omega \\ n &\mapsto \max\{BA(\mathcal{F}) : \mathcal{F} \subseteq \mathcal{S}, |\mathcal{F}| = n\}\end{aligned}$$

Counting Boolean atoms

Observe that $\pi_{\mathcal{S}}^*(n) \leq 2^n$ for all n .

Counting Boolean atoms

Observe that $\pi_{\mathcal{S}}^*(n) \leq 2^n$ for all n .

If $\pi_{\mathcal{S}}^*(d) < 2^d$ for some d then $\pi_{\mathcal{S}}^*(n) = O(n^{d-1})$ (Sauer's Lemma).

Counting Boolean atoms

Observe that $\pi_{\mathcal{S}}^*(n) \leq 2^n$ for all n .

If $\pi_{\mathcal{S}}^*(d) < 2^d$ for some d then $\pi_{\mathcal{S}}^*(n) = O(n^{d-1})$ (Sauer's Lemma).

In this case we call \mathcal{S} a *VC class* (VC=Vapnik–Chervonenkis).

Counting Boolean atoms

Observe that $\pi_{\mathcal{S}}^*(n) \leq 2^n$ for all n .

If $\pi_{\mathcal{S}}^*(d) < 2^d$ for some d then $\pi_{\mathcal{S}}^*(n) = O(n^{d-1})$ (Sauer's Lemma).

In this case we call \mathcal{S} a *VC class* (VC=Vapnik–Chervonenkis).

Definition

The VC-codensity of a VC class \mathcal{S} is

$$\text{vc}^*(\mathcal{S}) = \inf\{r \in \mathbb{R}^{\geq 0} : \pi_{\mathcal{S}}^*(n) = O(n^r)\}.$$

Counting Boolean atoms

Observe that $\pi_{\mathcal{S}}^*(n) \leq 2^n$ for all n .

If $\pi_{\mathcal{S}}^*(d) < 2^d$ for some d then $\pi_{\mathcal{S}}^*(n) = O(n^{d-1})$ (Sauer's Lemma).

In this case we call \mathcal{S} a *VC class* (VC=Vapnik–Chervonenkis).

Definition

The VC-codensity of a VC class \mathcal{S} is

$$\text{vc}^*(\mathcal{S}) = \inf\{r \in \mathbb{R}^{\geq 0} : \pi_{\mathcal{S}}^*(n) = O(n^r)\}.$$

E.g. If $\mathcal{S} = \{\text{intervals in } \mathbb{R}\}$ then $\pi_{\mathcal{S}}^*(n) = 2n$ and $\text{vc}^*(\mathcal{S}) = 1$.

Definition

A family of sets \mathcal{S} is *(downward) directed* if, for any **finite** subfamily $\mathcal{F} \subseteq \mathcal{S}$, there exists some $S \in \mathcal{S}$ such that $S \subseteq \cap \mathcal{F}$.


Equivalently, \mathcal{S} is directed if, for any $S_1, S_2 \in \mathcal{S}$ there exists $S \in \mathcal{S}$ with $S \subseteq S_1 \cap S_2$.

E.g.

- In a poset (P, \leq) the sets $\{a \in P : a \leq b\}$ for $b \in P$.
- The neighborhoods of a given point in a topological spaces.
- Any filter basis.


Definition

A family of sets \mathcal{S} has a finite transversal $T = \{a_1, \dots, a_n\}$ if every $S \in \mathcal{S}$ satisfies that $S \cap T \neq \emptyset$.



Definition


A family of sets \mathcal{S} has a finite transversal $T = \{a_1, \dots, a_n\}$ if every $S \in \mathcal{S}$ satisfies that $S \cap T \neq \emptyset$.



Definition

A family of sets \mathcal{S} has a finite transversal $T = \{a_1, \dots, a_n\}$ if every $S \in \mathcal{S}$ satisfies that $S \cap T \neq \emptyset$.


We care about the property that \mathcal{S} does **not** have a finite transversal.



Definition

A family of sets \mathcal{S} has a finite transversal $T = \{a_1, \dots, a_n\}$ if every $S \in \mathcal{S}$ satisfies that $S \cap T \neq \emptyset$.

We care about the property that \mathcal{S} does **not** have a finite transversal.



Is there a connection between directedness and the dual shatter function?


Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.


\mathcal{S} could be directed.



Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

\mathcal{S} could be directed.




Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

\mathcal{S} could be directed.


$$\pi_{\mathcal{S}}^*(3) \geq 5$$



Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .




Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .

$$\pi_{\mathcal{S}}^*(3) \geq 8$$




Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .

$$\pi_{\mathcal{S}}^*(3) \geq 6$$




Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .

$$\pi_{\mathcal{S}}^*(3) \geq 6$$




Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .

$$\pi_{\mathcal{S}}^*(3) \geq 8$$




Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .

$$BA(S_1, S_2) \geq 3$$




Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .

$$BA(S_1, S_2) \geq 3$$




Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .

$$BA(S_1, S_2, S_3) \geq 3 + 2 = 5$$




Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .

$$BA(S_1, S_2, S_3) \geq 3 + 1 + \textcolor{red}{1}.$$




Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .

$$F = S_1 \cap S_2 \cap S_3$$



Is there a connection between directedness and the dual shatter function?

Fix a family of sets \mathcal{S} that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that \mathcal{S} is **not** directed, witnessed by S_1 and S_2 .

$$F = S_1 \cap S_2 \cap S_3$$



Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.


$$F = S_1 \cap S_2 \cap S_3$$



Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.


$F \cap \mathcal{S}$ could be directed.



Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.


$F \cap \mathcal{S}$ could be directed.



Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.

$F \cap \mathcal{S}$ could be directed.




Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.

Suppose that $F \cap \mathcal{S}$ is **not** directed, witnessed by S_4 and S_5 .

$$BA(S_1, S_2, S_3, S_4, S_5) \geq BA(S_1, S_2, S_3) + 3$$




Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.

Suppose that $F \cap \mathcal{S}$ is **not** directed, witnessed by S_4 and S_5 .

$$BA(S_1, S_2, S_3, S_4, S_5) \geq BA(S_1, S_2, S_3) + 3$$




Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.

Suppose that $F \cap \mathcal{S}$ is **not** directed, witnessed by S_4 and S_5 .

$$BA(S_1, S_2, S_3, S_4, S_5) \geq BA(S_1, S_2, S_3) + 3$$




Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.

Suppose that $F \cap \mathcal{S}$ is **not** directed, witnessed by S_4 and S_5 .

$$BA(S_1, \dots, S_6) \geq BA(S_1, S_2, S_3) + 3 + 2.$$




Is there a connection between directedness and the dual shatter function?

Consider the family of sets $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$.

Suppose that $F \cap \mathcal{S}$ is **not** directed, witnessed by S_4 and S_5 .

$$BA(S_1, \dots, S_6) \geq BA(S_1, S_2, S_3) + 3 + 2 + 1.$$



Is there a connection between directedness and the dual shatter function?

We continue this process, and one of two things can happen:

- We find a finite intersection F of sets in \mathcal{S} such that $F \cap \mathcal{S}$ is directed.
- There is a sequence $(S_i)_i$ of sets in \mathcal{S} such that

$$BA(B_i : i \leq 3n) \geq \sum_{j=1}^n j.$$

Meaning that

$$\pi_{\mathcal{S}}^*(3n) \geq \frac{n^2 + n}{2}.$$

Lemma A (A.G. '23)

Let \mathcal{S} be a family of sets with $\text{vc}^*(\mathcal{S}) < 2$. Then at least one holds:

- (a) There exists a finite intersection F of sets in \mathcal{S} such that $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ is directed.
- (b) \mathcal{S} has a finite transversal.

Lemma A (A.G. '23)

Let \mathcal{S} be a family of sets with $\text{vc}^*(\mathcal{S}) < 2$. Then at least one holds:

- (a) There exists a finite intersection F of sets in \mathcal{S} such that $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ is directed.
- (b) \mathcal{S} has a finite transversal.

In fact we can improve (b) in the above lemma.

Lemma A (A.G. '23)

Let \mathcal{S} be a family of sets with $\text{vc}^*(\mathcal{S}) < 2$. Then at least one holds:

- (a) There exists a finite intersection F of sets in \mathcal{S} such that $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ is directed.
- (b) \mathcal{S} has a finite transversal.

In fact we can improve (b) in the above lemma.

Better Lemma A (A.G. '23)

Let \mathcal{S} be a family of sets with $\text{vc}^*(\mathcal{S}) < 2$. Then at least one holds:

- (a) There exists a finite intersection F of sets in \mathcal{S} such that $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ is directed.
- (b) There exists F_1, \dots, F_m , finite non-empty Boolean combinations of sets in \mathcal{S} such that, for every $S \in \mathcal{S}$ there is some $i \leq m$ such that $F_i \subseteq S$.

Now let us talk compressibility.

Our two main references are:

- Martin Bays, Itay Kaplan and Pierre Simon. *Density of compressible types and some consequences*.
ArXiv: 2107.05197, 2022.
- Xi Chen, Yu Cheng, and Bo Tang. *On the recursive teaching dimension of VC classes*.
Advances in Neural Information Processing Systems 29:
Annual Conference on Neural Information Processing
Systems, 2016, Barcelona, Spain, p 2164–2171.

Model-theoretic framework

Let T be a theory and $\varphi(x, y)$ be a (partitioned) formula. For any $M \models T$ and $b \in M^y$ let $\varphi(M, b) = \{a \in M^x : M \models \varphi(a, b)\}$. We identify $\varphi(x, y)$ with the family of sets $\{\varphi(M, b) : b \in M^y\}$:

- $\varphi(x, y)$ may be directed.
- It has dual shatter function π_φ^* .
- It has VC-codensity ...

(All these are independent of M .)

Model-theoretic framework

Let T be a theory and $\varphi(x, y)$ be a (partitioned) formula. For any $M \models T$ and $b \in M^y$ let $\varphi(M, b) = \{a \in M^x : M \models \varphi(a, b)\}$. We identify $\varphi(x, y)$ with the family of sets $\{\varphi(M, b) : b \in M^y\}$:

- $\varphi(x, y)$ may be directed.
- It has dual shatter function π_φ^* .
- It has VC-codensity ...

(All these are independent of M .)

A formula φ is *NIP* if it is a VC class.

A theory T is *NIP* if every formula φ is NIP.

Model-theoretic framework

Let T be a theory and $\varphi(x, y)$ be a (partitioned) formula. For any $M \models T$ and $b \in M^y$ let $\varphi(M, b) = \{a \in M^x : M \models \varphi(a, b)\}$. We identify $\varphi(x, y)$ with the family of sets $\{\varphi(M, b) : b \in M^y\}$:

- $\varphi(x, y)$ may be directed.
- It has dual shatter function π_φ^* .
- It has VC-codensity ...

(All these are independent of M .)

A formula φ is *NIP* if it is a VC class.

A theory T is *NIP* if every formula φ is NIP.

A theory T is of *VC-density one* if every formula $\varphi(x, y)$ with $|x| = 1$ has VC-codensity less than 2.

$$\text{VC-density one} \subseteq \text{dp-minimal} \subsetneq \text{NIP}$$

Model-theoretic framework

Let T be a theory and $\varphi(x, y)$ be a (partitioned) formula. For any $M \models T$ and $b \in M^y$ let $\varphi(M, b) = \{a \in M^x : M \models \varphi(a, b)\}$. We identify $\varphi(x, y)$ with the family of sets $\{\varphi(M, b) : b \in M^y\}$:

- $\varphi(x, y)$ may be directed.
- It has dual shatter function π_φ^* .
- It has VC-codensity ...

(All these are independent of M .)

A formula φ is *NIP* if it is a VC class.

A theory T is *NIP* if every formula φ is NIP.

A theory T is of *VC-density one* if every formula $\varphi(x, y)$ with $|x| = 1$ has VC-codensity less than 2.

Conjecture

VC-density one = dp-minimal \subsetneq NIP

Theorem (Baldwin-Saxl '76). *Let G be a group definable in an NIP structure. Let \mathcal{H} be a (uniformly) definable family of subgroups of G . Then there is $k < \omega$ such that, for any finite family $\mathcal{F} \subseteq \mathcal{H}$, there is a subfamily $\mathcal{C} \subseteq \mathcal{F}$ with $|\mathcal{C}| \leq k$ such that $\bigcap \mathcal{C} = \bigcap \mathcal{F}$.*

Definition

Let $k < \omega$. A family of sets \mathcal{S} is *k-compressible* if, for any **finite** subfamily $\mathcal{F} \subseteq \mathcal{S}$, there exists a subfamily $\mathcal{G} \subseteq \mathcal{S}$ of size **at most** k such that $\cap \mathcal{G} \subseteq \cap \mathcal{F}$.

Equivalently, \mathcal{S} is *k-compressible* if, for any subfamily $\mathcal{F} \subseteq \mathcal{S}$ of size $k + 1$, there exists a subfamily $\mathcal{G} \subseteq \mathcal{S}$ of size **at most** k such that $\cap \mathcal{G} \subseteq \cap \mathcal{F}$.

Definition

Let $k < \omega$. A family of sets \mathcal{S} is *k-compressible* if, for any **finite** subfamily $\mathcal{F} \subseteq \mathcal{S}$, there exists a subfamily $\mathcal{G} \subseteq \mathcal{S}$ of size **at most** k such that $\cap \mathcal{G} \subseteq \cap \mathcal{F}$.

Equivalently, \mathcal{S} is *k-compressible* if, for any subfamily $\mathcal{F} \subseteq \mathcal{S}$ of size $k + 1$, there exists a subfamily $\mathcal{G} \subseteq \mathcal{S}$ of size **at most** k such that $\cap \mathcal{G} \subseteq \cap \mathcal{F}$.


Note: Directed = 1-compressible.

“This terminology is inspired by the terminology around compression schemes in the statistical learning literature.”


— Bays-Kaplan-Simon

- The set of real intervals in 0^+ is 1-compressible.
- The set of unbounded real intervals in 0^+ is 2-compressible.
- In \mathbb{R}^k , the set of “stripes” of the form $\{(x_1, \dots, x_k) : a < x_i < b\}$ for $a \leq 0 < b$ and $i \leq n$, is k -compressible.


- The set of real intervals in 0^+ is 1-compressible.
- The set of unbounded real intervals in 0^+ is 2-compressible.
- In \mathbb{R}^k , the set of “stripes” of the form $\{(x_1, \dots, x_k) : a < x_i < b\}$ for $a \leq 0 < b$ and $i \leq n$, is k -compressible.




- The set of real intervals in 0^+ is 1-compressible.
- The set of unbounded real intervals in 0^+ is 2-compressible.
- In \mathbb{R}^k , the set of “stripes” of the form $\{(x_1, \dots, x_k) : a < x_i < b\}$ for $a \leq 0 < b$ and $i \leq n$, is k -compressible.



- The set of real intervals in 0^+ is 1-compressible.
- The set of unbounded real intervals in 0^+ is 2-compressible.
- In \mathbb{R}^k , the set of “stripes” of the form $\{(x_1, \dots, x_k) : a < x_i < b\}$ for $a \leq 0 < b$ and $i \leq n$, is k -compressible.



- The set of real intervals in 0^+ is 1-compressible.
- The set of unbounded real intervals in 0^+ is 2-compressible.
- In \mathbb{R}^k , the set of “stripes” of the form $\{(x_1, \dots, x_k) : a < x_i < b\}$ for $a \leq 0 < b$ and $i \leq n$, is k -compressible.



- The set of real intervals in 0^+ is 1-compressible.
- The set of unbounded real intervals in 0^+ is 2-compressible.
- In \mathbb{R}^k , the set of “stripes” of the form $\{(x_1, \dots, x_k) : a < x_i < b\}$ for $a \leq 0 < b$ and $i \leq n$, is k -compressible.

A k -compressible family of sets is consistent iff it is k -consistent (every k sets have non-empty intersection).

So for k -compressible **definable** families of sets consistency is expressible with a single first order sentence.

Let T be a theory and $M \models T$.

For a formula $\varphi(x, y)$ let $\varphi^1(x, y) = \varphi(x, y)$ and $\varphi^0(x, y) = \neg\varphi(x, y)$.

Let T be a theory and $M \models T$.

For a formula $\varphi(x, y)$ let $\varphi^1(x, y) = \varphi(x, y)$ and $\varphi^0(x, y) = \neg\varphi(x, y)$.

By a φ -type $p(x)$ over $B \subseteq M^y$ we mean a maximal consistent collection of formulas of the form $\varphi^i(x, b)$, for $b \in B$ and $i \in \{0, 1\}$.

Let T be a theory and $M \models T$.

For a formula $\varphi(x, y)$ let $\varphi^1(x, y) = \varphi(x, y)$ and $\varphi^0(x, y) = \neg\varphi(x, y)$.

By a φ -type $p(x)$ over $B \subseteq M^y$ we mean a maximal consistent collection of formulas of the form $\varphi^i(x, b)$, for $b \in B$ and $i \in \{0, 1\}$.

We identify types $p(x)$ with the family of sets defined in M by the formulas in it.

Let T be a theory and $M \models T$.

For a formula $\varphi(x, y)$ let $\varphi^1(x, y) = \varphi(x, y)$ and $\varphi^0(x, y) = \neg\varphi(x, y)$.

By a φ -type $p(x)$ over $B \subseteq M^y$ we mean a maximal consistent collection of formulas of the form $\varphi^i(x, b)$, for $b \in B$ and $i \in \{0, 1\}$.

We identify types $p(x)$ with the family of sets defined in M by the formulas in it.

Definition

A φ -type $p(x)$ is k -compressible if for any **finite** partial type $q(x) \subseteq p(x)$ there exists $r(x) \subseteq p(x)$ with $|r(x)| \leq k$ such that

$$r(x) \vdash q(x) \quad (\text{i.e. } r(M) \subseteq q(M)).$$

Better Lemma A (A.G. '23)

Let $\varphi(x, y)$ be a formula in a theory T and $M \models T$. Let $B \subseteq M^y$ and $p(x)$ be a φ -type over B with $\text{vc}^*(p(x)) < 2$.

Then at least one holds:

- (a) There exists a finite conjunction $\psi(x)$ of formulas in $p(x)$ such that $\psi(x) \wedge p(x) = \{\psi(x) \wedge \sigma(x) : \sigma(x) \in p(x)\}$ is directed (i.e. 1-compressible).

In particular $p(x)$ is k -compressible for some $k < \omega$.

- (b) There exist $q_1(x), \dots, q_m(x)$ finite (partial) φ -types such that, for every $\sigma(x) \in p(x)$ there exists some $i \leq m$ such that $q_i(x) \vdash \sigma(x)$.

Better Lemma A* (A.G. and Johnson '23)

Let $\varphi(x, y)$ be a formula with $|x| = 1$ in a dp-minimal theory T and $M \models T$. Let $B \subseteq M^y$ and $p(x)$ be a φ -type over B with $\text{ve}^*(p(x)) \leq 2$. Then at least one holds:

- (a) There exists a finite conjunction $\psi(x)$ of formulas in $p(x)$ such that $\psi(x) \wedge p(x) = \{\psi(x) \wedge \sigma(x) : \sigma(x) \in p(x)\}$ is directed (i.e. 1-compressible).

In particular $p(x)$ is k -compressible for some $k < \omega$.

- (b) There exist $q_1(x), \dots, q_m(x)$ finite (partial) φ -types such that, for every $\sigma(x) \in p(x)$ there exists some $i \leq m$ such that $q_i(x) \vdash \sigma(x)$.

Better Lemma A* (A.G. and Johnson '23)

Let $\varphi(x, y)$ be a formula with $|x| = 1$ in a dp-minimal theory T and $M \models T$. Let $B \subseteq M^y$ and $p(x)$ be a φ -type over B with $\text{ve}^*(p(x)) \leq 2$. Then at least one holds:

- (a) There exists a finite conjunction $\psi(x)$ of formulas in $p(x)$ such that $\psi(x) \wedge p(x) = \{\psi(x) \wedge \sigma(x) : \sigma(x) \in p(x)\}$ is directed (i.e. 1-compressible).

In particular $p(x)$ is k -compressible for some $k < \omega$.

- (b) There exist $q_1(x), \dots, q_m(x)$ finite (partial) φ -types such that, for every $\sigma(x) \in p(x)$ there exists some $i \leq m$ such that $q_i(x) \vdash \sigma(x)$.

So the type $p(x)$ is either k -compressible (some $k = k(p)$) or it can be partitioned into finitely many types that are finitely axiomatisable (in particular k -compressible).

Can we say something similar to this for all formulas in NIP theories?

Can we say something similar to this for all formulas in NIP theories?

Yes (thanks to Bays, Kaplan and Simon).

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$.

There exists $k, n < \omega$ such that, for every $B \subseteq M^y$ and φ -type $p(x)$ over B , there exist n many k -compressible φ -types over B ,

$$q_1(x), \dots, q_n(x),$$

such that any formula $\varphi(x, b)$, $b \in B$, belongs in $p(x)$ if and only if it belongs in more than $n/2$ of the types $q_i(x)$:

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > n/2, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$.

There exists $k, n < \omega$ such that, for every $B \subseteq M^y$ and φ -type $p(x)$ over B , there exist n many k -compressible φ -types over B ,

$$q_1(x), \dots, q_n(x),$$

such that any formula $\varphi(x, b)$, $b \in B$, belongs in $p(x)$ if and only if it belongs in more than $n/2$ of the types $q_i(x)$:

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > n/2, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

The type $p(x)$ is the *rounded average* of the k -compressible types $q_1(x), \dots, q_n(x)$.

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$.

There exists $k, n < \omega$ such that, for every $B \subseteq M^y$ and φ -type $p(x)$ over B , there exist n many k -compressible φ -types over B ,

$$q_1(x), \dots, q_n(x),$$

such that any formula $\varphi(x, b)$, $b \in B$, belongs in $p(x)$ if and only if it belongs in more than $n/2$ of the types $q_i(x)$:

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > n/2, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

The type $p(x)$ is the *rounded average* of the k -compressible types $q_1(x), \dots, q_n(x)$.

The n and k depend only on the dual VC-dimension of φ (the maximum d such that $\pi_\varphi^*(d) = 2^d$).

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$. Fix $1/2 \leq \alpha < 1$.

There exists $k, n < \omega$ such that, for every $B \subseteq M^y$ and φ -type $p(x)$ over B , there exist n many k -compressible φ -types over B ,

$$q_1(x), \dots, q_n(x),$$

such that

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > \alpha n, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

The $n/2$ in the theorem can be improved to αn for any fixed $1/2 \leq \alpha < 1$.

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$. Fix $1/2 \leq \alpha < 1$.

There exists $k, n < \omega$ such that, for every $B \subseteq M^y$ and φ -type $p(x)$ over B , there exist n many k -compressible φ -types over B ,

$$q_1(x), \dots, q_n(x),$$

such that

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > \alpha n, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

The $n/2$ in the theorem can be improved to αn for any fixed $1/2 \leq \alpha < 1$.

Theorem B was used to prove existence of uniform honest definitions for NIP formulas.

Two considerations around Theorem B

- Does VC-codensity play a role in refining Theorem B?
- If the type $p(x)$ is definable, can the k -compressible types $q_1(x), \dots, q_n(x)$ be chosen definable too?

Does VC-codensity play a role in refining Theorem B?

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$. Then there exists $k, n < \omega$ such that every φ -type over any $B \subseteq M^y$ is the rounded average of n many k -compressible φ -types over B .

Does VC-codensity play a role in refining Theorem B?

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$. Then there exists $k, n < \omega$ such that every φ -type over any $B \subseteq M^y$ is the rounded average of n many k -compressible φ -types over B .

Proposition C (A.G. '23) - *Uniform Better Lemma A*

Let $\varphi(x, y)$ be a formula with $\mathbf{vc}^*(\varphi) < 2$ in a theory T . Let $M \models T$. Then there exist $k, n < \omega$ such, for every φ -type $p(x)$ over any $B \subseteq M^y$, at least one holds:

- (a) $p(x)$ is k -compressible.
- (b) There are n many partial φ -types $q_1(x), \dots, q_n(x)$, with $\mathbf{vc}^*(q_i) = 0$ for every $i \leq n$, such that every formula in $p(x)$ is implied by **at least one** of the $q_i(x)$.

Does VC-codensity play a role in refining Theorem B?

Theorem B (Bays-Kaplan-Simon '22)

Let $\varphi(x, y)$ be an NIP formula in a theory T and $M \models T$. Then there exists $k, n < \omega$ such that every φ -type over any $B \subseteq M^y$ is the rounded average of n many k -compressible φ -types over B .

Proposition C (A.G. '23) - *Uniform Better Lemma A*

Let $\varphi(x, y)$ be a formula with $\mathbf{vc}^*(\varphi) < 2$ in a theory T . Let $M \models T$. Then there exist $k, n < \omega$ such, for every φ -type $p(x)$ over any $B \subseteq M^y$, at least one holds:

- (a) $p(x)$ is k -compressible.
- (b) There are n many partial φ -types $q_1(x), \dots, q_n(x)$, with $\mathbf{vc}^*(q_i) = 0$ for every $i \leq n$, such that every formula in $p(x)$ is implied by **at least one** of the $q_i(x)$.

Does Proposition C have a generalization to $\mathbf{vc}^*(\varphi) < n$?

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

Why do we care?

Because we are interested in further understanding the density of k -compressible types.

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

Why do we care?

Because we are interested in further understanding the density of k -compressible types.

Corollary D (of Theorem B)

Let M be a distal structure, $\varphi(x, y)$ a formula, and $p(x)$ a φ -type. Then there exists another formula $\psi(x, z)$ and a directed (i.e. 1-compressible) ψ -type $q(x)$ such that

$$q(x) \vdash p(x).$$

($q(x)$ can be chosen over the same parameters as $p(x)$)

This corollary is a strong “uniform” density result for k -compressible types.

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

Question E

In Corollary D, if the φ -type $p(x)$ extends to a definable type over M (in the full language), can the directed ψ -type $q(x)$ satisfying $q(x) \vdash p(x)$ be chosen definable?

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

Question E

In Corollary D, if the φ -type $p(x)$ extends to a definable type over M (in the full language), can the directed ψ -type $q(x)$ satisfying $q(x) \vdash p(x)$ be chosen definable?

A positive answer to our question about Theorem B, would yield a positive answer to Question E.

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?



This one.

Question E

In Corollary D, if the φ -type $p(x)$ extends to a definable type over M (in the full language), can the directed ψ -type $q(x)$ satisfying $q(x) \vdash p(x)$ be chosen definable?

A positive answer to our question about Theorem B, would yield a positive answer to Question E.

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

↑

This one.

Question E

In Corollary D, if the φ -type $p(x)$ extends to a definable type over M (in the full language), can the directed ψ -type $q(x)$ satisfying $q(x) \vdash p(x)$ be chosen definable?

A positive answer to our question about Theorem B, would yield a positive answer to Question E.

Using Proposition C (uniform Better Lemma A) we can show that Question E has a positive answer in distal theories of VC-density one.

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

Theorem (A.G. '23)

Let M be a distal structure of VC-density one, $\varphi(x, y)$ be a formula, and $p(x)$ a φ -type. If $p(x)$ extends to a **definable** type in $S_x(M)$, then there exists another formula $\psi(x, z)$ and a directed **definable** ψ -type $q(x)$ such that

$$q(x) \vdash p(x).$$

(dp -minimal version to come)

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

↑

This one.

But what about our question?

If the type $p(x)$ is definable, can the k -compressible types $q_i(x)$ be chosen definable too?

Itay Kaplan <[REDACTED]>



Para: Pablo Andujar Guerrero

Mar 07/03/2023 18:57

I talked about it with Martin and Pierre.

It seems that there might be a counterexample but maybe it's not so easy to verify.

The idea is to take a meet-tree with a height function. So let's take M to be $(\omega^{\omega}, \omega, h)$ where h goes from $w^{<w}$ to w and gives the level. The language has the tree order, the ω order, h (and the meet). Now if you take a point s in the tree, you can look at the type saying $x \geq s$ and x is incomparable to any other point $\geq s$ (the generic type of a "closed ball") and $h(x) > M$ (height = infinity).

Then this type is the average of compressible types by taking 3 different branches $\geq s$, say B_1, B_2, B_3 , and looking at the types saying $x \geq B_i, h(x) > M$, but these branches won't be definable.

In order to verify that this actually works, one would need to understand what are definable types, etc.

Itay

In the 2-sorted structure $(\omega^{<\omega}, \omega; <, <_\omega, h)$, with $h : \omega^{<\omega} \rightarrow \omega$ the level function, consider the formulas

$$\varphi_1(x, y_1) := "(x = y_1) \vee (x < y_1) \vee (x > y_1)" \text{ for } y_1 \in \omega^{<\omega},$$

$$\varphi_2(x, y_2) := "y_2 <_\omega h(x)" \text{ for } y_2 \in \omega.$$

In the 2-sorted structure $(\omega^{<\omega}, \omega; <, <_\omega, h)$, with $h : \omega^{<\omega} \rightarrow \omega$ the level function, consider the formulas

$$\begin{aligned}\varphi_1(x, y_1) &:= "(x = y_1) \vee (x < y_1) \vee (x > y_1)" \text{ for } y_1 \in \omega^{<\omega}, \\ \varphi_2(x, y_2) &:= "y_2 <_\omega h(x)" \text{ for } y_2 \in \omega.\end{aligned}$$

Consider the $\{\varphi_1, \varphi_2\}$ -type $p(x)$ given by

$$\begin{aligned}\varphi_1(x, ()), \\ \neg\varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\}, \\ b_2 <_\omega h(x) \text{ for every } b_2 \in \omega.\end{aligned}$$

The type $p(x)$ is clearly definable.

In the 2-sorted structure $(\omega^{<\omega}, \omega; <, <_\omega, h)$, with $h : \omega^{<\omega} \rightarrow \omega$ the level function, consider the formulas

$$\begin{aligned}\varphi_1(x, y_1) &:= "(x = y_1) \vee (x < y_1) \vee (x > y_1)" \text{ for } y_1 \in \omega^{<\omega}, \\ \varphi_2(x, y_2) &:= "y_2 <_\omega h(x)" \text{ for } y_2 \in \omega.\end{aligned}$$

Consider the $\{\varphi_1, \varphi_2\}$ -type $p(x)$ given by

$$\begin{aligned}\varphi_1(x, ()), \\ \neg\varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\}, \\ b_2 <_\omega h(x) \text{ for every } b_2 \in \omega.\end{aligned}$$

The type $p(x)$ is clearly definable.

(I believe that we can apply Theorem B to $p(x)$.)

Consider the $\{\varphi_1, \varphi_2\}$ -type $p(x)$ given by

$$\begin{aligned} &\varphi_1(x, ()), \\ &\neg\varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\}, \\ &b_2 <_\omega h(x) \text{ for every } b_2 \in \omega. \end{aligned}$$

The type $p(x)$ is clearly definable.

(I believe that we can apply Theorem B to $p(x)$.)

Consider the $\{\varphi_1, \varphi_2\}$ -type $p(x)$ given by

$$\begin{aligned} &\varphi_1(x, ()), \\ &\neg\varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\}, \\ &b_2 <_\omega h(x) \text{ for every } b_2 \in \omega. \end{aligned}$$

The type $p(x)$ is clearly definable.

(I believe that we can apply Theorem B to $p(x)$.)

Pick 3 disjoint branches B_1, B_2, B_3 . For each $i \in \{1, 2, 3\}$ let $q_i(x)$ be the $\{\varphi_1, \varphi_2\}$ -type:

$$\begin{aligned} &\varphi_1(x, b_1) \text{ iff } b_1 \in B_i, \\ &b_2 <_\omega h(x) \text{ for every } b_2 \in \omega. \end{aligned}$$

Consider the $\{\varphi_1, \varphi_2\}$ -type $p(x)$ given by

$$\begin{aligned} &\varphi_1(x, ()), \\ &\neg\varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\}, \\ &b_2 <_\omega h(x) \text{ for every } b_2 \in \omega. \end{aligned}$$

The type $p(x)$ is clearly definable.

(I believe that we can apply Theorem B to $p(x)$.)

Pick 3 disjoint branches B_1, B_2, B_3 . For each $i \in \{1, 2, 3\}$ let $q_i(x)$ be the $\{\varphi_1, \varphi_2\}$ -type:

$$\begin{aligned} &\varphi_1(x, b_1) \text{ iff } b_1 \in B_i, \\ &b_2 <_\omega h(x) \text{ for every } b_2 \in \omega. \end{aligned}$$

The type $p(x)$ is the rounded average of the types $q_1(x), q_2(x), q_3(x)$. Moreover these are 2-compressible. However they are *probably* not definable.

Email (Bays-Kaplan-Simon)

The $\{\varphi_1, \varphi_2\}$ -type $p(x)$ is definable but might not be the rounded average of finitely many k -compressible (for any k) **definable** $\{\varphi_1, \varphi_2\}$ -types.

In particular $p(x)$ is the rounded average of the 2-compressible types $q_1(x)$, $q_2(x)$ and $q_3(x)$, but these are probably not definable.

The big picture: what do we know about NIP types?

1. Few types over finite sets.
2. Indiscernible sequences converge to types.
3. Equivalence of invariance, non-dividing and non-forking over models.
4. Existence of uniform honest definitions.
5. Described by **compressible types** (Theorem B).

The big picture: what do we know about NIP types?

1. Few types over finite sets.
2. Indiscernible sequences converge to types.
3. Equivalence of invariance, non-dividing and non-forking over models.
4. Existence of uniform honest definitions.
5. Described by **compressible types** (Theorem B).

Opinion: VC-(co)density is conspicuously absent.


The bigger picture: VC meets NIP.

- 1968 – 1972: Vapnik-Chervonenkis, Shelah-Perles and Sauer all prove independently Sauer's Lemma.

The bigger picture: VC meets NIP.

- 1968 – 1972: Vapnik-Chervonenkis, Shelah-Perles and Sauer all prove independently Sauer's Lemma.
 - statistics
 - Erdős question
 - classification theory

The bigger picture: VC meets NIP.


- 1968 – 1972: Vapnik-Chervonenkis, Shelah-Perles and Sauer all prove independently Sauer's Lemma. 

→ statistics

→ Erdős question

→ classification theory
- 1992: Laskowski “discovers” the connection between VC learning and NIP model theory. *He uses model theory to point out new VC classes (e.g. families of sets definable in o-minimal structures).*

The bigger picture: VC meets NIP.


- 1968 – 1972: Vapnik-Chervonenkis, Shelah-Perles and Sauer all prove independently Sauer's Lemma. 

→ statistics

→ Erdős question

→ classification theory
- 1992: Laskowski “discovers” the connection between VC learning and NIP model theory. *He uses model theory to point out new VC classes (e.g. families of sets definable in o-minimal structures).*
- 2020: Bays, Kaplan, Simon improve a result from VC theory on recursive teaching dimension to characterize NIP types as averages of k -compressible types.

The bigger picture: VC meets NIP.

- 1968 – 1972: Vapnik-Chervonenkis, Shelah-Perles and Sauer all prove independently Sauer's Lemma. 

statistics

Erdős question

classification theory
- 1992: Laskowski “discovers” the connection between VC learning and NIP model theory. *He uses model theory to point out new VC classes (e.g. families of sets definable in o-minimal structures).*
- 2020: Bays, Kaplan, Simon improve a result from VC theory on recursive teaching dimension to characterize NIP types as averages of k -compressible types.

We see fruitful outcomes (every 20+ years) from the communication between VC machine learning and NIP model theory.