

# NIP compressible types

Pablo Andújar Guerrero

University of Leeds

[anguempa.github.io/handout.pdf](https://anguempa.github.io/handout.pdf)

**Theorem:** For any NIP formula  $\varphi(x, y)$  there exists  $n, k < \omega$  such any  $\varphi$ -type  $p(x)$  is a rounded average of  $n$  many  $k$ -compressible  $\varphi$ -types.

**Theorem:** For any NIP formula  $\varphi(x, y)$  there exists  $n, k < \omega$  such any  $\varphi$ -type  $p(x)$  is a rounded average of  $n$  many  $k$ -compressible  $\varphi$ -types.


## Counting Boolean atoms

A *Boolean atom* of a finite family of sets  $\mathcal{F}$  is a maximal non-empty Boolean combination of sets in  $\mathcal{F}$ .

## Counting Boolean atoms

A *Boolean atom* of a finite family of sets  $\mathcal{F}$  is a maximal non-empty Boolean combination of sets in  $\mathcal{F}$ .


$$\mathcal{F} = \{S_1, S_2, S_3\}$$



## Counting Boolean atoms

A *Boolean atom* of a finite family of sets  $\mathcal{F}$  is a maximal non-empty Boolean combination of sets in  $\mathcal{F}$ .


$$S_1 \setminus (S_2 \cup S_3)$$



## Counting Boolean atoms

A *Boolean atom* of a finite family of sets  $\mathcal{F}$  is a maximal non-empty Boolean combination of sets in  $\mathcal{F}$ .

$$S_3 \setminus (S_1 \cup S_2)$$



## Counting Boolean atoms

A *Boolean atom* of a finite family of sets  $\mathcal{F}$  is a maximal non-empty Boolean combination of sets in  $\mathcal{F}$ .

(In model-theoretic terminology a Boolean atom can be called a type.)

Let  $BA(\mathcal{F})$  denote the number of Boolean atoms of  $\mathcal{F}$ .




## Counting Boolean atoms

A *Boolean atom* of a finite family of sets  $\mathcal{F}$  is a maximal non-empty Boolean combination of sets in  $\mathcal{F}$ .

Let  $BA(\mathcal{F})$  denote the number of Boolean atoms of  $\mathcal{F}$ .

$$BA(\mathcal{F}) = 5$$




## Counting Boolean atoms

A *Boolean atom* of a finite family of sets  $\mathcal{F}$  is a maximal non-empty Boolean combination of sets in  $\mathcal{F}$ .

Let  $BA(\mathcal{F})$  denote the number of Boolean atoms of  $\mathcal{F}$ .

$$BA(\mathcal{F}) = 2^3 = 8$$



# Counting Boolean atoms

A *Boolean atom* of a finite family of sets  $\mathcal{F}$  is a maximal non-empty Boolean combination of sets in  $\mathcal{F}$ .

Let  $BA(\mathcal{F})$  denote the number of Boolean atoms of  $\mathcal{F}$ .

## Definition

Let  $\mathcal{S}$  be a family of sets. The *dual shatter function* of  $\mathcal{S}$  is the function

$$\begin{aligned}\pi_{\mathcal{S}}^* : \omega &\rightarrow \omega \\ n &\mapsto \max\{BA(\mathcal{F}) : \mathcal{F} \subseteq \mathcal{S}, |\mathcal{F}| = n\}\end{aligned}$$

## Counting Boolean atoms

Observe that  $\pi_{\mathcal{S}}^*(n) \leq 2^n$  for all  $n$ .

# Counting Boolean atoms

Observe that  $\pi_{\mathcal{S}}^*(n) \leq 2^n$  for all  $n$ .

If  $\pi_{\mathcal{S}}^*(d) < 2^d$  for some  $d$  then  $\pi_{\mathcal{S}}^*(n) = O(n^{d-1})$  (Sauer's Lemma).

## Counting Boolean atoms

Observe that  $\pi_{\mathcal{S}}^*(n) \leq 2^n$  for all  $n$ .

If  $\pi_{\mathcal{S}}^*(d) < 2^d$  for some  $d$  then  $\pi_{\mathcal{S}}^*(n) = O(n^{d-1})$  (Sauer's Lemma).

In this case we call  $\mathcal{S}$  a *VC class* (VC=Vapnik–Chervonenkis).

# Counting Boolean atoms

Observe that  $\pi_{\mathcal{S}}^*(n) \leq 2^n$  for all  $n$ .

If  $\pi_{\mathcal{S}}^*(d) < 2^d$  for some  $d$  then  $\pi_{\mathcal{S}}^*(n) = O(n^{d-1})$  (Sauer's Lemma).

In this case we call  $\mathcal{S}$  a *VC class* (VC=Vapnik–Chervonenkis).

## Definition

The VC-codensity of a VC class  $\mathcal{S}$  is

$$\text{vc}^*(\mathcal{S}) = \inf\{r \in \mathbb{R}^{\geq 0} : \pi_{\mathcal{S}}^*(n) = O(n^r)\}.$$

# Counting Boolean atoms

Observe that  $\pi_{\mathcal{S}}^*(n) \leq 2^n$  for all  $n$ .

If  $\pi_{\mathcal{S}}^*(d) < 2^d$  for some  $d$  then  $\pi_{\mathcal{S}}^*(n) = O(n^{d-1})$  (Sauer's Lemma).

In this case we call  $\mathcal{S}$  a *VC class* (VC=Vapnik–Chervonenkis).

## Definition

The VC-codensity of a VC class  $\mathcal{S}$  is

$$\text{vc}^*(\mathcal{S}) = \inf\{r \in \mathbb{R}^{\geq 0} : \pi_{\mathcal{S}}^*(n) = O(n^r)\}.$$

E.g. If  $\mathcal{S} = \{\text{intervals in } \mathbb{R}\}$  then  $\pi_{\mathcal{S}}^*(n) = 2n$  and  $\text{vc}^*(\mathcal{S}) = 1$ .



## Definition

A family of sets  $\mathcal{S}$  is *(downward) directed* if, for any **finite** subfamily  $\mathcal{F} \subseteq \mathcal{S}$ , there exists some  $S \in \mathcal{S}$  such that  $S \subseteq \cap \mathcal{F}$ .


Equivalently,  $\mathcal{S}$  is directed if, for any  $S_1, S_2 \in \mathcal{S}$  there exists  $S \in \mathcal{S}$  with  $S \subseteq S_1 \cap S_2$ .

E.g.

- In a poset  $(P, \leq)$  the sets  $\{a \in P : a \leq b\}$  for  $b \in P$ .
- The neighborhoods of a given point in a topological spaces.
- Any filter basis.


## Definition

A family of sets  $\mathcal{S}$  has a finite transversal  $T = \{a_1, \dots, a_n\}$  if every  $S \in \mathcal{S}$  satisfies that  $S \cap T \neq \emptyset$ .



## Definition


A family of sets  $\mathcal{S}$  has a finite transversal  $T = \{a_1, \dots, a_n\}$  if every  $S \in \mathcal{S}$  satisfies that  $S \cap T \neq \emptyset$ .



## Definition

A family of sets  $\mathcal{S}$  has a finite transversal  $T = \{a_1, \dots, a_n\}$  if every  $S \in \mathcal{S}$  satisfies that  $S \cap T \neq \emptyset$ .


We care about the property that  $\mathcal{S}$  does **not** have a finite transversal.



## Definition

A family of sets  $\mathcal{S}$  has a finite transversal  $T = \{a_1, \dots, a_n\}$  if every  $S \in \mathcal{S}$  satisfies that  $S \cap T \neq \emptyset$ .

We care about the property that  $\mathcal{S}$  does **not** have a finite transversal.



Is there a connection between directedness and the dual shatter function?


**Is there a connection between directedness and the dual shatter function?**

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

$\mathcal{S}$  could be directed.






## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

$\mathcal{S}$  could be directed.




## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

$\mathcal{S}$  could be directed.


$$\pi_{\mathcal{S}}^*(3) \geq 5$$



## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that  $\mathcal{S}$  is **not** directed, witnessed by  $S_1$  and  $S_2$ .




## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that  $\mathcal{S}$  is **not** directed, witnessed by  $S_1$  and  $S_2$ .

$$\pi_{\mathcal{S}}^*(3) \geq 8$$




## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that  $\mathcal{S}$  is **not** directed, witnessed by  $S_1$  and  $S_2$ .

$$\pi_{\mathcal{S}}^*(3) \geq 6$$




## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that  $\mathcal{S}$  is **not** directed, witnessed by  $S_1$  and  $S_2$ .

$$\pi_{\mathcal{S}}^*(3) \geq 6$$




## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that  $\mathcal{S}$  is **not** directed, witnessed by  $S_1$  and  $S_2$ .

$$\pi_{\mathcal{S}}^*(3) \geq 8$$




## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that  $\mathcal{S}$  is **not** directed, witnessed by  $S_1$  and  $S_2$ .

$$BA(S_1, S_2) \geq 3$$






## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that  $\mathcal{S}$  is **not** directed, witnessed by  $S_1$  and  $S_2$ .

$$BA(S_1, S_2) \geq 3$$




## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that  $\mathcal{S}$  is **not** directed, witnessed by  $S_1$  and  $S_2$ .

$$BA(S_1, S_2, S_3) \geq 3 + 2 = 5$$




## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that  $\mathcal{S}$  is **not** directed, witnessed by  $S_1$  and  $S_2$ .

$$BA(S_1, S_2, S_3) \geq 3 + 1 + \textcolor{red}{1}.$$




## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that  $\mathcal{S}$  is **not** directed, witnessed by  $S_1$  and  $S_2$ .

$$F = S_1 \cap S_2 \cap S_3$$



## Is there a connection between directedness and the dual shatter function?

Fix a family of sets  $\mathcal{S}$  that is consistent (=has the finite intersection property). Suppose it does **not** have a finite transversal.

Now suppose that  $\mathcal{S}$  is **not** directed, witnessed by  $S_1$  and  $S_2$ .

$$F = S_1 \cap S_2 \cap S_3$$



Is there a connection between directedness and the dual shatter function?

Consider the family of sets  $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ .


$$F = S_1 \cap S_2 \cap S_3$$



**Is there a connection between directedness and the dual shatter function?**

Consider the family of sets  $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ .


$F \cap \mathcal{S}$  could be directed.



**Is there a connection between directedness and the dual shatter function?**

Consider the family of sets  $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ .

$F \cap \mathcal{S}$  could be directed.






**Is there a connection between directedness and the dual shatter function?**

Consider the family of sets  $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ .

$F \cap \mathcal{S}$  could be directed.




Is there a connection between directedness and the dual shatter function?

Consider the family of sets  $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ .

Suppose that  $F \cap \mathcal{S}$  is **not** directed, witnessed by  $S_4$  and  $S_5$ .

$$BA(S_1, S_2, S_3, S_4, S_5) \geq BA(S_1, S_2, S_3) + 3$$




Is there a connection between directedness and the dual shatter function?

Consider the family of sets  $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ .

Suppose that  $F \cap \mathcal{S}$  is **not** directed, witnessed by  $S_4$  and  $S_5$ .

$$BA(S_1, S_2, S_3, S_4, S_5) \geq BA(S_1, S_2, S_3) + 3$$




Is there a connection between directedness and the dual shatter function?

Consider the family of sets  $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ .

Suppose that  $F \cap \mathcal{S}$  is **not** directed, witnessed by  $S_4$  and  $S_5$ .

$$BA(S_1, S_2, S_3, S_4, S_5) \geq BA(S_1, S_2, S_3) + 3$$




Is there a connection between directedness and the dual shatter function?

Consider the family of sets  $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ .

Suppose that  $F \cap \mathcal{S}$  is **not** directed, witnessed by  $S_4$  and  $S_5$ .

$$BA(S_1, \dots, S_6) \geq BA(S_1, S_2, S_3) + 3 + 2.$$




Is there a connection between directedness and the dual shatter function?

Consider the family of sets  $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$ .

Suppose that  $F \cap \mathcal{S}$  is **not** directed, witnessed by  $S_4$  and  $S_5$ .

$$BA(S_1, \dots, S_6) \geq BA(S_1, S_2, S_3) + 3 + 2 + 1.$$



## Is there a connection between directedness and the dual shatter function?

We continue this process, and one of two things can happen:

- We find a finite intersection  $F$  of sets in  $\mathcal{S}$  such that  $F \cap \mathcal{S}$  is directed.
- There is a sequence  $(S_i)_i$  of sets in  $\mathcal{S}$  such that

$$BA(B_i : i \leq 3n) \geq \sum_{j=1}^n j.$$

Meaning that

$$\pi_{\mathcal{S}}^*(3n) \geq \frac{n^2 + n}{2}.$$

## Lemma A (A.G. '23)

Let  $\mathcal{S}$  be a family of sets with  $\text{vc}^*(\mathcal{S}) < 2$ . Then at least one holds:

- (a) There exists a finite intersection  $F$  of sets in  $\mathcal{S}$  such that  $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$  is directed.
- (b)  $\mathcal{S}$  has a finite transversal.



## Lemma A (A.G. '23)

Let  $\mathcal{S}$  be a family of sets with  $\text{vc}^*(\mathcal{S}) < 2$ . Then at least one holds:

- (a) There exists a finite intersection  $F$  of sets in  $\mathcal{S}$  such that  $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$  is directed.
- (b)  $\mathcal{S}$  has a finite transversal.

In fact we can improve (b) in the above lemma.

### Lemma A (A.G. '23)

Let  $\mathcal{S}$  be a family of sets with  $\text{vc}^*(\mathcal{S}) < 2$ . Then at least one holds:

- (a) There exists a finite intersection  $F$  of sets in  $\mathcal{S}$  such that  $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$  is directed.
- (b)  $\mathcal{S}$  has a finite transversal.

In fact we can improve (b) in the above lemma.

### Better Lemma A (A.G. '23)

Let  $\mathcal{S}$  be a family of sets with  $\text{vc}^*(\mathcal{S}) < 2$ . Then at least one holds:

- (a) There exists a finite intersection  $F$  of sets in  $\mathcal{S}$  such that  $F \cap \mathcal{S} = \{F \cap S : S \in \mathcal{S}\}$  is directed.
- (b) There exists  $F_1, \dots, F_m$ , finite non-empty Boolean combinations of sets in  $\mathcal{S}$  such that, for every  $S \in \mathcal{S}$  there is some  $i \leq m$  such that  $F_i \subseteq S$ .

Now let us talk compressibility.

Our two main references are:

- Martin Bays, Itay Kaplan and Pierre Simon. *Density of compressible types and some consequences*.  
ArXiv: 2107.05197, 2022.
- Xi Chen, Yu Cheng, and Bo Tang. *On the recursive teaching dimension of VC classes*.  
Advances in Neural Information Processing Systems 29:  
Annual Conference on Neural Information Processing  
Systems, 2016, Barcelona, Spain, p 2164–2171.

## Model-theoretic framework

Let  $T$  be a theory and  $\varphi(x, y)$  be a (partitioned) formula. For any  $M \models T$  and  $b \in M^y$  let  $\varphi(M, b) = \{a \in M^x : M \models \varphi(a, b)\}$ . We identify  $\varphi(x, y)$  with the family of sets  $\{\varphi(M, b) : b \in M^y\}$ :

- $\varphi(x, y)$  may be directed.
- It has dual shatter function  $\pi_\varphi^*$ .
- It has VC-codensity ...

(All these are independent of  $M$ .)

## Model-theoretic framework

Let  $T$  be a theory and  $\varphi(x, y)$  be a (partitioned) formula. For any  $M \models T$  and  $b \in M^y$  let  $\varphi(M, b) = \{a \in M^x : M \models \varphi(a, b)\}$ . We identify  $\varphi(x, y)$  with the family of sets  $\{\varphi(M, b) : b \in M^y\}$ :

- $\varphi(x, y)$  may be directed.
- It has dual shatter function  $\pi_\varphi^*$ .
- It has VC-codensity ...

(All these are independent of  $M$ .)

A formula  $\varphi$  is *NIP* if it is a VC class.

A theory  $T$  is *NIP* if every formula  $\varphi$  is NIP.

## Model-theoretic framework

Let  $T$  be a theory and  $\varphi(x, y)$  be a (partitioned) formula. For any  $M \models T$  and  $b \in M^y$  let  $\varphi(M, b) = \{a \in M^x : M \models \varphi(a, b)\}$ . We identify  $\varphi(x, y)$  with the family of sets  $\{\varphi(M, b) : b \in M^y\}$ :

- $\varphi(x, y)$  may be directed.
- It has dual shatter function  $\pi_\varphi^*$ .
- It has VC-codensity ...

(All these are independent of  $M$ .)

A formula  $\varphi$  is *NIP* if it is a VC class.

A theory  $T$  is *NIP* if every formula  $\varphi$  is NIP.

A theory  $T$  is of *VC-density one* if every formula  $\varphi(x, y)$  with  $|x| = 1$  has VC-codensity less than 2.

$$\text{VC-density one} \subseteq \text{dp-minimal} \subsetneq \text{NIP}$$

## Model-theoretic framework

Let  $T$  be a theory and  $\varphi(x, y)$  be a (partitioned) formula. For any  $M \models T$  and  $b \in M^y$  let  $\varphi(M, b) = \{a \in M^x : M \models \varphi(a, b)\}$ . We identify  $\varphi(x, y)$  with the family of sets  $\{\varphi(M, b) : b \in M^y\}$ :

- $\varphi(x, y)$  may be directed.
- It has dual shatter function  $\pi_\varphi^*$ .
- It has VC-codensity ...

(All these are independent of  $M$ .)

A formula  $\varphi$  is *NIP* if it is a VC class.

A theory  $T$  is *NIP* if every formula  $\varphi$  is NIP.

A theory  $T$  is of *VC-density one* if every formula  $\varphi(x, y)$  with  $|x| = 1$  has VC-codensity less than 2.

Conjecture

VC-density one = dp-minimal  $\subsetneq$  NIP

**Theorem** (Baldwin-Saxl '76). *Let  $G$  be a group definable in an NIP structure. Let  $\mathcal{H}$  be a definable family of subgroups of  $G$ . Then there is  $k < \omega$  such that, for any finite subfamily  $\mathcal{F} \subseteq \mathcal{H}$ , there is a subfamily  $\mathcal{C} \subseteq \mathcal{F}$  with  $|\mathcal{C}| \leq k$  such that  $\cap \mathcal{C} = \cap \mathcal{F}$ .*



## Definition

Let  $k < \omega$ . A family of sets  $\mathcal{S}$  is *k-compressible* if, for any **finite** subfamily  $\mathcal{F} \subseteq \mathcal{S}$ , there exists a subfamily  $\mathcal{G} \subseteq \mathcal{S}$  of size **at most**  $k$  such that  $\cap \mathcal{G} \subseteq \cap \mathcal{F}$ .

Equivalently,  $\mathcal{S}$  is *k-compressible* if, for any subfamily  $\mathcal{F} \subseteq \mathcal{S}$  of size  $k + 1$ , there exists a subfamily  $\mathcal{G} \subseteq \mathcal{S}$  of size **at most**  $k$  such that  $\cap \mathcal{G} \subseteq \cap \mathcal{F}$ .

## Definition

Let  $k < \omega$ . A family of sets  $\mathcal{S}$  is *k-compressible* if, for any **finite** subfamily  $\mathcal{F} \subseteq \mathcal{S}$ , there exists a subfamily  $\mathcal{G} \subseteq \mathcal{S}$  of size **at most**  $k$  such that  $\cap \mathcal{G} \subseteq \cap \mathcal{F}$ .

Equivalently,  $\mathcal{S}$  is *k-compressible* if, for any subfamily  $\mathcal{F} \subseteq \mathcal{S}$  of size  $k + 1$ , there exists a subfamily  $\mathcal{G} \subseteq \mathcal{S}$  of size **at most**  $k$  such that  $\cap \mathcal{G} \subseteq \cap \mathcal{F}$ .


**Note:** Directed = 1-compressible.

“This terminology is inspired by the terminology around compression schemes in the statistical learning literature.”


— Bays-Kaplan-Simon

- The set of real intervals in  $0^+$  is 1-compressible.
- The set of unbounded real intervals in  $0^+$  is 2-compressible.
- In  $\mathbb{R}^k$ , the set of “stripes” of the form  $\{(x_1, \dots, x_k) : a < x_i < b\}$  for  $a \leq 0 < b$  and  $i \leq n$ , is  $k$ -compressible.


- The set of real intervals in  $0^+$  is 1-compressible.
- The set of unbounded real intervals in  $0^+$  is 2-compressible.
- In  $\mathbb{R}^k$ , the set of “stripes” of the form  $\{(x_1, \dots, x_k) : a < x_i < b\}$  for  $a \leq 0 < b$  and  $i \leq n$ , is  $k$ -compressible.




- The set of real intervals in  $0^+$  is 1-compressible.
- The set of unbounded real intervals in  $0^+$  is 2-compressible.
- In  $\mathbb{R}^k$ , the set of “stripes” of the form  $\{(x_1, \dots, x_k) : a < x_i < b\}$  for  $a \leq 0 < b$  and  $i \leq n$ , is  $k$ -compressible.



- The set of real intervals in  $0^+$  is 1-compressible.
- The set of unbounded real intervals in  $0^+$  is 2-compressible.
- In  $\mathbb{R}^k$ , the set of “stripes” of the form  $\{(x_1, \dots, x_k) : a < x_i < b\}$  for  $a \leq 0 < b$  and  $i \leq n$ , is  $k$ -compressible.



- The set of real intervals in  $0^+$  is 1-compressible.
- The set of unbounded real intervals in  $0^+$  is 2-compressible.
- In  $\mathbb{R}^k$ , the set of “stripes” of the form  $\{(x_1, \dots, x_k) : a < x_i < b\}$  for  $a \leq 0 < b$  and  $i \leq n$ , is  $k$ -compressible.



- The set of real intervals in  $0^+$  is 1-compressible.
- The set of unbounded real intervals in  $0^+$  is 2-compressible.
- In  $\mathbb{R}^k$ , the set of “stripes” of the form  $\{(x_1, \dots, x_k) : a < x_i < b\}$  for  $a \leq 0 < b$  and  $i \leq n$ , is  $k$ -compressible.

A  $k$ -compressible family of sets is consistent iff it is  $k$ -consistent (every  $k$  sets have non-empty intersection).

So for  $k$ -compressible **definable** families of sets consistency is expressible with a single first order sentence.



Let  $T$  be a theory and  $M \models T$ .

For a formula  $\varphi(x, y)$  let  $\varphi^1(x, y) = \varphi(x, y)$  and  $\varphi^0(x, y) = \neg\varphi(x, y)$ .

Let  $T$  be a theory and  $M \models T$ .

For a formula  $\varphi(x, y)$  let  $\varphi^1(x, y) = \varphi(x, y)$  and  $\varphi^0(x, y) = \neg\varphi(x, y)$ .

By a  $\varphi$ -type  $p(x)$  over  $B \subseteq M^y$  we mean a maximal consistent collection of formulas of the form  $\varphi^i(x, b)$ , for  $b \in B$  and  $i \in \{0, 1\}$ .

Let  $T$  be a theory and  $M \models T$ .

For a formula  $\varphi(x, y)$  let  $\varphi^1(x, y) = \varphi(x, y)$  and  $\varphi^0(x, y) = \neg\varphi(x, y)$ .

By a  $\varphi$ -type  $p(x)$  over  $B \subseteq M^y$  we mean a maximal consistent collection of formulas of the form  $\varphi^i(x, b)$ , for  $b \in B$  and  $i \in \{0, 1\}$ .

We identify types  $p(x)$  with the family of sets defined in  $M$  by the formulas in it.

Let  $T$  be a theory and  $M \models T$ .

For a formula  $\varphi(x, y)$  let  $\varphi^1(x, y) = \varphi(x, y)$  and  $\varphi^0(x, y) = \neg\varphi(x, y)$ .

By a  $\varphi$ -type  $p(x)$  over  $B \subseteq M^y$  we mean a maximal consistent collection of formulas of the form  $\varphi^i(x, b)$ , for  $b \in B$  and  $i \in \{0, 1\}$ .

We identify types  $p(x)$  with the family of sets defined in  $M$  by the formulas in it.

### Definition

A  $\varphi$ -type  $p(x)$  is  $k$ -compressible if for any **finite** type  $q(x) \subseteq p(x)$  there exists  $r(x) \subseteq p(x)$  with  $|r(x)| \leq k$  such that

$$r(x) \vdash q(x) \quad (\text{i.e. } r(M) \subseteq q(M)).$$

## Better Lemma A (A.G. '23)

Let  $\varphi(x, y)$  be a formula in a theory  $T$  and  $M \models T$ . Let  $B \subseteq M^y$  and  $p(x)$  be a  $\varphi$ -type over  $B$  with  $\text{vc}^*(p(x)) < 2$ . Then at least one holds:

- (a) There exists a finite conjunction  $\psi(x)$  of formulas in  $p(x)$  such that  $\psi(x) \wedge p(x) = \{\psi(x) \wedge \sigma(x) : \sigma(x) \in p(x)\}$  is directed (i.e. 1-compressible).

In particular  $p(x)$  is  $k$ -compressible for some  $k < \omega$ .

- (b) There exist  $q_1(x), \dots, q_m(x)$  finite  $\varphi$ -types such that, for every  $\sigma(x) \in p(x)$  there exists some  $i \leq m$  such that  $q_i(x) \vdash \sigma(x)$  (i.e.  $q_i(M) \subseteq \sigma(M)$ ).

## Better Lemma A\* (A.G. and Johnson '23)

Let  $\varphi(x, y)$  be a formula with  $|x| = 1$  in a dp-minimal theory  $T$  and  $M \models T$ . Let  $B \subseteq M^y$  and  $p(x)$  be a  $\varphi$ -type over  $B$  with  $\text{ve}^*(p(x)) < 2$ . Then at least one holds:

- (a) There exists a finite conjunction  $\psi(x)$  of formulas in  $p(x)$  such that  $\psi(x) \wedge p(x) = \{\psi(x) \wedge \sigma(x) : \sigma(x) \in p(x)\}$  is directed (i.e. 1-compressible).

In particular  $p(x)$  is  $k$ -compressible for some  $k < \omega$ .

- (b) There exist  $q_1(x), \dots, q_m(x)$  finite  $\varphi$ -types such that, for every  $\sigma(x) \in p(x)$  there exists some  $i \leq m$  such that  $q_i(x) \vdash \sigma(x)$  (i.e.  $q_i(M) \subseteq \sigma(M)$ ).

## Better Lemma A\* (A.G. and Johnson '23)

Let  $\varphi(x, y)$  be a formula with  $|x| = 1$  in a dp-minimal theory  $T$  and  $M \models T$ . Let  $B \subseteq M^y$  and  $p(x)$  be a  $\varphi$ -type over  $B$  with  $\text{ve}^*(p(x)) \leq 2$ . Then at least one holds:

- (a) There exists a finite conjunction  $\psi(x)$  of formulas in  $p(x)$  such that  $\psi(x) \wedge p(x) = \{\psi(x) \wedge \sigma(x) : \sigma(x) \in p(x)\}$  is directed (i.e. 1-compressible).

In particular  $p(x)$  is  $k$ -compressible for some  $k < \omega$ .

- (b) There exist  $q_1(x), \dots, q_m(x)$  finite  $\varphi$ -types such that, for every  $\sigma(x) \in p(x)$  there exists some  $i \leq m$  such that  $q_i(x) \vdash \sigma(x)$  (i.e.  $q_i(M) \subseteq \sigma(M)$ ).

So the type  $p(x)$  is either  $k$ -compressible (some  $k = k(p)$ ) or it can be partitioned into finitely many types each of which extends to a finitely axiomatizable (in particular  $k$ -compressible) type.

Can we say something similar to this for **all** NIP formulas?



Can we say something similar to this for **all** NIP formulas?

Yes (thanks to Bays, Kaplan and Simon).

## Theorem B (Bays-Kaplan-Simon '22)

Let  $\varphi(x, y)$  be an NIP formula in a theory  $T$  and  $M \models T$ .

There exists  $k, n < \omega$  such that, for every  $B \subseteq M^y$  and  $\varphi$ -type  $p(x)$  over  $B$ , there exist  $n$  many  $k$ -compressible  $\varphi$ -types over  $B$ ,

$$q_1(x), \dots, q_n(x),$$

such that any formula  $\varphi(x, b)$ ,  $b \in B$ , belongs in  $p(x)$  if and only if it belongs in more than  $n/2$  of the types  $q_i(x)$ :

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > n/2, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

## Theorem B (Bays-Kaplan-Simon '22)

Let  $\varphi(x, y)$  be an NIP formula in a theory  $T$  and  $M \models T$ .

There exists  $k, n < \omega$  such that, for every  $B \subseteq M^y$  and  $\varphi$ -type  $p(x)$  over  $B$ , there exist  $n$  many  $k$ -compressible  $\varphi$ -types over  $B$ ,

$$q_1(x), \dots, q_n(x),$$

such that any formula  $\varphi(x, b)$ ,  $b \in B$ , belongs in  $p(x)$  if and only if it belongs in more than  $n/2$  of the types  $q_i(x)$ :

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > n/2, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

The type  $p(x)$  is the *rounded average* of the  $k$ -compressible types  $q_1(x), \dots, q_n(x)$ .

## Theorem B (Bays-Kaplan-Simon '22)

Let  $\varphi(x, y)$  be an NIP formula in a theory  $T$  and  $M \models T$ .

There exists  $k, n < \omega$  such that, for every  $B \subseteq M^y$  and  $\varphi$ -type  $p(x)$  over  $B$ , there exist  $n$  many  $k$ -compressible  $\varphi$ -types over  $B$ ,

$$q_1(x), \dots, q_n(x),$$

such that any formula  $\varphi(x, b)$ ,  $b \in B$ , belongs in  $p(x)$  if and only if it belongs in more than  $n/2$  of the types  $q_i(x)$ :

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > n/2, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

The type  $p(x)$  is the *rounded average* of the  $k$ -compressible types  $q_1(x), \dots, q_n(x)$ .

The  $n$  and  $k$  depend only on the dual VC-dimension of  $\varphi$  (the maximum  $d$  such that  $\pi_\varphi^*(d) = 2^d$ ).

## Theorem B (Bays-Kaplan-Simon '22)

Let  $\varphi(x, y)$  be an NIP formula in a theory  $T$  and  $M \models T$ . Fix  $1/2 \leq \alpha < 1$ .

There exists  $k, n < \omega$  such that, for every  $B \subseteq M^y$  and  $\varphi$ -type  $p(x)$  over  $B$ , there exist  $n$  many  $k$ -compressible  $\varphi$ -types over  $B$ ,

$$q_1(x), \dots, q_n(x),$$

such that

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > \alpha n, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

The  $n/2$  in the theorem can be improved to  $\alpha n$  for any fixed  $1/2 \leq \alpha < 1$ .

## Theorem B (Bays-Kaplan-Simon '22)

Let  $\varphi(x, y)$  be an NIP formula in a theory  $T$  and  $M \models T$ . Fix  $1/2 \leq \alpha < 1$ .

There exists  $k, n < \omega$  such that, for every  $B \subseteq M^y$  and  $\varphi$ -type  $p(x)$  over  $B$ , there exist  $n$  many  $k$ -compressible  $\varphi$ -types over  $B$ ,

$$q_1(x), \dots, q_n(x),$$

such that

$$p(x) = \{\varphi^i(x, b) : b \in B, i \in \{0, 1\}, \exists I \subseteq \{1, \dots, n\}, \\ |I| > \alpha n, \varphi^i(x, b) \in q_i(x) \forall i \in I\}.$$

The  $n/2$  in the theorem can be improved to  $\alpha n$  for any fixed  $1/2 \leq \alpha < 1$ .

Theorem B was used to prove existence of uniform honest definitions for NIP formulas.

## Two considerations around Theorem B

- Does VC-codensity play a role in refining Theorem B?
- If the type  $p(x)$  is definable, can the  $k$ -compressible types  $q_1(x), \dots, q_n(x)$  be chosen definable too?

## Does VC-codensity play a role in refining Theorem B?

Theorem B (Bays-Kaplan-Simon '22)

Let  $\varphi(x, y)$  be an NIP formula in a theory  $T$  and  $M \models T$ . Then there exists  $k, n < \omega$  such that every  $\varphi$ -type over any  $B \subseteq M^y$  is the rounded average of  $n$  many  $k$ -compressible  $\varphi$ -types over  $B$ .



# Does VC-codensity play a role in refining Theorem B?

## Theorem B (Bays-Kaplan-Simon '22)

Let  $\varphi(x, y)$  be an NIP formula in a theory  $T$  and  $M \models T$ . Then there exists  $k, n < \omega$  such that every  $\varphi$ -type over any  $B \subseteq M^y$  is the rounded average of  $n$  many  $k$ -compressible  $\varphi$ -types over  $B$ .

## Proposition C (A.G. '23) - *Uniform Better Lemma A*

Let  $\varphi(x, y)$  be a formula with  $\mathbf{vc}^*(\varphi) < 2$  in a theory  $T$ . Let  $M \models T$ . Then there exist  $k, m < \omega$  such, for every  $\varphi$ -type  $p(x)$  over any  $B \subseteq M^y$ , at least one holds:

- (a)  $p(x)$  is  $k$ -compressible.
- (b) There are  $m$  many  $\varphi$ -types  $q_1(x), \dots, q_n(x)$ , with  $\mathbf{vc}^*(q_i) = 0$  for every  $i \leq m$ , such that every formula in  $p(x)$  is implied by **at least one** of the  $q_i(x)$ .

# Does VC-codensity play a role in refining Theorem B?

## Theorem B (Bays-Kaplan-Simon '22)

Let  $\varphi(x, y)$  be an NIP formula in a theory  $T$  and  $M \models T$ . Then there exists  $k, n < \omega$  such that every  $\varphi$ -type over any  $B \subseteq M^y$  is the rounded average of  $n$  many  $k$ -compressible  $\varphi$ -types over  $B$ .

## Proposition C (A.G. '23) - *Uniform Better Lemma A*

Let  $\varphi(x, y)$  be a formula with  $\mathbf{vc}^*(\varphi) < 2$  in a theory  $T$ . Let  $M \models T$ . Then there exist  $k, m < \omega$  such, for every  $\varphi$ -type  $p(x)$  over any  $B \subseteq M^y$ , at least one holds:

- (a)  $p(x)$  is  $k$ -compressible.
- (b) There are  $m$  many  $\varphi$ -types  $q_1(x), \dots, q_n(x)$ , with  $\mathbf{vc}^*(q_i) = 0$  for every  $i \leq m$ , such that every formula in  $p(x)$  is implied by **at least one** of the  $q_i(x)$ .

Does Proposition C have a generalization to  $\mathbf{vc}^*(\varphi) < n$ ?

If the type  $p(x)$  is definable, can the  $k$ -compressible types  $q_i(x)$  be chosen definable too?

If the type  $p(x)$  is definable, can the  $k$ -compressible types  $q_i(x)$  be chosen definable too?

Why do we care?

Because we are interested in further understanding the density of  $k$ -compressible types.

If the type  $p(x)$  is definable, can the  $k$ -compressible types  $q_i(x)$  be chosen definable too?

Why do we care?

Because we are interested in further understanding the density of  $k$ -compressible types.

#### Corollary D (of Theorem B)

Let  $M$  be a distal structure,  $\varphi(x, y)$  a formula, and  $p(x)$  a  $\varphi$ -type. Then there exists another formula  $\psi(x, z)$  and a directed (i.e. 1-compressible)  $\psi$ -type  $q(x)$  such that

$$q(x) \vdash p(x).$$

( $q(x)$  can be chosen over the same parameters as  $p(x)$ )

This corollary is a strong “uniform” density result for  $k$ -compressible types.

If the type  $p(x)$  is definable, can the  $k$ -compressible types  $q_i(x)$  be chosen definable too?

### Question E

In Corollary D, if the  $\varphi$ -type  $p(x)$  extends to a definable type over  $M$  (in the full language), can the directed  $\psi$ -type  $q(x)$  satisfying  $q(x) \vdash p(x)$  be chosen definable?

If the type  $p(x)$  is definable, can the  $k$ -compressible types  $q_i(x)$  be chosen definable too?

### Question E

In Corollary D, if the  $\varphi$ -type  $p(x)$  extends to a definable type over  $M$  (in the full language), can the directed  $\psi$ -type  $q(x)$  satisfying  $q(x) \vdash p(x)$  be chosen definable?

A positive answer to our question about Theorem B, would yield a positive answer to Question E.

If the type  $p(x)$  is definable, can the  $k$ -compressible types  $q_i(x)$  be chosen definable too?



This one.

### Question E

In Corollary D, if the  $\varphi$ -type  $p(x)$  extends to a definable type over  $M$  (in the full language), can the directed  $\psi$ -type  $q(x)$  satisfying  $q(x) \vdash p(x)$  be chosen definable?

A positive answer to our question about Theorem B, would yield a positive answer to Question E.



If the type  $p(x)$  is definable, can the  $k$ -compressible types  $q_i(x)$  be chosen definable too?



This one.

### Question E

In Corollary D, if the  $\varphi$ -type  $p(x)$  extends to a definable type over  $M$  (in the full language), can the directed  $\psi$ -type  $q(x)$  satisfying  $q(x) \vdash p(x)$  be chosen definable?

A positive answer to our question about Theorem B, would yield a positive answer to Question E.

Using Proposition C (uniform Better Lemma A) we can show that Question E has a positive answer in distal theories of VC-density one.

If the type  $p(x)$  is definable, can the  $k$ -compressible types  $q_i(x)$  be chosen definable too?

Theorem (A.G. '23)

Let  $M$  be a distal structure of VC-density one,  $\varphi(x, y)$  be a formula, and  $p(x)$  a  $\varphi$ -type. If  $p(x)$  extends to a **definable** type in  $S_x(M)$ , then there exists another formula  $\psi(x, z)$  and a directed **definable**  $\psi$ -type  $q(x)$  such that

$$q(x) \vdash p(x).$$

(dp-minimal version to come)

If the type  $p(x)$  is definable, can the  $k$ -compressible types  $q_i(x)$  be chosen definable too?

↑

This one.

But what about our question?

If the type  $p(x)$  is definable, can the  $k$ -compressible types  $q_i(x)$  be chosen definable too?

Itay Kaplan <[REDACTED]>



Para: Pablo Andujar Guerrero

Mar 07/03/2023 18:57

I talked about it with Martin and Pierre.

It seems that there might be a counterexample but maybe it's not so easy to verify.

The idea is to take a meet-tree with a height function. So let's take  $M$  to be  $(\omega^{\omega}, \omega, h)$  where  $h$  goes from  $w^{<w}$  to  $w$  and gives the level. The language has the tree order, the  $\omega$  order,  $h$  (and the meet). Now if you take a point  $s$  in the tree, you can look at the type saying  $x \geq s$  and  $x$  is incomparable to any other point  $\geq s$  (the generic type of a "closed ball") and  $h(x) > M$  (height = infinity).

Then this type is the average of compressible types by taking 3 different branches  $\geq s$ , say  $B_1, B_2, B_3$ , and looking at the types saying  $x \geq B_i, h(x) > M$ , but these branches won't be definable.

In order to verify that this actually works, one would need to understand what are definable types, etc.

Itay

In the 2-sorted structure  $(\omega^{<\omega}, \omega; <, <_\omega, h)$ , with  $h : \omega^{<\omega} \rightarrow \omega$  the level function, consider the formulas

$$\varphi_1(x, y_1) := "(x = y_1) \vee (x < y_1) \vee (x > y_1)" \text{ for } y_1 \in \omega^{<\omega},$$

$$\varphi_2(x, y_2) := "y_2 <_\omega h(x)" \text{ for } y_2 \in \omega.$$

In the 2-sorted structure  $(\omega^{<\omega}, \omega; <, <_\omega, h)$ , with  $h : \omega^{<\omega} \rightarrow \omega$  the level function, consider the formulas

$$\begin{aligned}\varphi_1(x, y_1) &:= "(x = y_1) \vee (x < y_1) \vee (x > y_1)" \text{ for } y_1 \in \omega^{<\omega}, \\ \varphi_2(x, y_2) &:= "y_2 <_\omega h(x)" \text{ for } y_2 \in \omega.\end{aligned}$$

Consider the  $\{\varphi_1, \varphi_2\}$ -type  $p(x)$  given by

$$\begin{aligned}\varphi_1(x, ()), \\ \neg\varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\}, \\ b_2 <_\omega h(x) \text{ for every } b_2 \in \omega.\end{aligned}$$

The type  $p(x)$  is clearly definable.

In the 2-sorted structure  $(\omega^{<\omega}, \omega; <, <_\omega, h)$ , with  $h : \omega^{<\omega} \rightarrow \omega$  the level function, consider the formulas

$$\begin{aligned}\varphi_1(x, y_1) &:= "(x = y_1) \vee (x < y_1) \vee (x > y_1)" \text{ for } y_1 \in \omega^{<\omega}, \\ \varphi_2(x, y_2) &:= "y_2 <_\omega h(x)" \text{ for } y_2 \in \omega.\end{aligned}$$

Consider the  $\{\varphi_1, \varphi_2\}$ -type  $p(x)$  given by

$$\begin{aligned}\varphi_1(x, ()), \\ \neg\varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\}, \\ b_2 <_\omega h(x) \text{ for every } b_2 \in \omega.\end{aligned}$$

The type  $p(x)$  is clearly definable.

(I believe that we can apply Theorem B to  $p(x)$ .)

Consider the  $\{\varphi_1, \varphi_2\}$ -type  $p(x)$  given by

$$\begin{aligned} &\varphi_1(x, ()), \\ &\neg\varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\}, \\ &b_2 <_\omega h(x) \text{ for every } b_2 \in \omega. \end{aligned}$$

The type  $p(x)$  is clearly definable.

(I believe that we can apply Theorem B to  $p(x)$ .)



Consider the  $\{\varphi_1, \varphi_2\}$ -type  $p(x)$  given by

$$\begin{aligned} &\varphi_1(x, ()), \\ &\neg\varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\}, \\ &b_2 <_\omega h(x) \text{ for every } b_2 \in \omega. \end{aligned}$$

The type  $p(x)$  is clearly definable.

(I believe that we can apply Theorem B to  $p(x)$ .)

Pick 3 disjoint branches  $B_1, B_2, B_3$ . For each  $i \in \{1, 2, 3\}$  let  $q_i(x)$  be the  $\{\varphi_1, \varphi_2\}$ -type:

$$\begin{aligned} &\varphi_1(x, b_1) \text{ iff } b_1 \in B_i, \\ &b_2 <_\omega h(x) \text{ for every } b_2 \in \omega. \end{aligned}$$

Consider the  $\{\varphi_1, \varphi_2\}$ -type  $p(x)$  given by

$$\begin{aligned} &\varphi_1(x, ()), \\ &\neg\varphi_1(x, b_1) \text{ for every } b_1 \in \omega^{<\omega} \setminus \{()\}, \\ &b_2 <_\omega h(x) \text{ for every } b_2 \in \omega. \end{aligned}$$

The type  $p(x)$  is clearly definable.

(I believe that we can apply Theorem B to  $p(x)$ .)

Pick 3 disjoint branches  $B_1, B_2, B_3$ . For each  $i \in \{1, 2, 3\}$  let  $q_i(x)$  be the  $\{\varphi_1, \varphi_2\}$ -type:

$$\begin{aligned} &\varphi_1(x, b_1) \text{ iff } b_1 \in B_i, \\ &b_2 <_\omega h(x) \text{ for every } b_2 \in \omega. \end{aligned}$$

The type  $p(x)$  is the rounded average of the types  $q_1(x), q_2(x), q_3(x)$ . Moreover these are 2-compressible. However they are *probably* not definable.

## Email (Bays-Kaplan-Simon)

The  $\{\varphi_1, \varphi_2\}$ -type  $p(x)$  is definable but might not be the rounded average of finitely many  $k$ -compressible (for any  $k$ ) **definable**  $\{\varphi_1, \varphi_2\}$ -types.

In particular  $p(x)$  is the rounded average of the 2-compressible types  $q_1(x)$ ,  $q_2(x)$  and  $q_3(x)$ , but these are probably not definable.

# The big picture: what do we know about NIP types?

1. Few types over finite sets.
2. Indiscernible sequences converge to types.
3. Equivalence of invariance, non-dividing and non-forking over models.
4. Existence of uniform honest definitions.
5. Described by **compressible types** (Theorem B).

# The big picture: what do we know about NIP types?

1. Few types over finite sets.
2. Indiscernible sequences converge to types.
3. Equivalence of invariance, non-dividing and non-forking over models.
4. Existence of uniform honest definitions.
5. Described by **compressible types** (Theorem B).

**Opinion:** VC-(co)density is conspicuously absent.


## The bigger picture: VC meets NIP.

- 1968 – 1972: Vapnik-Chervonenkis, Shelah-Perles and Sauer all prove independently Sauer's Lemma.

## The bigger picture: VC meets NIP.

- 1968 – 1972: Vapnik-Chervonenkis, Shelah-Perles and Sauer all prove independently Sauer's Lemma.
  - statistics
  - Erdős question
  - classification theory

# The bigger picture: VC meets NIP.

- 1968 – 1972: Vapnik-Chervonenkis, Shelah-Perles and Sauer all prove independently Sauer's Lemma. 


→ statistics

→ Erdős question

→ classification theory
- 1992: Laskowski “discovers” the connection between VC learning and NIP model theory. *He uses model theory to point out new VC classes (e.g. families of sets definable in o-minimal structures).*



# The bigger picture: VC meets NIP.


- 1968 – 1972: Vapnik-Chervonenkis, Shelah-Perles and Sauer all prove independently Sauer's Lemma. 

→ statistics

→ Erdős question

→ classification theory
- 1992: Laskowski “discovers” the connection between VC learning and NIP model theory. *He uses model theory to point out new VC classes (e.g. families of sets definable in o-minimal structures).*
- 2020: Bays, Kaplan, Simon improve a result from VC theory on recursive teaching dimension to characterize NIP types as averages of  $k$ -compressible types.

# The bigger picture: VC meets NIP.

- 1968 – 1972: Vapnik-Chervonenkis, Shelah-Perles and Sauer all prove independently Sauer's Lemma. 

statistics

Erdős question

classification theory
- 1992: Laskowski “discovers” the connection between VC learning and NIP model theory. *He uses model theory to point out new VC classes (e.g. families of sets definable in o-minimal structures).*
- 2020: Bays, Kaplan, Simon improve a result from VC theory on recursive teaching dimension to characterize NIP types as averages of  $k$ -compressible types.

We see fruitful outcomes (every 20+ years) from the communication between VC machine learning and NIP model theory.