

My beautiful dissertation

My Name

November 8, 2021

*Thesis submitted for the degree of
Doctor of Philosophy*

in

Applied Mathematics

at The University of Adelaide

Faculty of Engineering, Computer and Mathematical Sciences

School of Mathematical Sciences



THE UNIVERSITY
of ADELAIDE

Contents

Signed Statement	ix
Acknowledgements	xi
Dedication	xiii
Abstract	xv
1 Introduction	1
1.1 Fluid queues	2
1.2 Matrix-exponential distributions	4
1.3 QBD-RAPs	6
2 A stochastic modelling approach to approximating fluid queues	9
2.1 Inspiration and motivation	10
2.2 Time to exit an interval	12
2.2.1 Modelling the residual time to exit on no change of phase	12
2.2.2 On a change of phase from $i \in \mathcal{S}_+$ to $j \in \mathcal{S}_+$	15
2.2.3 On a change of phase from $i \in \mathcal{S}_+$ to $j \in \mathcal{S}_-$	17
2.2.4 Upon exiting $\mathcal{D}_{k,i}$	22
2.3 The association of $j \in \mathcal{S}_0$ with \mathcal{S}_+ or \mathcal{S}_-	22
2.4 The dynamics of the QBD-RAP approximation	24
2.5 Boundary conditions	29
2.6 Initial conditions	31
2.7 At time t – closing operators	32
3 Weak convergence of the QBD-RAP scheme	37
3.1 Characterisations on no change of level	39
3.1.1 Characterising the fluid queue	39
3.1.2 Characterising the QBD-RAP	42
3.2 Laplace transforms with respect to time on no change of level	43
3.3 Convergence on no change of level	50

3.3.1	Convergence at the first change of level	61
3.4	At the n th change of level	63
3.5	Between the n th and $n + 1$ th change of level	68
3.6	To global convergence	70
3.6.1	First, another domination condition	70
3.6.2	Global convergence	74
3.7	Extension to arbitrary (but fixed) discretisation structures	78
A	Technical results for convergence	81
A.1	Some bounds for integrating against matrix exponential distributions. . . .	81
A.1.1	One integral	81
A.1.2	Many integrals.	93
A.2	Properties of closing operators	108
A.3	Kronecker properties	112
B	second appendix	119
	Bibliography	121

List of Tables

List of Figures

2.1	The density function for a <i>concentrated matrix exponential of order 21</i> from (Horváth et al. 2020) (blue) and corresponding density functions of the residual lives, $R_{0.3}$ (red), $R_{0.6}$ (blue). Observe how the density function of the Z_i (blue) to approximate a point mass at $\Delta = 1$, while the density functions of $R_{0.3}$ (red) and $R_{0.6}$ (blue) approximate point masses at 0.7 and 0.4, respectively.	14
3.1	Sample paths corresponding to the summands in (3.10).	41
3.2	Sample paths corresponding to the Laplace transforms (3.29). $z_1 = \Delta - x_1 - (x - y_{\ell_0})$, $z_2 = \Delta - x_2 - x_1$	45

Signed Statement

I certify that this work contains no material which has been accepted for the award of any other degree or diploma in my name in any university or other tertiary institution and, to the best of my knowledge and belief, contains no material previously published or written by another person, except where due reference has been made in the text. In addition, I certify that no part of this work will, in the future, be used in a submission in my name for any other degree or diploma in any university or other tertiary institution without the prior approval of the University of Adelaide and where applicable, any partner institution responsible for the joint award of this degree.

I give consent to this copy of my thesis, when deposited in the University Library, being made available for loan and photocopying, subject to the provisions of the Copyright Act 1968.

I also give permission for the digital version of my thesis to be made available on the web, via the University's digital research repository, the Library Search and also through web search engines, unless permission has been granted by the University to restrict access for a period of time.

Signed: Date:

Acknowledgements

Dedication

Abstract

Chapter 1 introduces

Chapter 1

Introduction

A fluid queue is a two-dimensional stochastic process $\{(X(t), \varphi(t))\}_{t \geq 0}$. The phase process, also known as the driving process, $\{\varphi(t)\}_{t \geq 0}$, is a continuous-time Markov chain (CTMC). The level process, $\{X(t)\}_{t \geq 0}$, is a real-valued, continuous, and piecewise linear.

Stochastic fluid queues have found a variety of applications such as telecommunications (see Anick et al. (1982) as a canonical application in this area), power systems (Bean et al. 2010), risk processes (Badescu et al. 2005) and environmental modelling (Wurm 2020). Fluid queues are relatively well studied. Largely, the analysis of fluid queues falls into two categories, matrix-analytic methods (Ahn et al. 2003, Ahn & Ramaswami 2003, 2004, Bean et al. 2005a,b, 2009a,b, da Silva Soares 2005, Latouche & Nguyen 2019), and differential equation-based methods (Anick et al. 1982, Karandikar & Kulkarni 1995, Bean et al. 2021).

More recently, Bean & O'Reilly (2014) extended fluid queues to *so-called* stochastic fluid-fluid queues. In a fluid-fluid queue there is a second level process, $\{Y(t)\}_{t \geq 0}$ which is itself driven by a fluid queue, $\{(X(t), \varphi(t))\}_{t \geq 0}$. The analysis, (Bean & O'Reilly 2014), is in principal similar to the matrix-analytic methods of (Bean et al. 2005b), and derives results about the second level process $\{Y(t)\}_{t \geq 0}$ in terms of the infinitesimal generator (a differential operator) of the fluid queue, $\{(X(t), \varphi(t))\}_{t \geq 0}$. For practical computation of the results of Bean & O'Reilly (2014), a matrix-discretisation of the infinitesimal generator of the fluid queue can be used. To this end, to date, two possible discretisation have been suggested. Taking a differential equations-based approach, Bean et al. (2021) use the discontinuous Galerkin (DG) method to discretise this operator, while Bean & O'Reilly (2013) take a stochastic modelling and matrix-analytic methods approach to approximate the fluid-queue by a quasi-birth-and-death (QBD) process. Both approaches are insightful and offer different tools and perspectives with which to analyse the resulting approximations. It turns out that the latter approach is a sub-class of the former; the QBD can be viewed as the simplest DG scheme where the operator is projected onto a basis of piecewise constant functions.

In the context of approximating fluid queues, one advantage of the QBD discretisa-

tion and, equivalently, DG schemes with constant basis functions, is that they guarantee probabilities computed from the approximation are positive (Koltai 2011, Section 3.3), see also (Hesthaven & Warburton 2007) and references therein. One justification for the positivity preserving property is from the interpretation of the discretisation as a stochastic process thereby ensuring positivity. For higher order DG schemes there is no such interpretation. Moreover, higher-order DG approximation schemes may produce negative, or highly oscillatory solutions, particularly when discontinuities or steep gradients are present. Methods to navigate the problem of negative or highly oscillatory solutions have been developed, such as slope limiters, and filtering (see (Hesthaven & Warburton 2007, Section 6.5) and references therein). Slope limiting effectively alters the discretised operator in regions where oscillations are detected and lowers the order of the approximation in these regions. Filtering is a post-hoc method which looks to recover an accurate solution, given an oscillatory approximation.

1.1 Fluid queues

A fluid queue is a two-dimensional stochastic process $\{(X(t), \varphi(t))\}_{t \geq 0}$ where $\{\varphi(t)\}_{t \geq 0}$ is known as the phase or driving process, and $\{X(t)\}_{t \geq 0}$ is known as the level process or buffer. The phase process $\{\varphi(t)\}_{t \geq 0}$, is an irreducible continuous-time Markov chain (CTMC) with finite state space, which we assume to be $\mathcal{S} = \{1, 2, \dots, N\}$ without loss of generality, and infinitesimal generator $\mathbf{T} = [T_{ij}]_{i,j \in \mathcal{S}}$. We assume that \mathbf{T} is *conservative*. Associated with states $i \in \mathcal{S}$ are real-valued *rates* $c_i \in \mathbb{R}$.

Partition the state space \mathcal{S} into $\mathcal{S}_+ = \{i \in \mathcal{S} \mid c_i > 0\}$, $\mathcal{S}_- = \{i \in \mathcal{S} \mid c_i < 0\}$ and $\mathcal{S}_0 = \{i \in \mathcal{S} \mid c_i = 0\}$. We assume, without loss of generality, that the generator \mathbf{T} is partitioned into sub-matrices

$$\mathbf{T} = \begin{bmatrix} \mathbf{T}_{++} & \mathbf{T}_{+-} & \mathbf{T}_{+0} \\ \mathbf{T}_{-+} & \mathbf{T}_{--} & \mathbf{T}_{-0} \\ \mathbf{T}_{0+} & \mathbf{T}_{0-} & \mathbf{T}_{00} \end{bmatrix},$$

where $\mathbf{T}_{mn} = [T_{ij}]_{i \in \mathcal{S}_m, j \in \mathcal{S}_n}$, $m, n \in \{+, -, 0\}$. Also define the diagonal matrices

$$\mathbf{C} = \begin{bmatrix} \mathbf{C}_+ & & \\ & \mathbf{C}_- & \\ & & \mathbf{0} \end{bmatrix}, \quad \mathbf{C}_+ = \text{diag}(c_i, i \in \mathcal{S}_+), \quad \mathbf{C}_- = \text{diag}(|c_i|, i \in \mathcal{S}_-),$$

where $\text{diag}(a_i, i \in \mathcal{I})$ denotes a diagonal matrix with entries a_i down the diagonal.

When no boundary conditions are imposed, the level process is given by

$$X(t) = X(0) + \int_{s=0}^t c_{\varphi(s)} \, ds.$$

Sample paths of $\{X(t)\}$ are continuous and piecewise linear, with $\frac{d}{dt}X(t) = c_\varphi(t)$, when $X(t)$ is differentiable. Given sample paths of $\{\varphi(t)\}$, then $\{X(t)\}$ is deterministic, and in this sense, $\{\varphi(t)\}$ is the only stochastic element of the fluid queue.

Often, boundary conditions are imposed. Here, we consider a mixture of *regulated* and *reflecting* boundary conditions. Upon hitting a boundary we suppose that, with probability p_{ij} , $i, j \in \mathcal{S}$, the phase process instantaneously transitions from phase i to phase j (note that we might have $i = j$ i.e. no transition). At a lower boundary, if $j \in \mathcal{S}_0 \cup \mathcal{S}_-$, then $\frac{d}{dt}X(t) = 0$, and the phase process continues to evolve according to the sub-generator

$$\begin{bmatrix} \mathbf{T}_{--} & \mathbf{T}_{-0} \\ \mathbf{T}_{0-} & \mathbf{T}_{00} \end{bmatrix},$$

until such a time that $\varphi(t)$ transitions to a phase $k \in \mathcal{S}_+$, at which time $X(t)$ leaves the boundary. Similarly, at an upper boundary if $j \in \mathcal{S}_0 \cup \mathcal{S}_+$, then $\frac{d}{dt}X(t) = 0$ and the phase process continues to evolve according to the sub-generator

$$\begin{bmatrix} \mathbf{T}_{++} & \mathbf{T}_{+0} \\ \mathbf{T}_{0+} & \mathbf{T}_{00} \end{bmatrix},$$

until such a time that $\varphi(t)$ transitions to a phase $k \in \mathcal{S}_-$ at which time $X(t)$ leaves the boundary. Without loss of generality, we assume the lower and upper boundaries (when present) are at $x = 0$ and $x = M > 0$, respectively.

In summary, the evolution of the level can be expressed as

$$\frac{d}{dt}X(t) = \begin{cases} c_{\varphi(t)}, & \text{if } X(t) > 0, \\ \max\{0, c_{\varphi(t)}\}, & \text{if } X(t) = 0, \\ \min\{0, c_{\varphi(t)}\}, & \text{if } X(t) = M. \end{cases}$$

Let $\mathbf{f}(x, t) = [f_i(x, t)]_{i \in \mathcal{S}}$ be a row-vector function where $f_i(x, t)$ is the density of $\mathbb{P}(X(t) \leq x, \varphi(t) = i)$, assuming it exists. When a differentiable density exists, the system of partial differential equation which describes the evolution of the densities $\mathbf{f}(x, t)$ is

$$\frac{\partial}{\partial t}\mathbf{f}(x, t) = \mathbf{f}(x, t)\mathbf{T} - \frac{\partial}{\partial x}\mathbf{f}(x, t)\mathbf{C}. \quad (1.1)$$

The initial condition is the initial distribution of the fluid queue, $f_i(x, 0)$. Often a differentiable density function does not exist and therefore the partial differential equation (1.1) is not well-defined. For example, for a fluid queue with a regulated boundary, if the initial distribution of the fluid queue is a point mass at any point $x_0 \geq 0$ and in phase $i \in \mathcal{S}_+ \cup \mathcal{S}_0$, then a density function $f_i(x, t)$ will not exist for any finite t . Specifically,

a point mass will persist along the ray $x_0 + c_i t$, $t \geq 0$. In such situations, it is the *weak solution* to (1.1) that we seek. A weak solution satisfies

$$-\int_x \int_t \mathbf{f}(x, t) \frac{\partial}{\partial t} \boldsymbol{\psi}(x, t) dt dx = \int_x \int_t \mathbf{f}(x, t) \mathbf{T} \boldsymbol{\psi}(x, t) dt dx + \int_x \int_t \mathbf{f}(x, t) \mathbf{C} \frac{\partial}{\partial x} \boldsymbol{\psi}(x, t) dt dx, \quad (1.2)$$

for every vector of test functions, $\boldsymbol{\psi}(x, t)$, which are smooth and have compact support.

Boundary conditions may also be imposed on (1.1).

Discretisation methods approximate the operator on the right-hand side of (1.1) by a matrix, in our case, by the generator of a QBD-RAP.

1.2 Matrix-exponential distributions

Here we recount some facts about matrix exponential distributions. See (Bladt & Nielsen 2017) for a more detailed exposition. A random variable, Z , is said to have a matrix-exponential distribution if it has a distribution function of the form $1 - \boldsymbol{\alpha} e^{\mathbf{S}x} (-\mathbf{S})^{-1} \mathbf{s}$, where $\boldsymbol{\alpha}$ is a $1 \times p$ initial vector, \mathbf{S} a $p \times p$ matrix, and \mathbf{s} a $p \times 1$ closing vector, and $e^{\mathbf{S}x} := \sum_{n=0}^{\infty} \frac{(\mathbf{S}x)^n}{n!}$ is the matrix exponential. The density function of Z is given by $f_Z(x) = \boldsymbol{\alpha} e^{\mathbf{S}x} \mathbf{s}$. The only restrictions on the parameters $(\boldsymbol{\alpha}, \mathbf{S}, \mathbf{s})$ are that $\boldsymbol{\alpha} e^{\mathbf{S}x} \mathbf{s}$ be a valid density function, i.e. $\boldsymbol{\alpha} e^{\mathbf{S}x} \mathbf{s} \geq 0$, for all $x \geq 0$ and $\lim_{x \rightarrow \infty} 1 - \boldsymbol{\alpha} e^{\mathbf{S}x} (-\mathbf{S})^{-1} \mathbf{s} = 1$. There is the possibility of an *atom* (a point mass) at 0, but here we do not consider this possibility. These conditions that $\boldsymbol{\alpha} e^{\mathbf{S}x} \mathbf{s}$ be a valid density imposes some properties on representations $(\boldsymbol{\alpha}, \mathbf{S}, \mathbf{s})$. However, in general there is no way to determine whether, given a triplet $(\boldsymbol{\alpha}, \mathbf{S}, \mathbf{s})$ whether it is a representation of a matrix-exponential distribution, or not. Nonetheless, some properties of a triplet $(\boldsymbol{\alpha}, \mathbf{S}, \mathbf{s})$ are known, such as the following, which is used in the characterisation of QBD-RAPs.

Theorem 1.1 (Theorem 4.1.3, Bladt & Nielsen (2017)). *The density function of a matrix-exponential distribution with representation $(\boldsymbol{\alpha}, \mathbf{S}, \mathbf{s})$ can be expressed in terms of real-valued constants as*

$$\begin{aligned} \psi(x) = & \sum_{j=1}^{m_1} \sum_{k=1}^{p_j} c_{jk} \frac{x^{k-1}}{(k-1)!} e^{\mu_j x} + \sum_{j=1}^{m+2} \sum_{k=1}^{q_j} d_{jk} \frac{x^{k-1}}{(k-1)!} e^{\eta_j x} \cos(\sigma_j x) \\ & + \sum_{j=1}^{m_2} \sum_{k=1}^{q_j} e_{jk} \frac{x^{k-1}}{(k-1)!} e^{\eta_j x} \sin(\sigma_j x), \end{aligned} \quad (1.3)$$

for integers m_1, m_2, p_j , and q_j and some real constants $c_{jk}, d_{jk}, e_{jk}, \mu_j, \eta_j$, and σ_j . Here $\mu_j, j = 1, \dots, m_1$ are the real eigenvalues of \mathbf{S} , while $\eta_j + i\sigma_j, \eta_j - i\sigma_j, j = 1, \dots, m_2$ denote

its complex eigenvalues, which come in conjugate pairs. Thus $m_1 + 2m_2$ is the total number of eigenvalues, while the dimension of the representation is given by $p = \sum_{j=1}^{m+1} p_j + 2 \sum_{j=1}^{m_2} q_j$.

Theorem 1.2 (Theorem 4.1.4, Bladt & Nielsen (2017)). *Consider the nonvanishing terms of the matrix exponential density (1.3), i.e., the terms for which $c_{jk} = 0$, $d_{jk} = 0$, or $e_{jk} = 0$. Among the corresponding eigenvalues λ_j , there is a real dominating eigenvalue κ , say. That is, κ is real, $\kappa \geq \operatorname{Re}(\lambda_j)$ for all j , and the multiplicity of κ is at least the multiplicity of every other eigenvalue with real part κ .*

Corollary 1.3 (Corollary 4.1.5, Bladt & Nielsen (2017)). *If $(\boldsymbol{\alpha}, \mathbf{S}, \mathbf{s})$ is a representation for a matrix-exponential distribution, then \mathbf{S} has a real dominating eigenvalue.*

Theorem 1.4 (Theorem 4.1.6, Bladt & Nielsen (2017)). *Let Z be a matrix-exponentially distributed random variable with density (1.3). Then the dominant real eigenvalue κ of Theorem 1.2 is strictly negative.*

We define $\operatorname{dev}(\mathbf{S})$ to be the real dominating eigenvalue of \mathbf{S} , that is $\operatorname{dev}(\mathbf{S}) = \kappa$ in Theorem 1.2.

The class of matrix-exponential distribution is characterised as the class of probability distributions which have a rational Laplace transform. That is, $\int_{x=0}^{\infty} e^{-\lambda x} \boldsymbol{\alpha} e^{\mathbf{S}x} \mathbf{s} dx$ is a ratio of two polynomial functions in λ . Matrix exponential distributions are an extension of Phase-type distributions, where for the latter, \mathbf{S} must be a sub-generator matrix of a CTMC, $\mathbf{s} = -\mathbf{S}\mathbf{e}$ where \mathbf{e} is a $1 \times p$ vector of ones, and $\boldsymbol{\alpha}$ is a discrete probability distribution.

A representation of a matrix exponential distribution is a triplet $(\boldsymbol{\alpha}, \mathbf{S}, \mathbf{s})$, and we write $Z \sim ME(\boldsymbol{\alpha}, \mathbf{S}, \mathbf{s})$ to denote that Z has a matrix-exponential distribution with this representation. The order of the representation is the dimension of the square matrix \mathbf{S} , i.e. if \mathbf{S} is $p \times p$, then the matrix exponential distribution is said to be of order p . Representations of matrix-exponential distributions are not unique (Bladt & Nielsen 2017). A representation is called *minimal* when \mathbf{S} has the smallest possible dimension. Throughout this work, we assume that the representation of any matrix exponential distribution is minimal. Let \mathbf{e}_i be a vector with a 1 in the i th position and zeros elsewhere. We assume that $\mathbf{s} = -\mathbf{S}\mathbf{e}$, and that $(\mathbf{e}_i, \mathbf{S}, \mathbf{s})$ for $i = 1, \dots, p$ are representations of matrix exponential distributions. It is always possible to find such a representation (Bladt & Nielsen 2017, Theorem 4.5.17, Corollary 4.5.18). As such, we abbreviate our notation $Z \sim ME(\boldsymbol{\alpha}, \mathbf{S}, \mathbf{s})$ to $Z \sim ME(\boldsymbol{\alpha}, \mathbf{S})$. Further, given $\mathbf{s} = -\mathbf{S}\mathbf{e}$ then observe that $\int_{x=0}^{\infty} e^{\mathbf{S}x} \mathbf{s} dx = (-\mathbf{S})^{-1} \mathbf{s} = \mathbf{e}$.

For a given $p \times p$ matrix \mathbf{S} , denote by $\mathcal{A} \subset \mathbb{R}^p$ the space of all possible vectors \mathbf{a} such that (\mathbf{a}, \mathbf{S}) is a valid representation of a matrix exponential distribution.

1.3 QBD-RAPs

As RAPs are an extension of Markovian arrival processes to include matrix-exponential inter-arrival times, QBD-RAPs are extensions of QBDs to include matrix-exponential times between level changes. To define a QBD-RAP we first need a Batch (Marked) RAP.

Let $\mathcal{K} \subset \mathbb{Z}$ be a set of *marks*. Let N be a point process, and $Y_0 = 0 < Y_1 < Y_2 \dots$ be event times of N . Let $\{N(t)\}$ be the counting process associated with N such that $N(t)$ returns the number of events by time t . Associated with the n th event is a mark M_n . For $i \in \mathcal{K}$, let N_i be simple point processes associated with events with marks of type i only, and let $\{N_i(t)\}_{t \geq 0}$ be the associated counting processes of events of mark i . For a matrix \mathbf{B} let $\text{dev}(\mathbf{B})$ denote the real part of the dominant eigenvalue of \mathbf{B} (the one with maximal real part). Denote by $f_{N,n}(y_1, m_1, y_2, m_2, \dots, y_n, m_n)$ the joint density, probability mass function of the first n inter-arrival times, $Y_1, Y_2 - Y_1, \dots, Y_n - Y_{n-1}$, and the associated marks M_n . From Bean and Nielsen (Bean & Nielsen 2010, Theorem 1) we have the following.

Theorem 1.5. *A process N is a Marked RAP if there exist matrices $\mathbf{S}, \mathbf{D}_i, i \in \mathcal{K}$, and a row vector $\boldsymbol{\alpha}$ such that $\text{dev}(\mathbf{S}) < 0$, $\text{dev}(\mathbf{S} + \mathbf{D}) = 0$, $(\mathbf{S} + \mathbf{D})\mathbf{e} = 0$, $\mathbf{D} = \sum_{i \in \mathcal{K}} \mathbf{D}_i$, and*

$$f_{N,n}(y_1, m_1, y_2, m_2, \dots, y_n, m_n) = \boldsymbol{\alpha} e^{\mathbf{S}y_1} \mathbf{D}_{m_1} e^{\mathbf{S}y_2} \mathbf{D}_{m_2} \dots e^{\mathbf{S}y_n} \mathbf{D}_{m_n} \mathbf{e}. \quad (1.4)$$

Conversely, if a point process has the property (1.4) then it is a Marked RAP.

Denote such a process $N \sim \text{BRAP}(\boldsymbol{\alpha}, \mathbf{S}, \mathbf{D}_i, i \in \mathcal{K})$.

Also from Bean & Nielsen (2010), associated with a Marked RAP is a row-vector-valued *orbit* process, $\{\mathbf{A}(t)\}_{t \geq 0}$,

$$\mathbf{A}(t) = \frac{\boldsymbol{\alpha} \left(\prod_{i=1}^{N(t)} e^{\mathbf{S}(Y_i - Y_{i-1})} \mathbf{D}_{M_i} \right) e^{\mathbf{S}(t - Y_{N(t)})}}{\boldsymbol{\alpha} \left(\prod_{i=1}^{N(t)} e^{\mathbf{S}(Y_i - Y_{i-1})} \mathbf{D}_{M_i} \right) e^{\mathbf{S}(t - Y_{N(t)})} \mathbf{e}}.$$

Thus, $\{\mathbf{A}(t)\}$ is a piecewise-deterministic Markov process where, in between events $\{\mathbf{A}(t)\}$ evolves deterministically according to

$$\mathbf{A}(t) = \frac{\mathbf{A}(Y_{N(t)}^-) e^{\mathbf{S}(t - Y_{N(t)})}}{\mathbf{A}(Y_{N(t)}^-) e^{\mathbf{S}(t - Y_{N(t)})} \mathbf{e}},$$

where $\mathbf{A}(Y_{N(t)}^-) = \lim_{u \rightarrow 0^+} \mathbf{A}(Y_{N(t)} - u)$. The process $\{\mathbf{A}(t)\}$ "jumps" at event times of N (the process may not always jump at these times, but typically the dynamics

change discontinuously at this point). At time t the intensity with which $\{\mathbf{A}(t)\}$ has a jump is $\mathbf{A}(t)\mathbf{D}\mathbf{e}$. Upon an event the event is associated with mark i with probability $\mathbf{A}(t)\mathbf{D}_i\mathbf{e}/\mathbf{A}(t)\mathbf{D}\mathbf{e}$. Upon an event at time t with mark i , the new position of the orbit is $\mathbf{A}(t) = \mathbf{A}(t^-)\mathbf{D}_i/\mathbf{A}(t^-)\mathbf{D}_i\mathbf{e}$. It is important to note that the jumps of the orbit process are *linear* transformations of the orbit process immediately before the time of the jump.

Marked RAPs are an extension of Marked Markovian arrival processes to include matrix-exponential inter-arrival times. For Marked MAPs, the vector $\mathbf{A}(t)$ is a vector of posterior probabilities of a continuous-time Markov chain.

Intuitively, $\mathbf{A}(t)$ encodes all of the information about the event times of the Marked RAP and associated marks up to time t that is needed to determine the future behaviour of the point process. Let \mathcal{F}_t be the σ -algebra generated by $N(u), u \in [0, t]$. Then $N \mid \mathcal{F}_t \equiv N \mid \mathbf{A}(t) \sim BRAP(\mathbf{A}(t), \mathbf{S}, \mathbf{D}_i, i \in \mathcal{K})$. In words, the future of the point process after time t given all of the information about the process up to and including time t , is distributed as a Marked RAP with initial vector $\mathbf{A}(t)$.

Now consider a Marked RAP, $N \sim BRAP(\boldsymbol{\alpha}, \mathbf{S}, \mathbf{D}_i, i \in \{-1, 0, +1\})$. The process $\{(L(t), \mathbf{A}(t))\}_{t \geq 0}$ formed by letting $L(t) = N_{+1}(t) - N_{-1}(t)$ is a QBD-RAP.

Chapter 2

A stochastic modelling approach to approximating fluid queues

In this chapter we develop a new discretisation of a fluid queue using a *quasi-birth-and-death process with rational arrival process components* (QBD-RAP). In the following chapter, Chapter 3 we prove the scheme converges weakly. To improve upon the discretisation of Bean & O'Reilly (2013), we argue that the shift from QBD to QBD-RAP is necessary. The QBD construction of Bean & O'Reilly (2013) uses Erlang distributions to approximate the deterministic time that the fluid queue spends in an interval on the event that the phase is constant – it is well known that the Erlang distribution is the distribution with the smallest variance in the class of Phase-type distributions of a given order Aldous & Shepp (1987), and in this sense is the best approximation to the deterministic behaviour we want to capture.

By choosing a matrix-exponential distribution with sufficiently small variance, the QBD-RAP approximation can be made arbitrarily accurate. Due to its stochastic interpretation, the QBD-RAP discretisation ensures solutions are positive, even when discontinuities are present. Unlike slope limiters and post-hoc filtering, for the QBD-RAP the positivity preserving nature is a property of the discretised operator.

The structure of this chapter is as follows. Section 2.1 motivates the idea using the QBD approximation of Bean & O'Reilly (2014). Sections 2.2 and 2.3 describe the modelling of certain events of the fluid queue with matrix exponential distributions to ultimately construct the behaviour of the QBD-RAP on the event that the QBD-RAP remains in a given level. Section 2.4 describes the dynamics of the construction and also constructs the level process of the QBD-RAP. Section 2.5 deals with modelling boundary behaviour. Section 2.6 describes how to model initial conditions of the fluid queue. Section 2.7 introduces the concept of a *closing operator* which maps the state space of the QBD-RAP to estimates of the density of the fluid queue.

2.1 Inspiration and motivation

The inspiration for our approach stems from the QBD-based discretisation of Bean & O'Reilly (2013). A QBD can be seen as a two dimensional CTMC, $\{(L(t), \phi(t))\}_{t \geq 0}$, where $\{L(t)\}$ is the discrete level, and $\{\phi(t)\}$ is the phase. Bean & O'Reilly (2013) discretise the state space of the fluid into small intervals of width Δ ; denote these by $\mathcal{D}_k = [k\Delta, (k+1)\Delta]$. Their approximation captures the dynamics of the phase process exactly, but discretises the level process of the fluid queue. Their approximation supposes that when $L(t) = k$, $\phi(t) = i$, then $X(t) \approx k\Delta$, $\phi(t) = i$.

When in level k and phase i , the approximation sees events at rate $|T_{ii}| + |c_i|\Delta$. Upon an event, with probability $T_{ij}/(|T_{ii}| + |c_i|\Delta)$ a change of phase from i to j is observed and the approximation remains in level k ; with probability $|c_i|\Delta/(|T_{ii}| + |c_i|\Delta)$ a change of level occurs, to $k+1$ if $c_i > 0$ and $k-1$ if $c_i < 0$. The generator of their QBD approximation (for an unbounded fluid queue) is

$$B = \begin{bmatrix} \ddots & \ddots & \ddots & & \\ B_{-1}(\Delta) & B_0(\Delta) & B_{+1}(\Delta) & & \\ & B_{-1}(\Delta) & B_0(\Delta) & B_{+1}(\Delta) & \\ & \ddots & \ddots & \ddots & \ddots \end{bmatrix}$$

$$B_0 = T - C\Delta, \quad B_{-1} = \begin{bmatrix} 0 & & \\ & C_{-\Delta} & \\ & & 0 \end{bmatrix}, \quad B_{+1} = \begin{bmatrix} C_{+\Delta} & & \\ & 0 & \\ & & 0 \end{bmatrix}. \quad (2.1)$$

Bean and O'Reilly then take $\Delta \rightarrow 0$ and show that the approximation converges weakly to the fluid queue. It seems that this is the best we can do if we want to keep the interpretation of the approximation as a QBD. We now elaborate on this point.

For a given Δ , the discretisation of Bean and O'Reilly supposes that, when the phase is i and on the event of no change of phase, the sojourn time in a given level has an exponential distribution with rate $|c_i|\Delta$. However, the corresponding sojourn time of the fluid queue is deterministic: given $X(t) = x \in \mathcal{D}_k$, then $\{X(t)\}$ will leave \mathcal{D}_k in exactly $((k+1)\Delta - x)/|c_i|$ units of time if $c_i > 0$, and $(x - k\Delta)/|c_i|$ units of time if $c_i < 0$, on the event that the phase does not change before this time. We can extend the QBD model of Bean and O'Reilly to model this determinism more accurately by supposing that the sojourn times in each interval have Erlang distributions (rather than exponential). However, we effectively realise the same QBD approximation as the original, but on a finer discretisation. For example, using Erlang distributions of order 2 we would get a

generator of the QBD approximation

$$B = \begin{bmatrix} \ddots & \ddots & \ddots & \\ B_{-1}(\Delta) & B_0(\Delta) & B_{+1}(\Delta) & \\ & B_{-1}(\Delta) & B_0(\Delta) & B_{+1}(\Delta) \\ & \ddots & \ddots & \ddots \end{bmatrix}$$

$$B_0 = T \otimes I_2 - \begin{bmatrix} \begin{bmatrix} -\Delta/2 & \Delta/2 \\ 0 & -\Delta/2 \end{bmatrix} \otimes C_+ & \begin{bmatrix} -\Delta/2 & 0 \\ \Delta/2 & -\Delta/2 \end{bmatrix} \otimes C_- \\ & \mathbf{0} \end{bmatrix}, \quad (2.2)$$

$$B_{-1} = \begin{bmatrix} \mathbf{0} & & \\ & C_- \otimes \begin{bmatrix} 0 & \Delta/2 \\ 0 & 0 \end{bmatrix} & \\ & & \mathbf{0} \end{bmatrix}, \quad B_{+1} = \begin{bmatrix} C_+ \otimes \begin{bmatrix} 0 & 0 \\ \Delta/2 & 0 \end{bmatrix} & & \\ & \mathbf{0} & \\ & & \mathbf{0} \end{bmatrix}. \quad (2.3)$$

But this is the same discretisation we would get if we were to use intervals of width $\Delta/2$ in the QBD approximation of Bean & O'Reilly (2013), with the rows/columns reordered. Indeed, this appears to be the best we can do with a QBD approximation as the Erlang distribution is known to be the least variable Phase-type distribution for a given order (Aldous & Shepp 1987) and therefore gives the closest approximation to the deterministic behaviour we are trying to approximate. This necessitates the extension to QBD-RAPs, whereby we can find more concentrated distributions than any Phase-type and retain enough stochastic interpretation and matrix-analytic tools.

Here, the construction of the approximating QBD-RAP is developed intuitively, from a stochastic modelling perspective. The key observation is that, given $\{\varphi(t)\}$ is constant, then $\{X(t)\}$ moves deterministically. Upon discretising the state space of the level process into intervals of width Δ , then, given $X(t)$, the distribution of time it takes for $\{X(t)\}$ to leave a given interval on the event that $\{\varphi(t)\}$ remains in the same phase, is deterministic. We model this deterministic behaviour approximately by concentrated matrix exponential distributions (matrix exponential distributions with low-variance) (Élteto et al. 2006, Horváth et al. 2016, Élteto et al. 2006, Horváth et al. 2020, Mészáros & Telek 2021). We construct a level process for the QBD-RAP and correspond it to a discretisation the level process of the fluid queue. This is similar to Bean & O'Reilly (2013) where the level process of their QBD corresponds a discretisation of the fluid level. Unlike Bean & O'Reilly (2013), however, here we do not take $\Delta \rightarrow 0$. Rather, we suppose that the variance of the concentrated matrix exponential distributions we use to approximate deterministic events gets small.

It turns out that the corresponding orbit process $\{\mathbf{A}(t)\}$ can be seen to “track”, approximately, how far $X(t)$ is from the left of a given interval when the phase is in \mathcal{S}_+ , or how far from the right of the interval $X(t)$ is when the phase is in \mathcal{S}_- . This leaves us with two problems. 1) on a transition from \mathcal{S}_+ to \mathcal{S}_- or \mathcal{S}_- to \mathcal{S}_+ how must $\mathbf{A}(t)$ jump to retain this information about where $X(t)$ is within a given interval. 2) how to use the orbit position at time t , $\mathbf{A}(t)$, to obtain an approximation for the density of $X(t)$. Answers to both problems can be derived from the *residual time* of a matrix exponential distribution. Let us now formalise these concepts some more.

2.2 Time to exit an interval

Consider the partition of the state space of the level of a fluid queue, $[0, M]$ into K intervals of width $\Delta = M/(K + 1)$, specifically $\mathcal{D}_{k,i} = [k\Delta, (k + 1)\Delta)$ if $i \in \mathcal{S}_+$ and $\mathcal{D}_{k,i} = (k\Delta, (k + 1)\Delta]$ if $i \in \mathcal{S}_-$, $k = 0, \dots, K + 1$. The distinction between $\mathcal{D}_{k,i}$ for $i \in \mathcal{S}_+$ and \mathcal{S}_- is a technical one and will be discussed later. For now, one reason we might want to specify the discretisation in this way, is that it ensures the stochastic process

$$\left\{ \sum_{k \in \mathcal{K}} \sum_{i \in \mathcal{S}} k 1(X(t) \in \mathcal{D}_{k,i}) \right\}_{t \geq 0} \quad (2.4)$$

is right-continuous with left limits (cádlág). The expression (2.4) is the discrete process of the fluid-queue which the level process of the QBD-RAP will approximate. Let $y_k = k\Delta$, $k = 0, \dots, K + 1$ and $\mathcal{D}_k = [y_k, y_{k+1}]$. We now look to model the behaviour of $\{X(t)\}$ on an interval \mathcal{D}_k over the time $[t, t + u]$ given information about the phase process over this time.

2.2.1 Modelling the residual time to exit on no change of phase

We first consider no change of phase in $[t, t + u]$. Suppose that $\varphi(t) = i \in \mathcal{S}_+$ and $X(t) = x \in \mathcal{D}_{k,i}$. Observe that on no change of phase, $\varphi(t + s) = i$, $s \in [0, u]$, and at time $t + s$ the level of the fluid queue is given by $X(t + s) = x + c_i s$, $s \in [0, u]$. Further, for $u > (y_{k+1} - x)/c_i$ the level process $\{X(t)\}$ leaves the interval $\mathcal{D}_{k,i}$ at exactly time $t + (y_{k+1} - x)/c_i$. Similarly, for $i \in \mathcal{S}_-$, $\{X(t)\}$ leaves the interval $\mathcal{D}_{k,i}$ at exactly time $t + (x - y_k)/|c_i|$ on the event that there is no change of phase on $[t, t + u]$ for $u > (x - y_k)/|c_i|$.

To approximate the deterministic behaviour observed above, consider a matrix-exponential distribution $Z \sim ME(\boldsymbol{\alpha}, \mathbf{S}, \mathbf{s})$ with mean Δ and low variance. As we shall see later, the assumption that Z has low variance will allow us to claim that Z can be used to approximate deterministic behaviour. Let $Z_i \sim ME(\boldsymbol{\alpha}, \mathbf{S}_i, \mathbf{s}_i)$ where $\mathbf{S}_i = |c_i| \mathbf{S}$, $\mathbf{s}_i = -\mathbf{S}_i \mathbf{e}$, $i \in \mathcal{S}$. The random variable Z_i has mean $\Delta/|c_i|$.

Suppose first that $X(t) = y_k$ and $\varphi(t) = i \in \mathcal{S}_+$, so that $X(t)$ will leave $\mathcal{D}_{k,i}$ in exactly $\Delta/|c_i|$ units of time on the event that the phase process of the QBD-RAP remains in i for

at least this amount of time. A sensible approximation for this deterministic event is to choose the value of the orbit process at time t to be α , $\mathbf{A}(t) = \alpha$. With this choice, the distribution of time until the next event of the QBD-RAP, on the event that the phase process remains constant will be the distribution of Z_i . Observe,

$$\begin{aligned} \mathbb{P}(Z_i \in ((\Delta - \varepsilon)/|c_i|, (\Delta + \varepsilon)/|c_i|)) &= \mathbb{P}(Z \in (\Delta - \varepsilon, \Delta + \varepsilon)) \\ &\geq 1 - \frac{\text{Var}(Z)}{\varepsilon^2}, \end{aligned}$$

since, by Chebyshev's inequality, $\mathbb{P}(Z \in (\Delta - \varepsilon, \Delta + \varepsilon)) \geq 1 - \frac{\text{Var}(Z)}{\varepsilon^2}$. Choosing $\varepsilon = \text{Var}(Z)^{1/3}$, then

$$\mathbb{P}(R_i(u) \in ((\Delta - \varepsilon)/|c_i| - u, (\Delta + \varepsilon)/|c_i|)) \geq 1 - \text{Var}(Z)^{1/3} \approx 1,$$

when $\text{Var}(Z)$ is small. Hence, when the variance of Z (equivalently Z_i) is low, then with high probability the time until the next event of the QBD-RAP, given the phase process remains constant will occur at approximately $\Delta/|c_i|$.

Now, let $R_i(u)$ be the residual time, $R_i(u) = (Z_i - u)1\{Z_i - u > 0\}$, $i \in \mathcal{S}$. After $u \in [0, \Delta/|c_i|)$ amount of time has elapsed, on the event that $\varphi(t+s) = i$, for all $s \in [0, u)$, then $X(t+u) = y_k + c_i u$. If the phase remains i for a further $\Delta/|c_i| - u$ amount of time, then $\{X(t)\}$ will leave $\mathcal{D}_{k,i}$ at exactly this time. Also at time $t+u$, given $Z_i > u$, the distribution of the residual time, $R_i(u)$, has density

$$f_{R_i(u)}(r) = \frac{\alpha e^{\mathbf{S}_i(u+r)} \mathbf{s}_i}{\alpha e^{\mathbf{S}_i u} \mathbf{e}} = \mathbf{A}(t+u) e^{\mathbf{S}_i r} \mathbf{s}_i.$$

Here, the event $Z_i > u$ approximates the event $X(t+u) \in \mathcal{D}_{k,i}$ and we want the residual time, $R_i(u)$, to approximate the time until $\{X(t)\}$ leaves $\mathcal{D}_{k,i}$. That is, we want $R_i(u)$ to approximate a deterministic random variable at $\Delta/|c_i| - u$. To this end, we observe that for $\varepsilon > 0$ and $u < (\Delta - \varepsilon)/|c_i|$, then

$$\begin{aligned} \mathbb{P}(R_i(u) \in ((\Delta - \varepsilon)/|c_i| - u, (\Delta + \varepsilon)/|c_i| - u)) \\ &= \mathbb{P}(Z_i \in ((\Delta - \varepsilon)/|c_i|, (\Delta + \varepsilon)/|c_i|)) \\ &= \mathbb{P}(Z \in (\Delta - \varepsilon, \Delta + \varepsilon)) \\ &\geq 1 - \frac{\text{Var}(Z)}{\varepsilon^2}. \end{aligned}$$

Choosing $\varepsilon = \text{Var}(Z)^{1/3}$, then

$$\mathbb{P}(R_i(u) \in ((\Delta - \varepsilon)/|c_i| - u, (\Delta + \varepsilon)/|c_i|)) \geq 1 - \text{Var}(Z)^{1/3} \approx 1,$$




Figure 2.1: The density function for a *concentrated matrix exponential of order 21* from (Horváth et al. 2020) (blue) and corresponding density functions of the residual lives, $R_{0.3}$ (red), $R_{0.6}$ (blue). Observe how the density function of the Z_i (blue) to approximate a point mass at $\Delta = 1$, while the density functions of $R_{0.3}$ (red) and $R_{0.6}$ (blue) approximate point masses at 0.7 and 0.4, respectively.

when $\text{Var}(Z)$ is small. That is, when the variance of Z (equivalently Z_i) is low, the residual time $R_i(u)$ will be concentrated around $\Delta/|c_i| - u$, as required. Figure 2.1 gives an example of a density function for a Z_i with mean $\Delta = 1$ and $c_i = 1$, as well as the density function of $R_i(0.3)$ conditional on $Z_i > 0.3$ and $R_i(0.6)$ conditional on $Z_i > 0.6$, for comparison. Observe that the density of the residual life $R_i(0.3)$ conditional on $Z_i > 0.3$ is concentrated around $\Delta - 0.3 = 0.7$ and, similarly the density of the residual life $R_i(0.6)$ conditional in $Z_i > 0.6$ is concentrated around $\Delta - 0.6 = 0.4$.

For notational convenience, define the row-vector-valued function $\mathbf{k}(t) : \mathbb{R} \rightarrow \mathcal{A}$,

$$\mathbf{k}(t) = \frac{\boldsymbol{\alpha} e^{St}}{\boldsymbol{\alpha} e^{St} \mathbf{e}}, \quad (2.5)$$

and the operator $\mathbf{K}(t) : \mathcal{A} \rightarrow \mathbb{R}$, $t \geq 0$,

$$\mathbf{a}\mathbf{K}(t) = \frac{\mathbf{a}e^{\mathbf{S}t}}{\mathbf{a}e^{\mathbf{S}t}\mathbf{e}}, \quad (2.6)$$

for any $\mathbf{a} \in \mathcal{A}$.

We interpret the position of the orbit $\mathbf{A}(t+u) = \mathbf{k}(|c_i|u)$ as corresponding to the fact that $X(t+u)$ is $|c_i|u$ units from the left-hand boundary of the interval $\mathcal{D}_{k,i}$. This gives a heuristic argument as to how we can model the sojourn times in a given interval $\mathcal{D}_{k,i}$ on the event that the phase does not change.

We can apply analogous arguments to heuristically develop a model for the sojourn time of the fluid queue in an interval, $\mathcal{D}_{k,i}$, $i \in \mathcal{S}_-$, on the event that the phase does not change, given the fluid is $X(t) = y_{k+1}$. For $i \in \mathcal{S}_-$, the orbit position $\mathbf{A}(t+u) = \mathbf{k}(|c_i|u)$ is interpreted as corresponding to $X(t)$ being $|c_i|u$ units from the right-hand boundary of the interval $\mathcal{D}_{k,i}$, y_{k+1} .

2.2.2 On a change of phase from $i \in \mathcal{S}_+$ to $j \in \mathcal{S}_+$

Let E be the event that, at time t the orbit of the QBD-RAP is $\mathbf{A}(t) = \boldsymbol{\alpha}$, the phase process is $\varphi(t) = i \in \mathcal{S}_+$, and there are no events of the QBD-RAP before time $t+u$. On the event E the orbit position at time $t+u$ will be

$$\mathbf{A}(t+u) = \mathbf{k}(|c_i|u) = \frac{\boldsymbol{\alpha}e^{\mathbf{S}_i u}}{\boldsymbol{\alpha}e^{\mathbf{S}_i u}\mathbf{e}}. \quad (2.7)$$

Correspondingly, on the event that at time t the level process of the fluid queue is $X(t) = y_k$, the phase is $\varphi(t) = i$ and there are no change of phase by time $t+u$, then $X(t+u) = y_k + c_i u$. That is, $X(t+u)$ is $c_i u$ units from the left-hand edge of \mathcal{D}_k .

If, at time $t+u$ there is a change of phase from $i \in \mathcal{S}_+$ to $j \in \mathcal{S}_+$, then we need to map the orbit position $\mathbf{A}(t+u^-) = \mathbf{k}(|c_i|u)$ to an orbit position which corresponds to the being $c_i u$ units from the left-hand edge of \mathcal{D}_k in phase j . Call this mapping $\mathbf{D}(i, j)(\cdot)$. As noted in Section 1.3, the mapping must be linear, $\mathbf{D}(i, j)(\mathbf{a}) = \mathbf{a}\mathbf{D}(i, j)$, $\mathbf{a} \in \mathcal{A}$, for some matrix $\mathbf{D}(i, j)$. The matrix $\mathbf{D}(i, j)$ is a modelling choice.

So that the process is a QBD-RAP, the matrix $\mathbf{D}(i, j)$ must have the property that, for any $\mathbf{a} \in \mathcal{A}$, $(\mathbf{a}\mathbf{D}(i, j), \mathbf{S})$ is a representation of a matrix exponential distribution. We would also like the matrix $\mathbf{D}(i, j)$ to have the property $\mathbf{D}(i, j)\mathbf{e} = \mathbf{e}$ as this will mean that the rate at which a change of phase from i to j of the QBD-RAP occurs proportional to

$$\mathbf{A}(t+u)\mathbf{D}(i, j)\mathbf{e} = 1,$$

which is constant and therefore the distribution of time until a change from phase i to j is exponential. We can use this to show, for certain models, the distribution of the phase

process of the fluid queue and the distribution of the phase process of the QBD-RAP share the same distribution.

An orbit position which corresponds to the level process of the QBD-RAP beign $c_i u$ units to the right of y_k in phase $j \in \mathcal{S}_+$ is

$$\mathbf{A}(t+u) = \frac{\boldsymbol{\alpha} e^{\mathbf{S}_j(|c_i|u/|c_j|)}}{\boldsymbol{\alpha} e^{\mathbf{S}_j(|c_i|u/|c_j|)u} \mathbf{e}}.$$

To see this, consider the event that at time t the orbit process of the QBD-RAP is $\mathbf{A}(t) = \boldsymbol{\alpha}$, the phase process is $\varphi(t) = j \in \mathcal{S}_+$, and there are no events of the QBD-RAP before time $t + |c_i|u/|c_j|$. On this event, the orbit position at time $t + u$ will be

$$\mathbf{A}(t+u) = \frac{\boldsymbol{\alpha} e^{\mathbf{S}_j(|c_i|u/|c_j|)}}{\boldsymbol{\alpha} e^{\mathbf{S}_j(|c_i|u/|c_j|)u} \mathbf{e}}.$$

Correspondingly, on the event that at time t the level process of the fluid queue is $X(t) = y_k$, the phase is $\varphi(t) = j$ and there are no change of phase by time $t + |c_i|u/|c_j|$, then $X(t+u) = y_k + c_j|c_i|u/|c_j| = y_k + c_i u$. That is, $X(t+u)$ is $c_i u$ units from the left-hand edge of \mathcal{D}_k .

Now, observe

$$\frac{\boldsymbol{\alpha} e^{\mathbf{S}_j(|c_i|u/|c_j|)}}{\boldsymbol{\alpha} e^{\mathbf{S}_j(|c_i|u/|c_j|)u} \mathbf{e}} = \frac{\boldsymbol{\alpha} e^{\mathbf{S}_i u}}{\boldsymbol{\alpha} e^{\mathbf{S}_i u} \mathbf{e}} = \mathbf{k}(|c_i|u),$$

which is exactly (2.7), the orbit position on the event E . Hence, it is sensible to choose $\mathbf{D}(i, j) = \mathbf{I}$.

Moreover, the residual time $R_j(|c_i|u/|c_j|)$ has density

$$f_{R_j(|c_i|u/|c_j|)}(r) = \frac{\boldsymbol{\alpha} e^{\mathbf{S}_i u}}{\boldsymbol{\alpha} e^{\mathbf{S}_i u} \mathbf{e}} e^{\mathbf{S}_j r} \mathbf{s}_j,$$

and

$$\begin{aligned} \mathbb{P}(R_j(|c_i|u/|c_j|) \in ((\Delta - |c_i|u - \varepsilon)/|c_j|, (\Delta - |c_i|u + \varepsilon)/|c_j|)) \\ = \mathbb{P}(Z_j \in ((\Delta - \varepsilon)/|c_j|, (\Delta + \varepsilon)/|c_j|)) \end{aligned} \quad (2.8)$$

$$= \mathbb{P}(Z \in (\Delta - \varepsilon, \Delta + \varepsilon)) \quad (2.9)$$

$$\geq 1 - \text{Var}(Z)^{1/3} \approx 1. \quad (2.10)$$

Hence, when the variance of Z is low, the residual time, $R_j(|c_i|u/|c_j|)$, is concentrated around $(\Delta - |c_i|u)/|c_j|$, as required.

Analogous arguments suggest that the same applies for changes of phase from $i \in \mathcal{S}_-$ to $j \in \mathcal{S}_-$.

2.2.3 On a change of phase from $i \in \mathcal{S}_+$ to $j \in \mathcal{S}_-$

Now suppose $X(t) = y_k$, $\varphi(t) = i \in \mathcal{S}_+$, and the phase remains in state i until there is a change of phase from $i \in \mathcal{S}_+$ to $j \in \mathcal{S}_-$ at time $t + u$, $u \in [0, \Delta/|c_i|)$. As before, we need to find a matrix $\mathbf{D}(i, j)$ to map the orbit position on the event E when there is a change of phase from $i \in \mathcal{S}_+$ to $j \in \mathcal{S}_-$ at time $t + u$, from

$$\mathbf{A}(t + u^-) = \mathbf{k}(|c_i|u^-),$$

to

$$\mathbf{A}(t + u) = \frac{\mathbf{k}(u^-)\mathbf{D}(i, j)}{\mathbf{k}(u^-)\mathbf{D}(i, j)\mathbf{e}} = \frac{\boldsymbol{\alpha}e^{\mathbf{S}_i u}\mathbf{D}(i, j)}{\boldsymbol{\alpha}e^{\mathbf{S}_i u}\mathbf{D}(i, j)\mathbf{e}}$$

in such a way that the orbit position after the jump corresponds, in some way, to the fluid level being a distance of $|c_i|u$ from y_k in phase j .

Again, the matrix $\mathbf{D}(i, j)$ is a modelling choice. We first discuss how we might choose the matrix $\mathbf{D}(i, j)$ for when the matrix-exponential Z is a phase-type distribution.

The Phase-type case If $Z \sim ME(\boldsymbol{\alpha}, \mathbf{S})$ is chosen to be a phase-type distribution then Z has the interpretation as the distribution of time until absorption of a finite-state continuous-time Markov chain with transient states $\{1, \dots, p\}$ and a single absorbing state. The sub-generator matrix describing the dynamics of the Markov chain on transient states is \mathbf{S} , and $\boldsymbol{\alpha}$ is an initial probability distribution over the transient states. Let $\{J(t)\}$ be the Markov chain associated with the phase-type distribution.

In the discussions above, we have relied on the relationship between the event E and the orbit position $\mathbf{k}(|c_i|u)$. The relationship allows us to associate this orbit position with the level of the fluid queue. For phase-type distributions the vector $\mathbf{k}(|c_i|u)$ is the vector of posterior probabilities $[\mathbb{P}(J(u) = k \mid E)]_{k \in \{1, \dots, p\}}$. We can use this vector of posterior probabilities in the same way as we do for matrix-exponential distributions and QBD-RAPs. However, the orbit interpretation forgoes the presence of the phase process $\{J(t)\}$. Here, we will use the phase process $\{J(t)\}$ to derive a choice of $\mathbf{D}(i, j)$.

Recall the position of the orbit process can be used to determine the residual time. This idea can be replicated when we are given the phase of the phase-type rather than the value of the orbit process. Given the phase of the phase-type distribution is $k \in \{1, \dots, p\}$, the distribution of the residual time is

$$\mathbb{P}(R_i \leq r \mid \text{phase} = k) = 1 - \mathbf{e}_k e^{\mathbf{S}_i r} \mathbf{e}.$$

Notice that this is independent of the time since the phase-type distribution was initialised. We call the time since the phase-type distribution was initialised the *age*. As noted by Hautphenne et al. (2017), the distribution of the age given the phase of the phase-type is $k \in \{1, \dots, p\}$ depends on the sampling scheme which determines the observation time. Here, on the event that at time t the phase of the QBD-RAP is i , a change of phase occurs

after an exponential amount of time with rate $-T_{ii}$. Proposition 4.1, Hautphenne et al. (2017) states that the distribution function of the age, given the phase of the phase-type is k and the process is observed after an exponential time with rate $-T_{ii}$ is

$$\mathbb{P}(\text{age} \leq u \mid \text{phase} = k) = 1 - \frac{\alpha e^{(\mathbf{S}_i + T_{ii}\mathbf{I})u} (-\mathbf{S}_i - T_{ii}\mathbf{I})^{-1} \mathbf{e}_k}{\alpha (-\mathbf{S}_i - T_{ii}\mathbf{I})^{-1} \mathbf{e}_k}.$$

Let $\widehat{\mathbf{S}}_i(T_{ii}) = \text{diag}(\boldsymbol{\nu})^{-1} \mathbf{S}'_i \text{diag}(\boldsymbol{\nu})$, where $\boldsymbol{\nu} = \alpha (-\mathbf{S}_i - T_{ii}\mathbf{I})^{-1} / (\alpha (-\mathbf{S}_i - T_{ii}\mathbf{I})^{-1} \mathbf{e})$. Algebraic manipulations show

$$1 - \frac{\alpha e^{(\mathbf{S}_i + T_{ii}\mathbf{I})u} (-\mathbf{S}_i - T_{ii}\mathbf{I})^{-1} \mathbf{e}_k}{\alpha (-\mathbf{S}_i - T_{ii}\mathbf{I})^{-1} \mathbf{e}_k} = 1 - \mathbf{e}'_k e^{(\widehat{\mathbf{S}}_i(T_{ii}) + T_{ii}\mathbf{I})u} \mathbf{e}, \quad (2.11)$$

which is of Phase-type.

Recall that in the matrix-exponential case we associated the value of the orbit process $\mathbf{k}(|c_i|u)$ with being $|c_i|u$ units to the right of y_k . In contrast, for the phase-type case if we use the phase process $\{J(t)\}$ to inform our modelling of the fluid queue (as opposed to the value of the orbit process), then we associate phase $k \in \{1, \dots, p\}$ of $\{J(t)\}$ with being a *random* distance to the right of y_k where this distance has distribution

$$1 - \mathbf{e}'_k e^{(\widehat{\mathbf{S}}_i(T_{ii}) + T_{ii}\mathbf{I})u} \mathbf{e}. \quad (2.12)$$

Therefore, on the event E , upon a change from of phase of the QBD-RAP from $i \in \mathcal{S}_+$ to $j \in \mathcal{S}_-$ at time $t + u$, it seems reasonable to want the distribution of time until the next event of the QBD-RAP to be

$$1 - \mathbf{e}'_k e^{(\widehat{\mathbf{S}}_i(T_{ii}) + T_{ii}\mathbf{I})|c_j|u/|c_i| + T_{jj}Iu} \mathbf{e}. \quad (2.13)$$

The factor $|c_j|u/|c_i|$ arises as a conversion between the speed at which the fluid level moves in phase j compared to phase i . The rate T_{jj} in the exponent is the rate at which the next change of phase occurs.

While this does achieve what we want, it is not quite satisfactory for the purpose of our approximation scheme due to dependence on the sample path of $\{\varphi(t)\}$. Specifically, the evolution of the QBD-RAP from time $t + u$ until the next event depends on the phase immediately before the change of phase at time $t + u$, $\varphi(t + u^-) = i$. This increases the size of the approximating QBD-RAP as we need a separate model for each $\varphi(t + u^-) \in \mathcal{S}_+$. Furthermore, we have not yet considered how to model any further changes from \mathcal{S}_- to \mathcal{S}_+ or beyond, which further complicates matters.

A solution is to suppose that, rather than observing the phase-type random variable at an exponential time with rate $-T_{ii}$, we instead observe the process uniformly randomly on the lifetime of length Z_i . Let $\boldsymbol{\Pi} = \text{diag}(\boldsymbol{\pi})$, where $[\pi_k]_{k \in \{1, \dots, p\}} =: \boldsymbol{\pi} = \alpha (-\mathbf{S})^{-1} / m$ and $m = \alpha (-\mathbf{S})^{-1} \mathbf{e}$. There is a time-reverse representation of a Phase-type distribution

given by $(\tilde{\alpha}, \tilde{\mathbf{S}}, \tilde{\mathbf{s}})$, where $\tilde{\alpha} = m\mathbf{s}'\mathbf{\Pi}$, $\tilde{\mathbf{S}} = \mathbf{\Pi}^{-1}\mathbf{S}'\mathbf{\Pi}$ and $\tilde{\mathbf{s}} = \mathbf{\Pi}^{-1}\alpha'/m$ (Asmussen 2008, Page 91). In the case where the phase is observed randomly on the lifetime of Z_i , the distribution function of the age, given the phase is k is (Hautphenne et al. 2017, Lemma 3.1)

$$\mathbb{P}(\text{age} \leq u \mid \text{phase} = k) = 1 - \frac{\alpha e^{\mathbf{S}_i u} (-\mathbf{S}_i)^{-1} \mathbf{e}_k}{\alpha (-\mathbf{S}_i)^{-1} \mathbf{e}_k} = 1 - \mathbf{e}_k e^{\tilde{\mathbf{S}}_i u} \mathbf{e}. \quad (2.14)$$

With this interpretation, we may associate phase $k \in \{1, \dots, p\}$ of $\{J(t)\}$ with being a *random* distance, with distribution (2.14), to the right of y_k .

Therefore, on the event E and on a change of phase from $i \in \mathcal{S}_+$ to $j \in \mathcal{S}_-$ at time $t+u$, a reasonable model for the time until the next event of the QBD-RAP has distribution

$$1 - \mathbf{e}_k e^{\tilde{\mathbf{S}}_i u |c_j|/|c_i| + T_{jj} \mathbf{I} u} \mathbf{e} = 1 - \mathbf{e}_k e^{\tilde{\mathbf{S}}_j u + T_{jj} \mathbf{I} u} \mathbf{e}. \quad (2.15)$$

This suggests that, at a jump from \mathcal{S}_+ to \mathcal{S}_- , the state of $\{J(t)\}$ does not change, but begins to evolve according to the time-reverse generator at an appropriate speed. Since the time-reverse of $\tilde{\mathbf{S}}$ is \mathbf{S} , then upon a jump back to \mathcal{S}_+ from \mathcal{S}_- , the phase of the phase-type random variable remains k but begins to evolve according to \mathbf{S} . This suggests that we use the representation (α, \mathbf{S}) when in phases in \mathcal{S}_+ , and use the time-reverse representation $(\tilde{\alpha}, \tilde{\mathbf{S}})$ when in phases in \mathcal{S}_- . The matrices $\mathbf{D}(i, j) = \mathbf{I}$ for all $i, j \in \mathcal{S}$.

With this construction, and choosing $Z \sim \text{Erlang}(p, \Delta/p)$, we recover the discretisation of Bean & O'Reilly (2013) with discretisation parameter Δ/p .

The matrix-exponential case For matrix exponential distributions we cannot rely on the phase process $\{J(t)\}$ as we did in the phase-type case. Recall that we want to find a matrix $\mathbf{D}(i, j)$ to map the orbit position on the event E when there is a change of phase from $i \in \mathcal{S}_+$ to $j \in \mathcal{S}_-$ at time $t+u$, from $\mathbf{A}(t+u^-) = \mathbf{k}(|c_i|u)$, to

$$\mathbf{A}(t+u) = \frac{\mathbf{k}(|c_i|u) \mathbf{D}(i, j)}{\mathbf{k}(|c_i|u) \mathbf{D}(i, j) \mathbf{e}} = \frac{\alpha e^{\mathbf{S}_i u} \mathbf{D}(i, j)}{\alpha e^{\mathbf{S}_i u} \mathbf{D}(i, j) \mathbf{e}}$$

in such a way that the orbit position after the jump corresponds, in some way, to the fluid level being a distance of $|c_i|u$ from y_k in phase j .

Given our interpretation of the orbit position, $\mathbf{k}(x)$, a solution would be find a linear map which takes $\mathbf{k}(x)$ and maps it directly to

$$\mathbf{k}(\Delta - x) = \frac{\alpha e^{\mathbf{S}(\Delta - x)}}{\alpha e^{\mathbf{S}(\Delta - x)} \mathbf{e}}.$$

However, we have been unsuccessful in finding such a mapping. The main hurdle being that the map must be linear. Instead, we approximate as follows.

Recall that we use $R_i(u)$ to approximate the distance of the fluid level to the left of y_{k+1} . Suppose we are given $R_i(u) = r$ which corresponds, approximately, to the fluid level being $|c_i|r$ units to the left of y_{k+1} . The position of the orbit process which corresponds to a distance of $|c_i|r$ to the left of y_{k+1} in phase j is $\mathbf{k}(|c_i|r)$. Hence, on $R_i(u) = r$, the event E and on the event that a change of phase from $i \in \mathcal{S}_+$ to $j \in \mathcal{S}_-$ occurs at time $t + u$, the orbit position should jump from $\mathbf{k}(|c_i|u)$ to $\mathbf{k}(|c_i|r)$. However, at time $t + u$, $R_i(u)$ is a random variable about the future of the process and therefore not known, so instead, we take the expected initial vector

$$\mathbb{E}[\mathbf{k}(|c_i|R_i(u))] = \int_{r=0}^{\infty} \frac{\boldsymbol{\alpha} e^{\mathbf{S}_i u}}{\boldsymbol{\alpha} e^{\mathbf{S}_i u} \mathbf{e}} e^{\mathbf{S}_i r} \mathbf{s}_i \mathbf{k}(|c_i|r) \mathrm{d}r = \frac{\boldsymbol{\alpha} e^{\mathbf{S}_i u}}{\boldsymbol{\alpha} e^{\mathbf{S}_i u} \mathbf{e}} \int_{r=0}^{\infty} e^{\mathbf{S}_i r} \mathbf{s}_i \frac{\boldsymbol{\alpha} e^{\mathbf{S}_i r}}{\boldsymbol{\alpha} e^{\mathbf{S}_i r} \mathbf{e}} \mathrm{d}r.$$

After a change of variables $x = |c_i|r$ we get

$$\frac{\boldsymbol{\alpha} e^{\mathbf{S}_i u}}{\boldsymbol{\alpha} e^{\mathbf{S}_i u} \mathbf{e}} \int_{x=0}^{\infty} e^{\mathbf{S}^x |c_i|} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}^x}}{\boldsymbol{\alpha} e^{\mathbf{S}^x} \mathbf{e}} \mathrm{d}x / |c_i| = \mathbf{A}(t + u^-) \int_{x=0}^{\infty} e^{\mathbf{S}^x} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}^x}}{\boldsymbol{\alpha} e^{\mathbf{S}^x} \mathbf{e}} \mathrm{d}x,$$

since at time $t + u^-$, the orbit position is $\mathbf{A}(t + u^-) = \boldsymbol{\alpha} e^{\mathbf{S}_i u} / \boldsymbol{\alpha} e^{\mathbf{S}_i u} \mathbf{e}$. Thus, we find

$$\mathbf{D}(i, j) = \int_{x=0}^{\infty} e^{\mathbf{S}^x} \mathbf{s} \mathbf{k}(x) \mathrm{d}x =: \mathbf{D}.$$

Observe that $\mathbf{D} \mathbf{e} = \int_{x=0}^{\infty} e^{\mathbf{S}^x} \mathbf{s} \mathbf{k}(x) \mathrm{d}x \mathbf{e} = \int_{x=0}^{\infty} e^{\mathbf{S}^x} \mathbf{s} \mathrm{d}x = \mathbf{e}$, since $\mathbf{k}(x) \mathbf{e} = 1$ for all $x \geq 0$. Further, since \mathcal{A} is closed and convex (Bladt & Nielsen 2017), then $(\mathbf{a} \mathbf{D}, \mathbf{S}, \mathbf{s})$ is a representation of a matrix exponential distribution for any $\mathbf{a} \in \mathcal{A}$.

We pose the choice of the matrix \mathbf{D} as a modelling choice. Other choices are possible, for example, $\mathbf{D} = \int_{x=0}^{\Delta} e^{\mathbf{S}^x} \mathbf{s} \mathbf{k}(x) \mathrm{d}x$ is another sensible choice. It may also be possible to construct other matrices \mathbf{D} , perhaps via geometric arguments.

Computing \mathbf{D} In practice we use the class of concentrated matrix exponential distributions (CMEs) found numerically in (Horváth et al. 2020). For this class of CMEs, we take the index p to be the order of the representation. Moreover, we take p to be odd. The justification for considering representations of odd orders only is that the variance of CME representations of orders $2p$ and $2p - 1$ are relatively similar and therefore have similar abilities to represent the delta function (Horváth et al. 2020). Hence, if we construct a QBD-RAP approximation with a representation of order $2p$ we expect it to perform only marginally better than an approximation constructed with representations of order $2p - 1$. However, the computational cost of the latter is lower, so we opt for the order $2p - 1$ representation.

For a given CME with odd order, p , and representation $(\boldsymbol{\alpha}, \mathbf{S})$, the matrix \mathbf{S} has one real eigenvalue, and $p - 1$ complex eigenvalues and all eigenvalues have the same real part.

We numerically evaluate the matrix \mathbf{D} where

$$\mathbf{D} = \int_{t=0}^{\infty} e^{\mathbf{S}t} \mathbf{s} \cdot \mathbf{k}(t) dt$$

using a trapezoidal rule as follows. For the CMEs considered here, the vector function $\mathbf{k}(t)$ is periodic with period $\rho = 2\pi/\omega$ where $\omega = \min_i(|\Im(\lambda_i)|)$, λ_i are the eigenvalues of \mathbf{S} and $\Im(z)$ is the imaginary component of a complex number z . Let $\mathbf{f}(t) = e^{\mathbf{S}t} \mathbf{s}$. Then $\mathbf{f}(t)e^{-\lambda t}$ where $\lambda = \Re(\lambda_i)$ is the real part of the eigenvalues of \mathbf{S} (they all share the same real part), is also periodic with the same period. Hence we can simplify the integral to a finite one;

$$\begin{aligned} \mathbf{D} &= \int_{t=0}^{\infty} \mathbf{f}(t) \cdot \mathbf{k}(t) dt \\ &= \sum_{k=0}^{\infty} \int_{k\rho}^{(k+1)\rho} e^{\lambda t} e^{-\lambda t} \mathbf{f}(t) \cdot \mathbf{k}(t) dt \\ &= \sum_{k=0}^{\infty} \int_0^{\rho} e^{\lambda(k\rho+t)} e^{-\lambda(k\rho+t)} \mathbf{f}(k\rho+t) \cdot \mathbf{k}(k\rho+t) dt. \end{aligned} \quad (2.16)$$

By periodicity, then $e^{-\lambda(k\rho+t)} \mathbf{f}(k\rho+t) \cdot \mathbf{k}(k\rho+t) = e^{-\lambda t} \mathbf{f}(t) \cdot \mathbf{k}(t)$, hence (2.16) is equal to

$$\sum_{k=0}^{\infty} (e^{\lambda\rho})^k \int_0^{\rho} e^{\lambda t} e^{-\lambda t} \mathbf{f}(t) \cdot \mathbf{k}(t) dt = \frac{1}{1 - e^{\lambda\rho}} \int_0^{\rho} \mathbf{f}(t) \cdot \mathbf{k}(t) dt, \quad (2.17)$$

where the sum converges as it is a geometric series and $\lambda < 0$, $\rho > 0$.

To approximate (2.17) numerically, we first partition $[0, \rho]$ into N equal-width intervals $[t_n, t_{n+1})$, where $t_n = (n-1)\rho/N$, $n = 1, 2, \dots, N+1$. On $[t_n, t_{n+1})$ we approximate the orbit $\mathbf{k}(t)$ by a constant $\mathbf{k}(t) \approx \mathbf{k}_n := \frac{1}{2}(\mathbf{k}(t_n) + \mathbf{k}(t_{n+1}))$, $t \in [t_n, t_{n+1})$. Substituting this approximation into the expression for \mathbf{D} gives

$$\begin{aligned} \mathbf{D} &\approx \frac{1}{1 - e^{\lambda\rho}} \sum_{n=1}^N \int_{t_n}^{t_{n+1}} \mathbf{f}(t) \cdot \mathbf{k}_n dt \\ &= \frac{1}{1 - e^{\lambda\rho}} \sum_{n=1}^N [e^{\mathbf{S}t_{n+1}} - e^{\mathbf{S}t_n}] \mathbf{e} \cdot \mathbf{k}_n. \end{aligned}$$

This approximation preserves the property that $\mathbf{D}\mathbf{e} = \mathbf{e}$.

Computationally efficient expressions for $e^{\mathbf{S}t} \mathbf{e}$ and $\mathbf{k}(t)$ are provided in (Horváth et al. 2020).

2.2.4 Upon exiting $\mathcal{D}_{k,i}$

Suppose that upon exiting $\mathcal{D}_{k,i}$ at time t the phase is $\varphi(t) = i \in \mathcal{S}_+$. At this time $X(t) = y_{k+1}$ which is the left-hand endpoint of $\mathcal{D}_{k+1,i}$. Hence, we restart the model of the sojourn time with the initial condition $\mathbf{A}(t) = \boldsymbol{\alpha}$. Similarly, upon exiting $\mathcal{D}_{k,i}$ at time t in phase $\varphi(t) = i \in \mathcal{S}_-$, then $X(t) = y_k$, which is the right-hand endpoint of $\mathcal{D}_{k,i}$, and so we restart the model of the sojourn time with the initial condition $\mathbf{A}(t) = \boldsymbol{\alpha}$.

2.3 The association of $j \in \mathcal{S}_0$ with \mathcal{S}_+ or \mathcal{S}_-

Let $X(0) = y_\ell$ and consider the event where $\{\varphi(t)\}$ transitions from $j_0 \rightarrow j_1 \rightarrow j_2$ where $j_0 \in \mathcal{S}_+$, $j_1 \in \mathcal{S}_0$ and $j_2 \in \mathcal{S}_-$, before there is a change of level, i.e.

$$\varphi(t) = \begin{cases} j_0 & t \in [0, t_1), \\ j_1 & t \in [t_1, t_2), \\ j_2 & t \in [t_2, t_3), \end{cases}$$

and $X(t) \in \mathcal{D}_\ell$, $t \in [0, t_3)$. On approximating this event, the initial orbit position is $\mathbf{A}(0) = \boldsymbol{\alpha}$. The corresponding sample path of the orbit process on $t \in [0, t_3)$, is

$$\begin{aligned} \mathbf{A}(t) &= \begin{cases} \frac{\boldsymbol{\alpha} e^{(\mathbf{S}_{j_0} + T_{j_0 j_0} \mathbf{I})t}}{\boldsymbol{\alpha} e^{(\mathbf{S}_{j_0} + T_{j_0 j_0} \mathbf{I})t} \mathbf{e}} & t \in [0, t_1), \\ \frac{\boldsymbol{\alpha} e^{(\mathbf{S}_{j_0} + T_{j_0 j_0} \mathbf{I})t_1} \mathbf{D}(j_0, j_1) e^{T_{j_1 j_1}(t-t_1)}}{\boldsymbol{\alpha} e^{(\mathbf{S}_{j_0} + T_{j_0 j_0} \mathbf{I})t_1} \mathbf{D}(j_0, j_1) e^{T_{j_1 j_1}(t-t_1)} \mathbf{e}} & t \in [t_1, t_2), \\ \frac{\boldsymbol{\alpha} e^{(\mathbf{S}_{j_0} + T_{j_0 j_0} \mathbf{I})t_1} \mathbf{D}(j_0, j_1) e^{T_{j_1 j_1}(t_2-t_1)} \mathbf{D}(j_1, j_2) e^{(\mathbf{S}_{j_2} + T_{j_2 j_2} \mathbf{I})(t-t_2)}}{\boldsymbol{\alpha} e^{(\mathbf{S}_{j_0} + T_{j_0 j_0} \mathbf{I})t_1} \mathbf{D}(j_0, j_1) e^{T_{j_1 j_1}(t_2-t_1)} \mathbf{D}(j_1, j_2) e^{(\mathbf{S}_{j_2} + T_{j_2 j_2} \mathbf{I})(t-t_2)} \mathbf{e}} & t \in [t_2, t_3), \end{cases} \\ &= \begin{cases} \frac{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t}}{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t} \mathbf{e}} & t \in [0, t_1), \\ \frac{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t_1} \mathbf{D}(j_0, j_1)}{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t_1} \mathbf{D}(j_0, j_1) \mathbf{e}} & t \in [t_1, t_2), \\ \frac{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t_1} \mathbf{D}(j_0, j_1) \mathbf{D}(j_1, j_2) e^{\mathbf{S}_{j_2}(t-t_2)}}{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t_1} \mathbf{D}(j_0, j_1) \mathbf{D}(j_1, j_2) e^{\mathbf{S}_{j_2}(t-t_2)} \mathbf{e}} & t \in [t_2, t_3), \end{cases} \end{aligned}$$

for some matrices $\mathbf{D}(j_0, j_1)$ and $\mathbf{D}(j_1, j_2)$. Notice that $\{\mathbf{A}(t)\}$ is constant on $t \in [t_1, t_2)$. The matrix product $\mathbf{D}(j_0, j_1) \mathbf{D}(j_1, j_2)$ is there to capture the change in direction due to the change from \mathcal{S}_+ to \mathcal{S}_- . Hence $\mathbf{D}(j_0, j_1) \mathbf{D}(j_1, j_2)$ should be equal to \mathbf{D} . These types of sample paths are the reason we need to associate states $j_1 \in \mathcal{S}_0$ with either \mathcal{S}_+ or \mathcal{S}_- .

Associating j_1 with \mathcal{S}_+ , amounts to choosing $\mathbf{D}(j_0, j_1) = \mathbf{I}$ and $\mathbf{D}(j_1, j_2) = \mathbf{D}$; associating j_1 with \mathcal{S}_- , amounts to choosing $\mathbf{D}(j_0, j_1) = \mathbf{D}$ and $\mathbf{D}(j_1, j_2) = \mathbf{I}$.

There are some consequences to this choice. Let $k_2 \in \mathcal{S}_+$. Consider an event where the phase process of the fluid queue transitions from $j_0 \rightarrow j_1 \rightarrow k_2$ and there is no change of level. If j_1 is associated with \mathcal{S}_+ , then $\mathbf{D}(j_0, j_1) = \mathbf{D}(j_1, k_2) = \mathbf{I}$ and the corresponding orbit process, given $\mathbf{A}(0) = \boldsymbol{\alpha}$, is

$$\mathbf{A}(t) = \begin{cases} \frac{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t}}{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t} \mathbf{e}} & t \in [0, t_1), \\ \frac{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t_1}}{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t_1} \mathbf{e}} & t \in [t_1, t_2), \\ \frac{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t_1} e^{\mathbf{S}_{k_2}(t-t_2)}}{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t_1} e^{\mathbf{S}_{k_2}(t-t_2)} \mathbf{e}} & t \in [t_2, t_3). \end{cases}$$

Notice that there is no matrix \mathbf{D} in this expression.

Compare this to if j_1 is associated with \mathcal{S}_- . In this case $\mathbf{D}(j_0, j_1) = \mathbf{D}(j_1, k_2) = \mathbf{D}$ and the corresponding orbit process of the approximation, given $\mathbf{A}(0) = \boldsymbol{\alpha}$ and there are no change of level in $[t_1, t_3)$, is

$$\mathbf{A}(t) = \begin{cases} \frac{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t}}{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t} \mathbf{e}} & t \in [0, t_1), \\ \frac{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t_1} \mathbf{D}}{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t_1} \mathbf{D} \mathbf{e}} & t \in [t_1, t_2), \\ \frac{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t_1} \mathbf{D} \mathbf{D} e^{\mathbf{S}_{k_2}(t-t_2)}}{\boldsymbol{\alpha} e^{\mathbf{S}_{j_0} t_1} \mathbf{D} \mathbf{D} e^{\mathbf{S}_{k_2}(t-t_2)} \mathbf{e}} & t \in [t_2, t_3). \end{cases}$$

Ideally $\mathbf{D}^2 = \mathbf{I}$, however this is not the case here. Recall that a jump according to \mathbf{D} corresponds to approximating the residual life by an expectation. With this interpretation as an approximation, it suggests that we might want to minimise the number of jumps according to \mathbf{D} which occur. Therefore, for $j_1 \in \mathcal{S}_0$, if transitions $\mathcal{S}_+ \rightarrow j_1 \rightarrow \mathcal{S}_+$ occur with high probability compared to transition $\mathcal{S}_- \rightarrow j_1 \rightarrow \mathcal{S}_-$, then this suggests we might want to associate j_1 with \mathcal{S}_+ . Which association is chosen will depend on the parameters of the fluid queue and on which aspects of the model we wish to approximate. Although this advice is based on intuition only, numerical results in Section TBC suggest that it is reasonable.

Augmented state-space schemes

Another way to approach the problem is to augment the state space of the phase process by duplicating \mathcal{S}_0 and associating one copy of \mathcal{S}_0 with \mathcal{S}_+ and one copy of \mathcal{S}_0 with \mathcal{S}_- . Let $\{\varphi^*(t)\}$ be the augmented CTMC with state space \mathcal{S}^* and generator \mathbf{T}^* . Let \mathcal{S}_+ and \mathcal{S}_- be as before and $\mathcal{S}_{m0} = \{(m, i) \mid i \in \mathcal{S}_m\}$, $m \in \{+, -\}$, then $\mathcal{S}^* = \mathcal{S}_+ \cup \mathcal{S}_- \cup \mathcal{S}_{+0} \cup \mathcal{S}_{-0}$.

The generator of $\varphi^*(t)$ can be written as

$$\mathbf{T}^* = \begin{bmatrix} \mathbf{T}_{++} & \mathbf{T}_{+0} & \mathbf{T}_{+-} & 0 \\ \mathbf{T}_{0+} & \mathbf{T}_{00} & \mathbf{T}_{0-} & 0 \\ \mathbf{T}_{-+} & 0 & \mathbf{T}_{--} & \mathbf{T}_{-0} \\ \mathbf{T}_{0+} & 0 & \mathbf{T}_{0-} & \mathbf{T}_{00} \end{bmatrix}.$$

Also define a fluid level $\{X^*(t)\}$ using $\{\varphi^*(t)\}$, with rates $c_i^* = c_i$ for $i \in \mathcal{S}_+ \cup \mathcal{S}_-$ and $c_{(m,i)}^* = 0$ for $(m,i) \in \mathcal{S}_{+0} \cup \mathcal{S}_{-0}$. The process $\{\varphi(t)\}$ is imbedded within $\{\varphi^*(t)\}$ and is recovered by marginalising over \mathcal{S}_{+0} and \mathcal{S}_{-0} . On the event $X^*(0) = X(0)$, the fluid levels $X^*(t)$ and $X(t)$ match exactly. Hence, by approximating $\{(X^*(t), \varphi^*(t))\}$, we can recover an approximation to $\{(X(t), \varphi(t))\}$. This construction removes the problem of having to choose how to associate states $j \in \mathcal{S}_0$ with either \mathcal{S}_+ or \mathcal{S}_- .

The generator for the QBD-RAP approximation to the augmented fluid process (ignoring boundaries) is

$$\mathbf{B} = \begin{bmatrix} \ddots & \ddots & \ddots & & & \\ & \mathbf{B}_{-1} & \mathbf{B}_0 & \mathbf{B}_{+1} & & \\ & & \mathbf{B}_{-1} & \mathbf{B}_0 & \mathbf{B}_{+1} & \\ & & & \ddots & \ddots & \ddots \end{bmatrix},$$

where,

$$\mathbf{B}_0 = \begin{bmatrix} \mathbf{C}_+ \otimes \mathbf{S} + \mathbf{T}_{++} \otimes \mathbf{I} & \mathbf{T}_{+0} \otimes \mathbf{I} & \mathbf{T}_{+-} \otimes \mathbf{D} & \mathbf{0} \\ \mathbf{T}_{0+} \otimes \mathbf{I} & \mathbf{T}_{00} \otimes \mathbf{I} & \mathbf{T}_{0-} \otimes \mathbf{D} & \mathbf{0} \\ \mathbf{T}_{-+} \otimes \mathbf{D} & \mathbf{0} & \mathbf{C}_- \otimes \mathbf{S} + \mathbf{T}_{--} \otimes \mathbf{I} & \mathbf{T}_{-0} \otimes \mathbf{I} \\ \mathbf{T}_{0+} \otimes \mathbf{D} & \mathbf{0} & \mathbf{T}_{0-} \otimes \mathbf{I} & \mathbf{T}_{00} \otimes \mathbf{I} \end{bmatrix},$$

$$\mathbf{B}_{-1} = \begin{bmatrix} \mathbf{0} & & & \\ & \mathbf{0} & & \\ & & \mathbf{C}_- \otimes (s\alpha) & \\ & & & \mathbf{0} \end{bmatrix}, \quad \mathbf{B}_{+1} = \begin{bmatrix} \mathbf{C}_+ \otimes (s\alpha) & & & \\ & \mathbf{0} & & \\ & & \mathbf{0} & \\ & & & \mathbf{0} \end{bmatrix}.$$

With this construction, jumps according to the matrix \mathbf{D} occur only on transitions from $\mathcal{S}_m \rightarrow \mathcal{S}_n$, or $\mathcal{S}_m \rightarrow \mathcal{S}_{0m} \rightarrow \mathcal{S}_n$, $m, n \in \{+,-\}$, $m \neq n$.

2.4 The dynamics of the QBD-RAP approximation

We now have all the elements we need to describe the dynamics of the QBD-RAP approximation. Since the phase dynamics are a CTMC, we choose to use an alternate notation

to the standard presented in Section 1.3. Let $\{(L(t), \mathbf{A}(t), \phi(t))\}_{t \geq 0}$ be the QBD-RAP approximation of a fluid queue, where $\{L(t)\}$ is the level, $\{\mathbf{A}(t)\}$ is the orbit process and $\{\phi(t)\}$ is the phase process. We use this representation to make explicit how the approximation captures the phase dynamics of the fluid queue. We will show later that $\phi(t)$ captures the phase dynamics of the fluid queue exactly, provided the phase process is independent of the fluid level $\{X(t)\}$. The level $L(t) = \ell$ approximates which band \mathcal{D}_ℓ that $X(t)$ is in at time t , and the orbit $\mathbf{A}(t)$ can be used to obtain an approximation of where $X(t)$ is within the interval \mathcal{D}_ℓ .

We now proceed to describe the evolution of the orbit and phase processes, before introducing the level variable later.

The orbit process and phase dynamics On $\phi(t) = i$, between event epochs, the process $\{\mathbf{A}(t)\}$ evolves deterministically according to the differential equation

$$\frac{d}{dt}\mathbf{A}(t) = \mathbf{A}(t)(\mathbf{S}_i - \mathbf{A}(t)(\mathbf{S}_i + T_{ii}\mathbf{I}_p)\mathbf{e})\mathbf{I}_p. \quad (2.18)$$

Let $\mathbf{a} \in \mathcal{A}$ be an arbitrary vector in \mathcal{A} . On the event that no events occur before time $t + u$, $\mathbf{A}(u) = \mathbf{a} \in \mathcal{A}$ and $\phi(u) = i$, the solution to (2.18) states that $\mathbf{A}(t + u)$ evolves deterministically according to

$$\mathbf{A}(t + u) = \frac{\mathbf{a}e^{(\mathbf{S}_i + T_{ii}\mathbf{I}_p)t}}{\mathbf{a}e^{(\mathbf{S}_i + T_{ii}\mathbf{I}_p)t}\mathbf{e}} = \frac{\mathbf{a}e^{\mathbf{S}_i t}}{\mathbf{a}e^{\mathbf{S}_i t}\mathbf{e}}.$$

At time t an event occurs at rate

$$\mathbf{A}(t)(\mathbf{S}_i - T_{ii}\mathbf{I}_p)\mathbf{e} = \mathbf{A}(t)\mathbf{s}_i - T_{ii}.$$

More precisely, an event corresponding to a change in phase for $\phi(t)$ occurs at rate $-\mathbf{A}(t)T_{ii}\mathbf{e} = -T_{ii}$ and an event corresponding to a *change of level* occurs at rate $-\mathbf{A}(t)\mathbf{S}_i\mathbf{e} = \mathbf{A}(t)\mathbf{s}_i$. Later, we will make clear why we say that the latter event corresponds to a change of level. Upon an event occurring at time t , with probability $-T_{ii}/(-T_{ii} + \mathbf{A}(t^-)\mathbf{s}_i)$ the event corresponds to a change of phase and with probability $\mathbf{A}(t^-)\mathbf{s}_i/(-T_{ii} + \mathbf{A}(t^-)\mathbf{s}_i)$ the event corresponds to a change of level.

Upon an event corresponding to a change of level occurring at time t the process $\{\mathbf{A}(t)\}$ jumps to $\mathbf{A}(t) = \boldsymbol{\alpha}$.

Upon an event corresponding to a change of phase from i to $j \neq i$ occurring at time t , there are two possibilities; either $\text{sign}(c_i) = \text{sign}(c_j)$, or $\text{sign}(c_i) \neq \text{sign}(c_j)$. As discussed earlier, for states $i \in \mathcal{S}_0$ we must specify some association with either \mathcal{S}_+ or \mathcal{S}_- . Here we choose the augmented state space approach and duplicate \mathcal{S}_0 and associate one copy with $i \in \mathcal{S}_+$ and one copy with \mathcal{S}_- ; call these \mathcal{S}_{+0} and \mathcal{S}_{-0} , respectively. We take $\text{sign}(c_i) = +$ for $i \in \mathcal{S}_{+0}$ and $\text{sign}(c_i) = -$ for $i \in \mathcal{S}_{-0}$.

Upon an event corresponding to a change of phase from i to $j \neq i$ occurring at time t , in the case $\text{sign}(c_i) = \text{sign}(c_j)$, at the time of the event $\mathbf{A}(t)$ is unchanged but immediately begins to evolve according to (2.18) with i replaced by j , while $\{\phi(t)\}$ jumps from $\phi(t^-) = i$ to $\phi(t) = j$.

Upon an event corresponding to a change of phase from i to $j \neq i$ occurring at time t , in the case $\text{sign}(c_i) \neq \text{sign}(c_j)$, the process $\{\mathbf{A}(t)\}$ jumps to

$$\mathbf{A}(t) = \frac{\mathbf{A}(t^-)T_{ij}\mathbf{D}}{\mathbf{A}(t^-)T_{ij}\mathbf{D}\mathbf{e}} = \frac{\mathbf{A}(t^-)T_{ij}\mathbf{D}}{T_{ij}} = \mathbf{A}(t^-)\mathbf{D}$$

and then immediately proceeds to evolve according to (2.18) with i replaced by j , and $\{\phi(t)\}$ jumps from $\phi(t^-) = i$ to $\phi(t) = j$.

The two scenarios, $\text{sign}(c_i) \neq \text{sign}(c_j)$ and $\text{sign}(c_i) = \text{sign}(c_j)$, can be written succinctly by stating that, at the time of an event corresponding to a change of phase from i to j , $\{\mathbf{A}(t)\}$ jumps to $\mathbf{A}(t)\mathbf{D}^{1(\text{sign}(c_i) \neq \text{sign}(c_j))}$ and begins to evolve according to (2.18) with i replaced by j , meanwhile $\{\phi(t)\}$ jumps from $\phi(t^-) = i$ to $\phi(t) = j$.

The following result states that $\{\phi(t)\}$ has the same distribution as $\{\varphi(t)\}$ when the fluid queue is unbounded, the latter being the phase process of the fluid queue.

Theorem 2.1. *Let Θ_i be the time at which the first jump of the phase process of the unbounded QBD-RAP, $\{\phi(t)\}$, occurs given $\phi(0) = i$. For any initial orbit $\mathbf{a} \in \mathcal{A}$, then Θ_i has an exponential distribution with rate parameter $|T_{ii}|$. Furthermore, given $\phi(t)$ leaves state i , it jumps to state j with probability $T_{ij}/|T_{ii}|$. Hence $\phi(t)$ and $\varphi(t)$ have the same probability law.*

Proof. Let $\{\tau_n\}_{n \geq 0}$ with $\tau_0 = 0$ and τ_n the time of the n -th change of level of the QBD-RAP. Consider partitioning $\{\Theta_i > t\}$ with respect to $\{\tau_{n-1} < t \leq \tau_n\}$, $n = 1, 2, \dots$. For $n = 1$ we can write

$$\mathbb{P}(\Theta_i > t, \tau_0 < t \leq \tau_1 \mid \mathbf{A}(0) = \mathbf{a}) = \mathbf{a}e^{(\mathbf{S}_i + T_{ii}\mathbf{I}_p)t}\mathbf{e}$$

and since $T_{ii}\mathbf{I}_p$ commutes with \mathbf{S}_i and $e^{T_{ii}t}$ is a scalar, then this is equal to

$$\mathbf{a}e^{\mathbf{S}_i t}e^{T_{ii}\mathbf{I}_p t}\mathbf{e} = \mathbf{a}e^{\mathbf{S}_i t}\mathbf{e}e^{T_{ii}t} = \mathbb{P}(\tau_0 < t \leq \tau_1)e^{T_{ii}t}.$$

For $n > 1$, by partitioning on the times of the first $n - 1$ level changes, τ_1, \dots, τ_n , we get

$$\begin{aligned} & \mathbb{P}(\Theta_i > t, \tau_{n-1} < t \leq \tau_n \mid \mathbf{A}(0) = \mathbf{a}) \\ &= \int_{t_1=0}^t \int_{t_2=t_1}^t \dots \int_{t_{n-1}=t_{n-2}}^t \mathbb{P}(\Theta_i > t, t \leq \tau_n, \tau_{n-1} \in dt_{n-1}, \dots, \tau_2 \in dt_2, \tau_1 \in dt_1 \mid \mathbf{A}(0) = \mathbf{a}) \\ &= \int_{t_1=0}^t \int_{t_2=t_1}^t \dots \int_{t_{n-1}=t_{n-2}}^t \mathbf{a}e^{(\mathbf{S}_i + T_{ii}\mathbf{I}_p)t_1}\mathbf{s}_i \left(\prod_{k=2}^{n-1} \alpha e^{(\mathbf{S}_i + T_{ii}\mathbf{I}_p)(t_k - t_{k-1})}\mathbf{s}_i \right) \end{aligned}$$

$$\times \alpha e^{(\mathbf{S}_i + T_{ii} \mathbf{I}_p)(t - t_{n-1})} \mathbf{e} dt_{n-1} dt_{n-2} \dots dt_1.$$

Since $T_{ii} \mathbf{I}_p$ commutes with \mathbf{S}_i , $e^{T_{ii} t_k}$, $k = 1, \dots, n-1$ are scalars, and $t_1 + (t_2 - t_1) + \dots + (t_{n-1} - t_{n-2}) + (t - t_{n-1}) = t$, then this is equal to

$$\begin{aligned} & \int_{t_1=0}^t \int_{t_2=t_1}^t \dots \int_{t_{n-1}=t_{n-2}}^t \alpha e^{\mathbf{S}_i t_1} \mathbf{s}_i \left(\prod_{k=2}^{n-1} \alpha e^{\mathbf{S}_i(t_k - t_{k-1})} \mathbf{s}_i \right) \alpha e^{\mathbf{S}_i(t - t_{n-1})} \mathbf{e} \times e^{T_{ii} t} dt_{n-1} dt_{n-2} \dots dt_1 \\ &= \mathbb{P}(\tau_{n-1} < t \leq \tau_n) e^{T_{ii} t}. \end{aligned}$$

Hence, by the law of total probability,

$$\begin{aligned} \mathbb{P}(\Theta_i > t) &= \sum_{n=1}^{\infty} \mathbb{P}(\Theta_i > t, \tau_{n-1} < t \leq \tau_n) \\ &= \sum_{n=1}^{\infty} \mathbb{P}(\tau_{n-1} < t \leq \tau_n) e^{T_{ii} t} \\ &= e^{T_{ii} t}, \end{aligned}$$

and therefore Θ_i has an exponential distribution with rate $|T_{ii}|$.

Upon leaving state i at time t , $\phi(t)$ transitions to state j with probability

$$\frac{\left(\frac{\mathbf{A}(t) \mathbf{D}^{1(\text{sign}(c_i) \neq \text{sign}(c_j))} T_{ij} \mathbf{e}}{\sum_{j \in \mathcal{S}} \mathbf{A}(t) \mathbf{D}^{1(\text{sign}(c_i) \neq \text{sign}(c_j))} T_{ij} \mathbf{e} + \mathbf{A}(t) \mathbf{s}_i} \right)}{\left(\frac{\sum_{j \in \mathcal{S}} \mathbf{A}(t) \mathbf{D}^{1(\text{sign}(c_i) \neq \text{sign}(c_j))} T_{ij} \mathbf{e}}{\sum_{j \in \mathcal{S}} \mathbf{A}(t) \mathbf{D}^{1(\text{sign}(c_i) \neq \text{sign}(c_j))} T_{ij} \mathbf{e} + \mathbf{A}(t) \mathbf{s}_i} \right)} = \frac{\mathbf{A}(t) \mathbf{e} T_{ij}}{\sum_{j \in \mathcal{S}} \mathbf{A}(t) \mathbf{e} T_{ij}} = \frac{T_{ij}}{-T_{ii}}.$$

Therefore the process $\{\phi(t)\}$ has the same probability law as $\{\varphi(t)\}$. \square

Remark 2.2. *The same result can be shown for a regulated boundary. For boundary conditions which interact with the phase dynamics, such as a reflecting boundary, the result does not hold. The cause is the fact that the phase dynamics are level dependent – we may see a forced change of phase upon a boundary being hit – and the QBD-RAP can only approximate the level process of the fluid queue. However, until a boundary is hit (by either the fluid queue or QBD-RAP) then the phase processes match. We show later that, in the limit as the variance of the the matrix exponential distribution used in the construction of the QBD-RAP goes to zero, then the dynamics of the level process of the fluid queue, $X(t)$, are captured by the QBD-RAP, and boundary behaviour which interacts with the phase dynamics can be captured too.*

Since the phase processes $\{\phi(t)\}$ and $\{\varphi(t)\}$ have the same law when boundaries are not present, henceforth, we shall assume $\{\phi(t)\}$ and $\{\varphi(t)\}$ are coupled when possible (they share the same sample path). Specifically, in Section 3.1, we will analyse the QBD-RAP on the event that it remains in the same level, ℓ , say, and we compare this to the fluid queue on the event that the level remains in the band \mathcal{D}_ℓ . No boundary behaviour is involved in this calculation, so we treat $\{\phi(t)\}$ and $\{\varphi(t)\}$ as coupled and use the latter notation. When we must distinguish the two processes, we use $\phi(t)$ for the phase of the QBD-RAP and $\varphi(t)$ for the phase of the fluid.

The level process To event epochs of $\{(\mathbf{A}(t), \varphi(t))\}_{t \geq 0}$ we associate marks $\{-1, 0, +1\}$ in the following way.

- To events epochs corresponding to a change of phase of $\varphi(t)$ we associate the mark 0.
- To event epochs at time t which correspond to a change in level and for which $\varphi(t^-) = i \in \mathcal{S}_-$ we associate the mark -1 .
- To event epochs at time t which correspond to a change in level and for which $\varphi(t^-) = i \in \mathcal{S}_+$ we associate the mark $+1$.

Now define $N_+(t)$ ($N_-(t)$) as the simple point process which counts the number of event epochs with marks $+1$ (-1) which have occurred in the time up to and including time t . The level process of the QBD-RAP is given by $L(t) = N_+(t) - N_-(t)$. The process $\{(L(t), \mathbf{A}(t), \varphi(t))\}_{t \geq 0}$ forms a QBD with RAP components.

One way to specify the QBD with RAP components, $\{(L(t), \mathbf{A}(t), \varphi(t))\}$, is to describe its generator:

$$B = \begin{bmatrix} \ddots & \ddots & \ddots & & \\ & B_{-1} & B_0 & B_{+1} & \\ & & B_{-1} & B_0 & B_{+1} \\ & & & \ddots & \ddots & \ddots \end{bmatrix},$$

where,

$$B = \begin{bmatrix} B_{++} & B_{+-} \\ B_{-+} & B_{--} \end{bmatrix}$$

$$B_{-1} = \begin{bmatrix} 0 & & & \\ & 0 & & \\ & & C_- \otimes (s\alpha) & \\ & & & 0 \end{bmatrix}, \quad B_{+1} = \begin{bmatrix} C_+ \otimes (s\alpha) & & & \\ & 0 & & \\ & & 0 & \\ & & & 0 \end{bmatrix},$$

and

$$\begin{aligned} B_{++} &= \begin{bmatrix} C_+ \otimes S + T_{++} \otimes I & T_{+0} \otimes I \\ T_{0+} \otimes I & T_{00} \otimes I \end{bmatrix}, B_{+-} = \begin{bmatrix} T_{+-} \otimes D & 0 \\ T_{0-} \otimes D & 0 \end{bmatrix}, \\ B_{-+} &= \begin{bmatrix} T_{-+} \otimes D & 0 \\ T_{0+} \otimes D & 0 \end{bmatrix}, B_{--} = \begin{bmatrix} C_- \otimes S - T_{--} \otimes I & T_{-0} \otimes I \\ T_{0-} \otimes I & T_{00} \otimes I \end{bmatrix}. \end{aligned}$$

2.5 Boundary conditions

In the study and practical application of fluid queues, regulated, reflecting, sticky, or a mixture of these boundary conditions may be imposed. We now present the intuition as to how one may include such boundary conditions in the QBD-RAP approximation scheme.

Without loss of generality, assume that there is a boundary for the fluid level at $y_0 = 0$. This boundary can only be hit from above in phases $i \in \mathcal{S}_-$. Suppose that, upon hitting the boundary in phase $i \in \mathcal{S}_-$, the phase jumps from i to $j \in \mathcal{S}$ with probability p_{ij} . If $j \in \mathcal{S}_- \cup \mathcal{S}_0$ the process remains at the boundary and the phase process evolves among states in $\mathcal{S}_- \cup \mathcal{S}_0$ until the first transition to a state in \mathcal{S}_+ , at which point the level $\{X(t)\}$ immediately leaves the boundary – we will call this a *sticky* boundary. If $j \in \mathcal{S}_+$ the process immediately leaves the boundary – we will call this a *reflecting* boundary. We collect the probabilities p_{ij} into the matrices $\mathbf{P}_{-+} = [p_{ij}]_{i \in \mathcal{S}_-, j \in \mathcal{S}_+}$, $\mathbf{P}_{-0} = [p_{ij}]_{i \in \mathcal{S}_-, j \in \mathcal{S}_0}$, and $\mathbf{P}_{--} = [p_{ij}]_{i \in \mathcal{S}_-, j \in \mathcal{S}_-}$.

Adjacent to the boundary at $y_0 = 0$ is the band $\mathcal{D}_0 = [0, \Delta]$ which corresponds to level 0 for the QBD-RAP. We denote the boundary of the QBD-RAP as level -1 , which corresponds to $\{0\}$ for the fluid queue. Modelling the behaviour of the fluid queue at boundaries can be broken down into three components. 1) Modelling the time and phase when the fluid level hits the boundary. 2) Modelling the phase whilst the fluid level remains at the boundary. 3) Modelling the fluid level and phase at the exit from the boundary.

We claim that the event $\{L(t) = \ell, \phi(t) = i\}$ models the event $\{X(t) \in \mathcal{D}_{\ell,i}, \varphi(t) = i\}$, and further that instants u with $L(u^-) \neq L(u), \phi(u) = i$ model the events $\{X(u^-) \in \mathcal{D}_{\ell,i}, X(u) \notin \mathcal{D}_{\ell,i}, \varphi(u) = i\}$. With this in mind, we suppose that when the QBD-RAP is in level 0 in phase $i \in \mathcal{S}_-$ and there is a change of level, this corresponds to approximating the event that the fluid level $\{X(t)\}$ hits the boundary at 0. We refer to this as the QBD-RAP hitting the boundary also. We show later that, using matrix exponentials with sufficiently small variance in the construction of the QBD-RAP, then the distribution of time until the QBD-RAP first hits the boundary closely models the time until the fluid level hits the boundary. This addresses component 1).

For a sticky lower boundary at 0, given the fluid level reaches zero at time u in phase $i \in \mathcal{S}_-$, then the phase transitions to some phase $j \in \mathcal{S}_- \cup \mathcal{S}_0$ with probability p_{ij} , and the

distribution of the phase $\varphi(t)$ (on the event that the process has not left the boundary at or before time t) is given by the elements of the vector

$$p_{ij} \mathbf{e}_j \exp \left(\begin{bmatrix} \mathbf{T}_{--} & \mathbf{T}_{-0} \\ \mathbf{T}_{0-} & \mathbf{T}_{00} \end{bmatrix} (t - u) \right).$$

The distribution for the time until $\{X(t)\}$ leaves $x = 0$ for the first time after u is

$$1 - p_{ij} \mathbf{e}_j \exp \left(\begin{bmatrix} \mathbf{T}_{--} & \mathbf{T}_{-0} \\ \mathbf{T}_{0-} & \mathbf{T}_{00} \end{bmatrix} (t - u) \right) \mathbf{e}.$$

Therefore, in the case of a sticky boundary, we can capture the behaviour of the process at the boundary exactly. For a reflecting boundary there is no time spent at the boundary. This addresses component 2) of the modelling problem.

For both sticky and reflecting boundaries, given the QBD-RAP is in the correct phase at the instant upon which it hits the boundary, then upon exiting the boundary the phase can be captured exactly too. For a sticky boundary, given the fluid/QBD-RAP hits the boundary in phase $i \in \mathcal{S}_-$ at time t , it then jumps to phase $j \in \mathcal{S}_+ \cup \mathcal{S}_0$ and exits the boundary in phase $k \in \mathcal{S}_+$ at time $(t - u)$ with density

$$p_{ij} \mathbf{e}_j \exp \left(\begin{bmatrix} \mathbf{T}_{--} & \mathbf{T}_{-0} \\ \mathbf{T}_{0-} & \mathbf{T}_{00} \end{bmatrix} (t - u) \right) \begin{bmatrix} \mathbf{T}_{-+} \\ \mathbf{T}_{0+} \end{bmatrix} \mathbf{e}_k.$$

For a reflecting boundary, given the boundary is hit in phase $i \in \mathcal{S}_-$, the phase upon leaving the boundary is $j \in \mathcal{S}_+$ with probability p_{ij} . Upon leaving the boundary, the appropriate orbit position for the QBD-RAP is α , as this corresponds to the fluid level $X = 0$.

To summarise, at time t the rate at which probability mass accumulates at the sticky boundary in phase $j \in \mathcal{S}_- \cup \mathcal{S}_0$ and level -1 , upon hitting the boundary in phase $i \in \mathcal{S}_-$ and level 0 , is

$$c_i p_{ij} \mathbf{A}(t) \mathbf{s}.$$

The rate at which density from phase $i \in \mathcal{S}_-$ and level 0 transitions to density in phase $j \in \mathcal{S}_+$ and level 0 upon hitting a boundary is

$$c_i p_{ij} \mathbf{A}(t) \mathbf{s},$$

and upon this transition, the orbit jumps to α . The rate at which mass leaves the sticky boundary from phase $i \in \mathcal{S}_- \cup \mathcal{S}_0$ and level -1 into phases $j \in \mathcal{S}_+$ and level 0 is

$$T_{ij},$$

and upon leaving the boundary the orbit is α .

This information can be inscribed in the generator of the QBD-RAP. For example, using the augmented state-space scheme to account for phases \mathcal{S}_0 , the generator of the QBD-RAP described above has boundary conditions included as

$$\left[\begin{array}{cc|ccc|c} T_{--} & T_{0-} & T_{-+} \otimes \alpha & 0 & 0 & 0 & \\ T_{-0} & T_{00} & T_{0+} \otimes \alpha & 0 & 0 & 0 & \\ \hline 0 & 0 & & & & & \\ 0 & 0 & & & & & \\ (C_- P_{--}) \otimes s & (C_- P_{-0}) \otimes s & \check{B}_0 & & & & B_{+1} \\ 0 & 0 & & & & & \ddots \\ \hline & & B_{-1} & & & & B_0 \\ & & & & \ddots & & \ddots \end{array} \right],$$

where

$$\check{B}_0 = \left[\begin{array}{ccc|cc} C_+ \otimes S + T_{++} \otimes I & T_{+0} \otimes I & T_{+-} \otimes D & & \\ T_{0+} \otimes I & T_{00} \otimes I & T_{0-} \otimes D & & \\ (C_- P_{-+}) \otimes s\alpha + T_{-+} \otimes D & & C_- \otimes S + T_{--} \otimes I & T_{-0} \otimes I & \\ T_{0+} \otimes D & & T_{0-} \otimes I & T_{00} \otimes I & \end{array} \right].$$

The top-left block represents the point masses due to the sticky boundary. The bottom-left block is the transition of density into point masses due to the sticky boundary. The top-right block is the transition of point masses into density as the process leaves the boundary. The term $(C_- P_{-+}) \otimes s\alpha$ in \check{B}_0 incorporates the reflecting boundary behaviour.

Upper boundaries can be included in an analogous manner.

The orbit process $\{\mathbf{A}(t)\}$ is not required to model the behaviour at the sticky boundary. However, we suppose that $\mathbf{A}(t) = 1$ at the boundaries. This choice is arbitrary but allows us to generalise some notation later when we are describing the evolution of the QBD-RAP. Further, for boundaries at $y_0 = 0$ and $y_{K+1} = (K+1)\Delta$, we let the level process take the value $L(t) = -1$ at the lower boundary, and $L(t) = K+1$ at the upper boundary. Denote the set of levels (including boundary levels) by $\mathcal{K} = \{-1, 0, \dots, K, K+1\}$.

Remark 2.3. *It may be possible to extend the QBD-RAP approximation to model jumps into and out of the boundary, provided that the jumps into/out of the boundary happen at an intensity which can be described by a matrix exponential distribution.*

Furthermore, it may be possible to model more general boundary behaviours which can be approximated by a limit of matrix exponentials. For example, a boundary condition where the fluid queue spends a deterministic time at the boundary.

2.6 Initial conditions

We argued earlier that we can think of the orbit $\mathbf{k}(x)$ as corresponding to the fluid being a distance of x from the left boundary of an interval when $i \in \mathcal{S}_+$, or from the right

boundary of an interval when $i \in \mathcal{S}_-$. With this interpretation, an approximation to the initial condition $(X(0), \varphi(0)) = (x_0, i)$, $x_0 \in \mathcal{D}_{\ell, i}$, $i \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$ is to set the initial orbit to $\mathbf{k}(x_0 - y_\ell)$. Similarly, an approximation to the initial condition $(X(0), \varphi(0)) = (x_0, i)$, $x_0 \in \mathcal{D}_{\ell, i}$, $x_0 \neq y_{\ell+1}$, $i \in \mathcal{S}_- \cup \mathcal{S}_{-0}$ is to set the initial orbit position to $\mathbf{k}(y_{\ell+1} - x_0)$.

We use the notation $\mathbf{a}_{\ell, i}(x_0) = \mathbf{k}(x_0 - y_\ell)$ for the initial orbit position corresponding to x_0 when the initial phase is $i \in \mathcal{S}_+$ and $x_0 \in \mathcal{D}_{\ell, i}$. Similarly, for $i \in \mathcal{S}_-$ and $x_0 \in \mathcal{D}_{\ell, i}$ define the notation $\mathbf{a}_{\ell, i}(x_0) = \mathbf{k}(y_{\ell+1} - x_0)$.

More generally, given an initial measure $\mu_i(\cdot) := \mathbb{P}(X(0) \in \cdot, \varphi(0) = i)$, an approximation to this initial condition is to set the orbit to $\int_{x \in \mathcal{D}_{\ell, i}} \mathbf{a}_{\ell, i}(x) d\mu_i$, for $i \in \mathcal{S}$, $\ell \in \mathcal{K} \setminus \{-1, K+1\}$.

2.7 At time t – closing operators

So far we have described how we can use a QBD-RAP to approximate a fluid queue. The level process $\{L(t)\}$ of the QBD-RAP approximates the process

$$\left\{ \sum_{k \in \mathcal{K}} \sum_{i \in \mathcal{S}} k 1(X(t) \in \mathcal{D}_{k, i}) \right\}_{t \geq 0}. \quad (2.19)$$

This may be of interest in its own right. However, for some applications, this approximation may be too coarse – the level process tells us nothing about where in the intervals $\mathcal{D}_{k, i}$ the fluid queue might be.

We now describe how the orbit position $\mathbf{A}(t)$ can be used to approximate the density of the level of the fluid queue $X(t)$. This reconstruction is entirely post-hoc in the sense that it does not affect the dynamics of the QBD-RAP itself. For any time t we only take the information contained in $\mathbb{E}[A(t)1(L(t) = \ell, \varphi(t) = i)]$ and rewrite it in a descriptive way.

Suppose that the QBD-RAP is in level $L(t) = \ell$, phase $\phi(t) = j \in \mathcal{S}_+$, and the orbit is $\mathbf{A}(t) = \mathbf{a}$. If the QBD-RAP remains in phase j , the QBD-RAP approximation will transition out of level ℓ in the infinitesimal time interval $t + du$ with density $\mathbf{a}e^{|c_j|S^u}|c_j|\mathbf{s} du$. At the time of the change of level we estimate the position of $X(t + u)$ by $X(t + u) \approx y_{\ell+1}$. Tracing this back to time t , we estimate the position of $X(t)$ as $X(t) \approx y_{\ell+1} - |c_j|u$. Reverse engineering this logic, the approximation to the density of $X(t) \in dx$ is $\mathbf{a}e^{|c_j|S(y_{\ell+1}-x)/|c_j|}|c_j|\mathbf{s} dx/|c_j|$, since $dx = |c_j|du$ where dx is an infinitesimal interval in space and du is an infinitesimal interval with respect to time.

Similarly, for $j \in \mathcal{S}_-$, if the QBD-RAP remains in phase j then the QBD-RAP approximation will transition out of level ℓ in the infinitesimal time interval $t + du$ with density $\mathbf{a}e^{|c_j|S^u}|c_j|\mathbf{s} du$. At the time of the transition of level, we estimate the position of $X(t + u)$ by $X(t + u) \approx y_\ell$. Tracing this back to time t , we estimate the position of $X(t)$

as $X(t) \approx y_\ell + |c_j|u$. Reverse engineering this logic, the approximation to the density $X(t) \in dx$ is $\mathbf{a}e^{|c_j|\mathbf{S}(x-y_\ell)/|c_j|}|c_j|\mathbf{s} dx/|c_j|$, since $dx = |c_j|du$.

For phases $j \in \mathcal{S}_0$, if phase j is associated with \mathcal{S}_+ , we use the same approximation as if $j \in \mathcal{S}_+$, and if phase $j \in \mathcal{S}_0$ is associated with \mathcal{S}_- , we use the same approximation as if $j \in \mathcal{S}_-$.

This reasoning leads to an approximation of the distribution of the fluid at time t , $\mathbb{P}(X(t) \in dx, \varphi(t) = j \mid X(0) = x_0, \varphi(0) = i)$, as

$$\int_{\mathbf{a} \in \mathcal{A}} \mathbb{P}(L(t) = \ell, \mathbf{A}(t) \in d\mathbf{a}, \phi(t) = j \mid L(0) = \ell_0, \mathbf{A}(0) = \mathbf{a}_{\ell_0, i}(x_0), \phi(0) = i) \mathbf{a}e^{\mathbf{S}z} \mathbf{s} dx, \quad (2.20)$$

where $x \in \mathcal{D}_{\ell, j}$, $x_0 \in \mathcal{D}_{\ell_0, i}$ and

$$z = \begin{cases} x - y_\ell & i \in \mathcal{S}_- \\ y_{\ell+1} - x & i \in \mathcal{S}_+. \end{cases}$$

While the estimate (2.20) is appealing, it is, however, defective – it does not integrate to 1 for any finite-dimensional matrix exponential distribution. To see this, compute

$$\begin{aligned} & \sum_{\ell \in \mathcal{K}} \sum_{j \in \mathcal{S}} \int_{z \in \mathcal{D}_{\ell, j}} \int_{\mathbf{a} \in \mathcal{A}} \mathbb{P}(L(t) = \ell, \mathbf{A}(t) \in d\mathbf{a}, \phi(t) = j \mid L(0) = \ell_0, \mathbf{A}(0) = \mathbf{a}_{\ell, i}(x_0), \\ & \quad \phi(0) = i) \mathbf{a}e^{\mathbf{S}z} \mathbf{s} dz \\ &= \sum_{\ell \in \mathcal{K}} \sum_{j \in \mathcal{S}} \int_{\mathbf{a} \in \mathcal{A}} \mathbb{P}(L(t) = \ell, \mathbf{A}(t) \in d\mathbf{a}, \phi(t) = j \mid L(0) = \ell_0, \mathbf{A}(0) = \mathbf{a}_{\ell, i}(x_0), \\ & \quad \phi(0) = i) (1 - \mathbf{a}e^{\mathbf{S}\Delta} \mathbf{e}) \\ &< 1, \end{aligned}$$

since $\mathbf{a}e^{\mathbf{S}u} \mathbf{e} > 0$ for any $u > 0$. For an orbit position $\mathbf{A}(t) = \mathbf{a}$, the amount of mass missing from each level and phase is

$$\mathbb{P}(L(t) = \ell, \mathbf{A}(t) \in d\mathbf{a}, \phi(t) = j \mid L(0) = \ell_0, \mathbf{A}(0) = \mathbf{a}_{\ell, i}(x_0), \phi(0) = i) \mathbf{a}e^{\mathbf{S}\Delta} \mathbf{e} dx.$$

The problem is that the triple $(\mathbf{a}, \mathbf{S}, \mathbf{s})$ defines a distribution on $x \in [0, \infty)$, however, it only makes sense to take $x \in [0, \Delta)$ in the approximation scheme.

To partially rectify this we can instead approximate $\mathbb{P}(X(t) \in dx, \varphi(t) = j \mid X(0) = x_0, \varphi(0) = i)$ by

$$\int_{\mathbf{a} \in \mathcal{A}} \mathbb{P}(L(t) = \ell, \mathbf{A}(t) \in d\mathbf{a}, \phi(t) = j \mid L(0) = \ell_0, \mathbf{A}(0) = \mathbf{a}_{\ell, i}(x_0), \phi(0) = i) \mathbf{a}u_i^\ell(x) dx, \quad (2.21)$$

where

$$\mathbf{a}u_i^\ell(x) = \begin{cases} \mathbf{a} \left(e^{\mathbf{S}(y_{\ell+1}-x)} \mathbf{s} + e^{\mathbf{S}(2\Delta-(y_{\ell+1}-x))} \mathbf{s} \right), & i \in \mathcal{S}_+, \\ \mathbf{a} \left(e^{\mathbf{S}(x-y_\ell)} \mathbf{s} + e^{\mathbf{S}(2\Delta-(x-y_\ell))} \mathbf{s} \right), & i \in \mathcal{S}_-. \end{cases} \quad (2.22)$$

Intuitively, we take the density function $\mathbf{a}e^{\mathbf{S}x}\mathbf{s}$ on $x \in [0, 2\Delta)$, and ‘fold’ it back on itself around Δ , to create a density function on $[0, \Delta)$,

$$\mathbf{a}e^{\mathbf{S}x}\mathbf{s} + \mathbf{a}e^{\mathbf{S}(2\Delta-x)}\mathbf{s}, \quad x \in [0, \Delta).$$

The missing mass is now proportional to

$$\mathbf{a}e^{\mathbf{S}2\Delta}\mathbf{e},$$

which is less than the quantity $\mathbf{a}e^{\mathbf{S}\Delta}\mathbf{e}$ from the approximation scheme (2.20). There is nothing special about the choice of truncating the distribution at 2Δ in this construction, and we could choose any truncation value greater than Δ .

A third option is to approximate $\mathbb{P}(X(t) \in dx, \varphi(t) = j \mid X(0) = x_0, \varphi(0) = i)$ by

$$\int_{\mathbf{a} \in \mathcal{A}} \mathbb{P}(L(t) = \ell, \mathbf{A}(t) \in d\mathbf{a}, \phi(t) = j \mid L(0) = \ell_0, \mathbf{A}(0) = \mathbf{a}_{\ell,i}(x_0), \phi(0) = i) \mathbf{a}v_i^\ell(x) dx, \quad (2.23)$$

where

$$\mathbf{a}v_i^\ell(x) = \begin{cases} \mathbf{a} \left(e^{\mathbf{S}(y_{\ell+1}-x)} + e^{\mathbf{S}(2\Delta-(y_{\ell+1}-x))} \right) [I - e^{\mathbf{S}2\Delta}]^{-1} \mathbf{s}, & i \in \mathcal{S}_+, \\ \mathbf{a} \left(e^{\mathbf{S}(x-y_\ell)} + e^{\mathbf{S}(2\Delta-(x-y_\ell))} \right) [I - e^{\mathbf{S}2\Delta}]^{-1} \mathbf{s}, & i \in \mathcal{S}_-. \end{cases} \quad (2.24)$$

Here, we take the density function $\mathbf{a}e^{\mathbf{S}x}\mathbf{s}$, defined on $[0, \infty)$, and map it to a density function on $[0, \Delta)$ by $\Delta - |(x \bmod 2\Delta) - \Delta|$. The resulting density function is

$$\begin{aligned} v(x) &= \sum_{m=0}^{\infty} \mathbf{a} \left(e^{\mathbf{S}(x+2\Delta m)} + e^{\mathbf{S}(2\Delta-x+2\Delta m)} \right) \mathbf{s} \\ &= \mathbf{a} \left(e^{\mathbf{S}x} + e^{\mathbf{S}(2\Delta-x)} \right) \sum_{m=0}^{\infty} e^{\mathbf{S}2\Delta m} \mathbf{s} \\ &= \mathbf{a} \left(e^{\mathbf{S}x} + e^{\mathbf{S}(2\Delta-x)} \right) [I - e^{\mathbf{S}2\Delta}]^{-1} \mathbf{s}. \end{aligned}$$

The difference between the approximation schemes is the way in which the vector \mathbf{a} is used. We generalise this idea with the concept of *closing operators* which is a linear operator $V(x) : \mathcal{A} \rightarrow \mathbb{R}$, for each $x \in [0, \Delta)$. For example, the closing operator in Equations (2.20) is the operator $V(x)$, $x \in [0, \Delta)$ such that for any $\mathbf{a} \in \mathcal{A}$,

$$\mathbf{a}V(x) = \mathbf{a}e^{\mathbf{S}x}\mathbf{s}. \quad (2.25)$$

Similarly, in (2.21)

$$\mathbf{a}V(x) = \mathbf{a} \left(e^{\mathbf{S}x} \mathbf{s} + e^{\mathbf{S}(2\Delta-x)} \mathbf{s} \right), \quad (2.26)$$

and in (2.23)

$$\mathbf{a}V(x) = \mathbf{a} \left(e^{\mathbf{S}x} \mathbf{s} + e^{\mathbf{S}(2\Delta-x)} [I - e^{\mathbf{S}2\Delta}]^{-1} \mathbf{s} \right). \quad (2.27)$$

The operator $V(x)$ is a choice which forms part of the definition of the approximation scheme. As we shall see, given certain properties of $V(x)$, we can prove that the approximation scheme converges, and ensures positivity. In the cases above, all the closing operators (2.25)-(2.26) lead to an approximation which converges and, due to their interpretation as probability densities, ensure positivity.

Comments on linearity and normalisation We considered taking the closing operators to be

$$\mathbf{a}V(x) = \frac{\mathbf{a} \left(e^{\mathbf{S}x} \mathbf{s} + e^{\mathbf{S}(2\Delta-x)} \mathbf{s} \right)}{1 - \mathbf{a}e^{\mathbf{S}2\Delta}\mathbf{e}}, \quad (2.28)$$

which also ensures that the solution always preserves mass (the result of integrating (2.28) over $[0, \Delta)$ is 1). However, (2.28) is not a linear operator. The importance of this fact is that for a linear operator,

$$\mathbb{E}[\mathbf{A}(t)V(x)] = \mathbb{E}[\mathbf{A}(t)]V(x),$$

thus, the computation of $\mathbb{E}[\mathbf{A}(t)V(x)]$ can be achieved by first computing $\mathbb{E}[\mathbf{A}(t)]$ and then applying $V(x)$ to the result. In this sense, $\mathbb{E}[\mathbf{A}(t)]$ contains all of the information about the history of the process up to time t needed to compute $\mathbb{E}[\mathbf{A}(t)V(x)]$. This is much the same as in RAPs and QBD-RAPs where $\mathbb{E}[\mathbf{A}(t)]$ is all that is required to compute probabilities about the future of the processes from time t onwards.

When $V(x)$ is not linear then, in general,

$$\mathbb{E}[\mathbf{A}(t)V(x)] \neq \mathbb{E}[\mathbf{A}(t)]V(x).$$

For example, with (2.28),

$$\mathbb{E}[\mathbf{A}(t)V(x)] = \int_{\mathbf{a} \in \mathcal{A}} \mathbb{P}(\mathbf{A}(t) \in d\mathbf{a}) \frac{\mathbf{a} \left(e^{\mathbf{S}x} \mathbf{s} + e^{\mathbf{S}(2\Delta-x)} \mathbf{s} \right)}{1 - \mathbf{a}e^{\mathbf{S}2\Delta}\mathbf{e}}. \quad (2.29)$$

While the calculation in (2.29) is theoretically possible, practically it is not. The seemingly innocuous integral over $\mathbf{a} \in \mathcal{A}$ actually amounts to an integral over all possible sample paths of the QBD-RAP (i.e. all possible event times of the QBD-RAP before time t) and in general, there is no ‘nice’ way to do this computation.

In the linear case, computation of

$$\mathbb{E}[\mathbf{A}(t)] = \int_{\mathbf{a} \in \mathcal{A}} \mathbb{P}(\mathbf{A}(t) \in d\mathbf{a}) \mathbf{a},$$

amounts to a matrix-exponential calculation.

In practice we use

$$\frac{\mathbb{E}[\mathbf{A}(t)] (e^{\mathbf{S}x} \mathbf{s} + e^{\mathbf{S}(2\Delta-x)} \mathbf{s})}{1 - \mathbb{E}[\mathbf{A}(t)] e^{\mathbf{S}2\Delta} \mathbf{e}} \quad (2.30)$$

as a normalised version of (2.26), and this seems to work well. However, this is not the same as using the closing operator (2.28).

Chapter 3

Weak convergence of the QBD-RAP scheme

This chapter details convergence of the approximation scheme constructed in Chapter 2. The QBD-RAP is constructed using *matrix-exponential distributions*. The main result shows the convergence of the approximation scheme under the assumption that the variance of the matrix-exponential distribution(s) used in the construction tends to 0. Given that there is no known simple closed form for the transient distributions of a fluid queue, and that we make minimal assumptions about the matrix-exponential distributions used in the construction of the approximation, it is somewhat remarkable that we are able to establish a convergence result. The generality of the result with respect to the assumptions on the matrix-exponential distributions used in the construction of the approximation is necessitated by the fact that we use a class of *concentrated matrix-exponential distributions* found numerically in (Horváth et al. 2020), and for which there is relatively little known about their properties.

We suppose that we have a sequence $\{Z^{(p)}\}_{p \geq 1}$ of matrix exponential distributions, $Z^{(p)} \sim ME(\boldsymbol{\alpha}^{(p)}, \boldsymbol{S}^{(p)}, \boldsymbol{s}^{(p)})$, such that $\text{Var}(Z^{(p)}) \rightarrow 0$ as $p \rightarrow \infty$. Ultimately, we show weak convergence (in both space and time) of the QBD-RAP approximation scheme via convergence of various Laplace transforms with respect to time of the QBD-RAP to the corresponding Laplace transforms of the fluid queue. We use the superscript (p) to denote dependence on the underlying choice of matrix exponential distribution that is used in the construction of the QBD-RAP scheme. To simplify notation, we omit the super script (p) where possible. In the following we show error bounds for an arbitrary parameter $\varepsilon > 0$. However, keep in mind the ultimate intention is to show convergence, for which we choose this parameter to be $\varepsilon^{(p)} = \text{Var}(Z^{(p)})^{1/3}$. Other notations which have been defined so far which are defined by $Z^{(p)}$ and therefore also implicitly depend on p are $\boldsymbol{\alpha}^{(p)}, \boldsymbol{S}^{(p)}, \boldsymbol{s}^{(p)}, \boldsymbol{S}_i^{(p)}, \boldsymbol{s}_i^{(p)}, \boldsymbol{D}^{(p)}, \mathcal{A}^{(p)}$.

In the following we show various results which involve integrating a function g , or a

sequence of functions g_1, g_2, \dots . We make the following assumptions about such functions,

Assumptions 3.1. Let g be a function $g : [0, \infty) \rightarrow [0, \infty)$ which is

(i) non-negative,

$$g(x) \geq 0 \text{ for all } x \geq 0,$$

(ii) bounded,

$$g(x) \leq G < \infty \text{ for all } x \geq 0,$$

(iii) integrable,

$$\int_{x=0}^{\infty} g(x) dx \leq \widehat{G} < \infty,$$

(iv) and Lipschitz continuous

$$|g(x) - g(u)| \leq L|x - u| \text{ for all } x, u \geq 0, 0 < L < \infty. \quad (3.1)$$

We also need a corresponding sequence of closing operators which we denote by $V^{(p)}$. For the convergence results, we require the following properties of the closing operators $V^{(p)}(x)$, $x \in [0, \Delta)$.

Properties 3.2. Let $\{V^{(p)}(x)\}_{p \geq 1}$ be a sequence of closing operators.

(i) For $x \in [0, \Delta)$, $u \geq 0$,

$$\alpha^{(p)} e^{\mathbf{S}^{(p)} u} (-\mathbf{S}^{(p)})^{-1} V^{(p)}(x) \leq \alpha^{(p)} e^{\mathbf{S}^{(p)} u} \mathbf{e} G_V,$$

for some $0 \leq G_V < \infty$ independent of p for $p > p_0$ and some $p_0 < \infty$.

(ii) Let g be a function satisfying the Assumptions 3.1. For $u \leq \Delta - \varepsilon^{(p)}$, $v \in [0, \Delta)$, then

$$\left| \int_{x=0}^{\infty} \frac{\alpha^{(p)} e^{\mathbf{S}^{(p)}(u+x)}}{\alpha^{(p)} e^{\mathbf{S}^{(p)} u} \mathbf{e}} V^{(p)}(v) g(x) dx - g(\Delta - u - v) 1(u + v \leq \Delta - \varepsilon^{(p)}) \right| = |r_V^{(p)}(u, v)|,$$

where

$$\int_{u=0}^{\Delta} |r_V^{(p)}(u, v)| du \leq R_{V,1}^{(p)} \rightarrow 0$$

and

$$\int_{v=0}^{\Delta} |r_V^{(p)}(u, v)| dv \leq R_{V,2}^{(p)} \rightarrow 0$$

as $\text{Var}(Z^{(p)}) \rightarrow 0$.

For convenience, in the sequel, let us adopt the notation

$$\mathbb{P}(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in dx, \phi^{(p)}(t) = j \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i) \quad (3.2)$$

for an approximation to

$$\mathbb{P}(X(t) \in dx, \varphi(t) = j \mid X(0) = x_0, \varphi(0) = i), \quad (3.3)$$

$x \in \mathcal{D}_{\ell,j}, x_0 \in \mathcal{D}_{\ell_0,i}$.

In Appendix A.2 we provide results which show that the closing operators (2.25) - (2.26) have the properties 3.2.

Ultimately, we will apply the Extended Continuity Theorem for Laplace transforms (Feller 1957, Chapter XIII, Theorem 2a) to claim convergence. The Extended Continuity Theorem for Laplace transforms requires us to show convergence of the Laplace transform pointwise with respect to the transform parameter, λ . Therefore, we can fix $\lambda > 0$ in the following sections.

The structure of this chapter is as follows. First, in Sections 3.1 we analyse the behaviour of the fluid queue and QBD-RAP on the event that they stay in the same level. Then, in Section 3.2, we derive expressions for certain Laplace transforms with respect to time of the QBD-RAP and fluid queue on the event that there is no change of level. Section 3.3 provides error bounds between the Laplace transforms of the QBD-RAP and the fluid queue that were derived in Section 3.2 and also a domination condition used so that we may apply the Dominated Convergence Theorem. At this stage we have shown weak convergence of the approximation up to the first change of level. The subsequent sections generalise this to global convergence. We start by showing weak convergence of the approximation at the n th change of level, $n \geq 1$ in Section 3.4. Section 3.5 generalises this to show weak convergence between the n th and $n + 1$ th change of level. Via the Dominated Convergence Theorem (again), Section 3.6 proves the weak convergence of the QBD-RAP approximation to the fluid queue. So as to not obscure the logical flow of the presentation, technical proofs are reserved for Appendix A.

3.1 Characterisations on no change of level

3.1.1 Characterising the fluid queue

Let τ_1^X be the minimum of the time at which $\{X(t)\}$ hits a boundary or exits \mathcal{D}_{ℓ_0} , where $X(0) = x_0 \in \mathcal{D}_{\ell_0}$, or $\{X(t)\}$ exits a boundary. More precisely,

$$\tau_1^X = \min \left\{ \begin{array}{l} \inf \{t > 0 \mid X(t) = y_\ell, \ell \in \mathcal{K}\}, \\ \inf \{t > 0 \mid X(t) \neq 0, X(0) = 0\}, \\ \inf \{t > 0 \mid X(t) \neq y_{K+1}, X(0) = y_{K+1}\} \end{array} \right\}.$$

Consider the measures

$$\mu^{\ell_0}(t)(dx, j; x_0, i) := \mathbb{P}(X(t) \in dx, t < \tau_1^X, \varphi(t) = j \mid X(0) = x_0, \varphi(0) = i), \quad (3.4)$$

$\ell_0 \in \{0, \dots, K\}$, $x, x_0 \in \mathcal{D}_{\ell_0, i}$, $i, j \in \mathcal{S}, t \geq 0$. In words, this is the distribution of the fluid queue at time t on the event that the fluid level remains within \mathcal{D}_{ℓ_0} up to and including time t and is in phase j at time t , given that it started at $X(0) = x_0 \in \mathcal{D}_{\ell_0, i}$ in phase i . The QBD-RAP approximation to (3.4) is

$$(\mathbf{e}_i \otimes \mathbf{a}_{\ell_0, i}^{(p)}(x_0)) \exp\left(\mathbf{B}^{(p)}t\right) (\mathbf{e}_j \otimes V^{(p)}(y_{\ell_0+1} - x)) dx, j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}. \quad (3.5)$$

There is no simple expression for (3.4). There are expressions for the Laplace transform of (3.4) with respect to time. One is in terms of the of first return matrices $\Psi(\lambda)$ and $\Xi(\lambda)$ (Bean et al. 2009b). Here we opt for another expression for the Laplace transform which is obtained by partitioning as follows.

Suppose $i \in \mathcal{S}_+, j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$. The arguments for all other combinations of $i, j \in \mathcal{S}$ are analogous and notable differences will be pointed out where necessary. There are some issues pertaining to starting the QBD-RAP in phases $i \in \mathcal{S}_{+0}$, hence the restriction $\varphi(0) = i \in \mathcal{S}_+$ only. We will treat the case $i \in \mathcal{S}_0$ separately and in due course.

Denote by $\Sigma_m, m \geq 1$ the sequence of (stopping) times at which $\{\varphi(t)\}$ jumps from $\mathcal{S}_+ \cup \mathcal{S}_{+0}$ to \mathcal{S}_- for the m th time. Denote by $\Gamma_m, m \geq 1$ the sequence of (stopping) times at which $\{\varphi(t)\}$ jumps from $\mathcal{S}_- \cup \mathcal{S}_{-0}$ to \mathcal{S}_+ for the m th time. See Figure 3.1. More precisely, for sample paths with $\varphi(0) \in \mathcal{S}_+$, let $\Gamma_0 = 0$, then for $m \geq 1$,

$$\Sigma_m := \inf\{t > \Gamma_{m-1} \mid \varphi(t) \in \mathcal{S}_-\}, \quad (3.6)$$

$$\Gamma_m := \inf\{t > \Sigma_m \mid \varphi(t) \in \mathcal{S}_+\}. \quad (3.7)$$

For times t such that $\Gamma_m \leq t < \Sigma_{m+1}$, then $\varphi(t) \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$. For times t such that $\Sigma_{m+1} \leq t < \Gamma_{m+1}$, then $\varphi(t) \in \mathcal{S}_- \cup \mathcal{S}_{-0}$. The events $\{\Gamma_m \leq t < \Sigma_{m+1}\}$, and $\{\Sigma_{m+1} \leq t < \Gamma_{m+1}\}$, $m \geq 0$, partition the sample paths of (3.4) into periods where the fluid is either non-decreasing or non-increasing, respectively; see Figure 3.1. Similarly, for sample paths with $\varphi(0) \in \mathcal{S}_-$, let $\Sigma_0 = 0$, then for $m \geq 1$,

$$\Gamma_m := \inf\{t > \Sigma_{m-1} \mid \varphi(t) \in \mathcal{S}_+\}. \quad (3.8)$$

$$\Sigma_m := \inf\{t > \Gamma_m \mid \varphi(t) \in \mathcal{S}_-\}, \quad (3.9)$$

Using the law of total probability, we may write (3.4) with $i \in \mathcal{S}_+, j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$, as

$$\sum_{m=0}^{\infty} \mu_{m,+,+}^{\ell_0}(t)(dx, j; x_0, i), \quad (3.10)$$

where we define $\mu_{m,+,+}^{\ell_0}(t)(\cdot, j; x_0, i)$, $m \geq 0$, to be the measures on \mathcal{D}_{ℓ_0}

$$\mathbb{P}(X(t) \in \cdot, t < \tau_1^X, \varphi(t) = j, \Gamma_m \leq t < \Sigma_{m+1} \mid X(0) = x_0, \varphi(0) = i), \quad (3.11)$$




Figure 3.1: Sample paths corresponding to the summands in (3.10).

$x_0 \in \mathcal{D}_{\ell_0}$. For $m \geq 0$, define analogously

$$\begin{aligned} \mu_{m,+,-}^{\ell_0}(t)(\cdot, j; x_0, i) \\ = \mathbb{P}(X(t) \in \cdot, t < \tau_1^X, \varphi(t) = j, \Sigma_{m+1} \leq t < \Gamma_{m+1} \mid X(0) = x_0, \varphi(0) = i) \end{aligned} \quad (3.12)$$

for $i \in \mathcal{S}_+$, $j \in \mathcal{S}_- \cup \mathcal{S}_{-0}$,

$$\begin{aligned} \mu_{m,-,+}^{\ell_0}(t)(\cdot, j; x_0, i) \\ = \mathbb{P}(X(t) \in \cdot, t < \tau_1^X, \varphi(t) = j, \Gamma_{m+1} \leq t < \Sigma_{m+1} \mid X(0) = x_0, \varphi(0) = i) \end{aligned} \quad (3.13)$$

for $i \in \mathcal{S}_-$, $j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$, and

$$\mu_{m,-,-}^{\ell_0}(t)(\cdot, j; x_0, i) = \mathbb{P}(X(t) \in \cdot, t < \tau_1^X, \varphi(t) = j, \Sigma_m \leq t < \Gamma_{m+1} \mid X(0) = x_0, \varphi(0) = i) \quad (3.14)$$

for $i \in \mathcal{S}_-$, $j \in \mathcal{S}_- \cup \mathcal{S}_{-0}$. Furthermore, let

$$\mu_{+,+}^{\ell_0}(t)(dx, j; x_0, i) := \sum_{m=0}^{\infty} \mu_{m,+,-}^{\ell_0}(t)(dx, j; x_0, i) \quad i \in \mathcal{S}_+, j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}, \quad (3.15)$$

$$\mu_{+,-}^{\ell_0}(t)(dx, j; x_0, i) := \sum_{m=1}^{\infty} \mu_{m,+,-}^{\ell_0}(t)(dx, j; x_0, i) \quad i \in \mathcal{S}_+, j \in \mathcal{S}_- \cup \mathcal{S}_{-0}, \quad (3.16)$$

$$\mu_{-,+}^{\ell_0}(t)(dx, j; x_0, i) := \sum_{m=1}^{\infty} \mu_{m,-,+}^{\ell_0}(t)(dx, j; x_0, i) \quad i \in \mathcal{S}_-, j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}, \quad (3.17)$$

$$\mu_{-,-}^{\ell_0}(t)(dx, j; x_0, i) := \sum_{m=0}^{\infty} \mu_{m,-,-}^{\ell_0}(t)(dx, j; x_0, i) \quad i \in \mathcal{S}_-, j \in \mathcal{S}_- \cup \mathcal{S}_{-0}. \quad (3.18)$$

3.1.2 Characterising the QBD-RAP

Let $\tau_1^{(p)}$ be the random (stopping) time at which the QBD-RAP changes level, or hits the boundary, or exits a boundary, for the first time;

$$\tau_1^{(p)} = \inf \{t > 0 \mid L^{(p)}(t) \neq L^{(p)}(0)\}.$$

According to the approximation described in Chapter 2.1, the summands in (3.10) are approximated by

$$f_{m,+,+}^{\ell_0,(p)}(t)(x, j; x_0, i) \, dx,$$

for $i \in \mathcal{S}_+$, $j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$, $x_0 \in \mathcal{D}_{\ell_0,i}$, $x \in \mathcal{D}_{\ell_0,j}$, $t \geq 0$, $\ell_0 \in \{0, \dots, K\}$, $m \geq 0$, which are given by

$$\begin{aligned} & \int_{\mathbf{a} \in \mathcal{A}^{(p)}} \mathbb{P} \left(\mathbf{A}^{(p)}(t) \in d\mathbf{a}, t < \tau_1^{(p)}, \varphi(t) = j, \Gamma_m \leq t < \Sigma_{m+1} \mid \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0,i}^{(p)}(x_0), \varphi(0) = i \right) \\ & \quad \times \mathbf{a} V^{(p)}(y_{\ell_0+1} - x) \, dx \\ &= \int_{\sigma_1=0}^t (\mathbf{e}_i \otimes \mathbf{a}_{\ell_0,i}^{(p)}(x_0)) e^{\mathbf{B}_{++}^{(p)} \sigma_1} \mathbf{B}_{+-}^{(p)} \int_{\gamma_1=\sigma_1}^t e^{\mathbf{B}_{--}^{(p)}(\gamma_1-\sigma_1)} \mathbf{B}_{-+}^{(p)} \dots \int_{\gamma_m=\sigma_m}^t e^{\mathbf{B}_{--}^{(p)}(\gamma_m-\sigma_m)} \mathbf{B}_{-+}^{(p)} \\ & \quad \times e^{\mathbf{B}_{++}^{(p)}(t-\gamma_m)} (\mathbf{e}_j \otimes V^{(p)}(y_{\ell_0+1} - x)) \, dx \, d\sigma_1 \, d\gamma_1 \dots d\sigma_m \, d\gamma_m. \end{aligned} \quad (3.19)$$

Analogously, approximations to (3.12)-(3.14) are

$$\begin{aligned} & f_{m,+,-}^{\ell_0,(p)}(t)(x, j; x_0, i) \\ &:= \int_{\sigma_1=0}^t (\mathbf{e}_i \otimes \mathbf{a}_{\ell_0,i}^{(p)}(x_0)) e^{\mathbf{B}_{++}^{(p)} \sigma_1} \mathbf{B}_{+-}^{(p)} \int_{\gamma_1=\sigma_1}^t e^{\mathbf{B}_{--}^{(p)}(\gamma_1-\sigma_1)} \mathbf{B}_{-+}^{(p)} \dots \int_{\sigma_{m+1}=\gamma_m}^t e^{\mathbf{B}_{++}^{(p)}(\sigma_{m+1}-\gamma_m)} \mathbf{B}_{+-}^{(p)} \\ & \quad \times e^{\mathbf{B}_{--}^{(p)}(t-\sigma_{m+1})} (\mathbf{e}_j \otimes V^{(p)}(x - y_{\ell_0})) \, dx \, d\sigma_1 \, d\gamma_1 \dots d\sigma_m \, d\gamma_m \, d\sigma_{m+1} \end{aligned} \quad (3.20)$$

for $i \in \mathcal{S}_+$, $j \in \mathcal{S}_- \cup \mathcal{S}_{-0}$;

$$\begin{aligned} & f_{m,-,+}^{\ell_0,(p)}(t)(x, j; x_0, i) \\ &:= \int_{\gamma_1=0}^t (\mathbf{e}_i \otimes \mathbf{a}_{\ell_0,i}^{(p)}(x_0)) e^{\mathbf{B}_{--}^{(p)} \gamma_1} \mathbf{B}_{-+}^{(p)} \int_{\sigma_1=\gamma_1}^t e^{\mathbf{B}_{++}^{(p)}(\sigma_1-\gamma_1)} \mathbf{B}_{+-}^{(p)} \dots \int_{\gamma_{m+1}=\sigma_m}^t e^{\mathbf{B}_{--}^{(p)}(\gamma_{m+1}-\sigma_m)} \mathbf{B}_{-+}^{(p)} \\ & \quad \times e^{\mathbf{B}_{++}^{(p)}(t-\gamma_{m+1})} (\mathbf{e}_j \otimes V^{(p)}(y_{\ell_0+1} - x)) \, dx \, d\gamma_1 \, d\sigma_1 \dots d\gamma_m \, d\sigma_m \, d\gamma_{m+1} \end{aligned} \quad (3.21)$$

for $i \in \mathcal{S}_-$, $j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$;

$$\begin{aligned} & f_{m,-,-}^{\ell_0,(p)}(t)(x, j; x_0, i) \\ &:= \int_{\gamma_1=0}^t (\mathbf{e}_i \otimes \mathbf{a}_{\ell_0,i}^{(p)}(x_0)) e^{\mathbf{B}_{--}^{(p)} \gamma_1} \mathbf{B}_{-+}^{(p)} \int_{\sigma_1=\gamma_1}^t e^{\mathbf{B}_{++}^{(p)}(\sigma_1-\gamma_1)} \mathbf{B}_{+-}^{(p)} \dots \int_{\sigma_m=\gamma_m}^t e^{\mathbf{B}_{++}^{(p)}(\sigma_m-\gamma_m)} \mathbf{B}_{+-}^{(p)} \end{aligned}$$

$$\times e^{\mathbf{B}_{--}^{(p)}(t-\sigma_m)} (\mathbf{e}_j \otimes V^{(p)}(x - y_{\ell_0})) \, dx \, d\gamma_1 \, d\sigma_1 \dots d\gamma_m \, d\sigma_m \quad (3.22)$$

for $i \in \mathcal{S}_-$, $j \in \mathcal{S}_- \cup \mathcal{S}_{-0}$, respectively.

For sample paths with $\varphi(0) \in \mathcal{S}_+$, ($\varphi(0) \in \mathcal{S}_-$) the events $\{\Gamma_m \leq t < \Sigma_{m+1}\}$, and $\{\Sigma_{m+1} \leq t < \Gamma_{m+1}\}$, (respectively, $\{\Sigma_m \leq t < \Gamma_{m+1}\}$, and $\{\Gamma_{m+1} \leq t < \Sigma_{m+1}\}$) $m \geq 0$, form a partition of the sample paths of the QBD-RAP. With this partition the phase of the QBD-RAP changes from \mathcal{S}_+ to $j_m \in \mathcal{S}_-$ at times Σ_m and from \mathcal{S}_- to $k_m \in \mathcal{S}_+$ at times Γ_m .

Define

$$\begin{aligned} f_{+,+}^{\ell_0,(p)}(t)(x, j; x_0, i) \, dx &:= \sum_{m=0}^{\infty} f_{m,+,+}^{\ell_0,(p)}(t)(x, j; x_0, i) \, dx & i \in \mathcal{S}_+, j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}, \\ f_{+,-}^{\ell_0,(p)}(t)(x, j; x_0, i) \, dx &:= \sum_{m=1}^{\infty} f_{m,+,-}^{\ell_0,(p)}(t)(x, j; x_0, i) \, dx & i \in \mathcal{S}_+, j \in \mathcal{S}_0 \cup \mathcal{S}_{-0}, \\ f_{-,+}^{\ell_0,(p)}(t)(x, j; x_0, i) &:= \sum_{m=1}^{\infty} f_{m,-,+}^{\ell_0,(p)}(t)(x, j; x_0, i) \, dx & i \in \mathcal{S}_-, j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}, \\ f_{-,-}^{\ell_0,(p)}(t)(x, j; x_0, i) \, dx &:= \sum_{m=0}^{\infty} f_{m,-,-}^{\ell_0,(p)}(t)(x, j; x_0, i) \, dx & i \in \mathcal{S}_-, j \in \mathcal{S}_- \cup \mathcal{S}_{-0}. \end{aligned}$$

which are the corresponding approximations of (3.11)-(3.18).

3.2 Laplace transforms with respect to time on no change of level

For $\lambda \geq 0$, define the matrices

$$\begin{aligned} \mathbf{Q}_{+0}(\lambda) &= \mathbf{C}_+^{-1} \mathbf{T}_{+0} [\lambda \mathbf{I} - \mathbf{T}_{00}]^{-1}, \\ \mathbf{Q}_{-0}(\lambda) &= \mathbf{C}_-^{-1} \mathbf{T}_{-0} [\lambda \mathbf{I} - \mathbf{T}_{00}]^{-1}, \\ \mathbf{Q}_{++}(\lambda) &= \mathbf{C}_+^{-1} (\mathbf{T}_{++} - \lambda \mathbf{I} + \mathbf{T}_{+0} [\lambda \mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0+}), \\ \mathbf{Q}_{+-}(\lambda) &= \mathbf{C}_+^{-1} (\mathbf{T}_{+-} + \mathbf{T}_{+0} [\lambda \mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0-}), \\ \mathbf{Q}_{--}(\lambda) &= \mathbf{C}_-^{-1} (\mathbf{T}_{--} - \lambda \mathbf{I} + \mathbf{T}_{-0} [\lambda \mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0-}), \\ \mathbf{Q}_{-+}(\lambda) &= \mathbf{C}_-^{-1} (\mathbf{T}_{-+} + \mathbf{T}_{-0} [\lambda \mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0+}), \end{aligned}$$

and the functions,

$$\mathbf{H}^{++}(\lambda, x) := e^{\mathbf{Q}_{++}(\lambda)x} [\mathbf{C}_+^{-1} \quad \mathbf{Q}_{+0}(\lambda)] = [h_{ij}^{++}(\lambda, x)]_{i \in \mathcal{S}_+, j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}}, \quad (3.23)$$

$$\mathbf{H}^{--}(\lambda, x) := e^{\mathbf{Q}_{--}(\lambda)x} [\mathbf{C}_-^{-1} \quad \mathbf{Q}_{-0}(\lambda)] = [h_{ij}^{--}(\lambda, x)]_{i \in \mathcal{S}_-, j \in \mathcal{S}_- \cup \mathcal{S}_{-0}}, \quad (3.24)$$

$$\mathbf{H}^{+-}(\lambda, x) := e^{\mathbf{Q}_{++}(\lambda)x} \mathbf{Q}_{+-}(\lambda) = [h_{ij}^{+-}(\lambda, x)]_{i \in \mathcal{S}_+, j \in \mathcal{S}_-}, \quad (3.25)$$

$$\mathbf{H}^{-+}(\lambda, x) := e^{\mathbf{Q}_{--}(\lambda)x} \mathbf{Q}_{-+}(\lambda) = [h_{ij}^{-+}(\lambda, x)]_{i \in \mathcal{S}_-, j \in \mathcal{S}_+}, \quad (3.26)$$

for $x, \lambda \geq 0$. The function $h_{ij}^{++}(\lambda, x)$ ($h_{ij}^{--}(\lambda, x)$) is the Laplace transform with respect to time of the time taken for the fluid level to shift by an amount x whilst remaining in phases in $\mathcal{S}_+ \cup \mathcal{S}_{+0}$ ($\mathcal{S}_- \cup \mathcal{S}_{-0}$), given the phase was initially $i \in \mathcal{S}_+$ ($i \in \mathcal{S}_-$) (Bean et al. 2005b). The function $h_{ij}^{+-}(\lambda, x)$ ($h_{ij}^{-+}(\lambda, x)$) is the Laplace transform with respect to time of the time taken for the fluid level, $\{X(t)\}$ to shift by an amount x whilst remaining in phases in $\mathcal{S}_+ \cup \mathcal{S}_{+0}$ ($\mathcal{S}_- \cup \mathcal{S}_{-0}$), after which time the phase instantaneously changes to $j \in \mathcal{S}_-$ (\mathcal{S}_+), given the phase was initially $i \in \mathcal{S}_+$ (\mathcal{S}_-) (Bean et al. 2005b).

Let $\psi : [0, \Delta] \rightarrow \mathbb{R}$ be bounded and Lipschitz continuous and consider expectations with respect to measures (3.11)-(3.14). For example,

$$\int_{x=0}^{\Delta} \mu_{m,+,+}^{\ell_0}(t)(y_{\ell_0} + dx, j; x_0, i) \psi(x) dx. \quad (3.27)$$

Consider taking the Laplace transform with respect to time of (3.27);

$$\begin{aligned} & \int_{t=0}^{\infty} e^{-\lambda t} \int_{x=0}^{\Delta} \mu_{m,+,+}^{\ell_0}(t)(y_{\ell_0} + dx, j; x_0, i) \psi(x) dt \\ &= \int_{x=0}^{\Delta} \int_{t=0}^{\infty} e^{-\lambda t} \mu_{m,+,+}^{\ell_0}(t)(y_{\ell_0} + dx, j; x_0, i) dt \psi(x) \\ &= \int_{x=0}^{\Delta} \hat{\mu}_{m,+,+}^{\ell_0}(\lambda)(y_{\ell_0} + x, j; x_0, i) \psi(x) dx \end{aligned} \quad (3.28)$$

where we use $\hat{\mu}_{m,+,+}^{\ell_0}(\lambda)(y_{\ell_0} + dx, j; x_0, i)$ to denote the Laplace transform with respect to time of (3.11). We now proceed to derive an expression for the Laplace transform with respect to time, $\hat{\mu}_{m,+,+}^{\ell_0}(\lambda)(dx, j; x_0, i)$, from which an expression for (3.28) follows. We use the hat $\hat{}$ notation to denote Laplace transforms with respect to time.

From the stochastic interpretations of the Laplace transforms (3.23)-(3.26) given in (Bean et al. 2005b) and summarised above, the Laplace transforms with respect to time, $\hat{\mu}_{m,+,+}^{\ell_0}(\lambda)(x, j; x_0, i)$, of (3.11) are given by

$$\hat{\mu}_{0,+,+}^{\ell_0}(\lambda)(x, j; x_0, i) dx = h_{ij}^{++}(\lambda, x - x_0) 1(x \geq x_0) dx,$$

for $m = 0$, and

$$\begin{aligned} & \int_{x_1=0}^{\Delta - (x_0 - y_{\ell_0})} \mathbf{e}_i \mathbf{H}^{+-}(\lambda, \Delta - (x_0 - y_{\ell_0}) - x_1) \\ & \times \left[\prod_{r=1}^{m-1} \int_{x_{2r}=0}^{\Delta - x_{2r-1}} \mathbf{H}^{-+}(\lambda, \Delta - x_{2r} - x_{2r-1}) dx_{2r-1} \int_{x_{2r+1}=0}^{\Delta - x_{2r}} \mathbf{H}^{+-}(\lambda, \Delta - x_{2r+1} - x_{2r}) dx_{2r} \right] \end{aligned}$$




Figure 3.2: Sample paths corresponding to the Laplace transforms (3.29). $z_1 = \Delta - x_1 - (x - y_{l_0})$, $z_2 = \Delta - x_2 - x_1$.

$$\begin{aligned}
& \times \int_{x_{2m}=0}^{\Delta - x_{2m-1}} \mathbf{H}^{-+}(\lambda, \Delta - x_{2m-1} - x_{2m}) dx_{2m-1} \mathbf{H}^{++}(\lambda, \Delta - x_{2m} - (y_{l_0+1} - x)) \mathbf{e}_j \\
& \times 1(\Delta - x_{2m} - (y_{l_0+1} - x) \geq 0) dx_{2m} dx
\end{aligned} \tag{3.29}$$

for $m \geq 1$. Figure 3.2 shows an example of the sample paths to which these Laplace transforms correspond. Analogously, we can write down similar expressions for the Laplace transform with respect to time of (3.11)-(3.14).

Observe that $\hat{\mu}_{0,+,+}^{\ell_0}(\lambda)(x, j; x_0, i)$ is a smooth function in the variable $x > x_0$ (and $\lambda > 0$ too), even though the measure $\mu_{0,+,+}^{\ell_0}(t)(dx, j; x_0, i)$ may have point masses. For example, given $X(0) = x_0$ and $\varphi(0) = i$, then $\mu_{0,+,+}^{\ell_0}(t)(dx, j; x_0, i)$ has a point mass of size (at least) $e^{T_{ii}t} \delta(x - (x_0 + c_i t)) 1(j = i) 1(x \in \mathcal{D}_{\ell_0, i})$ which arises from the event that the phase remains in i until time t . Meanwhile, the Laplace transform $\hat{\mu}_{0,+,+}^{\ell_0}(\lambda)(dx, j; x_0, i)$ is given by $[e^{\mathbf{Q}_{++}(\lambda)(x-x_0)} 1(x \geq x_0)]_{ij}$, which is a smooth function for all x except at $x = x_0$.

Now consider expectations with respect to measures (3.19)-(3.22). For example,

$$\int_{x=0}^{\Delta} f_{m,+,+}^{\ell_0, (p)}(t)(y_{l_0} + dx, j; x_0, i) \psi(x). \tag{3.30}$$

Taking the Laplace transform with respect to time of (3.30);

$$\begin{aligned}
& \int_{t=0}^{\infty} e^{-\lambda t} \int_{x=0}^{\Delta} f_{m,+,+}^{\ell_0, (p)}(t)(y_{l_0} + x, j; x_0, i) \psi(x) dx dt \\
& = \int_{x=0}^{\Delta} \int_{t=0}^{\infty} e^{-\lambda t} f_{m,+,+}^{\ell_0, (p)}(t)(y_{l_0} + x, j; x_0, i) dt \psi(x) dx
\end{aligned}$$

$$= \int_{x=0}^{\Delta} \widehat{f}_{m,+,+}^{\ell_0,(p)}(\lambda)(y_{\ell_0} + x, j; x_0, i) \psi(x) dx \quad (3.31)$$

where we use $\widehat{f}_{m,+,+}^{\ell_0,(p)}(\lambda)(y_{\ell_0} + x, j; x_0, i)$ to denote the Laplace transform with respect to time of (3.19). We now proceed to derive an expression for the Laplace transform with respect to time, $\widehat{f}_{m,+,+}^{\ell_0,(p)}(\lambda)(dx, j; x_0, i)$, from which an expression for (3.31) follows.

Notice that (3.19) is a convolution. The Laplace transform with respect to time of (3.19) is

$$\begin{aligned} & \widehat{f}_{m,+,+}^{\ell_0,(p)}(\lambda)(x, j; x_0, i) \\ &= (\mathbf{e}_i \otimes \mathbf{a}_{\ell_0,i}^{(p)}(x_0)) \int_{t_1=0}^{\infty} e^{-\lambda t_1} e^{\mathbf{B}_{++}^{(p)} t_1} dt_1 \mathbf{B}_{+-}^{(p)} \int_{t_2=0}^{\infty} e^{-\lambda t_2} e^{\mathbf{B}_{--}^{(p)} t_2} dt_2 \mathbf{B}_{-+}^{(p)} \\ & \dots \int_{t_{2m}=0}^{\infty} e^{-\lambda t_{2m}} e^{\mathbf{B}_{--}^{(p)} t_{2m}} dt_{2m} \mathbf{B}_{-+}^{(p)} \int_{t=0}^{\infty} e^{-\lambda t} e^{\mathbf{B}_{++}^{(p)} t} dt (\mathbf{e}_j \otimes V^{(p)}(y_{\ell_0+1} - x)). \end{aligned} \quad (3.32)$$

Analogous expressions can be computed for the Laplace transforms with respect to time of (3.20)-(3.22).

In Corollary A.25 in Appendix A.3, we show the following relation

$$\begin{aligned} & [\mathbf{I}_{p|S_m|} \quad \mathbf{0}_{p|S_m| \times p|S_0|}] \int_{t=0}^{\infty} e^{-\lambda t} e^{\mathbf{B}_{mm}^{(p)} t} dt \mathbf{B}_{mn}^{(p)} \\ &= \int_{x=0}^{\infty} \mathbf{H}^{mn}(\lambda, x) \otimes e^{\mathbf{S}^{(p)} x} \mathbf{D}^{(p)} dx [\mathbf{I}_{p|S_n|} \quad \mathbf{0}_{p|S_n| \times p|S_0|}], \end{aligned} \quad (3.33)$$

for $m, n \in \{+, -\}$, $m \neq n$. Before we can apply this result, observe that we can write the initial vector in (3.32) as

$$((\mathbf{e}_i)_{1 \times |S_+ \cup S_0|} \otimes \mathbf{a}_{\ell_0,i}^{(p)}(x_0)) = ((\mathbf{e}_i)_{1 \times |S_+|} \otimes \mathbf{a}_{\ell_0,i}^{(p)}(x_0)) [\mathbf{I}_{p|S_+|} \quad \mathbf{0}_{p|S_+| \times p|S_0|}]$$

by the Mixed Product Rule. Now, applying (3.33) to the first integral in (3.32) transforms the expression to

$$\begin{aligned} & (\mathbf{e}_i \otimes \mathbf{a}_{\ell_0,i}^{(p)}(x_0)) \int_{x_1=0}^{\infty} \left(\mathbf{H}^{+-}(\lambda, x_1) \otimes e^{\mathbf{S}^{(p)} x} \mathbf{D}^{(p)} \right) dx_1 [\mathbf{I}_{p|S_-|} \quad \mathbf{0}_{p|S_-| \times p|S_0|}] \int_{t_2=0}^{\infty} e^{-\lambda t_2} e^{\mathbf{B}_{--}^{(p)} t_2} dt_2 \mathbf{B}_{-+}^{(p)} \\ & \dots \int_{t_{2m}=0}^{\infty} e^{-\lambda t_{2m}} e^{\mathbf{B}_{--}^{(p)} t_{2m}} dt_{2m} \mathbf{B}_{-+}^{(p)} \int_{t=0}^{\infty} e^{-\lambda t} e^{\mathbf{B}_{++}^{(p)} t} dt (\mathbf{e}_j \otimes V^{(p)}(y_{\ell_0+1} - x)). \end{aligned} \quad (3.34)$$

We may now apply (3.33) to the second integral, after which we can apply (3.33) to the third integral and so on. Ultimately, after applying (3.33) to all of the integrals in (3.32), we get

$$(\mathbf{e}_i \otimes \mathbf{a}_{\ell_0,i}^{(p)}(x_0)) \left(\int_{x_1=0}^{\infty} \mathbf{H}^{+-}(\lambda, x_1) \otimes e^{\mathbf{S}^{(p)} x_1} \mathbf{D}^{(p)} dx_1 \right)$$

$$\begin{aligned}
& \left[\prod_{r=1}^{m-1} \left(\int_{x_{2r}=0}^{\infty} \mathbf{H}^{-+}(\lambda, x_{2r}) \otimes e^{\mathbf{S}^{(p)} x_{2r}} \mathbf{D}^{(p)} dx_{2r} \right) \right. \\
& \quad \left. \left(\int_{x_{2r+1}=0}^{\infty} \mathbf{H}^{+-}(\lambda, x_{2r+1}) \otimes e^{\mathbf{S}^{(p)} x_{2r+1}} \mathbf{D}^{(p)} dx_{2r+1} \right) \right] \\
& \quad \left(\int_{x_{2m}=0}^{\infty} \mathbf{H}^{-+}(\lambda, x_{2m}) \otimes e^{\mathbf{S}^{(p)} x_{2m}} \mathbf{D}^{(p)} dx_{2m} \right) \\
& \quad \left(\int_{x_{2m+1}=0}^{\infty} \mathbf{H}^{++}(\lambda, x_{2m+1}) \otimes e^{\mathbf{S}^{(p)} x_{2m+1}} dx_{2m+1} \right) (\mathbf{e}_j \otimes V^{(p)}(y_{\ell_0+1} - x)) \\
& = \int_{x_1=0}^{\infty} \cdots \int_{x_{2m+1}=0}^{\infty} \mathbf{e}_i \mathbf{M}_{++}(\lambda, x_1, \dots, x_{2m+1}) \mathbf{e}_j \\
& \quad \times \boldsymbol{\alpha}_{\ell_0, i}^{(p)}(x_0) \mathbf{N}^{(p)}(\lambda, x_1, \dots, x_{2m+1}) V^{(p)}(y_{\ell_0+1} - x) dx_{2m+1} \dots dx_1, \tag{3.35}
\end{aligned}$$

where we define matrices

$$\begin{aligned}
& \mathbf{M}_{++}(\lambda, x_1, \dots, x_{2m+1}) \\
& = \mathbf{H}^{+-}(\lambda, x_1) \prod_{r=1}^{m-1} \mathbf{H}^{-+}(\lambda, x_{2r}) \mathbf{H}^{+-}(\lambda, x_{2r+1}) \mathbf{H}^{-+}(\lambda, x_{2m}) \mathbf{H}^{++}(\lambda, x_{2m+1})
\end{aligned}$$

and

$$\mathbf{N}^{(p)}(\lambda, x_1, \dots, x_{2m+1}) = e^{\mathbf{S}^{(p)} x_1} \mathbf{D}^{(p)} \prod_{r=1}^{m-1} e^{\mathbf{S}^{(p)} x_{2r}} \mathbf{D}^{(p)} e^{\mathbf{S}^{(p)} x_{2r+1}} \mathbf{D}^{(p)} e^{\mathbf{S}^{(p)} x_{2m}} \mathbf{D}^{(p)} e^{\mathbf{S}^{(p)} x_{2m+1}}.$$

The relation (3.33) is key to our analysis. It allows us to factorise the integrand of the Laplace transform (3.35) into one factor solely related to the orbit process $\{\mathbf{A}^{(p)}(t)\}$ and another factor solely related to the fluid queue.

Further, if we define matrices

$$\begin{aligned}
\mathbf{M}_{-+}(\lambda, x_1, \dots, x_{2m}) & = \mathbf{H}^{-+}(\lambda, x_1) \prod_{r=1}^{m-1} \mathbf{H}^{+-}(\lambda, x_{2r}) \mathbf{H}^{-+}(\lambda, x_{2r+1}) \mathbf{H}^{++}(\lambda, x_{2m}), \\
\mathbf{M}_{+-}(\lambda, x_1, \dots, x_{2m}) & = \mathbf{H}^{+-}(\lambda, x_1) \prod_{r=1}^{m-1} \mathbf{H}^{-+}(\lambda, x_{2r}) \mathbf{H}^{+-}(\lambda, x_{2r+1}) \mathbf{H}^{--}(\lambda, x_{2m}),
\end{aligned}$$

and

$$\begin{aligned}
& \mathbf{M}_{--}(\lambda, x_1, \dots, x_{2m+1}) \\
& = \mathbf{H}^{-+}(\lambda, x_1) \prod_{r=1}^{m-1} \mathbf{H}^{+-}(\lambda, x_{2r}) \mathbf{H}^{-+}(\lambda, x_{2r+1}) \mathbf{H}^{+-}(\lambda, x_{2m}) \mathbf{H}^{--}(\lambda, x_{2m+1}),
\end{aligned}$$

then analogous expressions can be shown for (3.20)-(3.22) in terms of these matrices; for $m \geq 0$,

$$\begin{aligned}
\widehat{f}_{m+1,-,+}^{\ell_0,(p)}(\lambda)(x,j;x_0,i) &= \int_{x_1=0}^{\infty} \cdots \int_{x_{2m+2}=0}^{\infty} \mathbf{e}_i \mathbf{M}_{-+}(\lambda, x_1, \dots, x_{2m+2}) \mathbf{e}_j \\
&\quad \times \mathbf{a}_{\ell_0,i}^{(p)}(x_0) \mathbf{N}^{(p)}(\lambda, x_1, \dots, x_{2m+2}) V^{(p)}(y_{\ell_0+1} - x) dx_{2m+2} \dots dx_1, \\
\widehat{f}_{m+1,+,-}^{\ell_0,(p)}(\lambda)(x,j;x_0,i) &= \int_{x_1=0}^{\infty} \cdots \int_{x_{2m+2}=0}^{\infty} \mathbf{e}_i \mathbf{M}_{+-}(\lambda, x_1, \dots, x_{2m+2}) \mathbf{e}_j \\
&\quad \times \mathbf{a}_{\ell_0,i}^{(p)}(x_0) \mathbf{N}^{(p)}(\lambda, x_1, \dots, x_{2m+2}) V^{(p)}(y_{\ell_0+1} - x) dx_{2m+2} \dots dx_1, \\
\widehat{f}_{m,-,-}^{\ell_0,(p)}(\lambda)(x,j;x_0,i) &= \int_{x_1=0}^{\infty} \cdots \int_{x_{2m+1}=0}^{\infty} \mathbf{e}_i \mathbf{M}_{--}(\lambda, x_1, \dots, x_{2m+2}) \mathbf{e}_j \\
&\quad \times \mathbf{a}_{\ell_0,i}^{(p)}(x_0) \mathbf{N}^{(p)}(\lambda, x_1, \dots, x_{2m+1}) V^{(p)}(y_{\ell_0+1} - x) dx_{2m+1} \dots dx_1.
\end{aligned}$$

For $\varphi(0) = k \in \mathcal{S}_{-0}$ (or $\varphi(0) = k \in \mathcal{S}_{+0}$) there is a slight complexity as in this case it is possible that, upon the phase process first leaving \mathcal{S}_{-0} (\mathcal{S}_{+0}) the phase transitions to a state in \mathcal{S}_{+} (\mathcal{S}_{-}). Since the orbit of the QBD-RAP is constant on $\varphi(t) \in \mathcal{S}_{-0}$ ($\varphi(t) \in \mathcal{S}_{+0}$), then upon a first transition out of \mathcal{S}_{-0} (\mathcal{S}_{+0}) and into \mathcal{S}_{+} (\mathcal{S}_{-}) the orbit jumps to $\mathbf{a}_{\ell_0,i}^{(p)}(x_0) \mathbf{D}^{(p)}$. For $k \in \mathcal{S}_{-0}$, the corresponding Laplace transform of the QBD-RAP is

$$\begin{aligned}
&\widehat{f}_{m,-0,+}^{\ell_0,(p)}(\lambda)(x,j;x_0,k) \\
&:= \int_{x_1=0}^{\infty} \cdots \int_{x_{2m+1}=0}^{\infty} \mathbf{e}_k [\mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0+} \mathbf{M}_{++}(\lambda, x_1, \dots, x_{2m+1}) \mathbf{e}_j \\
&\quad \mathbf{a}_{\ell_0,i}^{(p)}(x_0) \mathbf{D}^{(p)} \mathbf{N}^{(p)}(\lambda, x_1, \dots, x_{2m+1}) V^{(p)}(y_{\ell_0+1} - x) dx_{2m+1} \dots dx_2 dx_1 \\
&\quad + \int_{x_1=0}^{\infty} \cdots \int_{x_{2m+1}=0}^{\infty} \mathbf{e}_k [\mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0-} \mathbf{M}_{-+}(\lambda, x_1, \dots, x_{2m+2}) \mathbf{e}_j \\
&\quad \mathbf{a}_{\ell_0,i}^{(p)}(x_0) \mathbf{N}^{(p)}(\lambda, x_1, \dots, x_{2m+2}) V^{(p)}(y_{\ell_0+1} - x) dx_{2m+2} \dots dx_2 dx_1. \tag{3.36}
\end{aligned}$$

The second term in expression (3.36) is a linear combination of $\widehat{f}_{m+1,-,+}^{\ell_0,(p)}(\lambda)(x,j;x_0,k)$. The first term is a linear combination of terms similar to $\widehat{f}_{m,+,-}^{\ell_0,(p)}(\lambda)(x,j;x_0,k)$, except there is a different initial vector, $\mathbf{a}_{\ell_0,i}^{(p)}(x_0) \mathbf{D}^{(p)}$. The Laplace transform of the fluid queue corresponding to (3.36) is

$$\begin{aligned}
\widehat{\mu}_{m,-0,+}^{\ell_0}(\lambda)(x,j;x_0,k) &:= \sum_{i \in \mathcal{S}_{+}} \mathbf{e}_k [\lambda \mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0i} \widehat{\mu}_{m,+,-}^{\ell_0}(\lambda)(x,j;x_0,i) \\
&\quad + \sum_{i \in \mathcal{S}_{-}} \mathbf{e}_k [\lambda \mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0i} \widehat{\mu}_{m+1,-,+}^{\ell_0}(\lambda)(x,j;x_0,i). \tag{3.37}
\end{aligned}$$

Since the second term is a linear combination of $\widehat{f}_{m+1,-,+}^{\ell_0,(p)}(\lambda)(x, j; x_0, k)$, then once we argue that $\widehat{f}_{m+1,-,+}^{\ell_0,(p)}(\lambda)(x, j; x_0, k)$ converges to $\widehat{\mu}_{m,+}^{\ell_0}(\lambda)(x, j; x_0, i)$, then we get that the second term of (3.36) converges to the second term of (3.37). The first term of (3.36) creates a bit more work regarding the analysis. Similar comments apply for all sample paths starting in \mathcal{S}_{-0} of \mathcal{S}_{+0} .

Analogously, for $k \in \mathcal{S}_{-0}$ we also have

$$\begin{aligned} \widehat{f}_{m,-0,-}^{\ell_0,(p)}(\lambda)(x, j; x_0, k) &:= \int_{x_1=0}^{\infty} \cdots \int_{x_{2m+2}=0}^{\infty} \mathbf{e}_k [\mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0+} \mathbf{M}_{+-}(\lambda, x_1, \dots, x_{2m+2}) \mathbf{e}_j \\ &\quad \mathbf{a}_{\ell_0,i}^{(p)}(x_0) \mathbf{D}^{(p)} \mathbf{N}^{(p)}(\lambda, x_1, \dots, x_{2m+2}) V^{(p)}(y_{\ell_0+1} - x) dx_{2m+2} \cdots dx_1 \\ &\quad + \int_{x_1=0}^{\infty} \cdots \int_{x_{2m+1}=0}^{\infty} \mathbf{e}_k [\mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0-} \mathbf{M}_{--}(\lambda, x_1, \dots, x_{2m+1}) \mathbf{e}_j \\ &\quad \mathbf{a}_{\ell_0,i}^{(p)}(x_0) \mathbf{N}^{(p)}(\lambda, x_1, \dots, x_{2m+1}) V^{(p)}(y_{\ell_0+1} - x) dx_{2m+1} \cdots dx_1. \end{aligned}$$

For $k \in \mathcal{S}_{+0}$ we have

$$\begin{aligned} \widehat{f}_{m,+0,+}^{\ell_0,(p)}(\lambda)(x, j; x_0, k) &:= \int_{x_1=0}^{\infty} \cdots \int_{x_{2m+1}=0}^{\infty} \mathbf{e}_k [\mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0-} \mathbf{M}_{--}(\lambda, x_1, \dots, x_{2m+1}) \mathbf{e}_j \\ &\quad \mathbf{a}_{\ell_0,i}^{(p)}(x_0) \mathbf{D}^{(p)} \mathbf{N}^{(p)}(\lambda, x_1, \dots, x_{2m+1}) V^{(p)}(y_{\ell_0+1} - x) dx_{2m+1} \cdots dx_1 \\ &\quad + \int_{x_1=0}^{\infty} \cdots \int_{x_{2m+1}=0}^{\infty} \mathbf{e}_k [\mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0+} \mathbf{M}_{+-}(\lambda, x_1, \dots, x_{2m+2}) \mathbf{e}_j \\ &\quad \mathbf{a}_{\ell_0,i}^{(p)}(x_0) \mathbf{N}^{(p)}(\lambda, x_1, \dots, x_{2m+2}) V^{(p)}(y_{\ell_0+1} - x) dx_{2m+2} \cdots dx_1, \end{aligned}$$

and

$$\begin{aligned} \widehat{f}_{m,+0,-}^{\ell_0,(p)}(\lambda)(x, j; x_0, k) &:= \int_{x_1=0}^{\infty} \cdots \int_{x_{2m+1}=0}^{\infty} \mathbf{e}_k [\mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0-} \mathbf{M}_{--}(\lambda, x_1, \dots, x_{2m+1}) \mathbf{e}_j \\ &\quad \mathbf{a}_{\ell_0,i}^{(p)}(x_0) \mathbf{D}^{(p)} \mathbf{N}^{(p)}(\lambda, x_1, \dots, x_{2m+1}) V^{(p)}(y_{\ell_0+1} - x) dx_{2m+1} \cdots dx_1 \\ &\quad + \int_{x_1=0}^{\infty} \cdots \int_{x_{2m+2}=0}^{\infty} \mathbf{e}_k [\mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0+} \mathbf{M}_{+-}(\lambda, x_1, \dots, x_{2m+2}) \mathbf{e}_j \\ &\quad \mathbf{a}_{\ell_0,i}^{(p)}(x_0) \mathbf{N}^{(p)}(\lambda, x_1, \dots, x_{2m+2}) V^{(p)}(y_{\ell_0+1} - x) dx_{2m+2} \cdots dx_1. \end{aligned}$$

In general, for $p \in \{+, -\}$, $q, r \in \{+, -\}$, $q \neq r$,

$$\begin{aligned} \widehat{\mu}_{m,p0,q}^{\ell_0}(\lambda)(x, j; x_0, k) &:= \sum_{i \in \mathcal{S}_r} \mathbf{e}_k [\lambda \mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0i} \widehat{\mu}_{m,r,q}^{\ell_0}(\lambda)(x, j; x_0, i) \\ &\quad + \sum_{i \in \mathcal{S}_r} \mathbf{e}_k [\lambda \mathbf{I} - \mathbf{T}_{00}]^{-1} \mathbf{T}_{0i} \widehat{\mu}_{m+1,r,q}^{\ell_0}(\lambda)(x, j; x_0, i). \end{aligned}$$

Remark 3.3. For technical reasons we should not have point masses at $x_0 \in \partial\mathcal{D}_{\ell_0}$ when $\varphi(0) \in \mathcal{S}_{+0} \cup \mathcal{S}_{-0}$. Intuitively, if $\varphi(0) = k \in \mathcal{S}_{+0}$ and $x_0 = y_{\ell_0}$ then, if upon exiting \mathcal{S}_{+0} the phase process transitions to \mathcal{S}_{-} then the fluid queue will instantaneously leave the interval \mathcal{D}_{ℓ_0} upon this transition. On the same event, the orbit of the QBD-RAP will be $\alpha^{(p)} \mathbf{D}^{(p)}$ at the instant of the transition to \mathcal{S}_{-} . Roughly speaking $\mathbf{D}^{(p)}$ maps $\alpha^{(p)}$ to approximately $\frac{\alpha^{(p)} e^{\mathbf{S}^{(p)} \Delta}}{\alpha^{(p)} e^{\mathbf{S}^{(p)} \Delta} \mathbf{e}}$ (asymptotically). Our asymptotic arguments rely on Chebyshev's inequality, in the form of a bound in terms of the distance of the random variable $Z^{(p)} \sim ME(\alpha^{(p)}, \mathbf{S}^{(p)})$ from its mean Δ . However, we cannot use such a technique to bound terms such as $\frac{\alpha^{(p)} e^{\mathbf{S}^{(p)} \Delta}}{\alpha^{(p)} e^{\mathbf{S}^{(p)} \Delta} \mathbf{e}} e^{\mathbf{S}^{(p)} z} \mathbf{s}^{(p)}$.

In practice, it may be possible to avoid this issue by choosing the intervals $\{\mathcal{D}_\ell\}$ so that the boundaries do not align with any point masses. Another option is to append an ephemeral class of phases to the fluid queue as follows.

Suppose we wish to have a point mass at $x_0 = y_\ell$, $i \in \mathcal{S}_{+0}$, and without loss of generality, assume the point mass has total mass 1. Let \mathcal{S}_{+0}^* with rates $c_j = 0$, $j \in \mathcal{S}_{+0}^*$. These states represent a duplicated copy of \mathcal{S}_{+0} . The initial orbit representing the point mass is $\alpha^{(p)}$, $i \in \mathcal{S}_{+0}^*$. The dynamics of the QBD-RAP process regarding the ephemeral class \mathcal{S}_{+0}^* is as follows. The phase process evolves according to the sub-generator matrix \mathbf{T}_{00} whilst it remains in \mathcal{S}_{+0}^* . Meanwhile, the orbit and level remain constant. When in phase $k \in \mathcal{S}_{+0}^*$, at rate $[\mathbf{T}_{0+}]_{kj}$ the process transitions out of \mathcal{S}_{+0}^* and into phase $j \in \mathcal{S}_{+}$. Similarly, when in phase $k \in \mathcal{S}_{+0}^*$, at rate $[\mathbf{T}_{0-}]_{kj}$ the process transitions out of \mathcal{S}_{+0}^* and into phase $j \in \mathcal{S}_{-}$. Upon a transition from \mathcal{S}_{+0}^* to $j \in \mathcal{S}_{+}$ the orbit and level remain unchanged, and the process evolves as usual. Upon a transition from \mathcal{S}_{+0}^* to $j \in \mathcal{S}_{-}$ the orbit remains unchanged, but the level jumps to $\ell - 1$. Thus, the process evolves as if it has just entered level $\ell - 1$ in phase $j \in \mathcal{S}_{-}$.

3.3 Convergence on no change of level

We need the following properties of $h_{ij}^{++}(\lambda, x)$, $h_{ij}^{--}(\lambda, x)$, $h_{ij}^{+-}(\lambda, x)$, $h_{ij}^{-+}(\lambda, x)$ which follow from their interpretation as a Laplace transform of a probability distribution. Let $c_{\min} = \min_{i \in \mathcal{S}_{-} \cup \mathcal{S}_{+}} |c_i|$. For all $\lambda \geq 0$, there is some $0 \leq G < \infty$ such that

$$\begin{aligned} 0 &\leq h_{ij}^{++}(\lambda, x) \leq h_{ij}^{++}(0, x) \leq \max\{1/c_{\min}, 1\} \leq G, \quad i \in \mathcal{S}_{+}, j \in \mathcal{S}_{+} \cup \mathcal{S}_{+0}, \\ 0 &\leq h_{ij}^{--}(\lambda, x) \leq h_{ij}^{--}(0, x) \leq \max\{1/c_{\min}, 1\} \leq G, \quad i \in \mathcal{S}_{-}, j \in \mathcal{S}_{-} \cup \mathcal{S}_{-0}, \\ 0 &\leq h_{ij}^{+-}(\lambda, x) \leq h_{ij}^{+-}(0, x) \leq \max_{k, \ell} [\mathbf{Q}_{+-}(0)]_{k, \ell} \leq G, \quad i \in \mathcal{S}_{+}, j \in \mathcal{S}_{-}, \\ 0 &\leq h_{ij}^{-+}(\lambda, x) \leq h_{ij}^{-+}(0, x) \leq \max_{k, \ell} [\mathbf{Q}_{-+}(0)]_{k, \ell} \leq G, \quad i \in \mathcal{S}_{-}, j \in \mathcal{S}_{+}, \end{aligned}$$

Furthermore, there exists some $0 \leq \widehat{G} < \infty$ such that, for $i \in \mathcal{S}_+$, $j \in \mathcal{S}_{+0}$,

$$\int_{x=0}^{\infty} h_{ij}^{++}(\lambda, x) dx \leq \int_{x=0}^{\infty} h_{ij}^{++}(0, x) dx = [-\mathbf{Q}_{++}(0)^{-1} \mathbf{C}_+ \quad -\mathbf{Q}_{++}(0)^{-1} \mathbf{Q}_{+0}(0)]_{ij} \leq \widehat{G},$$

for $i \in \mathcal{S}_-$, $j \in \mathcal{S}_{-0}$,

$$\int_{x=0}^{\infty} h_{ij}^{--}(\lambda, x) dx \leq \int_{x=0}^{\infty} h_{ij}^{--}(0, x) dx = [-\mathbf{Q}_{--}(0)^{-1} \mathbf{C}_- \quad -\mathbf{Q}_{--}(0)^{-1} \mathbf{Q}_{-0}(0)]_{ij} \leq \widehat{G},$$

for $i \in \mathcal{S}_+$, $j \in \mathcal{S}_-$,

$$\int_{x=0}^{\infty} h_{ij}^{+-}(\lambda, x) dx \leq \int_{x=0}^{\infty} h_{ij}^{+-}(0, x) dx = [-\mathbf{Q}_{++}(0)^{-1} \mathbf{Q}_{+-}(0)]_{ij} \leq \widehat{G},$$

for $i \in \mathcal{S}_-$, $j \in \mathcal{S}_+$,

$$\int_{x=0}^{\infty} h_{ij}^{-+}(\lambda, x) dx \leq \int_{x=0}^{\infty} h_{ij}^{-+}(0, x) dx = [-\mathbf{Q}_{--}(0)^{-1} \mathbf{Q}_{-+}(0)]_{ij} \leq \widehat{G}.$$

Moreover, since $h_{ij}^{mn}(\lambda, x)$, $m, n \in \{+, -\}$, $i \in \mathcal{S}_m$, $j \in \mathcal{S}_n$, are matrix exponential functions with exponent matrix which is a sub-generator matrix, then for every $\lambda > 0$, $h_{ij}^{mn}(\lambda, x)$ is Lipschitz continuous with respect to x on $x \in [0, \infty)$. Therefore there exists some $0 < L < \infty$ such that $|h_{ij}^{mn}(\lambda, x) - h_{ij}^{mn}(\lambda, y)| \leq L|x - y|$, $m, n \in \{+, -\}$, $i \in \mathcal{S}_m$, $j \in \mathcal{S}_n \cup \mathcal{S}_{n0}$.

We now present a series of results which provide provide error bounds and/or show convergence of the Laplace transforms presented in the previous section.

Lemma 3.4. *Let $\psi : [0, \Delta) \rightarrow \mathbb{R}$ be bounded, $\psi(x) \leq F$, and Lipschitz. Then, for $x \in \mathcal{D}_{\ell_0, j}$, $\ell_0 \in \mathcal{K} \setminus \{-1, K + 1\}$, $\lambda > 0$,*

$$\left| \int_{x=0}^{\Delta} \widehat{f}_{0,+,+}^{\ell_0,(p)}(\lambda)(x, j; x_0, i) \psi(x) dx - \int_{x=0}^{\Delta} \widehat{\mu}_{0,+,+}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) dx \right| \leq R_{V,2}^{(p)} GF + \varepsilon^{(p)} GF, \quad (3.38)$$

and

$$\left| \int_{x=0}^{\Delta} \widehat{f}_{0,-,-}^{\ell_0,(p)}(\lambda)(x, j; x_0, i) \psi(x) dx - \int_{x=0}^{\Delta} \widehat{\mu}_{0,-,-}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) dx \right| \leq R_{V,2}^{(p)} GF + \varepsilon^{(p)} GF. \quad (3.39)$$

Corollary 3.5. *Let $\psi : \mathcal{D}_{\ell_0} \rightarrow \mathbb{R}$ be bounded, $|\psi(x)| \leq F$, and Lipschitz continuous. Then, for $x_0 \in (y_{\ell_0}, y_{\ell_0+1})$, $k \in \mathcal{S}_{0+}$, $i \in \mathcal{S}_-$, $j \in \mathcal{S}_- \cup \mathcal{S}_{-0}$, there exists $r_{10}^{(p)} \rightarrow 0$ as $p \rightarrow \infty$,*

$$\left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{0,-0,-}^{\ell_0,(p)}(x, i, j; j, x_0) \psi(x) dx \rightarrow \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{0,-0,-}^{\ell_0}(x, i, j; k, x_0) \psi(x) dx \right| \leq r_{10}^{(p)}. \quad (3.40)$$

Similarly, for $k \in \mathcal{S}_{0-}$, $i \in \mathcal{S}_+$, $j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$

$$\left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{0,+0,+}^{\ell_0,(p)}(x, i, j; j, x_0) \psi(x) dx - \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{0,+0,+}^{\ell_0}(x, i, j; k, x_0) \psi(x) dx \right| \leq r_{10}^{(p)}, \quad (3.41)$$

where $r_{10}^{(p)} \rightarrow 0$ as $p \rightarrow \infty$.

Proof. The result follows from (A.5) in Appendix A. \square

Define

$$w_n^{(p)}(x_0, x) = \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}^{(p)}(x_0) e^{\mathbf{S}^{(p)} x_1} dx_1 \mathbf{D}^{(p)} \left[\prod_{k=2}^{n-1} \int_{x_k=0}^{\infty} g_k(x_k) e^{\mathbf{S}^{(p)} x_k} dx_k \mathbf{D}^{(p)} \right] \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}^{(p)} x_n} dx_n V^{(p)}(x), \quad (3.42)$$

where g_1, g_2, \dots , are functions satisfying the Assumptions 3.1 and $V^{(p)}(x)$ is a closing operator with the Properties 3.2. Expressions such as (3.32) are specific forms of (3.42). In Appendix A.1, we prove the following results.

Corollary 3.6. *Let g_1, g_2, \dots , be functions satisfying Assumptions 3.1 and let $V(x)$, $x \in [0, \Delta)$, be a closing operator with Properties 3.2. Then, for $n \geq 2$, $x_0 \in [0, \Delta)$,*

$$\begin{aligned} & \left| \int_{x=0}^{\Delta} w_n^{(p)}(x_0, x) \psi(x) dx - \int_{x=0}^{\Delta} \int_{u_1=0}^{\Delta-x_0} g_1(\Delta - u_1 - x_0) \right. \\ & \quad \left. \left[\prod_{k=2}^{n-1} \int_{u_k=0}^{\Delta-u_{k-1}} g_k(\Delta - u_k - u_{k-1}) du_{k-1} \right] g_n(\Delta - x - u_{n-1}) \right. \\ & \quad \left. 1(\Delta - x - u_{n-1} \geq 0) du_{n-1} \psi(x) dx \right| \\ & \leq (|r_5^{(p)}(n)| + |r_6^{(p)}(n)| + (n-1)|r_4^{(p)}(n)|) \Delta F, \end{aligned} \quad (3.43)$$

where

$$\begin{aligned} |r_4^{(p)}(n)| &= \left(2\varepsilon^{(p)} + \frac{\text{Var}(Z^{(p)})}{\varepsilon^{(p)}} \right) \frac{1}{1 - \text{Var}(Z^{(p)})/(\Delta - x_0)} G \widehat{G}^{n-2} G, \\ |r_5(n)| &= O \left(\max \left\{ G^{n-1} \Delta^{n-2} \left(\frac{1}{2} \Delta |r_2| + 2\varepsilon^{(p)} G + \frac{1}{2} \Delta G \frac{\text{Var}(Z^{(p)})/(\varepsilon^{(p)})^2}{1 - \text{Var}(Z^{(p)})/(\varepsilon^{(p)})^2} \right), \right. \right. \\ & \quad \left. \left. G^{n-1} \Delta^{n-2} R_{V,1}^{(p)} \right\} \right) \\ |r_6^{(p)}(n)| &\leq (\varepsilon^{(p)})^{n-1} G^n. \end{aligned}$$

Corollary 3.7. *Let g_1, g_2, \dots , be functions satisfying Assumptions 3.1 and let $V^{(p)}(x)$, $x \in (0, \Delta)$, be a closing operator with Properties 3.2. For $x_0, x \in (0, \Delta)$, $n \geq 2$*

$$\left| \int_{x \in [0, \Delta)} \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}^{(p)}(x_0) \mathbf{D}^{(p)} e^{\mathbf{S}^{(p)} x_1} dx_1 \mathbf{D}^{(p)} \left[\prod_{n=2}^{k-1} \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}^{(p)} x_n} dx_n \mathbf{D}^{(p)} \right] \right. \\ \left. \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}^{(p)} x_n} dx_n V^{(p)}(x) \psi(x) dx - \int_{x \in [0, \Delta)} \int_{u_1=0}^{x_0} g_1(x_0 - u_1) \right. \\ \left. \left[\prod_{k=2}^{n-1} \int_{u_k=0}^{\Delta - u_{k-1}} g_k(\Delta - u_k - u_{k-1}) du_{k-1} \right] g_n(\Delta - x - u_{n-1}) \right. \\ \left. \times 1(\Delta - x - u_{n-1} \geq 0) du_{n-1} \psi(x) dx \right| \quad (3.44)$$

$$\leq \left(|r_8^{(p)}(n)| + |r_5^{(p)}(n)| + |r_6^{(p)}(n)| + (n-1)|r_4^{(p)}(n)| \right) F \Delta. \quad (3.45)$$

where

$$|r_8^{(p)}(n)| \leq \left(2|r_5^{(p)}(n)| + 2|r_6^{(p)}(n)| + 2(n-1)|r_4^{(p)}(n)| + \varepsilon^{(p)} G^{n-1} \Delta^{n-2} (G + L \Delta) \right) \\ + 2\hat{G}^{n-2} G G_V \frac{\text{Var}(Z^{(p)}) / (\varepsilon^{(p)})^2}{1 - \text{Var}(Z^{(p)}) / (\varepsilon^{(p)})^2}.$$

Upon choosing $\varepsilon^{(p)} = \text{Var}(Z^{(p)})^{1/3}$, for fixed $n \geq 2$, $|r_5^{(p)}(n)| + |r_6^{(p)}(n)| + (n-1)|r_4^{(p)}(n)| \rightarrow 0$ and $|r_8^{(p)}| + |r_5^{(p)}(n)| + |r_6^{(p)}(n)| + (n-1)|r_4^{(p)}(n)| \rightarrow 0$ as $p \rightarrow \infty$.

Our first main result towards proving the convergence of the QBD-RAP scheme to the fluid queue is the following bound.

Corollary 3.8. *Let $\psi : \mathcal{D}_{\ell_0} \rightarrow \mathbb{R}$ be bounded, $|\psi(x)| \leq F$, and Lipschitz continuous. For $x \in \mathcal{D}_{\ell_0, j}$, $x_0 \in \mathcal{D}_{\ell_0, i}$, $j_1, j_2, \dots \in \mathcal{S}_-$, $k_1, k_2, \dots \in \mathcal{S}_+$, $\ell_0 \in \mathcal{K} \setminus \{-1, K+1\}$, $m \geq 0$, $\lambda > 0$, then*

1. if $i \in \mathcal{S}_+$, $j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$,

$$\left| \int_{x \in \mathcal{D}_{\ell_0}} \hat{f}_{m,+,+}^{\ell_0, (p)}(\lambda)(x, j_1, k_1, \dots, j_m, k_m, j; x_0, i) \psi(x) dx \right. \\ \left. - \int_{x \in \mathcal{D}_{\ell_0}} \hat{\mu}_{m,+,+}^{\ell_0}(\lambda)(x, j_1, k_1, \dots, j_m, k_m, j; x_0, i) \psi(x) dx \right| \\ \leq \begin{cases} R_{V,2}^{(p)} G F + \varepsilon^{(p)} G F, & m = 0, \\ (|r_5^{(p)}(2m)| + |r_6^{(p)}(2m)| + (2m-1)|r_4^{(p)}(2m)|) \Delta F, & m \geq 1, \end{cases} \quad (3.46)$$

2. if $i \in \mathcal{S}_-$, $j \in \mathcal{S}_- \cup \mathcal{S}_{-0}$,

$$\begin{aligned} & \left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{m,-,-}^{\ell_0,(p)}(\lambda)(x, k_1, j_1, \dots, k_m, j_m, j; x_0, i) \psi(x) dx \right. \\ & \quad \left. - \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{m,-,-}^{\ell_0}(\lambda)(x, k_1, j_1, \dots, k_m, j_m, j; x_0, i) \psi(x) dx \right| \\ & \leq \begin{cases} R_{V,2}^{(p)} GF + \varepsilon^{(p)} GF, & m = 0, \\ (|r_5^{(p)}(2m)| + |r_6^{(p)}(2m)| + (2m-1)|r_4^{(p)}(2m)|) \Delta F, & m \geq 1, \end{cases} \end{aligned} \quad (3.47)$$

3. if $i \in \mathcal{S}_+$, $j \in \mathcal{S}_- \cup \mathcal{S}_{-0}$,

$$\begin{aligned} & \left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{m,+, -}^{\ell_0,(p)}(\lambda)(x, j_1, k_1, \dots, j_m, k_m, j_{m+1}, j; x_0, i) \psi(x) dx \right. \\ & \quad \left. - \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{m,+, -}^{\ell_0}(\lambda)(x, j_1, k_1, \dots, j_m, k_m, j_{m+1}, j; x_0, i) \psi(x) dx \right| \\ & \leq (|r_5^{(p)}(2m+1)| + |r_6^{(p)}(2m+1)| + 2m|r_4^{(p)}(2m+1)|) \Delta F, \quad m \geq 0, \end{aligned} \quad (3.48)$$

4. and if $i \in \mathcal{S}_-$, $j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$,

$$\begin{aligned} & \left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{m,-, +}^{\ell_0,(p)}(\lambda)(x, k_1, j_1, \dots, k_m, j_m, k_{m+1}, j; x_0, i) \psi(x) dx \right. \\ & \quad \left. - \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{m,-, +}^{\ell_0}(\lambda)(x, k_1, j_1, \dots, k_m, j_m, k_{m+1}, j; x_0, i) \psi(x) dx \right| \\ & \leq (|r_5^{(p)}(2m+1)| + |r_6^{(p)}(2m+1)| + 2m|r_4^{(p)}(2m+1)|) \Delta F, \quad m \geq 0. \end{aligned} \quad (3.49)$$

5. if $k \in \mathcal{S}_{-0}$, $i \in \mathcal{S}_+$, $j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$,

$$\begin{aligned} & \left| \int_{x=0}^{\Delta} \widehat{f}_{m,-0,+}^{\ell_0}(\lambda)(x, i, j_1, k_1, \dots, j_m, k_m, j; x_0, k) \psi(x) dx \right. \\ & \quad \left. - \int_{x=0}^{\Delta} \widehat{\mu}_{m,-0,+}^{\ell_0}(\lambda)(x, i, j_1, k_1, \dots, j_m, k_m, j; x_0, k) \psi(x) dx \right| \\ & \leq \begin{cases} |r_{10}^{(p)}|, & m = 0 \\ (|r_8^{(p)}(2m)| + |r_5^{(p)}(2m)| + |r_6^{(p)}(2m)| + (2m-1)|r_4^{(p)}(2m)|) \Delta F & m \geq 1 \end{cases} \end{aligned} \quad (3.50)$$

6. if $k \in \mathcal{S}_{-0}$, $i \in \mathcal{S}_+$, $j \in \mathcal{S}_- \cup \mathcal{S}_{-0}$,

$$\begin{aligned} & \left| \int_{x=0}^{\Delta} \widehat{f}_{m,-0,-}^{\ell_0,(p)}(\lambda)(x, i, j_1, k_1, \dots, k_m, j_{m+1}, j; x_0, k) \psi(x) dx \right. \\ & \quad \left. - \int_{x=0}^{\Delta} \widehat{\mu}_{m,-0,-}^{\ell_0}(\lambda)(x, i, j_1, k_1, \dots, k_m, j_{m+1}, j; x_0, k) \psi(x) dx \right| \\ & \leq (|r_8^{(p)}(2m+1)| + |r_5^{(p)}(2m+1)| + |r_6^{(p)}(2m+1)| + 2m|r_4^{(p)}(2m+1)|) \Delta F, \quad m \geq 0 \end{aligned} \quad (3.51)$$

7. if $k \in \mathcal{S}_{+0}$, $i \in \mathcal{S}_-$, $j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$,

$$\begin{aligned} & \left| \int_{x=0}^{\Delta} \widehat{f}_{m,+0,+}^{\ell_0,(p)}(\lambda)(x, i, k_1, j_1, \dots, j_m, k_{m+1}, j; x_0, k) \psi(x) dx \right. \\ & \quad \left. - \int_{x=0}^{\Delta} \widehat{\mu}_{m,+0,+}^{\ell_0}(\lambda)(x, i, k_1, j_1, \dots, j_m, k_{m+1}, j; x_0, k) \psi(x) dx \right| \\ & \leq (|r_8^{(p)}(2m+1)| + |r_5^{(p)}(2m+1)| + |r_6^{(p)}(2m+1)| + 2m|r_4^{(p)}(2m+1)|) \Delta F, \quad m \geq 0 \end{aligned} \quad (3.52)$$

8. and if $k \in \mathcal{S}_{+0}$, $i \in \mathcal{S}_-$, $j \in \mathcal{S}_- \cup \mathcal{S}_{-0}$,

$$\begin{aligned} & \left| \int_{x=0}^{\Delta} \widehat{f}_{m,+0,-}^{\ell_0,(p)}(\lambda)(x, i, k_1, j_1, \dots, k_m, j_m, j; x_0, k) \psi(x) dx \right. \\ & \quad \left. - \int_{x=0}^{\Delta} \widehat{\mu}_{m,+0,-}^{\ell_0}(\lambda)(x, i, k_1, j_1, \dots, k_m, j_m, j; x_0, k) \psi(x) dx \right| \\ & \leq \begin{cases} |r_{10}^{(p)}|, & m = 0 \\ (|r_8^{(p)}(2m)| + |r_5^{(p)}(2m)| + |r_6^{(p)}(2m)| + (2m-1)|r_4^{(p)}(2m)|) \Delta F & m \geq 1. \end{cases} \end{aligned} \quad (3.53)$$

Proof. For (3.46) and (3.47) and $m = 0$ apply Lemma 3.4 to the differences on the left-hand sides.

For (3.46) and (3.47) and $m \geq 1$ apply Corollary 3.6 to the differences on the left-hand sides.

For (3.48) and (3.49) apply Corollary 3.6 to the differences on the left-hand sides.

For (3.50) and (3.53) and $m = 0$ apply Corollary 3.5 to the differences on the left-hand sides.

For (3.50) and (3.53) and $m \geq 1$ apply Corollary 3.7 to the differences on the left-hand sides.

For (3.51) and (3.52) apply Corollary 3.7 to the differences on the left-hand sides. \square

These bounds are enough to show the weak convergence (in space and time) of the QBD-RAP scheme to the fluid queue on the event on the event that there is no change of level, and on a given partition as described in (??)-(??). Since the state space of the phases, \mathcal{S} , is finite, then the same bounds show convergence when we sum over all possible phases at times $\{\Sigma_m\}$ and $\{\Gamma_m\}$. We formalise this with Corollary 3.9, below.

Corollary 3.9. *For $q \in \{+, -, +0, -0\}$, $r \in \{+, -\}$, $i \in \mathcal{S}_q$, $j \in \mathcal{S}_r$, $x \in \mathcal{D}_{\ell_0, j}$, $x_0 \in \mathcal{D}_{\ell_0, i}$, $\ell_0 \in \mathcal{K}$, $\ell_0 \notin \{-1, K+1\}$, $m \geq 0$, $\lambda > 0$, then*

$$\left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{m, q, r}^{\ell_0, (p)}(\lambda)(x, j; x_0, i) \psi(x) dx - \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{m, q, r}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) dx \right| \rightarrow 0 \quad (3.54)$$

as $p \rightarrow \infty$.

Proof. We prove the result for $q = r = +$ only, with the proofs for the other cases being analogous.

By (??) and (??) and the triangle inequality,

$$\begin{aligned} & \left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{m, +, +}^{\ell_0, (p)}(\lambda)(x, j; x_0, i) \psi(x) dx - \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{m, +, +}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) dx \right| \\ & \leq \sum_{j_1 \in \mathcal{S}_-} \sum_{k_1 \in \mathcal{S}_+} \cdots \sum_{j_m \in \mathcal{S}_-} \sum_{k_m \in \mathcal{S}_+} \left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{m, +, +}^{\ell_0, (p)}(\lambda)(x, j_1, k_1, \dots, j_m, k_m, j; x_0, i) \psi(x) dx \right. \\ & \quad \left. - \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{m, +, +}^{\ell_0}(\lambda)(x, j_1, k_1, \dots, j_m, k_m, j; x_0, i) \psi(x) dx \right| \\ & \leq \begin{cases} R_{V,2}^{(p)} GF + \varepsilon^{(p)} GF & m = 0 \\ (|r_5^{(p)}(2m)| + |r_6^{(p)}(2m)| + 2m|r_4^{(p)}(2m)|) \Delta F |\mathcal{S}_+|^m |\mathcal{S}_-|^m & m \geq 1. \end{cases} \end{aligned}$$

since, by Corollary 3.8, each term in the sum is bounded by either $R_{V,2}^{(p)} GF + \varepsilon^{(p)} GF$ for $m = 0$ or by $(|r_5^{(p)}(2m)| + |r_6^{(p)}(2m)| + 2m|r_4^{(p)}(2m)|) \Delta F$ for $m \geq 1$. \square

For $q \in \{+, -, +0, -0\}$, $r \in \{+, -\}$, consider the Laplace transforms

$$\int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{q, r}^{\ell_0, (p)}(\lambda)(x, j; x_0, i) \psi(x) dx := \int_{x \in \mathcal{D}_{\ell_0}} \sum_{m=0}^{\infty} \widehat{f}_{m, q, r}^{\ell_0, (p)}(\lambda)(x, j; x_0, i) \psi(x) dx,$$

for $i \in \mathcal{S}_q$, $j \in \mathcal{S}_r \cup \mathcal{S}_{r0}$ and

$$\int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{q, r}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) dx := \int_{x \in \mathcal{D}_{\ell_0}} \sum_{m=0}^{\infty} \widehat{\mu}_{m, q, r}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) dx,$$

for $i \in \mathcal{S}_q$, $j \in \mathcal{S}_r \cup \mathcal{S}_{r0}$. The difference,

$$\left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{q,r}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) \, dx - \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{q,r}^{\ell_0, (p)}(\lambda)(x, j; x_0, i) \psi(x) \, dx \right|$$

is less than or equal to

$$\sum_{m=0}^{\infty} \left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{m,q,r}^{\ell_0, (p)}(\lambda)(x, j; x_0, i) \psi(x) \, dx - \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{m,q,r}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) \, dx \right|,$$

which is an infinite sum of terms of the form of the left-hand side of (3.54). The issue that remains is that the bounds in Corollary 3.9 are not necessarily summable (over m). In a step to resolving this issue, we have the following geometric domination condition.

Let $c_{\min} = \min_{i \in \mathcal{S}_+ \cup \mathcal{S}_-} |c_i|$.

Lemma 3.10. *For all $M \geq 0$, $x \in \mathcal{D}_{\ell_0, j}$, $x_0 \in \mathcal{D}_{\ell_0, i}$, $\ell_0 \in \mathcal{K}$, $\ell_0 \notin \{-1, K+1\}$, $\lambda > 0$, $q \in \{+, -, +0, -0\}$, $r \in \{+, -\}$, $i \in \mathcal{S}$, $j \in \mathcal{S}_r \cup \mathcal{S}_{r0}$,*

$$\sum_{m=M+1}^{\infty} \left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{m,q,r}^{\ell_0, (p)}(\lambda)(x, j; x_0, i) \psi(x) \, dx - \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{m,q,r}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) \, dx \right| \leq r_7^M \quad (3.55)$$

where

$$r_7^M = \Delta F G (1 + G_V) \left(\frac{q}{q + \lambda} \right)^{2M+2} \left(1 - \left(\frac{q}{q + \lambda} \right)^2 \right)^{-1}.$$

Note that the bound r_7^M is independent of p .

We prove the result for $q = r = +$ only, with the proof for the other cases following analogously. Essentially, this result follows from noting the probabilistic interpretation of the Laplace transforms $\widehat{f}_{m,+,+}^{\ell_0}(t)(x, j; x_0, i)$, as the probability that,

- there are m changes from \mathcal{S}_+ to \mathcal{S}_- and \mathcal{S}_- to \mathcal{S}_+ ,
- the orbit process $\{\mathbf{A}(t)\}$ evolves accordingly,
- and an exponential random variable with rate λ has not yet occurred.

We obtain an upper bound by ignoring the behaviour of the orbit process $\{\mathbf{A}(t)\}$, then, by a uniformisation argument, we bound the probability that there are m changes from \mathcal{S}_+ to \mathcal{S}_- and \mathcal{S}_- to \mathcal{S}_+ before an exponential random variable with rate λ occurs, by the event that there are m independent exponential events before an exponential random variable with rate λ occurs.

Similarly, the probabilistic interpretation of the Laplace transforms $\widehat{\mu}_{m,+,+}^{\ell_0}(\lambda)(x, j; x_0, i)$, as the probability that,

- there are m changes from \mathcal{S}_+ to \mathcal{S}_- and \mathcal{S}_- to \mathcal{S}_+ ,
- the fluid level $X(t)$ remains in \mathcal{D}_{ℓ_0} ,
- and an exponential random variable with rate λ has not yet occurred.

We obtain an upper bound by removing the requirement that the fluid level $X(t)$ remain in \mathcal{D}_{ℓ_0} , then applying the same uniformisation argument as we do for $\widehat{f}_{m,+,+}^{\ell_0}(t)(x, j; x_0, i)$.

Proof. The same arguments and results apply for all p , so let us drop the dependence on p .

Consider $i \in \mathcal{S}_+, j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$. By the triangle inequality,

$$\begin{aligned} & \sum_{m=M+1}^{\infty} \left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{m,+,+}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) dx - \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{m,+,+}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) dx \right| \\ & \leq \sum_{m=M+1}^{\infty} \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{m,+,+}^{\ell_0}(\lambda)(x, j; x_0, i) |\psi(x)| dx \\ & \quad + \sum_{m=M+1}^{\infty} \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{m,+,+}^{\ell_0}(\lambda)(x, j; x_0, i) |\psi(x)| dx, \end{aligned}$$

since all terms are non-negative.

Consider $\widehat{f}_{m,+,+}^{\ell_0}(\lambda)(x, j; x_0, i)$, which is given by the (i, j) th entry of

$$\begin{aligned} & \int_{x_1=0}^{\infty} \mathbf{H}^{+-}(\lambda, x_1) \int_{x_2=0}^{\infty} \mathbf{H}^{-+}(\lambda, x_2) \dots \int_{x_m=0}^{\infty} \mathbf{H}^{-+}(\lambda, x_m) \int_{x_{m+1}=0}^{\infty} \mathbf{H}^{++}(\lambda, x_{m+1}) \\ & \mathbf{a}_{\ell_0, i}(x_0) e^{\mathbf{S}x_1} dx_1 \mathbf{D} e^{\mathbf{S}x_2} dx_2 \mathbf{D} \dots e^{\mathbf{S}x_m} dx_m \mathbf{D} e^{\mathbf{S}x_{m+1}} V(y_{\ell_0+1} - x) dx_{m+1}. \end{aligned} \quad (3.56)$$

Substituting the upper bound $h_{j_m, j}^{++}(\lambda, x_{m+1}) \leq G$, then (3.56) is less than or equal to

$$\begin{aligned} & \int_{x_1=0}^{\infty} \mathbf{H}^{+-}(\lambda, x_1) \int_{x_2=0}^{\infty} \mathbf{H}^{-+}(\lambda, x_2) \dots \int_{x_m=0}^{\infty} \mathbf{H}^{-+}(\lambda, x_m) eG \\ & \int_{x_{m+1}=0}^{\infty} \mathbf{a}_{\ell_0, i}(x_0) e^{\mathbf{S}x_1} dx_1 \mathbf{D} e^{\mathbf{S}x_2} dx_2 \mathbf{D} \dots e^{\mathbf{S}x_m} dx_m \mathbf{D} e^{\mathbf{S}x_{m+1}} V(y_{\ell_0+1} - x) dx_{m+1} \\ & = \int_{x_1=0}^{\infty} \mathbf{H}^{+-}(\lambda, x_1) \int_{x_2=0}^{\infty} \mathbf{H}^{-+}(\lambda, x_2) \dots \int_{x_m=0}^{\infty} \mathbf{H}^{-+}(\lambda, x_m) eG \\ & \mathbf{a}_{\ell_0, i}(x_0) e^{\mathbf{S}x_1} dx_1 \mathbf{D} e^{\mathbf{S}x_2} dx_2 \mathbf{D} \dots e^{\mathbf{S}x_m} dx_m \mathbf{D} (-\mathbf{S})^{-1} V(y_{\ell_0+1} - x) \\ & \leq \int_{x_1=0}^{\infty} \mathbf{H}^{+-}(\lambda, x_1) \int_{x_2=0}^{\infty} \mathbf{H}^{-+}(\lambda, x_2) \dots \int_{x_m=0}^{\infty} \mathbf{H}^{-+}(\lambda, x_m) eG \\ & \mathbf{a}_{\ell_0, i}(x_0) e^{\mathbf{S}x_1} dx_1 \mathbf{D} e^{\mathbf{S}x_2} dx_2 \mathbf{D} \dots e^{\mathbf{S}x_m} dx_m eG_V, \end{aligned} \quad (3.57)$$

The last inequality holds since

$$\mathbf{D}(-\mathbf{S})^{-1}V(x) = \int_{u=0}^{\infty} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} (-\mathbf{S})^{-1}V(x) \, du \leq \int_{u=0}^{\infty} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} G_V = \mathbf{e} G_V,$$

which relies on Property 3.2(i).

Now, applying Lemma A.19 repeatedly gives the bound

$$\mathbf{a}_{\ell_0, i}(x_0) e^{\mathbf{S}x_1} \mathbf{D} e^{\mathbf{S}x_2} \mathbf{D} \dots e^{\mathbf{S}x_m} \mathbf{e} \leq 1.$$

Hence (3.57) is less than or equal to

$$\int_{x_1=0}^{\infty} \mathbf{H}^{+-}(\lambda, x_1) \, dx_1 \int_{x_2=0}^{\infty} \mathbf{H}^{-+}(\lambda, x_2) \, dx_2 \dots \int_{x_{2m}=0}^{\infty} \mathbf{H}^{-+}(\lambda, x_{2m}) \, dx_{2m} \mathbf{e} G_V G. \quad (3.58)$$

The stochastic interpretation of the i th element of the vector $\mathbf{H}^{+-}(\lambda, x)\mathbf{e}$ is that it is the probability density that the phase of the fluid queue changes from a positive to a negative phase at the time when the in-out fluid has increased by dx and before an exponential random variable with rate λ occurs, given the phase is initially i . There may be multiple changes of phase within $\mathcal{S}_+ \cup \mathcal{S}_{+0}$ before the first change from $\mathcal{S}_+ \cup \mathcal{S}_{+0}$ to \mathcal{S}_- . The first change of phase occurs at rate (with respect to the in-out level) $-T_{ii}/|c_i|$ and this is the lowest in-out fluid level at which it may be possible for us to see a transition from \mathcal{S}_+ to \mathcal{S}_- . Consider a uniformisation of the phase process of the in-out fluid with uniformisation parameter $q = \max_{i \in \mathcal{S}} -T_{ii}/|c_i|$. Then the first event of the uniformised phase process occurs at rate q and occurs at, or before, the first change of phase. Hence the first uniformisation event occurs at, or before, the first change from $\mathcal{S}_+ \cup \mathcal{S}_{+0}$ to \mathcal{S}_- . This gives the bound $\mathbf{H}^{+-}(\lambda, x)\mathbf{e} \leq qe^{-(\lambda+q)x}\mathbf{e}$.

Similarly for $\mathbf{H}^{-+}(\lambda, x)\mathbf{e} \leq qe^{-(\lambda+q)x}\mathbf{e}$.

From this stochastic interpretation, (3.58) is less than or equal to

$$\begin{aligned} & \mathbf{H}^{+-}(\lambda, x_1) \, dx_1 \int_{x_2=0}^{\infty} \mathbf{H}^{-+}(\lambda, x_2) \, dx_2 \dots \int_{x_m=0}^{\infty} qe^{(-q-\lambda)x_m} \, dx_{2m} G_V G \\ & \leq \mathbf{e} \int_{x_1=0}^{\infty} qe^{(-q-\lambda)x_1} \, dx_1 \int_{x_2=0}^{\infty} qe^{(-q-\lambda)x_2} \, dx_2 \dots \int_{x_{2m}=0}^{\infty} qe^{(-q-\lambda)x_{2m}} \, dx_{2m} G_V G \\ & = \mathbf{e} \left(\frac{q}{q+\lambda} \right)^{2m} G_V G. \end{aligned} \quad (3.59)$$

Hence,

$$\sum_{m=M+1}^{\infty} \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{m,+,+}^{\ell_0}(\lambda)(x, j; x_0, i) |\psi(x)| \, dx$$

$$\begin{aligned}
&\leq G_V G \sum_{m=M+1}^{\infty} \left(\frac{q}{q+\lambda} \right)^{2m} \int_{x \in \mathcal{D}_{\ell_0}} |\psi(x)| \, dx \\
&\leq G_V G \left(\frac{q}{q+\lambda} \right)^{2M+2} \left(1 - \left(\frac{q}{q+\lambda} \right)^2 \right)^{-1} \Delta F.
\end{aligned} \tag{3.60}$$

Now consider $\widehat{\mu}_{m,+,+}^{\ell_0}(\lambda)(x, j; x_0, i)$ which is given by the (i, j) th entry of

$$\begin{aligned}
&\int_{x_1=0}^{\Delta-(x_0-y_{\ell_0})} \mathbf{H}^{+-}(\lambda, \Delta - (x_0 - y_{\ell_0}) - x_1) \int_{x_2=0}^{\Delta-x_1} \mathbf{H}^{-+}(\lambda, \Delta - x_2 - x_1) \, dx_1 \\
&\quad \dots \int_{x_{2m}=0}^{\Delta-x_{m-1}} \mathbf{H}^{-+}(\lambda, \Delta - x_{2m-1} - x_{2m}) \, dx_{2m-1} \mathbf{H}^{++}(\lambda, \Delta - x_{2m} - (y_{\ell_0+1} - x)) \, dx_{2m} \\
&= \int_{x_1=(x_0-y_{\ell_0})}^{\Delta} \mathbf{H}^{+-}(\lambda, \Delta - x_1) \int_{x_2=x_1}^{\Delta} \mathbf{H}^{-+}(\lambda, \Delta - x_2) \, dx_1 \dots \\
&\quad \int_{x_{2m}=x_{2m-1}}^{\Delta} \mathbf{H}^{-+}(\lambda, \Delta - x_{2m}) \, dx_{2m-1} \mathbf{H}^{++}(\lambda, \Delta - x_{2m} - x_{2m-1} - (y_{\ell_0+1} - x)) \\
&\quad \, dx_{2m-1} \, dx_{2m}.
\end{aligned} \tag{3.61}$$

Once again, use the bound $h_{j_m, j}^{++}(\lambda, x_{m+1}) \leq G$. Hence (3.61) is less than or equal to

$$\begin{aligned}
&\int_{x_1=(x_0-y_{\ell_0})}^{\Delta} \mathbf{H}^{+-}(\lambda, \Delta - x_1) \int_{x_2=x_1}^{\Delta} \mathbf{H}^{-+}(\lambda, \Delta - x_2) \, dx_1 \\
&\quad \dots \int_{x_{2m}=x_{2m-1}}^{\Delta} \mathbf{H}^{-+}(\lambda, \Delta - x_{2m}) \, dx_{2m-1} \, dx_{2m} G.
\end{aligned} \tag{3.62}$$

The expression (3.62) differs from (3.58) only by a constant factor G_V and that the integrals in the former are finite, hence we may bound it in the same way as (3.58). Therefore,

$$\begin{aligned}
&\sum_{m=M+1}^{\infty} \int_{x \in \mathcal{D}_{\ell_0}} |\psi(x)| \, dx \widehat{\mu}_{m,+,+}^{\ell_0}(\lambda)(x, j; x_0, i) |\psi(x)| \, dx \\
&\leq G \left(\frac{q}{q+\lambda} \right)^{2M+2} \left(1 - \left(\frac{q}{q+\lambda} \right)^2 \right)^{-1} \Delta F.
\end{aligned} \tag{3.63}$$

Analogous arguments show the same bounds for any $i, j \in \mathcal{S}$. \square

Lemma 3.11. *For all $x \in \mathcal{D}_{\ell_0,j}$, $x_0 \in \mathcal{D}_{\ell_0,i}$, $i, j \in \mathcal{S}$, $\ell_0 \in \mathcal{K}$, $\ell_0 \notin \{-1, K+1\}$, $\lambda > 0$,*

$$\left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}^{\ell_0,(p)}(\lambda)(x, j; x_0, i) \psi(x) \, dx - \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) \, dx \right| \rightarrow 0 \quad (3.64)$$

as $p \rightarrow \infty$.

Proof. Consider $i \in \mathcal{S}_+$ and $j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$. By partitioning on the number of changes from $\mathcal{S}_- \rightarrow \mathcal{S}_+$, (3.64) can be written as

$$\begin{aligned} & \left| \sum_{m=0}^{\infty} \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{m,+,+}^{\ell_0,(p)}(\lambda)(x, j; x_0, i) \psi(x) \, dx - \sum_{m=0}^{\infty} \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{m,+,+}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) \, dx \right| \\ & \leq \sum_{m=0}^{\infty} \left| \int_{x \in \mathcal{D}_{\ell_0}} \widehat{f}_{m,+,+}^{\ell_0,(p)}(\lambda)(x, j; x_0, i) \psi(x) \, dx - \int_{x \in \mathcal{D}_{\ell_0}} \widehat{\mu}_{m,+,+}^{\ell_0}(\lambda)(x, j; x_0, i) \psi(x) \, dx \right|. \end{aligned} \quad (3.65)$$

By Corollary 3.9 each term in the sum (3.65) converges to 0 as $p \rightarrow \infty$. Lemma 3.10 gives a domination condition, so we can apply the Dominated Convergence Theorem which proves the stated convergence in (3.64).

Analogous arguments can be applied for any $i, j \in \mathcal{S}$. \square

Remark 3.12. *For a fixed $\lambda > 0$, convergence of*

$$\left| \widehat{f}^{\ell_0,(p)}(\lambda)(x, j; x_0, i) - \widehat{\mu}^{\ell_0}(\lambda)(x, j; x_0, i) \right| \quad (3.66)$$

actually holds point-wise for each $\ell_0 \in \mathcal{K} \setminus \{-1, K+1\}$, and each $i, j \in \mathcal{S}$, $x_0 \in \mathcal{D}_{\ell_0,i}$, $x \in \mathcal{D}_{\ell_0,j}$ except at the set of points where $x = x_0$. Specifically, the lack of point-wise convergence at this point occurs due terms with the index $m = 0$, that is, terms where there are no changes of phase from $\mathcal{S}_+ \rightarrow \mathcal{S}_-$ or $\mathcal{S}_- \rightarrow \mathcal{S}_+$. On these sample paths the relevant Laplace transforms of the fluid queue are discontinuous at this point. For example,

$$\widehat{\mu}_{0,+,+}^{\ell_0}(\lambda)(x, j; x_0, i) \, dx = h_{ij}^{++}(\lambda, x - x_0) 1(x \geq x_0) \, dx,$$

is discontinuous at $x = x_0$. If we were to insist on point-wise convergence, then we would need to enforce the error term $r_V(u, v)^{(p)} \rightarrow 0$ point-wise for each u, v . In the cases presented here $r_V^{(p)}(u, v)$ converges point-wise to 0 everywhere except $u + v = \Delta$.

3.3.1 Convergence at the first change of level

Before moving on to global convergence we need another result about convergence of the QBD-RAP scheme and the fluid queue at changes of level.

Let E^λ be an exponential random variable with rate λ . In the following we use the stochastic interpretation of the Laplace transform of a probability distribution with non-negative support: for a random variable W with distribution function $F_W(w) = \mathbb{P}(W < w)$, then $\int_{w=0}^{\infty} e^{-\lambda w} dF_W(w) = \mathbb{P}(W < E^\lambda)$. That is, the Laplace transform with parameter $\lambda > 0$ is the probability that W occurs before an exponential time with rate λ occurs.

Regarding Remark 3.12 we do require convergence at certain boundary points which correspond to changes of level of the QBD-RAP. To this end, we have the following result.

Corollary 3.13. *For $\ell, \ell_0 \in \mathcal{K} \setminus \{-1, K+1\}$, $x_0 \in \mathcal{D}_{\ell_0, i}$, $i \in \mathcal{S}$ then, for $j \in \mathcal{S}_+$,*

$$\begin{aligned} & \mathbb{P}(L^{(p)}(\tau_1^{(p)}) = \ell_0 + 1, \varphi(\tau_1^{(p)}) = j, \tau_1^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \\ & \quad \varphi(0) = i) \\ & \rightarrow \mathbb{P}(X(\tau_1^X) = y_{\ell_0+1}, \varphi(\tau_1^X) = j, \tau_1^X \leq E^\lambda \mid X(0) = x_0, \varphi(0) = i) \end{aligned} \quad (3.67)$$

and for $j \in \mathcal{S}_-$

$$\begin{aligned} & \mathbb{P}(L^{(p)}(\tau_1^{(p)}) = \ell_0 - 1, \varphi(\tau_1^{(p)}) = j, \tau_1^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \\ & \quad \varphi(0) = i) \\ & \rightarrow \mathbb{P}(X(\tau_1^X) = y_{\ell_0}, \varphi(\tau_1^X) = j, \tau_1^X \leq E^\lambda \mid X(0) = x_0, \varphi(0) = i). \end{aligned}$$

At a boundary, for $\ell_0 \in \{-1, K+1\}$,

$$\begin{aligned} & \mathbb{P}(L^{(p)}(\tau_1^{(p)}) = \ell, \varphi(\tau_1^{(p)}) = j, \tau_1^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = 1, \varphi(0) = i) \\ & = \mathbb{P}(X(\tau_1^X) = x, \varphi(\tau_1^X) = j, \tau_1 \leq E^\lambda \mid X(0) = 0, \varphi(0) = i), \end{aligned} \quad (3.68)$$

where $\ell = 0$, $i \in \mathcal{S}_- \cup \mathcal{S}_{-0}$ and $j \in \mathcal{S}_+$ if $\ell_0 = -1$, and $\ell = K$, $i \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$ and $j \in \mathcal{S}_-$ if $\ell_0 = K+1$.

Proof. The proof follows the same structure as the proof of Lemma 3.11 however changes are required in all the results used in the proof, as here we do not integrate a function ψ . Here we only give an outline of the proof.

Consider first $i \in \mathcal{S}_+, j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$. Partition the probability (3.67) on the times $\{\Sigma_n\}_{n \geq 1}$ and $\{\Gamma_n\}_{n \geq 1}$, specifically, partition on the event that there are exactly m events $\{\Sigma_n\}_{n=1}^m$ and exactly m events $\{\Gamma_n\}_{n=1}^m$. Partition further by the phases at these times $\varphi(\Sigma_n) = j_n \in \mathcal{S}_-, n = 1, \dots, m$ and $\varphi(\Gamma_n) = k_n \in \mathcal{S}_+, n = 1, \dots, m$. The resulting partitioned probabilities are

$$\int_{x_1=0}^{\infty} h_{i, j_1}^{+-}(\lambda, x_1) \mathbf{a}_{\ell_0, i}^{(p)}(x_0) e^{\mathbf{S}^{(p)} x_1} dx_1 \mathbf{D}^{(p)}$$

$$\left[\prod_{r=1}^{m-1} \int_{x_{2r}=0}^{\infty} h_{j_r, k_r}^{-+}(\lambda, x_{2r}) e^{\mathbf{S}^{(p)} x_{2r}} dx_{2r} \mathbf{D}^{(p)} \int_{x_{2r+1}=0}^{\infty} h_{k_r, j_{r+1}}^{+-}(\lambda, x_{2r+1}) e^{\mathbf{S}^{(p)} x_{2r+1}} dx_{2r+1} \mathbf{D}^{(p)} \right] \int_{x_{2m}=0}^{\infty} h_{j_m, k_m}^{-+}(\lambda, x_{2m}) e^{\mathbf{S}^{(p)} x_{2m}} dx_{2m} \mathbf{D}^{(p)} \int_{x_{2m+1}=0}^{\infty} h_{k_m, j}^{++}(\lambda, x_{2m+1}) e^{\mathbf{S}^{(p)} x_{2m+1}} dx_{2m+1} \mathbf{S}^{(p)} \quad (3.69)$$

and (3.67) is obtained by summing over $j_1, \dots, j_m, k_1, \dots, k_m, m \geq 0$.

Corollary A.2 and Lemma A.4 are analogous to Lemma 3.4 and Corollary 3.4 used in the proof of Lemma 3.11. They provide relevant bounds for the difference between terms like (3.69) for $m = 0$ and

$$\begin{aligned} & \mathbb{P}(X(\tau_1^X) = y_{\ell_0+1}, \varphi(\tau_1^X) = j, \tau_1^X \leq E^\lambda, \tau_1^X < \Sigma_1 \mid X(0) = x_0, \varphi(0) = i) \\ & = h_{ij}^{++}(\lambda, y_{\ell_0+1} - x_0), \end{aligned}$$

for example.

Corollaries A.11 and A.15 are analogous to Corollaries 3.6 and 3.7 used in the proof of Lemma 3.11. Corollaries A.11 and A.15 give error bounds for the difference between terms such as (3.69) for $m \geq 1$ and the corresponding probability for the fluid queue,

$$\begin{aligned} & \mathbb{P}(X(\tau_1^X) = y_{\ell_0+1}, \varphi(\tau_1^X) = j, \tau_1^X \leq E^\lambda, \Sigma_m \leq \tau_1^X < \Gamma_{m+1}, \varphi(\Sigma_\ell) = j_\ell, \varphi(\Gamma_\ell) = k_\ell, \\ & \ell = 1, \dots, m \mid X(0) = x_0, \varphi(0) = i) \end{aligned} \quad (3.70)$$

which is given by (3.29).

The error bounds on each term in the partition imply that all the terms converge to the corresponding terms for the fluid queue. Equation (3.59) provides a domination condition. Thus we can apply the Dominated Convergence Theorem to prove that the sum of the partitioned probabilities (3.69) converges as $p \rightarrow \infty$. The sum of the limits is

$$\mathbb{P}(X(\tau_1^X) = y_{\ell_0+1}, \varphi(\tau_1^X) = j, \tau_1 \leq E^\lambda \mid X(0) = x_0, \varphi(0) = i).$$

The result for all other cases $i, j \in \mathcal{S}$ follow analogously.

At a boundary we can model the fluid queue exactly, hence (3.68) holds. \square

3.4 At the n th change of level

So far we have shown error bounds for the QBD-RAP approximation on the event that the fluid queue and QBD-RAP remain in the same band/level (Lemma 3.11) and also immediately upon exiting the band (Corollary 3.13). In the next three subsections we extend this to global convergence of the approximation. To do so, we partition the approximation and the fluid queue by the number of level changes. The local convergence on a given level, which we just proved, is then used to claim that each term in the partition

converges. What remains is to argue that we can swap a limit and countable sum which we do by the Dominated Convergence Theorem.

Let $\{\tau_n^{(p)}\}_{n \geq 0}$, $\tau_0^{(p)} = 0$, and

$$\tau_n^{(p)} = \inf \left\{ t \geq \tau_{n-1}^{(p)} \mid L^{(p)}(t) \neq L^{(p)}(\tau_{n-1}^{(p)}) \right\},$$

be the (stopping) times at which $\{L^{(p)}(t)\}$, the level process of the QBD-RAP, changes, or the boundary is hit, or, if the process is at the boundary, the process leaves the boundary. To simplify notation, we may drop the superscript p where it is not explicitly needed.

The process $\{L^{(p)}(\tau_n^{(p)})\}_{n \geq 0}$ is the level process of the (continuous-time) QBD-RAP process observed at changes of level. When $\tau_n^{(p)}$ is a change of level, or the time when the process leaves a boundary, the value of orbit process at these times is $\mathbf{A}^{(p)}(\tau_n^{(p)}) = \boldsymbol{\alpha}^{(p)}$. At time $\tau_0^{(p)} = 0$, $\mathbf{A}^{(p)}(\tau_0^{(p)}) = \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0)$. The process $\{(L^{(p)}(\tau_n^{(p)}), \phi^{(p)}(\tau_n^{(p)}))\}_{n \geq 0}$ is a time-inhomogeneous discrete-time QBD. Further, the process $\{\phi^{(p)}(\tau_n^{(p)})\}_{n \geq 0}$ is a time-inhomogeneous discrete-time Markov chain on the state space \mathcal{S} . For $\mathbf{a}_{\ell_0, i}^{(p)}(x_0) = \boldsymbol{\alpha}^{(p)}$, or for index $n \geq 1$, then $\{L^{(p)}(\tau_n^{(p)}), \phi^{(p)}(\tau_n^{(p)})\}_{n \geq 1}$, and $\{\phi^{(p)}(\tau_n^{(p)})\}_{n \geq 1}$ are time-homogenous.

Let $\{\tau_n^X\}_{n \geq 0}$, be the sequence of (stopping) times with $\tau_0^X = 0$, and

$$\tau_{n+1}^X = \min \left\{ \begin{array}{l} \inf \{t > \tau_n^X \mid X(t) = y_\ell, \ell \in \mathcal{K}\}, \\ \inf \{t > \tau_n^X \mid X(t) \neq 0, X(0) = 0\}, \\ \inf \{t > \tau_n^X \mid X(t) \neq y_{K+1}, X(0) = y_{K+1}\} \end{array} \right\}.$$

For $n \geq 1$, τ_n^X is the time at which $X(t)$ either changes band, or hits a boundary, or the process leaves a boundary, for the n th time.

For $n \geq 1$, consider the Laplace transform

$$\begin{aligned} & \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} \in dt \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \\ & \quad \phi^{(p)}(0) = i) dt \\ &= \mathbb{P}(L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \\ & \quad \phi^{(p)}(0) = i), \end{aligned}$$

which is the Laplace transform of the time until the n th change of level of the QBD-RAP on the event that the level and phase at the n th change of level are ℓ and j_n , respectively, given that the initial level and phases are ℓ_0 and i , respectively and the initial orbit is $\mathbf{a}_{\ell_0, i}^{(p)}(x_0)$. Partitioning on the time of the first change of level, τ_1 , and the level and phase at this time gives

$$\sum_{j_1 \in \mathcal{S}} \sum_{\ell_1 \in \{\ell_0+1, \ell_0-1\} \cap \mathcal{K}} \mathbb{P}(L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} \leq E^\lambda \mid L^{(p)}(\tau_1^{(p)}) = \ell_1,$$

$$\begin{aligned}
& \phi^{(p)}(\tau_1^{(p)}) = j_1, \tau_1^{(p)} \leq E^\lambda \\
& \times \mathbb{P}(L^{(p)}(\tau_1^{(p)}) = \ell_1, \phi^{(p)}(\tau_1^{(p)}) = j_1, \tau_1^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \\
& \phi^{(p)}(0) = i).
\end{aligned} \tag{3.71}$$

An application of Corollary (3.13) to the expression on the second line of (3.71) states, for $i \in \mathcal{S}$, $j \in \mathcal{S}$, $\ell_0 \in \mathcal{K}$,

$$\begin{aligned}
& \lim_{p \rightarrow \infty} \mathbb{P}(L^{(p)}(\tau_1^{(p)}) = \ell_1, \phi^{(p)}(\tau_1^{(p)}) = j_1, \tau_1^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \\
& \phi^{(p)}(0) = i) \\
& \rightarrow \mathbb{P}(X(\tau_1^X) = y_{\ell_0+1}, \varphi(\tau_1^X) = j_1, \tau_1^X \leq E^\lambda \mid X(0) = x_0, \varphi(0) = i)
\end{aligned} \tag{3.72}$$

for $j_1 \in \mathcal{S}_+$, $\ell_1 = \ell_0 + 1$ and

$$\begin{aligned}
& \lim_{p \rightarrow \infty} \mathbb{P}(L^{(p)}(\tau_1^{(p)}) = \ell_1, \phi^{(p)}(\tau_1^{(p)}) = j_1, \tau_1^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \\
& \phi^{(p)}(0) = i) \\
& \rightarrow \mathbb{P}(X(\tau_1^X) = y_{\ell_0}, \varphi(\tau_1^X) = j_1, \tau_1^X \leq E^\lambda \mid X(0) = x_0, \varphi(0) = i)
\end{aligned} \tag{3.73}$$

for $j_1 \in \mathcal{S}_-$, $\ell_1 = \ell_0 - 1$.

Now, for a given j_1 and ℓ_1 consider

$$\begin{aligned}
& \mathbb{P}(L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} \leq E^\lambda \mid L^{(p)}(\tau_1^{(p)}) = \ell_1, \phi^{(p)}(\tau_1^{(p)}) = j_1, \tau_1^{(p)} \leq E^\lambda) \\
& = \mathbb{P}(L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_1, \phi^{(p)}(0) = j_1, \tau_1^{(p)} = 0)
\end{aligned} \tag{3.74}$$

by the time-homogeneous property of the QBD-RAP and the memoryless property of the exponential distribution. Equation (3.74) appears as the first factor in the summands of (3.71). Let

$$\mathcal{P}^n(\ell_0, \ell_n) = \{(\ell_1, \dots, \ell_{n-1}) \in \mathcal{K}^{n-1} \mid |\ell_{r-1} - \ell_r| = 1, r = 1, \dots, n\}. \tag{3.75}$$

The set $\mathcal{P}^n(\ell_0, \ell)$ contains all of the possible sequences of levels which $\{L(t)\}$ or $\{X(t)\}$ may visit on a sample path which start in level ℓ_0 , end in level ℓ and change level n times. By partitioning on the times τ_m , $m = 2, \dots, n-1$, and the phases and the levels at these times and using the strong Markov property of the QBD-RAP, then (3.74) is

$$\begin{aligned}
& \sum_{j_2, \dots, j_{n-1} \in \mathcal{S}} \sum_{(\ell_2, \dots, \ell_{n-1}) \in \mathcal{P}^{n-1}(\ell_1, \ell)} \prod_{m=2}^n \mathbb{P}(L^{(p)}(\tau_m^{(p)}) = \ell_m, \phi^{(p)}(\tau_m^{(p)}) = j_m, \tau_m^{(p)} \leq E^\lambda \mid \\
& L^{(p)}(\tau_{m-1}^{(p)}) = \ell_{m-1}, \phi^{(p)}(\tau_{m-1}^{(p)}) = j_{m-1}, \tau_{m-1}^{(p)} \leq E^\lambda),
\end{aligned} \tag{3.76}$$

where we define $\ell_n = \ell$. Each factor in the product is

$$\mathbb{P}(L^{(p)}(\tau_m^{(p)}) = \ell_m, \phi^{(p)}(\tau_m^{(p)}) = j_m, \tau_m^{(p)} \leq E^\lambda \mid L^{(p)}(\tau_{m-1}^{(p)}) = \ell_{m-1}, \phi^{(p)}(\tau_{m-1}^{(p)}) = j_{m-1},$$

$$\begin{aligned}
& \tau_{m-1}^{(p)} \leq E^\lambda) \\
& = \mathbb{P}(L^{(p)}(\tau_m^{(p)}) = \ell_m, \phi^{(p)}(\tau_m^{(p)}) = j_m, \tau_m^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_{m-1}, \phi^{(p)}(0) = j_{m-1}, \\
& \quad \tau_{m-1}^{(p)} = 0) \\
& = \mathbb{P}(L^{(p)}(\tau_1^{(p)}) = \ell_m, \phi^{(p)}(\tau_1^{(p)}) = j_m, \tau_1^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_{m-1}, \phi^{(p)}(0) = j_{m-1}) \quad (3.77)
\end{aligned}$$

by the time-homogeneous property of the QBD-RAP and the memoryless property of the exponential distribution. We can also apply Corollary (3.13) to the terms (3.77) and conclude

$$\begin{aligned}
& \mathbb{P}(L^{(p)}(\tau_1^{(p)}) = \ell_m, \phi^{(p)}(\tau_1^{(p)}) = j_m, \tau_1^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_{m-1}, \phi^{(p)}(0) = j_{m-1}) \\
& \rightarrow \mathbb{P}(X(\tau_1^X) = y_{\ell_{m-1}+1}, \varphi(\tau_1^X) = j_m, \tau_1^X \leq E^\lambda \mid X(0) = x_0, \varphi(0) = j_{m-1}) \quad (3.78)
\end{aligned}$$

for $j_m \in \mathcal{S}_+$, $\ell_m = \ell_{m-1} + 1$ and

$$\begin{aligned}
& \mathbb{P}(L^{(p)}(\tau_1^{(p)}) = \ell_m, \phi^{(p)}(\tau_1^{(p)}) = j_m, \tau_1^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_{m-1}, \phi^{(p)}(0) = j_{m-1}) \\
& \rightarrow \mathbb{P}(X(\tau_1^X) = y_{\ell_{m-1}}, \varphi(\tau_1^X) = j_m, \tau_1^X \leq E^\lambda \mid X(0) = x_0, \varphi(0) = j_{m-1}) \quad (3.79)
\end{aligned}$$

for $j_m \in \mathcal{S}_-$, $\ell_m = \ell_{m-1} - 1$.

By the time-homogeneous property of the fluid queue and the memoryless property of the exponential distribution,

$$\begin{aligned}
& \mathbb{P}(X(\tau_1^X) = y_{\ell_{m-1}+1}, \varphi(\tau_1^X) = j_m, \tau_1^X \leq E^\lambda \mid X(0) = x_0, \varphi(0) = j_{m-1}) \\
& = \mathbb{P}(X(\tau_m^X) = y_{\ell_{m-1}+1}, \varphi(\tau_m^X) = j_m, \tau_m^X \leq E^\lambda \mid X(0) = x_0, \varphi(0) = j_{m-1}, \tau_{m-1}^X = 0) \\
& = \mathbb{P}(X(\tau_m^X) = y_{\ell_{m-1}+1}, \varphi(\tau_m^X) = j_m, \tau_m^X \leq E^\lambda \mid X(\tau_{m-1}^X) = x_0, \varphi(\tau_{m-1}^X) = j_{m-1}, \\
& \quad \tau_{m-1}^X \leq E^\lambda). \quad (3.80)
\end{aligned}$$

Similarly,

$$\begin{aligned}
& \mathbb{P}(X(\tau_1^X) = y_{\ell_{m-1}}, \varphi(\tau_1^X) = j_m, \tau_1^X \leq E^\lambda \mid X(0) = x_0, \varphi(0) = j_{m-1}) \\
& = \mathbb{P}(X(\tau_m^X) = y_{\ell_{m-1}}, \varphi(\tau_m^X) = j_m, \tau_m^X \leq E^\lambda \mid X(\tau_{m-1}^X) = x_0, \varphi(\tau_{m-1}^X) = j_{m-1}, \tau_{m-1}^X \leq E^\lambda). \quad (3.81)
\end{aligned}$$

Now, taking the limit of (3.76) gives

$$\begin{aligned}
& \lim_{p \rightarrow \infty} \sum_{j_2, \dots, j_{n-1} \in \mathcal{S}} \sum_{(\ell_2, \dots, \ell_{n-1}) \in \mathcal{P}^{n-1}(\ell_1, \ell)} \prod_{m=2}^n \mathbb{P}(L^{(p)}(\tau_m^{(p)}) = \ell_m, \phi^{(p)}(\tau_m^{(p)}) = j_m, \tau_m^{(p)} \leq E^\lambda \\
& \quad \mid L^{(p)}(\tau_{m-1}^{(p)}) = \ell_{m-1}, \phi^{(p)}(\tau_{m-1}^{(p)}) = j_{m-1}, \tau_{m-1}^{(p)} \leq E^\lambda) \\
& = \sum_{j_2, \dots, j_{n-1} \in \mathcal{S}} \sum_{(\ell_2, \dots, \ell_{n-1}) \in \mathcal{P}^{n-1}(\ell_1, \ell)} \prod_{m=2}^n \lim_{p \rightarrow \infty} \mathbb{P}(L^{(p)}(\tau_m^{(p)}) = \ell_m, \phi^{(p)}(\tau_m^{(p)}) = j_m, \tau_m^{(p)} \leq E^\lambda)
\end{aligned}$$

$$| L^{(p)}(\tau_{m-1}^{(p)}) = \ell_{m-1}, \phi^{(p)}(\tau_{m-1}^{(p)}) = j_{m-1}, \tau_{m-1}^{(p)} \leq E^\lambda), \quad (3.82)$$

where we may swap the limit and the sums as they are finite, and we can swap the limit and the product since all the limits exist and the product is finite. Substituting the limits into (3.82) gives

$$\begin{aligned} & \sum_{j_2, \dots, j_{n-1} \in \mathcal{S}} \sum_{(\ell_2, \dots, \ell_{n-1}) \in \mathcal{P}^{n-1}(\ell_1, \ell)} \prod_{m=2}^n \mathbb{P}(X(\tau_m^X) = y_{\ell_{m-1}+1(j_m \in \mathcal{S}_+)}, \varphi(\tau_m^X) = j_m, \tau_m^X \leq E^\lambda \\ & \quad | X(\tau_{m-1}^X) = y_{\ell_{m-2}+1(j_{m-1} \in \mathcal{S}_+)}, \tau_{m-1}^X \leq E^\lambda), \\ & = \mathbb{P}(X(\tau_n^X) = y_{\ell+1(j_n \in \mathcal{S}_-)}, \varphi(\tau_n^X) = j_n, \tau_n^X \leq E^\lambda | X(0) = y_{\ell_0+1(j_1 \in \mathcal{S}_+)}, \varphi(0) = j_1, \\ & \quad \tau_1^X \leq E^\lambda). \end{aligned} \quad (3.83)$$

Therefore, taking the limit as $p \rightarrow \infty$ of (3.71)

$$\begin{aligned} & \lim_{p \rightarrow \infty} \sum_{j_1 \in \mathcal{S}} \sum_{\ell_1 \in \{\ell_0+1, \ell_0-1\} \cap \mathcal{K}} \mathbb{P}(L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} \leq E^\lambda | L^{(p)}(0) = \ell_1, \\ & \quad \phi^{(p)}(0) = j_1, \tau_1^{(p)} = 0) \\ & \mathbb{P}(L^{(p)}(\tau_1^{(p)}) = \ell_1, \phi^{(p)}(\tau_1^{(p)}) = j_1, \tau_1^{(p)} \leq E^\lambda | L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \\ & \quad \phi^{(p)}(0) = i) \\ & = \sum_{j_1 \in \mathcal{S}} \sum_{\ell_1 \in \{\ell_0+1, \ell_0-1\} \cap \mathcal{K}} \lim_{p \rightarrow \infty} \mathbb{P}(L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} \leq E^\lambda | L^{(p)}(0) = \ell_1, \\ & \quad \phi^{(p)}(0) = j_1, \tau_1^{(p)} = 0) \\ & \lim_{p \rightarrow \infty} \mathbb{P}(L^{(p)}(\tau_1^{(p)}) = \ell_1, \phi^{(p)}(\tau_1^{(p)}) = j_1, \tau_1^{(p)} \leq E^\lambda | L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \\ & \quad \phi^{(p)}(0) = i) \\ & = \sum_{j_1 \in \mathcal{S}} \sum_{\ell_1 \in \{\ell_0+1, \ell_0-1\} \cap \mathcal{K}} \mathbb{P}(X(\tau_n^X) = y_{\ell+1(j_n \in \mathcal{S}_-)}, \varphi(\tau_n^X) = j_n, \tau_n^X \leq E^\lambda | X(\tau_1^X) = y_{\ell_0+1(j_1 \in \mathcal{S}_+)}, \\ & \quad \varphi(\tau_1^X) = j_1, \tau_1^X \leq E^\lambda) \\ & \mathbb{P}(X(\tau_1^X) = y_{\ell_0+1(j_1 \in \mathcal{S}_+)}, \varphi(\tau_1^X) = j_1, \tau_1^X \leq E^\lambda | X(0) = x_0, \varphi(0) = i) \\ & = \mathbb{P}(X(\tau_n^X) = y_{\ell+1(j_n \in \mathcal{S}_-)}, \varphi(\tau_n^X) = j_n, \tau_n^X \leq E^\lambda | X(0) = x_0, \varphi(0) = i) \end{aligned} \quad (3.84)$$

where the swapping of limits and sums is justified as the sums are finite, and swapping limits and products is justified as the product is finite and all limits exist.

Therefore we have proved the following result

Lemma 3.14. *For all $\ell, \ell_0 \in \mathcal{K}$, $i, j_n \in \mathcal{S}$, $x_0 \in \mathcal{D}_{\ell_0, i}$, $n \geq 1$, then, as $p \rightarrow \infty$,*

$$\begin{aligned} & \mathbb{P}(L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} \leq E^\lambda | L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i), \\ & \rightarrow \mathbb{P}(X(\tau_n^X) = y_{\ell+1(j_n \in \mathcal{S}_-)}, \varphi(\tau_n^X) = j_n, \tau_n^X \leq E^\lambda | X(0) = x_0, \varphi(0) = i). \end{aligned} \quad (3.85)$$

3.5 Between the n th and $n + 1$ th change of level

Consider the Laplace transform

$$\int_{t=0}^{\infty} e^{-\lambda t} \int_{x \in \mathcal{D}_\ell} \mathbb{P}(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in dx, \phi^{(p)}(t) = j, \tau_n^{(p)} \leq t < \tau_{n+1}^{(p)} \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i) \psi(x) dx dt. \quad (3.86)$$

Partitioning on the time of the n th change of level and the phase and level at this time gives

$$\begin{aligned} & \int_{t=0}^{\infty} e^{-\lambda t} \int_{u_n=0}^t \sum_{j_n \in \mathcal{S}} \int_{x \in \mathcal{D}_\ell} \mathbb{P}(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in dx, \phi^{(p)}(t) = j, u_n < t \leq \tau_{n+1}^{(p)} \mid \\ & \quad L^{(p)}(u_n) = \ell, \phi^{(p)}(u_n) = j_n, \tau_n^{(p)} = u_n) \psi(x) dx \\ & \quad \mathbb{P}(L^{(p)}(u_n) = \ell, \phi^{(p)}(u_n) = j_n, \tau_n^{(p)} \in du_n \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i) \\ & \quad du_n dt \\ & = \sum_{j_n \in \mathcal{S}} \int_{x \in \mathcal{D}_\ell} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in dx, \phi^{(p)}(t) = j, \tau_n^{(p)} < t \leq \tau_{n+1}^{(p)} \mid \\ & \quad L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} = 0) dt \psi(x) dx \\ & \quad \mathbb{P}(L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i) \end{aligned} \quad (3.87)$$

by the time homogenous property of the QBD-RAP, the memoryless property of the exponential distribution, and the convolution theorem of Laplace transforms. We recognise the probability

$$\mathbb{P}(L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i) \quad (3.88)$$

as that appearing in Lemma (3.14). Hence (3.88) converges to

$$\mathbb{P}(X(\tau_n^X) = y_{\ell+1}(j_n \in \mathcal{S}_-), \varphi(\tau_n^X) = j_n, \tau_n^X \leq E^\lambda \mid X(0) = x_0, \varphi(0) = i) \quad (3.89)$$

as $p \rightarrow \infty$.

Now consider the expression

$$\begin{aligned} & \int_{x \in \mathcal{D}_\ell} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in dx, \phi^{(p)}(t) = j, \tau_n^{(p)} < t \leq \tau_{n+1}^{(p)} \mid L^{(p)}(\tau_n^{(p)}) = \ell, \\ & \quad \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} = 0) dt \psi(x) dx \end{aligned} \quad (3.90)$$

which appears as part of (3.87). We can rewrite (3.90) as

$$\begin{aligned} & \int_{x \in \mathcal{D}_\ell} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in dx, \phi^{(p)}(t) = j, t \leq \tau_1^{(p)} \mid L^{(p)}(0) = \ell, \\ & \quad \phi^{(p)}(0) = j_n) dt \psi(x) dx \\ &= \int_{x \in \mathcal{D}_\ell} \hat{f}^{\ell, (p)}(\lambda)(x, j; x_0, j_n) \psi(x) dx \end{aligned} \quad (3.91)$$

Applying Lemma 3.11, then (3.91) converges to

$$\begin{aligned} & \int_{x \in \mathcal{D}_\ell} \hat{\mu}^\ell(\lambda)(x, j; y_{\ell+1}(j_n \in \mathcal{S}_-), j_n) \psi(x) dx \\ &= \int_{x \in \mathcal{D}_\ell} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(X(t) \in dx, \varphi(t) = j, t \leq \tau_1^X \mid X(0) = y_{\ell+1}(j_n \in \mathcal{S}_-), \varphi(0) = j_n) \psi(x) dx. \end{aligned} \quad (3.92)$$

Since the fluid queue is time homogeneous then we can write (3.92) as

$$\begin{aligned} & \int_{x \in \mathcal{D}_\ell} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(X(t) \in dx, \varphi(t) = j, \tau_n^X < t \leq \tau_{n+1}^X \mid X(\tau_n^X) = y_{\ell+1}(j_n \in \mathcal{S}_-), \\ & \quad \varphi(\tau_n^X) = j_n, \tau_n^X \leq E^\lambda) \psi(x) dx \end{aligned} \quad (3.93)$$

Returning to right-hand side of (3.87) and taking the limit as $p \rightarrow \infty$,

$$\begin{aligned} & \lim_{p \rightarrow \infty} \sum_{j_n \in \mathcal{S}} \int_{x \in \mathcal{D}_\ell} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in dx, \phi^{(p)}(t) = j, \tau_n^{(p)} < t \leq \tau_{n+1}^{(p)} \mid \\ & \quad L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} = 0) dt \psi(x) dx \\ & \quad \mathbb{P}(L^{(p)}(\tau_n) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i) \\ &= \sum_{j_n \in \mathcal{S}} \lim_{p \rightarrow \infty} \int_{x \in \mathcal{D}_\ell} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in dx, \phi^{(p)}(t) = j, \tau_n^{(p)} < t \leq \tau_{n+1}^{(p)} \mid \\ & \quad L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} = 0) dt \psi(x) dx \\ & \quad \lim_{p \rightarrow \infty} \mathbb{P}(L^{(p)}(\tau_n^{(p)}) = \ell, \phi^{(p)}(\tau_n^{(p)}) = j_n, \tau_n^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \\ & \quad \phi^{(p)}(0) = i) \end{aligned}$$

since the sum is finite and the limits exist. Replacing the limits with their limiting values gives

$$\sum_{j_n \in \mathcal{S}} \int_{x \in \mathcal{D}_\ell} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(X(t) \in dx, \varphi(t) = j_n, \tau_n^X < t \leq \tau_{n+1}^X \mid X(\tau_n^X) = y_{\ell+1}(j_n \in \mathcal{S}_-),$$

$$\begin{aligned}
& \varphi(\tau_n^X) = j_n, \tau_n^X \leq E^\lambda) \psi(x) \, dx \\
& \mathbb{P}(X(\tau_n^X) = y_{\ell+1(j_n \in \mathcal{S}_-)}, \varphi(\tau_n^X) = j_n, \tau_n^X \leq E^\lambda \mid X(0) = x_0, \varphi(0) = i) \\
& = \int_{t=0}^{\infty} e^{-\lambda t} \int_{x \in \mathcal{D}_\ell} \mathbb{P}(X(t) \in dx, \varphi(t) = j_n, \tau_n^X < t \leq \tau_{n+1}^X \mid X(0) = x_0, \varphi(0) = i) \psi(x) \, dx \, dt
\end{aligned} \tag{3.94}$$

Hence we have shown the following result

Lemma 3.15. *For $\ell, \ell_0 \in \mathcal{K}$, $i, j \in \mathcal{S}$, $n \geq 0$, then, as $p \rightarrow \infty$,*

$$\begin{aligned}
& \int_{t=0}^{\infty} e^{-\lambda t} \int_{x \in \mathcal{D}_\ell} \mathbb{P}(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in dx, \phi^{(p)}(t) = j, \tau_n^{(p)} < t \leq \tau_{n+1}^{(p)} \mid L^{(p)}(0) = \ell_0, \\
& \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i) \psi(x) \, dx \, dt \\
& \rightarrow \int_{t=0}^{\infty} e^{-\lambda t} \int_{x \in \mathcal{D}_\ell} \mathbb{P}(X(t) \in dx, \varphi(t) = j_n, \tau_n^X < t \leq \tau_{n+1}^X \mid X(0) = x_0, \varphi(0) = i) \psi(x) \, dx \, dt.
\end{aligned} \tag{3.95}$$

3.6 To global convergence

3.6.1 First, another domination condition

To extend Lemma 3.15 to show a global convergence result we will use the Dominated Convergence Theorem. Here we show some geometric bounds on the probability of n level changes of the QBD-RAP and fluid queue, which ultimately serve as a dominating function in the Dominated Convergence Theorem.

Lemma 3.16. *For all $i \in \mathcal{S}_+ \cup \mathcal{S}_-$, and $n \geq 2$,*

$$\mathbb{P}(\tau_n \leq E^\lambda \mid \phi(\tau_{n-1}) = i, \tau_{n-1} \leq E^\lambda) \leq b, \tag{3.96}$$

where

$$b = \min \left\{ 1 - e^{-q(\Delta + \varepsilon)} [1 - e^{q\varepsilon - \lambda\Delta/|c_{\min}|}] + \frac{\text{Var}(Z)}{\varepsilon^2} + |r_1|, \frac{q}{q + \lambda} \right\}$$

and

$$|r_1| \leq 2G \frac{\text{Var}(Z)}{\varepsilon^2} + 2L\varepsilon.$$

Note that b and r_1 depend on p which has been suppressed to simplify notation. When explicitly needed, we use a superscript p to denote this dependence.

Proof. For the QBD-RAP, changes of level can only occur when $i \in \mathcal{S}_+ \cup \mathcal{S}_-$.

Suppose that the phase at time τ_{n-1} is $i \in \mathcal{S}_+$ and that at time τ_{n-1} the QBD-RAP is not at a boundary. The arguments for an initial phase $i \in \mathcal{S}_-$ are analogous. For the QBD-RAP to change level or hit a boundary or leave a boundary one of two things must happen, either;

1. the fluid remains in phase i until there is a change of level or boundary hit, or
2. the fluid changes phase before there is a change of level or a boundary is hit.

Hence, for sample paths which contribute to the Laplace transform, one of two things must happen, either;

1. the fluid remains in phase i until there is a change of level or a boundary is hit and E^λ does not occur before the change of level, or,
2. the fluid changes phase before there is a change of level or a boundary is hit and E^λ does not occur before the change of phase.

The probability of 1. is

$$\int_{x=0}^{\infty} \alpha e^{\mathbf{S}x} \mathbf{e} e^{(T_{ii}-\lambda)x/|c_i|} dx = e^{(T_{ii}-\lambda)\Delta/|c_i|} + r_1, \quad (3.97)$$

by Lemma A.1.

The probability of 2 is

$$\begin{aligned} \int_{x=0}^{\infty} \alpha e^{\mathbf{S}x} \mathbf{e} e^{(T_{ii}-\lambda)x/|c_i|} (-T_{ii}/|c_i|) dx &= \int_{x=0}^{\Delta+\varepsilon} \alpha e^{\mathbf{S}x} \mathbf{e} e^{(T_{ii}-\lambda)x/|c_i|} (-T_{ii}/|c_i|) dx \\ &+ \int_{x=\Delta+\varepsilon}^{\infty} \alpha e^{\mathbf{S}x} \mathbf{e} e^{(T_{ii}-\lambda)x/|c_i|} (-T_{ii}/|c_i|) dx. \end{aligned} \quad (3.98)$$

Now, since $\alpha e^{\mathbf{S}x} \mathbf{e} \leq 1$ for $x \leq \Delta + \varepsilon$ then the first term on the right-hand side of (3.98) is less than or equal to

$$\int_{x=0}^{\Delta+\varepsilon} e^{(T_{ii}-\lambda)x/|c_i|} (-T_{ii}/|c_i|) dx \leq \int_{x=0}^{\Delta+\varepsilon} e^{T_{ii}/|c_i|x} (-T_{ii}/|c_i|) dx = 1 - e^{T_{ii}/|c_i|(\Delta+\varepsilon)}.$$

By Chebyshev's inequality, $\alpha e^{\mathbf{S}x} \mathbf{e} \leq \frac{\text{Var}(Z)}{\varepsilon^2}$ for $x > \Delta + \varepsilon$, hence the second term on the right-hand side of (3.98) is less than or equal to

$$\int_{x=\Delta+\varepsilon}^{\infty} \frac{\text{Var}(Z)}{\varepsilon^2} e^{(T_{ii}-\lambda)x/|c_i|} (-T_{ii}/|c_i|) dx \leq \frac{\text{Var}(Z)}{\varepsilon^2}.$$

Putting these together, then the right-hand side of (3.98) is less than or equal to

$$1 - e^{T_{ii}(\Delta+\varepsilon)/|c_i|} + \frac{\text{Var}(Z)}{\varepsilon^2}. \quad (3.99)$$

Combining (3.97) and (3.99), then $\mathbb{P}(\tau_n \leq E^\lambda \mid \phi(\tau_{n-1}) = i, \tau_{n-1} \leq E^\lambda)$ is less than or equal to

$$\begin{aligned} & e^{(T_{ii}-\lambda)\Delta/|c_i|} + |r_1| + 1 - e^{T_{ii}(\Delta+\varepsilon)/|c_i|} + \frac{\text{Var}(Z)}{\varepsilon^2} \\ &= 1 - e^{T_{ii}(\Delta+\varepsilon)/|c_i|} \left[1 - e^{(-T_{ii}\varepsilon-\lambda\Delta)/|c_i|} \right] + \frac{\text{Var}(Z)}{\varepsilon^2} + |r_1| \\ &= 1 - e^{-q(\Delta+\varepsilon)} \left[1 - e^{q\varepsilon-\lambda\Delta/|c_{\min}|} \right] + \frac{\text{Var}(Z)}{\varepsilon^2} + |r_1|, \end{aligned} \quad (3.100)$$

since $-T_{ii}/|c_i| \leq q$ and $\lambda\Delta/|c_i| \leq \lambda\Delta/c_{\min}$ for all $i \in \mathcal{S}_+ \cup \mathcal{S}_-$.

Now consider the QBD-RAP at a boundary. To leave the boundary there must be at-least one change of phase before E^λ . By a uniformisation argument, the probability of at-least one change of phase before E^λ is $q/(q + \lambda)$. \square

Lemma 3.17. For $n \geq 2$, $i \in \mathcal{S}_+ \cup \mathcal{S}_-$,

$$\mathbb{P}(\tau_n \leq E^\lambda \mid \phi(\tau_1) = i, \tau_1 \leq E^\lambda) \leq b^{n-1}. \quad (3.101)$$

Proof. The proof is by induction.

For the base case, set $n = 2$ and apply Lemma 3.16.

Now, assume the induction hypothesis $\mathbb{P}(\tau_{n-1} \leq E^\lambda \mid \phi(\tau_1) = i, \tau_1 \leq E^\lambda) \leq b^{n-2}$ for arbitrary $n \geq 3$.

Since $\{\tau_{n-1} \leq E^\lambda\}$ is a subset of $\{\tau_n \leq E^\lambda\}$, then

$$\mathbb{P}(\tau_n \leq E^\lambda \mid \phi(\tau_1) = i, \tau_1 \leq E^\lambda) = \mathbb{P}(\tau_n \leq E^\lambda, \tau_{n-1} \leq E^\lambda \mid \phi(\tau_1) = i, \tau_1 \leq E^\lambda). \quad (3.102)$$

Now partition (3.102) on the phase at time τ_{n-1} ,

$$\begin{aligned} & \sum_{j_{n-1} \in \mathcal{S}} \mathbb{P}(\tau_n \leq E^\lambda, \tau_{n-1} \leq E^\lambda, \phi(\tau_{n-1}) = j_{n-1} \mid \phi(\tau_1) = i, \tau_1 \leq E^\lambda) \\ &= \sum_{j_{n-1} \in \mathcal{S}} \mathbb{P}(\tau_n \leq E^\lambda \mid \phi(\tau_{n-1}) = j_{n-1}, \tau_{n-1} \leq E^\lambda) \\ & \quad \times \mathbb{P}(\phi(\tau_{n-1}) = j_{n-1}, \tau_{n-1} \leq E^\lambda \mid \phi(\tau_1) = i, \tau_1 \leq E^\lambda), \end{aligned} \quad (3.103)$$

by the strong Markov property of the QBD-RAP and the fact that $\mathbf{A}(\tau_{n-1}) = \boldsymbol{\alpha}$.

By Lemma 3.16 (3.103) is less than or equal to

$$\sum_{j_{n-1} \in \mathcal{S}} b \mathbb{P}(\phi(\tau_{n-1}) = j_{n-1}, \tau_{n-1} \leq E^\lambda \mid \phi(\tau_1) = i, \tau_1 \leq E^\lambda)$$

$$\begin{aligned}
&= b\mathbb{P}(\tau_{n-1} \leq E^\lambda \mid \phi(\tau_1) = i, \tau_1 \leq E^\lambda) \\
&\leq b \cdot b^{n-2},
\end{aligned} \tag{3.104}$$

by the induction hypothesis, and this completes the proof. \square

Corollary 3.18.

$$\begin{aligned}
&\left| \int_{t=0}^{\infty} e^{-\lambda t} \int_{x \in \mathcal{D}_\ell} \mathbb{P}(L(t) = \ell, \bar{X}(t) \in dx, \phi(t) = j, \tau_n \leq t < \tau_{n+1} \mid L(0) = \ell_0, \right. \\
&\quad \left. \mathbf{A}(0) = \mathbf{a}_{\ell_0, i}(x_0), \phi(0) = i) \psi(x) dx dt \right| \\
&\leq \frac{F b^{n-1}}{\lambda}
\end{aligned} \tag{3.105}$$

Proof. First, since $|\psi(x)| \leq F$, then the left-hand side of (3.105) is less than or equal to

$$\begin{aligned}
&\int_{t=0}^{\infty} e^{-\lambda t} \int_{x \in \mathcal{D}_\ell} \mathbb{P}(L(t) = \ell, \bar{X}(t) \in dx, \phi(t) = j, \tau_n \leq t < \tau_{n+1} \mid L(0) = \ell_0, \\
&\quad \mathbf{A}(0) = \mathbf{a}_{\ell_0, i}(x_0), \phi(0) = i) F dx dt \\
&= \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(L(t) = \ell, \bar{X}(t) \in \mathcal{D}_{\ell, j}, \phi(t) = j, \tau_n \leq t < \tau_{n+1} \mid L(0) = \ell_0, \\
&\quad \mathbf{A}(0) = \mathbf{a}_{\ell_0, i}(x_0), \phi(0) = i) F dt.
\end{aligned}$$

Partitioning on the time, τ_1 , of the 1st change of level, and the phase and level at time τ_1 ,

$$\begin{aligned}
&\int_{t=0}^{\infty} e^{-\lambda t} \int_{u_1=0}^t \sum_{j_1 \in \mathcal{S}} \sum_{\ell_1 \in \{\ell_0+1, \ell_0-1\} \cap \mathcal{K}} \mathbb{P}(L(t) = \ell, \bar{X}(t) \in \mathcal{D}_{\ell, j}, \phi(t) = j, \tau_n \leq t < \tau_{n+1} \\
&\quad \mid L(\tau_1) = \ell_1, \phi(\tau_1) = j_1, \tau_1 = u_1) \\
&\quad \mathbb{P}(L(\tau_1) = \ell_1, \phi(\tau_1) = j_1, \tau_1 \in du_1 \mid L(0) = \ell_0, \mathbf{A}(0) = \mathbf{a}_{\ell_0, i}(x_0), \phi(0) = i) F dt \\
&= \sum_{j_1 \in \mathcal{S}} \sum_{\ell_1 \in \{\ell_0+1, \ell_0-1\} \cap \mathcal{K}} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(L(t) = \ell, \bar{X}(t) \in \mathcal{D}_{\ell, j}, \phi(t) = j, \tau_n \leq t < \tau_{n+1} \\
&\quad \mid L(0) = \ell_1, \phi(0) = j_1, \tau_1 = 0) dt \\
&\quad \mathbb{P}(L(\tau_1) = \ell_1, \phi(\tau_1) = j_1, \tau_1 \leq E^\lambda \mid L(0) = \ell_0, \mathbf{A}(0) = \mathbf{a}_{\ell_0, i}(x_0), \phi(0) = i) F, \tag{3.106}
\end{aligned}$$

by the time-homogeneous property of the QBD-RAP and the convolution theorem for Laplace transforms.

The expression

$$\int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(L(t) = \ell, \bar{X}(t) \in \mathcal{D}_{\ell, j}, \phi(t) = j, \tau_n \leq t < \tau_{n+1} \mid L(0) = \ell_1, \phi(0) = j_1, \tau_1 = 0) dt$$

$$\begin{aligned}
&\leq \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(\tau_n \leq t \mid L(0) = \ell_1, \phi(0) = j_1, \tau_1 = 0) dt \\
&\leq b^{n-1} \int_{t=0}^{\infty} e^{-\lambda t} dt \\
&= b^{n-1} \frac{1}{\lambda},
\end{aligned} \tag{3.107}$$

by Lemma 3.17.

Using the bound (3.107) in (3.106), gives the result. \square

3.6.2 Global convergence

Consider the Laplace transform of the QBD-RAP

$$\begin{aligned}
&\int_{t=0}^{\infty} e^{-\lambda t} \int_{x \in \mathcal{D}_{\ell,j}} \mathbb{P}(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in dx, \phi^{(p)}(t) = j \mid L^{(p)}(0) = \ell_0, \\
&\quad \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0,i}^{(p)}(x_0), \phi^{(p)}(0) = i) \psi(x) dx dt.
\end{aligned} \tag{3.108}$$

Partition on the number of level changes by time t ,

$$\begin{aligned}
&\int_{x \in \mathcal{D}_{\ell,j}} \int_{t=0}^{\infty} e^{-\lambda t} \sum_{n=0}^{\infty} \mathbb{P}(L^{(p)}(t) = \ell, \mathbf{A}^{(p)}(t) \in d\mathbf{a}^{(p)}, \phi^{(p)}(t) = j, \tau_n^{(p)} < t \leq \tau_{n+1}^{(p)} \mid \\
&\quad L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0,i}^{(p)}(x_0), \phi^{(p)}(0) = i) dt \psi(x) dx \\
&= \sum_{n=0}^{\infty} \int_{x \in \mathcal{D}_{\ell,j}} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(L^{(p)}(t) = \ell, \mathbf{A}^{(p)}(t) \in d\mathbf{a}^{(p)}, \phi^{(p)}(t) = j, \tau_n^{(p)} < t \leq \tau_{n+1}^{(p)} \mid \\
&\quad L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0,i}^{(p)}(x_0), \phi^{(p)}(0) = i) dt \psi(x) dx.
\end{aligned} \tag{3.109}$$

By Lemma (3.15), each term in the sum (3.109) converges. Furthermore, for $n \geq 1$, each term is dominated by $(b^{(p)})^{n-1} F/\lambda$, from Corollary 3.18. The dominating terms $(b^{(p)})^{n-1} F/\lambda$ depend on p and may not be summable. However, for p sufficiently large, there exists a $p_0 < \infty$ and a B with $B < 1$ such that $b^{(p)} < B$ for all $p > p_0$.

Therefore we can apply the Dominated Convergence Theorem,

$$\begin{aligned}
&\lim_{p \rightarrow \infty} \sum_{n=0}^{\infty} \int_{x \in \mathcal{D}_{\ell,j}} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(L^{(p)}(t) = \ell, \mathbf{A}^{(p)}(t) \in d\mathbf{a}^{(p)}, \phi^{(p)}(t) = j, \tau_n^{(p)} < t \leq \tau_{n+1}^{(p)} \mid \\
&\quad L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0,i}^{(p)}(x_0), \phi^{(p)}(0) = i) dt \psi(x) dx \\
&= \sum_{n=0}^{\infty} \int_{x \in \mathcal{D}_{\ell,j}} \lim_{p \rightarrow \infty} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(L^{(p)}(t) = \ell, \mathbf{A}^{(p)}(t) \in d\mathbf{a}^{(p)}, \phi^{(p)}(t) = j, \tau_n^{(p)} < t \leq \tau_{n+1}^{(p)} \mid
\end{aligned}$$

$$\begin{aligned}
& L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i) dt \psi(x) dx \\
& = \sum_{n=0}^{\infty} \int_{t=0}^{\infty} e^{-\lambda t} \int_{x \in \mathcal{D}_{\ell}} \mathbb{P}(X(t) \in dx, \varphi(t) = j, \tau_n < t \leq \tau_{n+1} \mid X(0) = x_0, \varphi(0) = i) \\
& \quad \times \psi(x) dx dt.
\end{aligned}$$

where the limit is given by Lemma 3.15. Swapping the sum and integrals and by the law of total probability,

$$\begin{aligned}
& \int_{t=0}^{\infty} e^{-\lambda t} \int_{x \in \mathcal{D}_{\ell}} \sum_{n=0}^{\infty} \mathbb{P}(X(t) \in dx, \varphi(t) = j, \tau_n < t \leq \tau_{n+1} \mid X(0) = x_0, \varphi(0) = i) \psi(x) dx dt. \\
& = \int_{t=0}^{\infty} e^{-\lambda t} \int_{x \in \mathcal{D}_{\ell}} \mathbb{P}(X(t) \in dx, \varphi(t) = j \mid X(0) = x_0, \varphi(0) = i) \psi(x) dx dt.
\end{aligned}$$

Thus, we have shown the following result

Lemma 3.19. *For all $\ell_0, \ell \in \mathcal{K}$, $i, j \in \mathcal{S}$, $x_0 \in \mathcal{D}_{\ell_0, i}$, as $p \rightarrow \infty$,*

$$\begin{aligned}
& \int_{t=0}^{\infty} e^{-\lambda t} \int_{x \in \mathcal{D}_{\ell, j}} \mathbb{P}(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in dx, \phi^{(p)}(t) = j \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \\
& \quad \phi^{(p)}(0) = i) \psi(x) dx dt \\
& \rightarrow \int_{t=0}^{\infty} e^{-\lambda t} \int_{x \in \mathcal{D}_{\ell, j}} \mathbb{P}(X(t) \in dx, \varphi(t) = j \mid X(0) = x_0, \varphi(0) = i) \psi(x) dx dt.
\end{aligned}$$

Corollary 3.20. *Let $\psi(x, j) : \mathbb{R} \times \mathcal{S} \rightarrow \mathbb{R}$ be Lipschitz continuous and bounded functions, $|\psi(x, j)| \leq F$. For each $x_0 \in \mathbb{R}$, $i \in \mathcal{S}$, $\ell_0 \in \mathcal{K}$,*

$$\begin{aligned}
& \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{E} \left[\psi(\bar{X}^{(p)}(t), \phi^{(p)}(t)) \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i \right] dt \\
& \rightarrow \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{E} [\psi(X(t), \phi(t)) \mid X(0) = x_0, \varphi(0) = i] dt.
\end{aligned}$$

Proof. Consider the left-hand side

$$\begin{aligned}
& \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{E} \left[\psi(\bar{X}^{(p)}(t), \phi^{(p)}(t)) \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i \right] dt \\
& = \int_{t=0}^{\infty} e^{-\lambda t} \sum_{\ell \in \mathcal{K}} \sum_{j \in \mathcal{S}} \mathbb{E} \left[\psi(\bar{X}^{(p)}(t), \phi^{(p)}(t)) 1(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in \mathcal{D}_{\ell, j}, \phi^{(p)}(t) = j) \right. \\
& \quad \left. \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i \right] dt \\
& = \sum_{\ell \in \mathcal{K}} \sum_{j \in \mathcal{S}} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{E} \left[\psi(\bar{X}^{(p)}(t), \phi^{(p)}(t)) 1(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in \mathcal{D}_{\ell, j}, \phi^{(p)}(t) = j) \mid L
\end{aligned}$$

$$^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i \Big] dt. \quad (3.110)$$

By Lemma 3.19, for each $\ell \in \mathcal{K}$, $j \in \mathcal{S}$, the terms

$$\int_{t=0}^{\infty} e^{-\lambda t} \mathbb{E} \left[\psi(\bar{X}^{(p)}(t), \phi^{(p)}(t)) 1(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in \mathcal{D}_{\ell, j}, \phi^{(p)}(t) = j) \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i \right] dt \quad (3.111)$$

converge to

$$\int_{t=0}^{\infty} e^{-\lambda t} \mathbb{E} \left[\psi(X(t), \phi(t)) 1(X(t) \in \mathcal{D}_{\ell, j}, \varphi(t) = j) \mid X(0) = x_0, \varphi(0) = i \right] dt.$$

If \mathcal{K} is finite, we are done upon taking the limit of (3.110) as $p \rightarrow \infty$ and swapping the limit and the sums.

If \mathcal{K} is countably infinite, then for a given $k \in \mathcal{K}$, since ψ is bounded

$$\begin{aligned} & \left| \sum_{j \in \mathcal{S}} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{E} \left[\psi(\bar{X}^{(p)}(t), \phi^{(p)}(t)) 1(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in \mathcal{D}_{\ell, j}, \phi^{(p)}(t) = j) \mid L^{(p)}(0) = \ell_0, \right. \right. \\ & \quad \left. \left. \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i \right] dt \right| \\ & \leq F \sum_{j \in \mathcal{S}} \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{E} \left[1(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in \mathcal{D}_{\ell, j}, \phi^{(p)}(t) = j) \mid L^{(p)}(0) = \ell_0, \right. \\ & \quad \left. \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i \right] dt \\ & \leq F \int_{t=0}^{\infty} e^{-\lambda t} \mathbb{P}(L^{(p)}(t) = \ell, \bar{X}^{(p)}(t) \in \mathcal{D}_{\ell} \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i) dt \\ & \leq F \mathbb{P}(\tau_{|\ell - \ell_0|}^{(p)} \leq E^\lambda \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i), \end{aligned} \quad (3.112)$$

since, to be in level ℓ after starting in level ℓ_0 , there must be at least $|\ell_0 - \ell|$ changes of level. By Lemma 3.17 then (3.112) is bounded by $(b^{(p)})^{|\ell - \ell_0| - 1}$ for $|\ell - \ell_0| \geq 2$ and by 1 otherwise. Now, choose p_0 sufficiently large so that $b^{(p)} < B < 1$ for all $p > p_0$. Therefore, for all $p > p_0$, the terms in (3.110) are dominated by $F \min\{B^{|\ell - \ell_0| - 1}, 1\}$. Moreover

$$\begin{aligned} F \sum_{\ell \in \mathcal{K}} \min\{B^{|\ell - \ell_0| - 1}, 1\} & \leq 2 \sum_{n=1}^{\infty} B^{n-1} + 1 \\ & = \frac{2}{1 - B} + 1 \\ & < \infty, \end{aligned}$$

hence the dominating terms are summable. Hence we may apply the Dominated Convergence Theorem to swap the necessary limits and sums. \square

The Extended Continuity Theorem for Laplace transforms (Feller 1957, Chapter XIII, Theorem 2a) can now be used to claim that the QBD-RAP approximation scheme converges weakly to the fluid queue.

Theorem 3.21 (Extended Continuity Theorem). *For $p = 1, 2, \dots$ let U_p be a measure with Laplace transform ω_p . If $\omega_p(\lambda) \rightarrow \omega(\lambda)$ for $\lambda > a \geq 0$, then ω is the Laplace transform of a measure U and $U_p \rightarrow U$.*

Conversely, if $U_p \rightarrow U$ and the sequence $\{\omega_p(a)\}$ is bounded, then $\omega_p(\lambda) \rightarrow \omega(\lambda)$ for $\lambda > a$.

Thus, by the Extended Continuity Theorem 3.21

$$\begin{aligned} & \mathbb{E} \left[\psi(\bar{X}^{(p)}(t), \phi^{(p)}(t)) \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i \right] \\ & \rightarrow \mathbb{E} [\psi(X(t), \phi(t)) \mid X(0) = x_0, \varphi(0) = i] \end{aligned}$$

weakly in t as $p \rightarrow \infty$. Now,

$$\mathbb{E} [\psi(X(t), \phi(t)) \mid X(0) = x_0, \varphi(0) = i]$$

is a continuous function of t (it is a Feller semi-group), moreover, for $p < \infty$,

$$\mathbb{E} \left[\psi(\bar{X}^{(p)}(t), \phi^{(p)}(t)) \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i \right]$$

is also continuous in t . However, the limit

$$\lim_{p \rightarrow \infty} \mathbb{E} \left[\psi(\bar{X}^{(p)}(t), \phi^{(p)}(t)) \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i \right]$$

may not be continuous in t . We can claim that

$$\begin{aligned} & \mathbb{E} \left[\psi(\bar{X}^{(p)}(t), \phi^{(p)}(t)) \mid L^{(p)}(0) = \ell_0, \mathbf{A}^{(p)}(0) = \mathbf{a}_{\ell_0, i}^{(p)}(x_0), \phi^{(p)}(0) = i \right] \\ & \rightarrow \mathbb{E} [\psi(X(t), \phi(t)) \mid X(0) = x_0, \varphi(0) = i] \end{aligned}$$

for almost all $t \geq 0$. At such values of t , since ψ is an arbitrary bounded, Lipschitz continuous function, then the Portmanteau Theorem states that the QBD-RAP approximation scheme converges in distribution to the fluid queue.

A sufficient condition to upgrade the convergence from weak to point-wise (in the variable t) is to show that for $t \geq 0$

$$\sup_p \mathbb{E} [\psi(\bar{X}(t), \phi(t)) \mid L(0) = \ell_0, \mathbf{A}(0) = \mathbf{a}_{\ell_0, i}(x_0), \phi(0) = i] \leq M(t) < \infty$$

and the sequence $\mathbb{E} [\psi(\bar{X}(t), \phi(t)) \mid L(0) = \ell_0, \mathbf{A}(0) = \mathbf{a}_{\ell_0, i}(x_0), \phi(0) = i]$ is eventually equicontinuous in t . That is, for every $\varepsilon > 0$ there exists a $\delta(t, \varepsilon) > 0$ and an $p_0(t, \varepsilon)$ such that $|t - u| < \delta(t, \varepsilon)$ implies that

$$\left| \mathbb{E} [\psi(\bar{X}(t), \phi(t)) \mid L(0) = \ell_0, \mathbf{A}(0) = \mathbf{a}_{\ell_0, i}(x_0), \phi(0) = i] - \mathbb{E} [\psi(\bar{X}(u), \phi(u)) \mid L(0) = \ell_0, \mathbf{A}(0) = \mathbf{a}_{\ell_0, i}(x_0), \phi(0) = i] \right| < \varepsilon \quad (3.113)$$

for all $p \geq p_0(t, \varepsilon)$. This is an area for future work.

Remark 3.22. *While the convergence result implies a loose bound on the rate of convergence, it applies to a wide class of QBD-RAP constructions. The main cause of the loose bound on the convergence rate is due to our reliance on Chebyshev's inequality. This is necessary as we try to make minimal assumptions about the matrix-exponential distributions used in the construction. In practice, we use a class of concentrated matrix-exponential distributions found numerically in (Horváth et al. 2020), and for which there is relatively little known about their properties. This necessitates the generality of the convergence result.*

3.7 Extension to arbitrary (but fixed) discretisation structures

Throughout, we have assumed that all intervals are of width Δ , i.e. $|y_{\ell+1} - y_\ell| = \Delta$, and that on every interval the dynamics of the fluid queue are modelled based on the same matrix exponential representation $(\boldsymbol{\alpha}, \mathbf{S}, \mathbf{s})$. These assumptions are, in fact, not necessary, but they do serve to simplify the presentation. The convergence results can be extended to use different sequences of matrix exponential representations on each interval, provided that for each sequence of matrix exponential distributions, the variance tends to 0. Moreover, we can extend the results to intervals of arbitrary width, provided that the width of the intervals is not arbitrarily small. Here we describe how one would prove such results.

The arguments which prove Lemma 3.11 are independent of all other levels/intervals, i.e. the hypotheses of the Lemma depend only on the interval \mathcal{D}_{ℓ_0} , and the sequence of matrix exponential distributions used to model the behaviour of the fluid queue on this interval and not on any other interval. Thus Lemma 3.11 holds independently on each interval, as does Corollary 3.13.

Let the width of an interval \mathcal{D}_{ℓ_0} be $\Delta_{\ell_0} = y_{\ell_0+1} - y_{\ell_0}$ and suppose that sequence of matrix exponential random variable used to model the dynamics of the fluid queue on the interval \mathcal{D}_{ℓ_0} is $Z_{\ell_0}^{(p)}$. Regarding Lemma 3.16, we can extend it the following version,

Lemma 3.23. Assume $\inf_{\ell_0} \Delta_{\ell_0} > 0$ and $\sup_{\ell_0} \text{Var}(Z_{\ell_0}) < \infty$ exist. Then, for all $i \in \mathcal{S}_+ \cup \mathcal{S}_-$, $\ell_0 \in \mathcal{K} \setminus \{-1, K+1\}$, and $n \geq 2$,

$$\mathbb{P}(\tau_n^{(p)} \leq E^\lambda \mid L^{(p)}(\tau_{n-1}^{(p)}) = \ell_0, \phi^{(p)}(\tau_{n-1}^{(p)}) = i, \tau_{n-1}^{(p)} \leq E^\lambda) \leq b_{\ell_0}^{(p)}, \quad (3.114)$$

where

$$b_{\ell_0}^{(p)} = 1 - e^{-q(\Delta_{\ell_0} + \varepsilon_{\ell_0}^{(p)})} \left[1 - e^{q\varepsilon_{\ell_0}^{(p)} - \lambda \Delta_{\ell_0} / |c_{\min}|} \right] + \frac{\text{Var}(Z_{\ell_0}^{(p)})}{(\varepsilon^{(p)})^2} + |r_{1, \ell_0}^{(p)}|$$

and

$$|r_{1, \ell_0}^{(p)}| \leq 2G \frac{\text{Var}(Z_{\ell_0}^{(p)})}{(\varepsilon_{\ell_0}^{(p)})^2} + 2L\varepsilon_{\ell_0}^{(p)}.$$

Hence, for all $i \in \mathcal{S}_+ \cup \mathcal{S}_-$, and $n \geq 2$,

$$\mathbb{P}(\tau_n^{(p)} \leq E^\lambda \mid \phi^{(p)}(\tau_{n-1}^{(p)}) = i, \tau_{n-1}^{(p)} \leq E^\lambda) \leq b^{(p)}, \quad (3.115)$$

where

$$b^{(p)} = \max \left\{ \sup_{\ell_0} b_{\ell_0}^{(p)}, \frac{q}{\lambda + q} \right\}.$$

Proof. For the proof of (3.114) follow the same arguments as in the proof of Lemma 3.16. The bound in equation follows by the assumptions that $\inf_{\ell_0} \Delta_{\ell_0}$ and $\sup_{\ell_0} \text{Var}(Z_{\ell_0}^{(p)})$ exist, then

$$\begin{aligned} & \mathbb{P}(\tau_n^{(p)} \leq E^\lambda \mid \phi^{(p)}(\tau_{n-1}^{(p)}) = i, \tau_{n-1}^{(p)} \leq E^\lambda) \\ & \leq \sup_{\ell_0} \mathbb{P}(\tau_n^{(p)} \leq E^\lambda \mid L^{(p)}(\tau_{n-1}^{(p)}) = \ell_0, \phi^{(p)}(\tau_{n-1}^{(p)}) = i, \tau_{n-1}^{(p)} \leq E^\lambda) \\ & \leq \max \left\{ \sup_{\ell_0} b_{\ell_0}^{(p)}, \frac{q}{\lambda + q} \right\}. \end{aligned}$$

□

Given Lemma 3.23, then an equivalent of Lemma 3.17 remains true, the proof of which follows verbatim except with the use of Lemma 3.16 replaced by Lemma 3.23. Corollary 3.18 remains true without modification. Lemma 3.19 and Corollary 3.20 remain true without modification provided that $\lim_{p \rightarrow \infty} \text{Var}(Z_\ell^{(p)}) \rightarrow 0$ for all ℓ .

Appendix A

Technical results for coverage

A.1 Some bounds for integrating against matrix exponential distributions.

This appendix proves some technical Lemmas about integrating functions with respect to matrix exponential distributions.

As with Section 3, we use the superscript (p) to denote dependence on the underlying choice of matrix exponential random variable $Z^{(p)}$. However, to simplify notation, we may omit the super script (p) where it is not explicitly needed. We show results for an arbitrary parameter $\varepsilon > 0$. Keep in mind the ultimate intention is to show convergence, for which we choose this parameter to be $\varepsilon^{(p)} = \text{Var} (Z^{(p)})^{1/3}$. Other notations, previously defined and which depend on p are $\alpha^{(p)}$, $\alpha_0^{i, \ell_0, (p)}(x_0)$, $\mathbf{S}^{(p)}$, $\mathbf{s}^{(p)}$, $\varepsilon^{(p)}$, $V^{(p)}(x)$, $R_{V,1}^{(p)}$, $R_{V,2}^{(p)}$, $\mathbf{D}^{(p)}$, $\mathcal{A}^{(p)}$.

A.1.1 One integral

Here we show a collection of results that show that integrating functions against a ME density function, or against the density function of a ME conditional on the ME-life-time surviving until some time $u < \Delta - \varepsilon$ where $\Delta = \mathbb{E}[Z]$ is the mean of the matrix exponential distribution, approximates integrating said function against a Kronecker delta situated at the Δ , provided the variance of the ME is sufficiently low.

Lemma A.1. *Let g be a function satisfying Assumptions 3.1, then, for $u \leq \Delta - \varepsilon$,*

$$\int_{x=0}^{\infty} g(x) \alpha e^{\mathbf{S}(x+u)} \mathbf{s} dx = g(\Delta - u) + r_1,$$

where

$$|r_1| \leq 2G \frac{\text{Var}(Z)}{\varepsilon^2} + 2L\varepsilon.$$

The proof follows closely that of (Horváth et al. 2020, Appendix A, Theorem 4).

Proof. By a change of variables,

$$\begin{aligned} & \left| \int_{x=0}^{\infty} g(x) \alpha e^{\mathbf{S}(x+u)} \mathbf{s} \, dx - g(\Delta - u) \right| \\ &= \left| \int_{x=u}^{\infty} g(x-u) \alpha e^{\mathbf{S}x} \mathbf{s} \, dx - g(\Delta - u) \right| \\ &= \left| \int_{x=u}^{\infty} g(x-u) \alpha e^{\mathbf{S}x} \mathbf{s} \, dx - \int_{x=u}^{\infty} g(\Delta - u) \alpha e^{\mathbf{S}x} \mathbf{s} \, dx - g(\Delta - u) (1 - \alpha e^{\mathbf{S}u} \mathbf{e}) \right|. \end{aligned}$$

By the triangle inequality this is less than or equal to

$$\begin{aligned} & \left| \int_{x=u}^{\infty} (g(x-u) - g(\Delta - u)) \alpha e^{\mathbf{S}x} \mathbf{s} \, dx \right| + |g(\Delta - u) (1 - \alpha e^{\mathbf{S}u} \mathbf{e})| \\ &= \left| \int_{x=u}^{\infty} (g(x-u) - g(\Delta - u)) \alpha e^{\mathbf{S}x} \mathbf{s} \, dx \right| + \left| \int_{x=0}^u g(\Delta - u) \alpha e^{\mathbf{S}x} \mathbf{s} \, dx \right| \\ &\leq d_1 + d_2 \end{aligned}$$

where

$$\begin{aligned} d_1 &= \left| \int_{x=0}^u g(\Delta - u) \alpha e^{\mathbf{S}x} \mathbf{s} \, dx \right| + \left| \int_{x=u}^{\Delta-\varepsilon} (g(x-u) - g(\Delta - u)) \alpha e^{\mathbf{S}x} \mathbf{s} \, dx \right| \\ &\quad + \left| \int_{x=\Delta+\varepsilon}^{\infty} (g(x-u) - g(\Delta - u)) \alpha e^{\mathbf{S}x} \mathbf{s} \, dx \right|, \\ d_2 &= \left| \int_{x=\Delta-\varepsilon}^{\Delta+\varepsilon} (g(t-u) - g(\Delta - u)) \alpha e^{\mathbf{S}x} \mathbf{s} \, dx \right|. \end{aligned}$$

Applying the triangle inequality to d_1 ,

$$\begin{aligned} d_1 &\leq \int_{x=u}^{\Delta-\varepsilon} |g(x-u) - g(\Delta - u)| \alpha e^{\mathbf{S}x} \mathbf{s} \, dx + \int_{x=\Delta+\varepsilon}^{\infty} |g(x-u) - g(\Delta - u)| \alpha e^{\mathbf{S}x} \mathbf{s} \, dx \\ &\quad + \left| \int_{x=0}^u g(\Delta - u) \alpha e^{\mathbf{S}x} \mathbf{s} \, dx \right|. \end{aligned} \tag{A.1}$$

Since $|g(x)| \leq G$, then (A.1) is less than or equal to

$$2G \left(\int_{x=u}^{\Delta-\varepsilon} \alpha e^{\mathbf{S}x} \mathbf{s} \, dx + \int_{x=\Delta+\varepsilon}^{\infty} \alpha e^{\mathbf{S}x} \mathbf{s} \, dx + \int_{x=0}^u \alpha e^{\mathbf{S}x} \mathbf{s} \, dx \right) = 2G \mathbb{P}(|Z - \Delta| > \varepsilon). \tag{A.2}$$

By Chebyshev's inequality,

$$2G\mathbb{P}(|Z - \Delta| > \varepsilon) \leq 2G \frac{\text{Var}(Z)}{\varepsilon^2}. \quad (\text{A.3})$$

For the term d_2 we have

$$\begin{aligned} d_2 &= \left| \int_{x=\Delta-\varepsilon}^{\Delta+\varepsilon} (g(x-u) - g(\Delta-u)) \boldsymbol{\alpha} e^{\mathbf{S}x} \mathbf{s} \, dx \right| \\ &\leq \int_{x=\Delta-\varepsilon}^{\Delta+\varepsilon} |g(x-u) - g(\Delta-u)| \boldsymbol{\alpha} e^{\mathbf{S}x} \mathbf{s} \, dx \\ &\leq \int_{x=\Delta-\varepsilon}^{\Delta+\varepsilon} 2L\varepsilon \boldsymbol{\alpha} e^{\mathbf{S}x} \mathbf{s} \, dx \\ &= 2L\varepsilon \mathbb{P}(Z \in (\Delta - \varepsilon, \Delta + \varepsilon)) \\ &\leq 2L\varepsilon, \end{aligned}$$

where the first inequality is the triangle inequality and the second inequality is from the Lipschitz property of g in Assumption 3.1(iv). Hence there is some r_1 such that

$$\left| \int_{x=0}^{\infty} g(x) \boldsymbol{\alpha} e^{\mathbf{S}(x+u)} \mathbf{s} \, dx - g(\Delta-u) \right| = |r_1| \leq 2G \frac{\text{Var}(Z)}{\varepsilon^2} + 2L\varepsilon,$$

and this completes the proof. \square

The error term $r_1^{(p)}$ depends on p , as it is defined by $Z^{(p)}$ and $\varepsilon^{(p)}$, but we have omitted the superscript p here. Note that, upon choosing $\varepsilon = \text{Var}(Z^{(p)})^{1/3}$, the error term $|r_1^{(p)}|$ is at most $O\left(\text{Var}(Z^{(p)})^{1/3}\right)$, which tends to 0 as $p \rightarrow \infty$.

Corollary A.2. *Let g be a function satisfying the Assumptions 3.1. For $u \leq \Delta - \varepsilon$,*

$$\int_{x=0}^{\infty} \frac{\boldsymbol{\alpha} e^{\mathbf{S}(x+u)} \mathbf{s}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} g(x) \, dx = g(\Delta-u) + r_2,$$

where

$$|r_2| \leq 3G \frac{\text{Var}(Z)}{\varepsilon^2} + 2L\varepsilon.$$

Proof. Observe that Chebyshev's inequality gives

$$\begin{aligned} \boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e} &= \mathbb{P}(Z > u) \\ &\geq \mathbb{P}(|Z - \Delta| \leq \varepsilon) \\ &\geq 1 - \frac{\text{Var}(Z)}{\varepsilon^2} \end{aligned}$$

$$=: 1 - \delta.$$

Now, since $1 - \delta \leq \alpha e^{S^u} e \leq 1$, then

$$\int_{x=0}^{\infty} \alpha e^{S(x+u)} s g(x) dx \leq \int_{x=0}^{\infty} \frac{\alpha e^{S(x+u)} s}{\alpha e^{S^u} e} g(x) dx \leq \frac{1}{1 - \delta} \int_{x=0}^{\infty} \alpha e^{S(x+u)} s g(x) dx.$$

By Lemma A.1 we have

$$g(\Delta - u) + r_1 \leq \int_{x=0}^{\infty} \frac{\alpha e^{S(x+u)} s}{\alpha e^{S^u} e} g(x) dx \leq \frac{g(\Delta - u) + r_1}{1 - \delta}.$$

Multiplying by $1 - \delta$, then subtracting $g(\Delta - u)$ and adding $\int_{x=0}^{\infty} \frac{\alpha e^{S(x+u)} s}{\alpha e^{S^u} e} g(x) dx \delta$ gives

$$\begin{aligned} & r_1(1 - \delta) - g(\Delta - u)\delta + \int_{x=0}^{\infty} \frac{\alpha e^{S(x+u)} s}{\alpha e^{S^u} e} g(x) dx \delta \\ & \leq \int_{x=0}^{\infty} \frac{\alpha e^{S(x+u)} s}{\alpha e^{S^u} e} g(x) dx - g(\Delta - u) \\ & \leq r_1 + \int_{x=0}^{\infty} \frac{\alpha e^{S(x+u)} s}{\alpha e^{S^u} e} g(x) dx \delta. \end{aligned}$$

The right-hand side is bounded above as

$$r_1 + \int_{x=0}^{\infty} \frac{\alpha e^{S(x+u)} s}{\alpha e^{S^u} e} g(x) dx \delta \leq r_1 + G\delta.$$

The left-hand side is bounded below as

$$r_1(1 - \delta) - g(\Delta - u)\delta + \int_{x=0}^{\infty} \frac{\alpha e^{S(x+u)} s}{\alpha e^{S^u} e} g(x) dx \delta \geq r_1(1 - \delta) - g(\Delta - u)\delta.$$

Hence,

$$\left| \int_{x=0}^{\infty} \frac{\alpha e^{S(x+u)} s}{\alpha e^{S^u} e} g(x) dx - g(\Delta - u) \right| \leq \max(r_1(1 - \delta) + g(\Delta - u)\delta, r_1 + G\delta) \quad (\text{A.4})$$

and therefore

$$\int_{x=0}^{\infty} \frac{\alpha e^{S(x+u)} s}{\alpha e^{S^u} e} g(x) dx = g(\Delta - u) + r_2, \quad (\text{A.5})$$

where

$$|r_2| \leq \max(r_1(1 - \delta) + g(\Delta - u)\delta, r_1 + G\delta)$$

$$\begin{aligned}
&\leq r_1 + G\delta \\
&= 3G \frac{\text{Var}(Z)}{\varepsilon^2} + 2L\varepsilon.
\end{aligned} \tag{A.6}$$

This completes the proof. \square

The error term $r_2^{(p)}$ also depends on p , as it is defined by $Z^{(p)}$ and $\varepsilon^{(p)}$, but we have omitted the superscript p here. Choosing $\varepsilon = \text{Var}(Z^{(p)})$, the error term $|r_2^{(p)}|$ is at most $O\left(\text{Var}(Z^{(p)})^{1/3}\right)$, which tends to 0 as $p \rightarrow \infty$.

Corollary A.3. *Let g be a function satisfying the Assumptions 3.1. For $u \leq \Delta - \varepsilon$, $v \geq 0$,*

$$\int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(x+u+v)} \mathbf{s}}{\alpha e^{\mathbf{S}u} \mathbf{e}} g(x) dx = g(\Delta - u - v) 1(u + v \leq \Delta - \varepsilon) + r_3(u + v),$$

where

$$|r_3(u + v)| \leq \begin{cases} r_2 & u + v \leq \Delta - \varepsilon, \\ G & u + v \in (\Delta - \varepsilon, \Delta + \varepsilon), \\ G \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2} & u + v \geq \Delta + \varepsilon. \end{cases}$$

Proof. For $u + v \leq \Delta - \varepsilon$ we use similar arguments those as in the proof of Corollary A.2. We have

$$\int_{x=0}^{\infty} \alpha e^{\mathbf{S}(x+u+v)} \mathbf{s} g(x) dx \leq \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(x+u+v)} \mathbf{s}}{\alpha e^{\mathbf{S}u} \mathbf{e}} g(x) dx \leq \frac{1}{1 - \delta} \int_{x=0}^{\infty} \alpha e^{\mathbf{S}(x+u+v)} \mathbf{s} g(x) dx.$$

By Lemma A.1

$$g(\Delta - u - v) + r_1 \leq \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(x+u+v)} \mathbf{s}}{\alpha e^{\mathbf{S}u} \mathbf{e}} g(x) dx \leq \frac{g(\Delta - u - v) + r_1}{1 - \delta}.$$

Multiplying by $1 - \delta$, then subtracting $g(\Delta - u - v)$ and adding $\int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(x+u+v)} \mathbf{s}}{\alpha e^{\mathbf{S}u} \mathbf{e}} g(x) dx \delta$ gives

$$\begin{aligned}
&r_1(1 - \delta) - g(\Delta - u - v)\delta + \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(x+u+v)} \mathbf{s}}{\alpha e^{\mathbf{S}u} \mathbf{e}} g(x) dx \delta \\
&\leq \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(x+u+v)} \mathbf{s}}{\alpha e^{\mathbf{S}u} \mathbf{e}} g(x) dx - g(\Delta - u - v) \\
&\leq r_1 + \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(x+u+v)} \mathbf{s}}{\alpha e^{\mathbf{S}u} \mathbf{e}} g(x) dx \delta.
\end{aligned}$$

The right-hand side is bounded above by

$$r_1 + \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(x+u+v)} \mathbf{s}}{\alpha e^{\mathbf{S}u} \mathbf{e}} g(x) dx \delta \leq r_1 + G\delta.$$

The left-hand side is bounded below by

$$r_1(1 - \delta) - g(\Delta - u - v)\delta + \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(x+u+v)} \mathbf{s}}{\alpha e^{\mathbf{S}u} \mathbf{e}} g(x) dx \delta \geq r_1(1 - \delta) - g(\Delta - u - v)\delta.$$

and ultimately,

$$\left| \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(x+u+v)} \mathbf{s}}{\alpha e^{\mathbf{S}u} \mathbf{e}} g(x) dx - g(\Delta - u - v) \right| \leq 3G \frac{\text{Var}(Z)}{\varepsilon^2} + 2L\varepsilon. \quad (\text{A.7})$$

For $u + v \in (\Delta - \varepsilon, \Delta + \varepsilon)$,

$$\int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(x+u+v)} \mathbf{s}}{\alpha e^{\mathbf{S}u} \mathbf{e}} g(x) dx \leq G\mathbb{P}(Z > u + v \mid Z > u) \leq G \quad (\text{A.8})$$

For $u + v \geq \Delta + \varepsilon$,

$$\int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(x+u+v)} \mathbf{s}}{\alpha e^{\mathbf{S}u} \mathbf{e}} g(x) dx \leq G \frac{\mathbb{P}(Z > u + v)}{\mathbb{P}(Z > u)} \leq G \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2}. \quad (\text{A.9})$$

□

The error term $r_3^{(p)}$ depends on p , as it is defined by $Z^{(p)}$ and $\varepsilon^{(p)}$, but we have omitted the superscript p here. Choosing $\varepsilon = \text{Var}(Z^{(p)})^{1/3}$ then, outside of the vanishingly small interval $u \in (\Delta - \varepsilon^{(p)}, \Delta + \varepsilon^{(p)})$, the error term $|r_3^{(p)}(u)|$ is bounded by $O(\text{Var}(Z^{(p)})^{1/3})$, which tends to 0 as $p \rightarrow \infty$. On $u \in (\Delta - \varepsilon^{(p)}, \Delta + \varepsilon^{(p)})$ the error term $|r_3^{(p)}(u)|$ is bounded by a constant which does not tend to 0 as $p \rightarrow \infty$. However, when we integrate a bounded function against $r_3^{(p)}(u)$, then the resulting integral tends to 0, i.e. for $|\psi(x)| \leq F$, $M < \infty$, $\int_0^M f(u)r_3^{(p)}(u) du \leq F\Delta|r_2^{(p)}| + 2GF\varepsilon^{(p)} + (M - \Delta)GF \frac{\text{Var}(Z^{(p)})/(\varepsilon^{(p)})^2}{1 - \text{Var}(Z^{(p)})/(\varepsilon^{(p)})^2} = O(\text{Var}(Z^{(p)})^{1/3}) \rightarrow 0$ as $p \rightarrow \infty$. This is the context in which we we apply Corollary A.3 and thus the error bound is sufficient. See, for example, Corollary A.17.

Proof of Lemma 3.4. Property 3.2(ii) states

$$\left| \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(u+x)} \mathbf{s}}{\alpha e^{\mathbf{S}u} \mathbf{e}} V(v)g(x) dx - g(\Delta - u - v)1(u + v \leq \Delta - \varepsilon) \right| = |r_V(u, v)|. \quad (\text{A.10})$$

Setting $g(x) = h_{ij}^{++}(\lambda, x)$,

$$\left| \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(u+x)}}{\alpha e^{\mathbf{S}u} \mathbf{e}} V(v) h_{ij}^{++}(\lambda, x) dx - h_{ij}^{++}(\lambda, \Delta - u - v) 1(u + v \leq \Delta - \varepsilon) \right| = |r_V(u, v)|. \quad (\text{A.11})$$

We recognise the left-most term as

$$\hat{f}_{0,+,+}^{\ell_0,(p)}(\lambda)(y_{\ell_0+1} - v, j; y_{\ell_0} + u, i) = \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(u+x)}}{\alpha e^{\mathbf{S}u} \mathbf{e}} V(v) h_{ij}^{++}(\lambda, x) dx.$$

Now consider

$$\begin{aligned} & \left| \int_{v=0}^{\Delta} \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(u+x)}}{\alpha e^{\mathbf{S}u} \mathbf{e}} V(v) h_{ij}^{++}(\lambda, x) dx \psi(v) dv \right. \\ & \quad \left. - \int_{v=0}^{\Delta} h_{ij}^{++}(\lambda, \Delta - u - v) 1(\Delta - u - v \geq 0) \psi(v) dv \right| \\ & \leq \int_{v=0}^{\Delta} \left| \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(u+x)}}{\alpha e^{\mathbf{S}u} \mathbf{e}} V(v) h_{ij}^{++}(\lambda, x) dx - h_{ij}^{++}(\lambda, \Delta - u - v) 1(\Delta - u - v \geq 0) \right| |\psi(v)| dv \\ & \leq \int_{v=0}^{\Delta} \left| \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(u+x)}}{\alpha e^{\mathbf{S}u} \mathbf{e}} V(v) h_{ij}^{++}(\lambda, x) dx - h_{ij}^{++}(\lambda, \Delta - u - v) 1(\Delta - u - v \geq \varepsilon) \right| |\psi(v)| dv \\ & \quad + \int_{v=0}^{\Delta} |h_{ij}^{++}(\lambda, \Delta - u - v) 1(\varepsilon \geq \Delta - u - v \geq 0)| |\psi(v)| dv. \end{aligned} \quad (\text{A.12})$$

Applying 3.2(ii) to the first term, then (A.12) is less than or equal to

$$\begin{aligned} & \int_{v=0}^{\Delta} |r_V(u, v)| |\psi(v)| dv + \int_{v=0}^{\Delta} |h_{ij}^{++}(\lambda, \Delta - u - v) 1(\varepsilon \geq \Delta - u - v \geq 0)| |\psi(v)| dv \\ & \leq R_{V,2} GF + \varepsilon GF. \end{aligned}$$

Finally, noting $h_{ij}^{++}(\lambda, \Delta - u - v) 1(\Delta - u - v \geq 0) = \hat{\mu}_{0,+,+}^{\ell_0}(\lambda)(y_{\ell_0+1} - v, j; y_{\ell_0} + u, i)$, then we have shown (3.38).

Using analogous arguments we can show (3.39). \square

Lemma A.4. *Let g satisfy the Assumptions 3.1 and $x_0 \in (2\varepsilon, \Delta - \varepsilon)$. Then*

$$\left| \int_{x=0}^{\infty} \mathbf{a}(x_0) \mathbf{D} e^{\mathbf{S}x} g(x) \mathbf{s} dx - g(x_0) \right| \leq \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2} 4G + 3L\varepsilon + 6G \frac{\text{Var}(Z)}{\varepsilon^2}. \quad (\text{A.13})$$

Proof. First rewrite the left-hand side as

$$\left| \int_{x=0}^{\infty} \mathbf{a}(x_0) \mathbf{D} e^{\mathbf{S}x} (g(x) - g(x_0)) \mathbf{s} \, dx \right| \leq \int_{x=0}^{\infty} \mathbf{a}(x_0) \mathbf{D} e^{\mathbf{S}x} |g(x) - g(x_0)| \mathbf{s} \, dx. \quad (\text{A.14})$$

Substituting the expression for \mathbf{D} gives,

$$\begin{aligned} & \int_{x=0}^{\infty} \mathbf{a}(x_0) \int_{u=0}^{\infty} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} \, du e^{\mathbf{S}x} |g(x) - g(x_0)| \mathbf{s} \, dx \\ &= \int_{x=0}^{\infty} \mathbf{a}(x_0) \int_{u=0}^{\Delta-\varepsilon} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} \, du e^{\mathbf{S}x} |g(x) - g(x_0)| \mathbf{s} \, dx \\ & \quad + \int_{x=0}^{\infty} \mathbf{a}(x_0) \int_{u=\Delta-\varepsilon}^{\infty} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} \, du e^{\mathbf{S}x} |g(x) - g(x_0)| \mathbf{s} \, dx \end{aligned} \quad (\text{A.15})$$

Since g is bounded, the second term is less than or equal to

$$\begin{aligned} \int_{x=0}^{\infty} \mathbf{a}(x_0) \int_{u=\Delta-\varepsilon}^{\infty} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} \, du e^{\mathbf{S}x} \mathbf{s} \, dx 2G &= \mathbf{a}(x_0) \int_{u=\Delta-\varepsilon}^{\infty} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} \, du 2G \\ &= \mathbf{a}(x_0) \int_{u=\Delta-\varepsilon}^{\infty} e^{\mathbf{S}u} \mathbf{s} \, du 2G \\ &= \frac{\mathbb{P}(Z \geq x_0 + \Delta - \varepsilon)}{\mathbb{P}(Z > x_0)} 2G \end{aligned} \quad (\text{A.16})$$

For $x_0 \in (2\varepsilon, \Delta - \varepsilon)$, then (A.16) is less than or equal to

$$\frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2} 2G.$$

As for the first term in (A.15), it can be written as

$$\begin{aligned} & \int_{x=0}^{\infty} \mathbf{a}(x_0) \int_{u=\Delta-x_0-\varepsilon}^{u=\Delta-x_0+\varepsilon} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} \, du e^{\mathbf{S}x} |g(x) - g(x_0)| \mathbf{s} \, dx \\ & \quad + \int_{x=0}^{\infty} \mathbf{a}(x_0) \int_{u=0}^{\Delta-x_0-\varepsilon} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} \, du e^{\mathbf{S}x} |g(x) - g(x_0)| \mathbf{s} \, dx \\ & \quad + \int_{x=0}^{\infty} \mathbf{a}(x_0) \int_{u=\Delta-x_0+\varepsilon}^{\Delta-\varepsilon} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} \, du e^{\mathbf{S}x} |g(x) - g(x_0)| \mathbf{s} \, dx. \end{aligned} \quad (\text{A.17})$$

Since g is bounded, then the last two terms in (A.17) are

$$2G \left(\int_{x=0}^{\infty} \mathbf{a}(x_0) \int_{u=0}^{\Delta-x_0-\varepsilon} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} \, du e^{\mathbf{S}x} \mathbf{s} \, dx \right.$$

$$\begin{aligned}
& + \int_{x=0}^{\infty} \mathbf{a}(x_0) \int_{u=\Delta-x_0+\varepsilon}^{\Delta-\varepsilon} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} du e^{\mathbf{S}x} \mathbf{s} dx \Bigg) \\
& = 2G \left(\mathbf{a}(x_0) \int_{u=0}^{\Delta-x_0-\varepsilon} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} du + \mathbf{a}(x_0) \int_{u=\Delta-x_0+\varepsilon}^{\Delta-\varepsilon} e^{\mathbf{S}u} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} du \right) \\
& = 2G \frac{\mathbb{P}(Z > x_0, Z \notin (\Delta - \varepsilon, \Delta + \varepsilon))}{\mathbb{P}(Z > x_0)} \\
& \leq 2G \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2}, \tag{A.18}
\end{aligned}$$

provided that $x_0 \in [0, \Delta - \varepsilon)$. Exchanging the order of integration for the first term in (A.17)

$$\int_{u=\Delta-x_0-\varepsilon}^{u=\Delta-x_0+\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \int_{x=0}^{\infty} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} e^{\mathbf{S}x} |g(x) - g(x_0)| \mathbf{s} dx du \tag{A.19}$$

from which we see that we can apply Corollary A.2 to the integral over x , implying that (A.19) is less than or equal to

$$\int_{u=\Delta-x_0-\varepsilon}^{u=\Delta-x_0+\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \left(|g(\Delta - u) - g(x_0)| + 6G \frac{\text{Var}(Z)}{\varepsilon^2} + 2L\varepsilon \right) du. \tag{A.20}$$

Since g is Lipschitz, then (A.20) is less than or equal to

$$\int_{u=\Delta-x_0-\varepsilon}^{u=\Delta-x_0+\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \left(L\varepsilon + 6G \frac{\text{Var}(Z)}{\varepsilon^2} + 2L\varepsilon \right) du \leq L\varepsilon + 6G \frac{\text{Var}(Z)}{\varepsilon^2} + 2L\varepsilon. \tag{A.21}$$

Putting all the bounds together proves the result. \square

Lemma A.5. Let $\psi : \mathcal{D}_{\ell_0} \rightarrow \mathbb{R}$ be bounded $|\psi(x)| \leq F$ and Lipschitz continuous function and let $\lambda > 0$. For $i \in \mathcal{S}_-, j \in \mathcal{S}_- \cup \mathcal{S}_{-0}$, $x_0 \in (0, \Delta)$,

$$\int_{x_1=0}^{\infty} \int_{x=0}^{\Delta} \mathbf{a}^{(p)}(x_0) \mathbf{D}^{(p)} e^{\mathbf{S}x_1} V^{(p)}(x) h_{ij}^{--}(\lambda, x_1) \psi(x) dx dx_1 \rightarrow \int_{x=0}^{x_0} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) dx, \tag{A.22}$$

as $p \rightarrow \infty$. Similarly, for $i \in \mathcal{S}_+, j \in \mathcal{S}_+ \cup \mathcal{S}_{+0}$

$$\begin{aligned}
& \int_{x_1=0}^{\infty} \int_{x=0}^{\Delta} \mathbf{a}^{(p)}(x_0) \mathbf{D}^{(p)} e^{\mathbf{S}^{(p)}x_1} V^{(p)}(x) h_{ij}^{++}(\lambda, x_1) \psi(\Delta - x) dx dx_1 \\
& \rightarrow \int_{x=\Delta-x_0}^{\Delta} h_{ij}^{++}(\lambda, x - x_0) \psi(x) dx, \tag{A.23}
\end{aligned}$$

Proof. Assume, without loss of generality $\ell_0 = 0$ so $\mathcal{D}_{\ell_0} = [0, \Delta]$. Substituting the definition of \mathbf{D} , then the left-hand side of (A.22) is

$$\begin{aligned}
& \left| \int_{x_1=0}^{\infty} \int_{x=0}^{\Delta} \int_{u=0}^{\infty} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \mathbf{a}(u) \, du e^{\mathbf{S}x_1} V(x) h_{ij}^{--}(\lambda, x_1) \psi(x) \, dx \, dx_1 \right. \\
& \quad \left. - \int_{x=0}^{x_0} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \right| \\
& \leq \left| \int_{x_1=0}^{\infty} \int_{x=0}^{\Delta} \int_{u=0}^{\Delta-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \mathbf{a}(u) \, du e^{\mathbf{S}x_1} V(x) h_{ij}^{--}(\lambda, x_1) \psi(x) \, dx \, dx_1 \right. \\
& \quad \left. - \int_{x=0}^{x_0} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \right| \\
& \quad + \left| \int_{x_1=0}^{\infty} \int_{x=0}^{\Delta} \int_{u=\Delta-\varepsilon}^{\infty} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \mathbf{a}(u) \, du e^{\mathbf{S}x_1} V(x) h_{ij}^{--}(\lambda, x_1) \psi(x) \, dx \, dx_1 \right| \quad (\text{A.24})
\end{aligned}$$

Since $|\phi(x)| \leq F$ and $|h_{ij}^{--}(\lambda, x)| \leq G$, the third term in (A.24) is less than or equal to

$$\int_{x_1=0}^{\infty} \int_{x=0}^{\Delta} \int_{u=\Delta-\varepsilon}^{\infty} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \mathbf{a}(u) \, du e^{\mathbf{S}x_1} V(x) \, dx \, dx_1 GF. \quad (\text{A.25})$$

Computing the integral over x_1 in (A.25) gives

$$\begin{aligned}
& \int_{x=0}^{\Delta} \int_{u=\Delta-\varepsilon}^{\infty} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \mathbf{a}(u) \, du (-\mathbf{S})^{-1} V(x) \, dx GF \\
& \leq \int_{x=0}^{\Delta} \int_{u=\Delta-\varepsilon}^{\infty} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \, du \, dx G_V GF \\
& = \int_{u=\Delta-\varepsilon}^{\infty} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \, du \Delta G_V GF. \quad (\text{A.26})
\end{aligned}$$

since, by property 3.2(i), $\mathbf{a}(u)(-\mathbf{S})^{-1} V(x) \leq \mathbf{a}(u) \mathbf{e} G_V = G_V$. The bound (A.26) is equal to

$$\Delta \mathbb{P}(Z > x_0 + \Delta - \varepsilon) \, dx G_V GF \leq \Delta \frac{\text{Var}(Z)}{(x_0 - \varepsilon)^2} G_V GF, \quad (\text{A.27})$$

by Chebyshev's inequality.

The first and second terms in (A.24) are

$$\left| \int_{x=0}^{\Delta} \int_{u=0}^{\Delta-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \int_{x_1=0}^{\infty} \mathbf{a}(u) e^{\mathbf{S}x_1} V(x) h_{ij}^{--}(\lambda, x_1) \, dx_1 \, du \psi(x) \, dx \right|$$

$$\begin{aligned}
& \left| - \int_{x=0}^{x_0} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \right| \\
&= \left| \int_{x=0}^{\Delta} \int_{u=0}^{\Delta-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} [h_{ij}^{--}(\lambda, \Delta - u - x) 1(u + x \leq \Delta - \varepsilon) + r_V(u, x)] \psi(x) \, du \, dx \right. \\
&\quad \left. - \int_{x=0}^{x_0} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \right| \\
&\leq \left| \int_{x=0}^{\Delta} \int_{u=0}^{\Delta-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} h_{ij}^{--}(\lambda, \Delta - u - x) 1(u + x \leq \Delta - \varepsilon) \psi(x) \, du \, dx \right. \\
&\quad \left. - \int_{x=0}^{x_0} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \right| + \left| \int_{x=0}^{\Delta} \int_{u=0}^{\Delta-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} r_V(u, x) \psi(x) \, du \, dx \right| \quad (\text{A.28})
\end{aligned}$$

where the first equality holds from Property 3.2(ii). The last term in (A.28) is less than or equal to

$$\begin{aligned}
\int_{u=0}^{\Delta-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \int_{x=0}^{\Delta} \mathbf{s} |r_V(u, x)| \, dx \, du F &\leq \int_{u=0}^{\Delta-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} R_{V,2} \, du F \\
&\leq R_{V,2} F, \quad (\text{A.29})
\end{aligned}$$

by Property 3.2(ii). The first two terms in (A.28) are

$$\begin{aligned}
& \left| \int_{x=0}^{\Delta} \int_{u=0}^{\Delta-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} h_{ij}^{--}(\lambda, \Delta - u - x) 1(u + x \leq \Delta - \varepsilon) \psi(x) \, du \, dx \right. \\
&\quad \left. - \int_{x=0}^{x_0} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \right| \\
&= \left| \int_{x=0}^{\Delta-\varepsilon} \int_{u=0}^{\Delta-x-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} h_{ij}^{--}(\lambda, \Delta - u - x) \psi(x) \, du \, dx \right. \\
&\quad \left. - \int_{x=0}^{x_0} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \right| \\
&\leq \left| \int_{x=0}^{\Delta-\varepsilon} \int_{u=\Delta-x_0-\varepsilon}^{\Delta-x_0+\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} h_{ij}^{--}(\lambda, \Delta - u - x) \psi(x) \, du 1(\Delta - x_0 + \varepsilon \leq \Delta - x - \varepsilon) \, dx \right. \\
&\quad \left. - \int_{x=0}^{x_0} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \right| \\
&\quad + \left| \int_{x=0}^{\Delta-\varepsilon} \int_{u=0}^{\min(\Delta-x_0-\varepsilon, \Delta-x-\varepsilon)} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} h_{ij}^{--}(\lambda, \Delta - u - x) \psi(x) \, du \, dx \right|
\end{aligned}$$

$$\begin{aligned}
& + \left| \int_{x=0}^{\Delta-\varepsilon} \int_{u=\Delta-x_0+\varepsilon}^{\Delta-x-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} h_{ij}^{--}(\lambda, \Delta - u - x) \psi(x) \, du \right. \\
& \quad \left. \times 1(\Delta - x_0 + \varepsilon \leq \Delta - x - \varepsilon) \, dx \right| \\
& + \left| \int_{x=0}^{\Delta-\varepsilon} \int_{u=\Delta-x_0-\varepsilon}^{\Delta-x-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} h_{ij}^{--}(\lambda, \Delta - u - x) \psi(x) \, du \right. \\
& \quad \left. \times 1(\Delta - x_0 - \varepsilon \leq \Delta - x - \varepsilon < \Delta - x_0 + \varepsilon) \, dx \right| \tag{A.30}
\end{aligned}$$

Since $|\psi| \leq F$, the second and third terms in (A.30) are less than or equal to

$$\begin{aligned}
& \int_{x=0}^{\Delta-\varepsilon} \int_{u=0}^{\min(\Delta-x_0-\varepsilon, \Delta-x-\varepsilon)} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \, du \, dx GF \\
& \quad + \int_{x=0}^{\Delta-\varepsilon} \int_{u=\Delta-x_0+\varepsilon}^{\Delta-x-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \, du 1(\Delta - x_0 + \varepsilon \leq \Delta - x - \varepsilon) \, dx GF \\
& \leq \int_{x=0}^{\Delta-\varepsilon} \int_{u=0}^{\Delta-x_0-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \, du \, dx GF \\
& \quad + \int_{x=0}^{\Delta-\varepsilon} \int_{u=\Delta-x_0+\varepsilon}^{\infty} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \, du 1(\Delta - x_0 + \varepsilon \leq \Delta - x - \varepsilon) \, dx GF \\
& \leq \Delta \int_{u=0}^{\Delta-x_0-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \, du \, dx GF + \Delta \int_{u=\Delta-x_0+\varepsilon}^{\infty} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \, du GF \\
& \leq \Delta \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2} GF. \tag{A.31}
\end{aligned}$$

Since $\int_{u=\Delta-x_0-\varepsilon}^{\Delta-x-\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \leq 1$, the last term in (A.30) is less than or equal to

$$\int_{x=0}^{\Delta-\varepsilon} GF \, du 1(\Delta - x_0 - \varepsilon \leq \Delta - x - \varepsilon < \Delta - x_0 + \varepsilon) \, dx = 2\varepsilon GF. \tag{A.32}$$

Now consider the difference in the first absolute value in (A.30),

$$\begin{aligned}
& \left| \int_{x=0}^{\Delta-\varepsilon} \int_{u=\Delta-x_0-\varepsilon}^{\Delta-x_0+\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} h_{ij}^{--}(\lambda, \Delta - u - x) \psi(x) \, du 1(x \leq x_0 - 2\varepsilon) \, dx \right. \\
& \quad \left. - \int_{x=0}^{x_0} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \right|
\end{aligned}$$

$$\begin{aligned}
&\leq \left| \int_{x=0}^{x_0-2\varepsilon} \int_{u=\Delta-x_0-\varepsilon}^{\Delta-x_0+\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} h_{ij}^{--}(\lambda, \Delta - u - x) \psi(x) \, du \, dx \right. \\
&\quad \left. - \int_{x=0}^{x_0-2\varepsilon} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \right| + \left| \int_{x=x_0-2\varepsilon}^{x_0} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \right| \\
&\leq \int_{x=0}^{x_0-2\varepsilon} \int_{u=\Delta-x_0-\varepsilon}^{\Delta-x_0+\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} \left| h_{ij}^{--}(\lambda, \Delta - u - x) - h_{ij}^{--}(\lambda, x_0 - x) \right| \, du \, |\psi(x)| \, dx \\
&\quad + \left| \int_{x=0}^{x_0-2\varepsilon} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \mathbb{P}(|Z - \Delta| > \varepsilon) \right| \\
&\quad + \left| \int_{x=x_0-2\varepsilon}^{x_0} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \right| \\
&\leq \int_{x=0}^{x_0-2\varepsilon} \int_{u=\Delta-x_0-\varepsilon}^{\Delta-x_0+\varepsilon} \mathbf{a}(x_0) e^{\mathbf{S}u} \mathbf{s} L \varepsilon \, du \, |\psi(x)| \, dx + \Delta G F \frac{\text{Var}(Z)}{\varepsilon^2} + 2\varepsilon G F \\
&\leq \Delta L \varepsilon F + \Delta G F \frac{\text{Var}(Z)}{\varepsilon^2} + 2\varepsilon G F. \tag{A.33}
\end{aligned}$$

In summary, we have shown

$$\begin{aligned}
&\left| \int_{x_1=0}^{\infty} \int_{x=0}^{\Delta} \mathbf{a}^{(p)}(x_0) \mathbf{D}^{(p)} e^{\mathbf{S}^{(p)} x_1} V^{(p)}(x) h_{ij}^{--}(\lambda, x_1) \psi(x) \, dx \, dx_1 \right. \\
&\quad \left. - \int_{x=0}^{x_0} h_{ij}^{--}(\lambda, x_0 - x) \psi(x) \, dx \right| \\
&\leq \Delta L \varepsilon^{(p)} F + \Delta G F \frac{\text{Var}(Z^{(p)})}{(\varepsilon^{(p)})^2} + 4\varepsilon^{(p)} G F + \Delta \frac{\text{Var}(Z^{(p)}) / (\varepsilon^{(p)})^2}{1 - \text{Var}(Z^{(p)}) / (\varepsilon^{(p)})^2} G F + R_{V,2}^{(p)} F \\
&\quad + \Delta \frac{\text{Var}(Z^{(p)})}{(x_0 - \varepsilon^{(p)})^2} G_V G F. \tag{A.34}
\end{aligned}$$

The result follows upon choosing $\varepsilon^{(p)} = \text{Var}(Z^{(p)})^{1/3}$ and letting $p \rightarrow \infty$. \square

A.1.2 Many integrals.

Define the column vectors

$$\mathcal{I}_{m,k}(u_k) = \left[\prod_{\ell=m}^{k-1} \int_{x_\ell=0}^{\infty} g_\ell(x_\ell) e^{\mathbf{S}x_\ell} \, dx_\ell \mathbf{D} \right] \int_{x_k=0}^{\infty} g_k(x_k) e^{\mathbf{S}x_k} \, dx_k e^{\mathbf{S}u_k} \mathbf{s} \tag{A.35}$$

for $m, k \in \{1, 2, \dots\}$, $m \leq k$, where a product over an empty set is equal to 1. Also define the row vectors

$$\mathcal{J}_{k+1,k+1}(u_k, x_{k+1}) := g_{k+1}(x_{k+1}) \frac{\alpha e^{S u_k}}{\alpha e^{S u_k} \mathbf{e}} e^{S x_{k+1}} \quad (\text{A.36})$$

and

$$\begin{aligned} \mathcal{J}_{k+1,n}(u_k, x_{k+1}) &:= g_{k+1}(x_{k+1}) \frac{\alpha e^{S u_k}}{\alpha e^{S u_k} \mathbf{e}} e^{S x_{k+1}} \mathbf{D} \left[\prod_{m=k+2}^{n-1} \int_{x_m=0}^{\infty} g_m(x_m) e^{S x_m} dx_m \mathbf{D} \right] \\ &\quad \times \int_{x_n=0}^{\infty} g_n(x_n) e^{S x_n} dx_n \end{aligned} \quad (\text{A.37})$$

for $k, n \in \{0, 1, 2, \dots\}$, $k+1 < n$. Also define the row vector function $\mathbf{a}(x) : [0, \infty) \rightarrow \mathcal{A} \subset \mathbb{R}^p$,

$$\mathbf{a}(x) = \frac{\alpha e^{S x}}{\alpha e^{S x} \mathbf{e}}.$$

Lemma A.6. *Let g_1, g_2, \dots , be functions satisfying Assumptions 3.1, then, for $k \in \{1, 2, \dots\}$,*

$$\mathbf{a}(x_0) \mathcal{I}_{1,k}(u_k) \leq \frac{1}{\alpha e^{S x_0} \mathbf{e}} G \hat{G}^{k-1} \alpha e^{S u_k} \mathbf{e}. \quad (\text{A.38})$$

Proof. Recall the definition of $\mathbf{D} := \int_{u=0}^{\infty} e^{S u} \mathbf{s} \frac{\alpha e^{S u}}{\alpha e^{S u} \mathbf{e}} du$ and substitute it into the left-hand side of (A.38),

$$\begin{aligned} \mathbf{a}(x_0) \mathcal{I}_{1,k}(u_k) &= \mathbf{a}(x_0) \int_{x_1=0}^{\infty} g_1(x_1) e^{S x_1} \mathbf{D} \mathcal{I}_{2,k}(u_k) \\ &= \mathbf{a}(x_0) \int_{x_1=0}^{\infty} g_1(x_1) e^{S x_1} \int_{u_1=0}^{\infty} e^{S u_1} \mathbf{s} \frac{\alpha e^{S u_1}}{\alpha e^{S u_1} \mathbf{e}} du_1 \mathcal{I}_{2,k}(u_k). \end{aligned}$$

Now, since $|g_1| \leq G$, then this is less than or equal to

$$\mathbf{a}(x_0) \int_{x_1=0}^{\infty} G e^{S x_1} \int_{u_1=0}^{\infty} e^{S u_1} \mathbf{s} \frac{\alpha e^{S u_1}}{\alpha e^{S u_1} \mathbf{e}} du_1 \mathcal{I}_{2,k}(u_k). \quad (\text{A.39})$$

Computing the integral with respect to x_1 in (A.39) gives

$$\begin{aligned} &G \mathbf{a}(x_0) (-\mathbf{S})^{-1} \int_{u_1=0}^{\infty} e^{S u_1} \mathbf{s} \frac{\alpha e^{S u_1}}{\alpha e^{S u_1} \mathbf{e}} du_1 \mathcal{I}_{2,k}(u_k) \\ &= G \mathbf{a}(x_0) \int_{u_1=0}^{\infty} e^{S u_1} \mathbf{e} \frac{\alpha e^{S u_1}}{\alpha e^{S u_1} \mathbf{e}} du_1 \mathcal{I}_{2,k}(u_k) \end{aligned}$$

$$= \frac{G}{\alpha e^{\mathbf{S}x_0} \mathbf{e}} \int_{u_1=0}^{\infty} \alpha e^{\mathbf{S}(x_0+u_1)} \mathbf{e} \frac{\alpha e^{\mathbf{S}u_1}}{\alpha e^{\mathbf{S}u_1} \mathbf{e}} du_1 \mathcal{I}_{2,k}(u_k), \quad (\text{A.40})$$

since $(-\mathbf{S})^{-1}$ and $e^{\mathbf{S}t}$ commute, $\mathbf{s} = -\mathbf{S}\mathbf{e}$ and $e^{\mathbf{S}(t+u)} = e^{\mathbf{S}t} e^{\mathbf{S}u}$.

Now, as $\alpha e^{\mathbf{S}(x_0+u_1)} \mathbf{e} \leq \alpha e^{\mathbf{S}u_1} \mathbf{e}$, then (A.40) is less than or equal to

$$G \frac{1}{\alpha e^{\mathbf{S}x_0} \mathbf{e}} \int_{u_1=0}^{\infty} \alpha e^{\mathbf{S}u_1} \mathbf{e} \frac{\alpha e^{\mathbf{S}u_1}}{\alpha e^{\mathbf{S}u_1} \mathbf{e}} du_1 \mathcal{I}_{2,k}(u_k) = G \frac{1}{\alpha e^{\mathbf{S}x_0} \mathbf{e}} \int_{u_1=0}^{\infty} \alpha e^{\mathbf{S}u_1} du_1 \mathcal{I}_{2,k}(u_k).$$

Now integrate with respect to u_1 and use the facts that $(-\mathbf{S})^{-1}$ and $e^{\mathbf{S}x}$ commute, and $\mathbf{s} = -\mathbf{S}\mathbf{e}$, to get

$$\begin{aligned} & G \frac{1}{\alpha e^{\mathbf{S}x_0} \mathbf{e}} \alpha (-\mathbf{S})^{-1} \mathcal{I}_{2,k}(u_k) \\ &= G \frac{1}{\alpha e^{\mathbf{S}x_0} \mathbf{e}} \alpha (-\mathbf{S})^{-1} \int_{x_2=0}^{\infty} g_2(x_2) e^{\mathbf{S}x_2} dx_2 \int_{u_2=0}^{\infty} e^{\mathbf{S}u_2} \mathbf{s} \frac{\alpha e^{\mathbf{S}u_2}}{\alpha e^{\mathbf{S}u_2} \mathbf{e}} du_2 \mathcal{I}_{3,k}(u_k) \end{aligned} \quad (\text{A.41})$$

$$= G \frac{1}{\alpha e^{\mathbf{S}x_0} \mathbf{e}} \int_{x_2=0}^{\infty} g_2(x_2) \alpha e^{\mathbf{S}x_2} dx_2 \int_{u_2=0}^{\infty} e^{\mathbf{S}u_2} \mathbf{e} \frac{\alpha e^{\mathbf{S}u_2}}{\alpha e^{\mathbf{S}u_2} \mathbf{e}} du_2 \mathcal{I}_{3,k}(u_k) \quad (\text{A.42})$$

Since $\alpha e^{\mathbf{S}x_2} e^{\mathbf{S}u_2} \mathbf{e} \leq \alpha e^{\mathbf{S}u_2} \mathbf{e}$, then (A.42) is less than or equal to

$$\begin{aligned} & G \frac{1}{\alpha e^{\mathbf{S}x_0} \mathbf{e}} \int_{x_2=0}^{\infty} g_2(x_2) dx_2 \int_{u_2=0}^{\infty} \alpha e^{\mathbf{S}u_2} \mathbf{e} \frac{\alpha e^{\mathbf{S}u_2}}{\alpha e^{\mathbf{S}u_2} \mathbf{e}} du_2 \mathcal{I}_{3,k}(u_k) \\ &= G \frac{1}{\alpha e^{\mathbf{S}x_0} \mathbf{e}} \widehat{G} \int_{u_2=0}^{\infty} \alpha e^{\mathbf{S}u_2} \mathbf{e} \frac{\alpha e^{\mathbf{S}u_2}}{\alpha e^{\mathbf{S}u_2} \mathbf{e}} du_2 \mathcal{I}_{3,k}(u_k) \\ &= G \frac{1}{\alpha e^{\mathbf{S}x_0} \mathbf{e}} \widehat{G} \int_{u_2=0}^{\infty} \alpha e^{\mathbf{S}u_2} \mathcal{I}_{3,k}(u_k) \\ &= G \frac{1}{\alpha e^{\mathbf{S}x_0} \mathbf{e}} \widehat{G} \alpha (-\mathbf{S})^{-1} \mathcal{I}_{3,k}(u_k). \end{aligned} \quad (\text{A.43})$$

Repeating the arguments which got us from (A.41) to (A.43) another $k-2$ times gives the result. \square

Lemma A.7. *Let g_1, g_2, \dots , be functions satisfying the Assumptions 3.1 and let $V(x)$ be a closing operator with the Properties 3.2, then, for $k, n \in \{1, 2, \dots\}$, $k+1 < n$,*

$$\mathcal{J}_{k+1,n}(u_k, x_{k+1})V(x) \leq g_{k+1}(x_{k+1})\widehat{G}^{n-k-2}GG_V.$$

Proof. Starting with the left-hand side, upon substituting \mathbf{D} ,

$$\mathcal{J}_{k+1,n}(u_k, x_{k+1})V(x)$$

$$\begin{aligned}
&= \mathcal{J}_{k+1,n-1}(u_k, x_{k+1}) \mathbf{D} \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} dx_n V(x) \\
&= \mathcal{J}_{k+1,n-1}(u_k, x_{k+1}) \int_{u_{n-1}=0}^{\infty} e^{\mathbf{S}u_{n-1}} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u_{n-1}}}{\boldsymbol{\alpha} e^{\mathbf{S}u_{n-1}} \mathbf{e}} du_{n-1} \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} dx_n V(x) \\
&\leq \mathcal{J}_{k+1,n-1}(u_k, x_{k+1}) \int_{u_{n-1}=0}^{\infty} e^{\mathbf{S}u_{n-1}} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u_{n-1}}}{\boldsymbol{\alpha} e^{\mathbf{S}u_{n-1}} \mathbf{e}} du_{n-1} \int_{x_n=0}^{\infty} G e^{\mathbf{S}x_n} dx_n V(x). \quad (\text{A.44})
\end{aligned}$$

By the Property 3.2(i) of $V(x)$, $\int_{x_n=0}^{\infty} \boldsymbol{\alpha} e^{\mathbf{S}u_{n-1}} e^{\mathbf{S}x_n} V(x) dx_n \leq \boldsymbol{\alpha} e^{\mathbf{S}u_{n-1}} \mathbf{e} G_V$. Therefore (A.44) is less than or equal to

$$\begin{aligned}
&\mathcal{J}_{k+1,n-1}(u_k, x_{k+1}) \int_{u_{n-1}=0}^{\infty} e^{\mathbf{S}u_{n-1}} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u_{n-1}} \mathbf{e}}{\boldsymbol{\alpha} e^{\mathbf{S}u_{n-1}} \mathbf{e}} du_{n-1} G G_V \\
&= \mathcal{J}_{k+1,n-1}(u_k, x_{k+1}) \int_{u_{n-1}=0}^{\infty} e^{\mathbf{S}u_{n-1}} \mathbf{s} du_{n-1} G G_V \\
&= \mathcal{J}_{k+1,n-1}(u_k, x_{k+1}) \mathbf{e} G G_V \quad (\text{A.45})
\end{aligned}$$

$$\begin{aligned}
&= \mathcal{J}_{k+1,n-2}(u_k, x_{k+1}) \int_{u_{n-2}=0}^{\infty} e^{\mathbf{S}u_{n-2}} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u_{n-2}}}{\boldsymbol{\alpha} e^{\mathbf{S}u_{n-2}} \mathbf{e}} du_{n-2} \int_{x_{n-1}=0}^{\infty} g_{n-1}(x_{n-1}) e^{\mathbf{S}x_{n-1}} dx_{n-1} \mathbf{e} \\
&\quad \times G G_V. \quad (\text{A.46})
\end{aligned}$$

Now, since $\boldsymbol{\alpha} e^{\mathbf{S}(x_{n-1}+u_{n-2})} \mathbf{e} \leq \boldsymbol{\alpha} e^{\mathbf{S}(u_{n-2})} \mathbf{e}$, then (A.46) is less than or equal to

$$\begin{aligned}
&\mathcal{J}_{k+1,n-2}(u_k, x_{k+1}) \int_{u_{n-2}=0}^{\infty} e^{\mathbf{S}u_{n-2}} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}u_{n-2}} \mathbf{e}}{\boldsymbol{\alpha} e^{\mathbf{S}u_{n-2}} \mathbf{e}} du_{n-2} \int_{x_{n-1}=0}^{\infty} g_{n-1}(x_{n-1}) dx_{n-1} G G_V \\
&= \mathcal{J}_{k+1,n-2}(u_k, x_{k+1}) \int_{u_{n-2}=0}^{\infty} e^{\mathbf{S}u_{n-2}} \mathbf{s} du_{n-2} \hat{G} G G_V \\
&= \mathcal{J}_{k+1,n-2}(u_k, x_{k+1}) \mathbf{e} \hat{G} G G_V. \quad (\text{A.47})
\end{aligned}$$

This is of the same form as (A.45), hence repeating the same arguments which got us from (A.45) to (A.47) another $n - k - 3$ more times gives

$$\begin{aligned}
\mathcal{J}_{k+1,k+1}(u_k, x_{k+1}) \mathbf{e} \hat{G}^{n-k-2} G G_V &\leq g_{k+1}(x_{k+1}) \frac{\boldsymbol{\alpha} e^{\mathbf{S}(u_k+x_{k+1})}}{\boldsymbol{\alpha} e^{\mathbf{S}u_k} \mathbf{e}} \mathbf{e} \hat{G}^{n-k-2} G G_V \\
&\leq g_{k+1}(x_{k+1}) \hat{G}^{n-k-2} G G_V.
\end{aligned}$$

□

Corollary A.8. Let g_1, g_2, \dots , be functions satisfying the Assumptions 3.1 and let $V(x)$ be a closing operator with the Properties 3.2, then,

$$\int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}(x_0) e^{\mathbf{S}x_1} dx_1 \mathbf{D} \left[\prod_{k=2}^{n-1} \int_{x_k=0}^{\infty} g_k(x_k) e^{\mathbf{S}x_k} dx_k \mathbf{D} \right] \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} dx_n V(x)$$

$$\leq \widehat{G}^{n-1} G G_V \quad (\text{A.48})$$

Proof. The left-hand side of (A.48) can be seen to be equivalent to $\mathcal{J}_{1,n+1}(x_0, x_1)$, with $g_1(x_1) = 1$, and the integrability condition on g_1 is not required to prove the bound. \square

Corollary A.9. *Let g_1, g_2, \dots , be functions satisfying the Assumptions 3.1 and let $V(x)$ be a closing operator with the Properties 3.2, then, for $k, n \in \{1, 2, \dots\}$, $k+1 \leq n$,*

$$\begin{aligned} & \int_{x_{k+1}=0}^{\infty} \int_{u_k=\Delta-\varepsilon}^{\infty} \mathbf{a}(x_0) \mathcal{I}_{1,k}(u_k) \mathcal{J}_{k+1,n}(u_k, x_{k+1}) V(x) \\ & \leq \left(2\varepsilon + \frac{\text{Var}(Z)}{\varepsilon} \right) \frac{1}{\boldsymbol{\alpha} e^{\mathbf{S}x_0} \mathbf{e}} G \widehat{G}^{n-2} G G_V =: |r_4(n)|. \end{aligned} \quad (\text{A.49})$$

Proof. Consider first $k+1 < n$. Combining Lemmas A.6 and A.7 the left-hand side of (A.49) is less than or equal to

$$\frac{1}{\boldsymbol{\alpha} e^{\mathbf{S}x_0} \mathbf{e}} G \widehat{G}^{k-1} \int_{x_{k+1}=0}^{\infty} \int_{u_k=\Delta-\varepsilon}^{\infty} \boldsymbol{\alpha} e^{\mathbf{S}u_k} \mathbf{e} g_{k+1}(x_{k+1}) du_k dx_{k+1} \widehat{G}^{n-k-2} G G_V \quad (\text{A.50})$$

$$= \frac{1}{\boldsymbol{\alpha} e^{\mathbf{S}x_0} \mathbf{e}} G \widehat{G}^{k-1} \int_{u_k=\Delta-\varepsilon}^{\infty} \boldsymbol{\alpha} e^{\mathbf{S}u_k} \mathbf{e} du_k \widehat{G}_{k+1} \widehat{G}^{n-k-2} G G_V. \quad (\text{A.51})$$

Now

$$\begin{aligned} \int_{u_k=\Delta-\varepsilon}^{\infty} \boldsymbol{\alpha} e^{\mathbf{S}u_k} \mathbf{e} du_k &= \int_{u_k=\Delta-\varepsilon}^{\Delta+\varepsilon} \mathbb{P}(Z > u_k) du_k + \int_{u_k=\Delta+\varepsilon}^{\infty} \mathbb{P}(Z > u_k) du_k \\ &\leq \int_{u_k=\Delta-\varepsilon}^{\Delta+\varepsilon} du_k + \int_{u_k=\Delta+\varepsilon}^{\infty} \frac{\text{Var}(Z)}{(u_k - \Delta)^2} du_k \\ &= 2\varepsilon + \frac{\text{Var}(Z)}{\varepsilon}, \end{aligned} \quad (\text{A.52})$$

where we have used Chebyshev's inequality to bound the tail probability,

$$\mathbb{P}(Z > u_k) \leq \mathbb{P}(|Z - \Delta| > |u_k - \Delta|) \leq \frac{\text{Var}(Z)}{(u_k - \Delta)^2},$$

for $u_k \geq \Delta + \varepsilon$. Hence (A.51) is less than or equal to

$$\frac{1}{\boldsymbol{\alpha} e^{\mathbf{S}x_0} \mathbf{e}} G \widehat{G}^{k-1} \left(2\varepsilon + \frac{\text{Var}(Z)}{\varepsilon} \right) \widehat{G}_{k+1} \widehat{G}^{n-k-2} G G_V.$$

Now consider $k+1 = n$. By Lemma A.6 we have

$$\frac{1}{\boldsymbol{\alpha} e^{\mathbf{S}x_0} \mathbf{e}} G \widehat{G}^{k-1} \int_{x_{k+1}=0}^{\infty} \int_{u_k=\Delta-\varepsilon}^{\infty} \boldsymbol{\alpha} e^{\mathbf{S}u_k} \mathbf{e} g_{k+1}(x_{k+1}) \frac{\boldsymbol{\alpha} e^{\mathbf{S}(u_k+x_{k+1})}}{\boldsymbol{\alpha} e^{\mathbf{S}u_k} \mathbf{e}} V(x) du_k dx_{k+1}. \quad (\text{A.53})$$

Since $g_{k+1} \leq G$, and upon integrating over x_{k+1} , then (A.53) is less than or equal to

$$\begin{aligned} & \frac{1}{\alpha e^{S_{x_0}} \mathbf{e}} G \widehat{G}^{k-1} \int_{u_k=\Delta-\varepsilon}^{\infty} G \alpha e^{S_{u_k}} (-\mathbf{S})^{-1} V(x) du_k \\ & \leq \frac{1}{\alpha e^{S_{x_0}} \mathbf{e}} G \widehat{G}^{k-1} \int_{u_k=\Delta-\varepsilon}^{\infty} G \alpha e^{S_{u_k}} \mathbf{e} G_V du_k, \end{aligned} \quad (\text{A.54})$$

where we have used Property 3.2(i) to get the upper bound on the right-hand side of (A.54). Using (A.52) again, then (A.54) is less than or equal to

$$\frac{1}{\alpha e^{S_{x_0}} \mathbf{e}} G \widehat{G}^{n-2} G G_V \left(2\varepsilon + \frac{\text{Var}(Z)}{\varepsilon} \right). \quad (\text{A.55})$$

This completes the proof. \square

The error term $r_4(n)$ depends on p and we write $r_4^{(p)}(n)$ when we need to make this dependence explicit, otherwise it is omitted from the notation. Upon choosing $\varepsilon = \text{Var}(Z^{(p)})^{1/3}$, then for fixed $n < \infty$ the error term $|r_4^{(p)}(n)| = O\left(\text{Var}(Z^{(p)})^{1/3}\right) \rightarrow 0$ as $p \rightarrow \infty$.

Define $\mathbf{D}(b) = \int_{u=0}^b e^{S_u} \mathbf{s} \frac{\alpha e^{S_u}}{\alpha e^{S_u} \mathbf{e}} du$. Also define

$$\begin{aligned} g_{2,n}^*(u_1, x) &= \int_{u_2=0}^{\Delta-u_1} g_2(\Delta - u_2 - u_1) du_2 \dots \int_{u_{n-1}=0}^{\Delta-u_{n-2}} g_{n-1}(\Delta - u_{n-1} - u_{n-2}) du_{n-2} \\ &\quad g_n(\Delta - x - u_{n-1}) 1(\Delta - x - u_{n-1} \geq 0) du_{n-1}, \end{aligned} \quad (\text{A.56})$$

and

$$g_{1,n}^*(x_0, x) = \int_{u_1=0}^{\Delta-x_0} g_1(\Delta - u_1 - x_0) g_{2,n}^*(u_1, x) du_1 \quad (\text{A.57})$$

and

$$\begin{aligned} g_{1,n}^{*,\varepsilon}(x_0, x) &= \int_{u_1=0}^{\Delta-\varepsilon-x_0} g_1(\Delta - u_1 - x_0) \int_{u_2=0}^{\Delta-\varepsilon-u_1} g_2(\Delta - u_2 - u_1) du_2 \\ &\quad \dots \int_{u_{n-1}=0}^{\Delta-\varepsilon-u_{n-2}} g_{n-1}(\Delta - u_{n-1} - u_{n-2}) du_{n-2} g_n(\Delta - x - u_{n-1}) \\ &\quad \times 1(\Delta - x - u_{n-1} \geq \varepsilon) \end{aligned} \quad (\text{A.58})$$

For later, observe that

$$g_{2,n}^*(u_1, x) = \int_{u_2=0}^{\Delta-u_1} g_2(\Delta - u_2 - u_1) du_2 \dots \int_{u_{n-1}=0}^{\Delta-u_{n-2}} g_{n-1}(\Delta - u_{n-1} - u_{n-2}) du_{n-2}$$

$$\begin{aligned}
& g_n(\Delta - x - u_{n-1})1(\Delta - x - u_{n-1} \geq 0) du_{n-1} \\
& \leq G^{n-1} \int_{u_2=0}^{\Delta-u_1} du_1 \dots \int_{u_{n-1}=0}^{\Delta-u_{n-2}} du_{n-1} \\
& \leq G^{n-1} \Delta^{n-2} := G_n^*.
\end{aligned} \tag{A.59}$$

Lemma A.10. Let g_1, g_2, \dots , be functions satisfying the Assumptions 3.1 and let $V(x)$ be a closing operator with the Properties 3.2. Then, for $n \geq 2$,

$$\begin{aligned}
& \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}(x_0) e^{\mathbf{S}x_1} dx_1 \mathbf{D}(\Delta - \varepsilon) \left[\prod_{k=2}^{n-1} \int_{x_k=0}^{\infty} g_k(x_k) e^{\mathbf{S}x_k} dx_k \right] \mathbf{D}(\Delta - \varepsilon) \\
& \quad \times \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} dx_n V(x) \\
& = g_{1,n}^*(x_0, x) + r_5(n) + r_6(n),
\end{aligned} \tag{A.60}$$

where

$$\begin{aligned}
|r_5(n)| &= O \left(\max \left\{ G^{n-1} \Delta^{n-2} \left(\frac{1}{2} \Delta |r_2| + 2\varepsilon G + \frac{1}{2} \Delta G \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2} \right), G^{n-1} \Delta^{n-2} R_{V,1} \right\} \right), \\
|r_6(n)| &\leq \varepsilon^{n-2} G^{n-1}
\end{aligned}$$

Proof. Rewrite the left-hand side of (A.60) as

$$\begin{aligned}
& \int_{u_1=0}^{\Delta-\varepsilon} \int_{x_1=0}^{\infty} \frac{\boldsymbol{\alpha} e^{\mathbf{S}(x_0+x_1+u_1)} \mathbf{s}}{\boldsymbol{\alpha} e^{\mathbf{S}x_0} \mathbf{e}} g_1(x_1) du_1 dx_1 \\
& \quad \times \left[\prod_{\ell=2}^{n-1} \int_{u_\ell=0}^{\Delta-\varepsilon} \int_{x_\ell=0}^{\infty} \frac{\boldsymbol{\alpha} e^{\mathbf{S}(u_{\ell-1}+x_\ell+u_\ell)} \mathbf{s}}{\boldsymbol{\alpha} e^{\mathbf{S}u_{\ell-1}} \mathbf{e}} g_\ell(x_\ell) du_\ell dx_\ell \right] \\
& \quad \times \int_{x_n=0}^{\infty} \frac{\boldsymbol{\alpha} e^{\mathbf{S}(u_{n-1}+x_n)} \mathbf{s}}{\boldsymbol{\alpha} e^{\mathbf{S}u_{n-1}} \mathbf{e}} V(x) g_n(x_n) du_{n-1} dx_n,
\end{aligned}$$

then we see that we can apply Corollary A.3 to all of the integrals over x_k , $k = 1, \dots, n-1$ and use Property 3.2(ii) of $V(x)$ to get

$$\begin{aligned}
& \int_{u_1=0}^{\Delta-\varepsilon} [g_1(\Delta - u_1 - x_0)1(u_1 + x_0 \leq \Delta - \varepsilon) + r_3(u_1 + x_0)] \\
& \quad \times \int_{u_2=0}^{\Delta-\varepsilon} [g_2(\Delta - u_2 - u_1)1(u_2 + u_1 \leq \Delta - \varepsilon) + r_3(u_2 + u_1)] du_1 \\
& \quad \dots \int_{u_{n-1}=0}^{\Delta-\varepsilon} [g_{n-1}(\Delta - u_{n-1} - u_{n-2})1(u_{n-1} + u_{n-2} \leq \Delta - \varepsilon) + r_3(u_{n-1} + u_{n-2})] du_{n-2}
\end{aligned}$$

$$\begin{aligned} & \times [g_n(\Delta - u_{n-1} - x)1(u_{n-1} + x \leq \Delta - \varepsilon) + r_V(u_{n-1}, x)] du_{n-1} \\ & = g_{1,n}^{*,\varepsilon}(x_0, x) + r_5(n) \end{aligned}$$

where $r_5(n)$ is an error term. The leading terms of $r_5(n)$ are of the form

$$\begin{aligned} & \int_{u_1=0}^{\Delta-\varepsilon-x_0} g_1(\Delta - u_1 - x_0) \int_{u_2=0}^{\Delta-\varepsilon-u_1} g_2(\Delta - u_2 - u_1) du_1 \\ & \quad \dots \int_{u_{k-1}=0}^{\Delta-\varepsilon-u_{k-2}} g_{k-1}(\Delta - u_{k-1} - u_{k-2}) du_{k-2} \int_{u_k=0}^{\Delta-\varepsilon} r_3(u_k + u_{k-1}) du_{k-1} \\ & \quad \times \int_{u_{k+1}=0}^{\Delta-\varepsilon-u_k} g_{k+1}(\Delta - u_{k+1} - u_k) du_k \dots \int_{u_{n-1}=0}^{\Delta-\varepsilon-u_{n-2}} g_{n-1}(\Delta - u_{n-1} - u_{n-2}) du_{n-2} \\ & \quad \times g_n(\Delta - u_{n-1} - x)1(u_{n-1} + x \leq \Delta - \varepsilon) du_{n-1} \\ & \leq G^{k-1} \Delta^{k-2} \int_{u_{k-1}=0}^{\Delta-\varepsilon} \int_{u_k=0}^{\Delta-\varepsilon} r_3(u_k + u_{k-1}) du_k du_{k-1} G^{n-k} \Delta^{n-k-1}, \end{aligned}$$

and

$$\begin{aligned} & \int_{u_1=0}^{\Delta-\varepsilon-x_0} g_1(\Delta - u_1 - x_0) \int_{u_2=0}^{\Delta-\varepsilon-u_1} g_2(\Delta - u_2 - u_1) du_1 \\ & \quad \dots \int_{u_{n-1}=0}^{\Delta-\varepsilon-u_{n-2}} g_{n-1}(\Delta - u_{n-1} - u_{n-2}) du_{n-2} r_V(u_{n-1}, x) du_{n-1} \\ & \leq G^{n-1} \Delta^{n-2} \int_{u_{n-1}=0}^{\Delta-\varepsilon} r_V(u_{n-1}, x) du_{n-1}. \end{aligned}$$

Now,

$$\begin{aligned} & \left| \int_{u_{k-1}=0}^{\Delta-\varepsilon} \int_{u_k=0}^{\Delta-\varepsilon} r_3(u_k + u_{k-1}) du_k du_{k-1} \right| \\ & \leq \int_{u_{k-1}=0}^{\Delta-\varepsilon} \left[\int_{u_k=u_{k-1}}^{\Delta-\varepsilon} |r_3(u_k)| du_k + \int_{u_k=\Delta-\varepsilon}^{\Delta+\varepsilon} |r_3(u_k)| du_k \right. \\ & \quad \left. + \int_{u_k=\Delta+\varepsilon}^{\Delta-\varepsilon+u_{k-1}} |r_3(u_k)| du_k 1(u_{k-1} > 2\varepsilon) \right] du_{k-1} \\ & \leq \left[\int_{u_{k-1}=0}^{\Delta-\varepsilon} (\Delta - \varepsilon - u_{k-1}) |r_2| + 2\varepsilon G + G \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2} (u_{k-1} - 2\varepsilon) 1(u_{k-1} > 2\varepsilon) \right] du_{k-1} \\ & \leq \frac{1}{2} \Delta^2 |r_2| + 2\Delta\varepsilon G + \frac{1}{2} \Delta^2 G \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2}, \end{aligned}$$

and, by Property 3.2(ii),

$$\int_{u_{n-1}=0}^{\Delta-\varepsilon} |r_V(u_{n-1}, x)| du_{n-1} \leq R_{V,1}$$

Therefore, the error term $|r_5(n)|$ is less than or equal to the larger of these two terms,

$$|r_5(n)| = O \left(\max \left\{ G^{n-1} \Delta^{n-2} \left(\frac{1}{2} \Delta |r_2| + 2\varepsilon G + \frac{1}{2} \Delta G \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2} \right), G^{n-1} \Delta^{n-2} R_{V,1} \right\} \right).$$

Now,

$$\begin{aligned} & \left| g_{1,n}^{*,\varepsilon}(x_0, x) - g_{1,n}^*(x_0, x) \right| \\ &= \int_{u_1=\Delta-\varepsilon-x_0}^{\Delta-x_0} g_1(\Delta - u_1 - x_0) \int_{u_2=\Delta-\varepsilon-u_1}^{\Delta-u_1} g_2(\Delta - u_2 - u_1) du_1 \\ & \quad \dots \int_{u_{n-1}=\Delta-\varepsilon-u_{n-2}}^{\Delta-u_{n-2}} g_{n-1}(\Delta - u_{n-1} - u_{n-2}) du_{n-2} g_n(\Delta - x - u_{n-1}) \\ & \quad \times 1(\Delta - x - u_{n-1} \geq 0) du_{n-1} \\ &\leq \int_{u_1=\Delta-\varepsilon-x_0}^{\Delta-x_0} G \int_{u_2=\Delta-\varepsilon-u_1}^{\Delta-u_1} G du_1 \dots \int_{u_{n-1}=\Delta-\varepsilon-u_{n-2}}^{\Delta-u_{n-2}} G du_{n-2} G du_{n-1} \\ &= \varepsilon^{n-1} G^n \end{aligned}$$

Therefore, the left-hand side of (A.60) is equal to

$$g_{1,n}^{*,\varepsilon}(x_0, x) + r_5(n) + r_6(n),$$

where $r_6(n) = g_{1,n}^*(x_0, x) - g_{1,n}^{*,\varepsilon}(x_0, x)$, and $|r_6(n)| \leq \varepsilon^{n-1} G^n$. \square

The error terms $r_5(n)$ depend on p as they are functions of $r_2^{(p)}$, $\varepsilon^{(p)}$, $\text{Var}(Z^{(p)})$ and $R_{V,1}^{(p)}$. We write $r_5^{(p)}(n)$ when this dependence is explicitly needed, otherwise this dependence is omitted from the notation. Choosing $\varepsilon = \text{Var}(Z^{(p)})^{1/3}$, the error term $|r_5^{(p)}(n)| = O(\text{Var}(Z^{(p)})^{1/3}) \rightarrow 0$ as $p \rightarrow \infty$. Similarly, $r_6(n)$ depends on p as it is a function of $\varepsilon^{(p)}$ and we write $r_6^{(p)}(n)$ when we need to denote this explicitly.

Corollary A.11. *Let g_1, g_2, \dots , be functions satisfying Assumptions 3.1 and let $V(x)$, $x \in [0, \Delta)$, be a closing operator with Properties 3.2. Then, for $n \geq 2$, $x_0 \in [0, \Delta)$,*

$$\left| \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}(x_0) e^{\mathbf{S}x_1} dx_1 \mathbf{D} \left[\prod_{k=2}^{n-1} \int_{x_k=0}^{\infty} g_k(x_k) e^{\mathbf{S}x_k} dx_k \mathbf{D} \right] \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} dx_n V(x) \right|$$

$$\begin{aligned}
& - \int_{u_1=0}^{\Delta-x_0} g_1(\Delta - u_1 - x_0) \left[\prod_{k=2}^{n-1} \int_{u_k=0}^{\Delta-u_{k-1}} g_k(\Delta - u_k - u_{k-1}) \, du_{k-1} \right] g_n(\Delta - x - u_{n-1}) \\
& \quad \left| 1(\Delta - x - u_{n-1} \geq 0) \, du_{n-1} \right| \\
& \leq |r_5(n)| + |r_6(n)| + (n-1)|r_4(n)|,
\end{aligned} \tag{A.61}$$

where

$$\begin{aligned}
|r_4(n)| &= \left(2\varepsilon + \frac{\text{Var}(Z)}{\varepsilon} \right) \frac{1}{1 - \text{Var}(Z)/(\Delta - x_0)} G \hat{G}^{n-2} G, \\
|r_5(n)| &= O \left(\max \left\{ G^{n-1} \Delta^{n-2} \left(\frac{1}{2} \Delta |r_2| + 2\varepsilon G + \frac{1}{2} \Delta G \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2} \right), G^{n-1} \Delta^{n-2} R_{V,1} \right\} \right) \\
|r_6(n)| &\leq \varepsilon^{n-1} G^n.
\end{aligned}$$

Proof. The left-most term on the left-hand side of (A.61) can be written as

$$\begin{aligned}
& \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}(x_0) e^{\mathbf{S}x_1} \, dx_1 \mathbf{D}(\Delta - \varepsilon) \left[\prod_{k=2}^{n-1} \int_{x_k=0}^{\infty} g_k(x_k) e^{\mathbf{S}x_k} \, dx_k \mathbf{D}(\Delta - \varepsilon) \right] \\
& \quad \times \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} \, dx_n V(x) + \sum_{k=1}^{n-1} \int_{x_{k+1}=0}^{\infty} \int_{u_k=\Delta-\varepsilon}^{\infty} \mathbf{a}(x_0) \mathcal{I}_{1,k}(u_k) \mathcal{J}_{k+1,n}(u_k, x_{k+1}) V(x).
\end{aligned} \tag{A.62}$$

Now, substitute this into the left-hand side of (A.61), apply the triangle inequality and Lemmas A.9 and A.10 to get the result. \square

Define

$$\begin{aligned}
w_n(x_0, x) &= \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}(x_0) e^{\mathbf{S}x_1} \, dx_1 \mathbf{D} \left[\prod_{k=2}^{n-1} \int_{x_k=0}^{\infty} g_k(x_k) e^{\mathbf{S}x_k} \, dx_k \mathbf{D} \right] \\
& \quad \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} \, dx_n V(x),
\end{aligned}$$

where g_1, g_2, \dots , are functions satisfying the Assumptions 3.1 and $V(x)$ is a closing operator with the Properties 3.2.

Corollary A.12. *Let g_1, g_2, \dots , be functions satisfying Assumptions 3.1 and let $V(x)$, $x \in [0, \Delta)$, be a closing operator with Properties 3.2. Then, for $n \geq 2$, $x_0 \in [0, \Delta)$,*

$$\left| \int_{x=0}^{\Delta} w_n(x_0, x) \psi(x) \, dx \right|$$

$$\begin{aligned}
& - \int_{x=0}^{\Delta} \int_{u_1=0}^{\Delta-x_0} g_1(\Delta - u_1 - x_0) \left[\prod_{k=2}^{n-1} \int_{u_k=0}^{\Delta-u_{k-1}} g_k(\Delta - u_k - u_{k-1}) \, du_{k-1} \right] g_n(\Delta - x - u_{n-1}) \\
& \quad \left| 1(\Delta - x - u_{n-1} \geq 0) \, du_{n-1} \psi(x) \, dx \right| \\
& \leq (|r_5(n)| + |r_6(n)| + (n-1)|r_4(n)|) \Delta F.
\end{aligned} \tag{A.63}$$

Proof. The left-hand side of (A.63) is less than or equal to

$$\begin{aligned}
& \int_{x=0}^{\Delta} \left| w_n(x_0, x) \right. \\
& \quad - \int_{u_1=0}^{\Delta-x_0} g_1(\Delta - u_1 - x_0) \left[\prod_{k=2}^{n-1} \int_{u_k=0}^{\Delta-u_{k-1}} g_k(\Delta - u_k - u_{k-1}) \, du_{k-1} \right] g_n(\Delta - x - u_{n-1}) \\
& \quad \left. 1(\Delta - x - u_{n-1} \geq 0) \, du_{n-1} \right| |\psi(x)| \, dx.
\end{aligned} \tag{A.64}$$

Apply Corollary A.11 to bound the first absolute value so that (A.64) is less than or equal to

$$\begin{aligned}
& \int_{x=0}^{\Delta} (|r_5(n)| + |r_6(n)| + (n-1)|r_4(n)|) |\psi(x)| \, dx \\
& \leq \int_{x=0}^{\Delta} (|r_5(n)| + |r_6(n)| + (n-1)|r_4(n)|) F \, dx \\
& = (|r_5(n)| + |r_6(n)| + (n-1)|r_4(n)|) \Delta F
\end{aligned} \tag{A.65}$$

□

At this point, we have all the results we need to prove weak convergence of the QBD-RAP approximation on the event that $\varphi(0) \notin \mathcal{S}_{+0} \cup \mathcal{S}_{-0}$. The difficulty that remains is to deal with sample paths of the QBD-RAP which start in $\phi(0) \in \mathcal{S}_{+0}$ (or \mathcal{S}_{-0}), and upon leaving this subset of phases jump to $i \in \mathcal{S}_-$ (\mathcal{S}_+). The orbit process at the time at which $\{\phi(t)\}$ first exits \mathcal{S}_{+0} (\mathcal{S}_{-0}) is $\mathbf{a}(x_0)\mathbf{D}$, thus we can treat this case simply as a change in the initial condition of the QBD-RAP. We do so by showing that asymptotically, the initial conditions $\mathbf{a}(\Delta - x_0)$ and $\mathbf{a}(x_0)\mathbf{D}$ produce results that are arbitrarily close to each other. We first show a Lipschitz-like condition in x_0 for $w_n(x_0, x)$.

Corollary A.13. For $x_0, x \in [0, \Delta)$, $n \geq 2$,

$$|w_n(x_0, x) - w_n(z_0, x)| \leq 2|r_5(n)| + 2|r_6(n)| + 2(n-1)|r_4(n)| + |x_0 - z_0|G_n^*(G + L\Delta), \tag{A.66}$$

Proof. By adding and subtracting both $\int_{u_1=0}^{\Delta-x_0} g_1(\Delta-u_1-x_0)g_{2,n}^*(u_1, x) du_1$ and $\int_{u_1=0}^{\Delta-z_0} g_1(\Delta-u_1-z_0)g_{2,n}^*(u_1, x) du_1$, we can write the left-hand side of (A.66) as

$$\begin{aligned} & \left| w_n(x_0, x) - \int_{u_1=0}^{\Delta-x_0} g_1(\Delta-u_1-x_0)g_{2,n}^*(u_1, x) du_1 \right. \\ & \quad - w_n(z_0, x) + \int_{u_1=0}^{\Delta-z_0} g_1(\Delta-u_1-z_0)g_{2,n}^*(u_1, x) du_1 \\ & \quad \left. + \int_{u_1=0}^{\Delta-x_0} g_1(\Delta-u_1-x_0)g_{2,n}^*(u_1, x) du_1 - \int_{u_1=0}^{\Delta-z_0} g_1(\Delta-u_1-z_0)g_{2,n}^*(u_1, x) du_1 \right| \end{aligned}$$

which, by the triangle inequality, is less than or equal to

$$\begin{aligned} & \left| w_n(x_0, x) - \int_{u_1=0}^{\Delta-x_0} g_1(\Delta-u_1-x_0)g_{2,n}^*(u_1, x) du_1 \right| \\ & + \left| w_n(z_0, x) - \int_{u_1=0}^{\Delta-z_0} g_1(\Delta-u_1-z_0)g_{2,n}^*(u_1, x) du_1 \right| \\ & + \left| \int_{u_1=0}^{\Delta-x_0} g_1(\Delta-u_1-x_0)g_{2,n}^*(u_1, x) du_1 - \int_{u_1=0}^{\Delta-z_0} g_1(\Delta-u_1-z_0)g_{2,n}^*(u_1, x) du_1 \right| \end{aligned} \quad (\text{A.67})$$

By Corollary A.11, the first two terms of (A.67) are less than or equal to $|r_5(n)| + |r_6(n)| + (n-1)|r_4(n)|$. As for the last term, adding and subtracting $\int_{u_1=0}^{\Delta-z_0} g_1(\Delta-u_1-x_0)g_{2,n}^*(u_1, x) du_1$ gives

$$\begin{aligned} & = \left| \int_{u_1=0}^{\Delta-x_0} g_1(\Delta-u_1-x_0)g_{2,n}^*(u_1, x) du_1 - \int_{u_1=0}^{\Delta-z_0} g_1(\Delta-u_1-x_0)g_{2,n}^*(u_1, x) du_1 \right. \\ & \quad \left. - \int_{u_1=0}^{\Delta-z_0} (g_1(\Delta-u_1-z_0) - g_1(\Delta-u_1-x_0))g_{2,n}^*(u_1, x) du_1 \right| \\ & \leq \left| \int_{u_1=\Delta-z_0}^{\Delta-x_0} g_1(\Delta-u_1-x_0)g_{2,n}^*(u_1, x) du_1 \right| \\ & \quad + \int_{u_1=0}^{\Delta-z_0} |g_1(\Delta-u_1-z_0) - g_1(\Delta-u_1-x_0)|g_{2,n}^*(u_1, x) du_1 \\ & \leq GG_n^*|x_0 - z_0| + \int_{u_1=0}^{\Delta-z_0} L|x_0 - z_0|G_n^* du_1 \end{aligned} \quad (\text{A.68})$$

since g_1 is Lipschitz by Assumption 3.1(iv) and $g_{2,n}^* \leq G_n^*$. Bounding the integral over u_1 by Δ , then (A.68) is less than or equal to

$$GG_n^*|x_0 - z_0| + \Delta L|x_0 - z_0|G_n^*. \quad (\text{A.69})$$

□

Corollary A.14. *Let g_1, g_2, \dots , be functions satisfying Assumptions 3.1 and let $V(x)$, $x \in [0, \Delta)$, be a closing operator with Properties 3.2. For $x_0, x \in [0, \Delta)$, $n \geq 2$,*

$$\begin{aligned} & \left| \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}(x_0) \mathbf{D} e^{\mathbf{S}x_1} dx_1 \mathbf{D} \left[\prod_{n=2}^{k-1} \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} dx_n \mathbf{D} \right] \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} dx_n V(x) \right. \\ & \quad \left. - w_n(\Delta - x_0, x) \right| \\ & = r_8(n), \end{aligned} \quad (\text{A.70})$$

where

$$\begin{aligned} |r_8(n)| & \leq (2|r_5(n)| + 2|r_6(n)| + 2(n-1)|r_4(n)| + \varepsilon G_n^*(G + L\Delta)) \\ & \quad + 2\hat{G}^{n-2}GG_V \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2}. \end{aligned}$$

Proof. Observe that

$$\begin{aligned} & \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}(x_0) \mathbf{D} e^{\mathbf{S}x_1} dx_1 \mathbf{D} \left[\prod_{n=2}^{k-1} \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} dx_n \mathbf{D} \right] \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} dx_n V(x) \\ & = \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}(x_0) \int_{z_0=0}^{\infty} e^{\mathbf{S}z_0} \mathbf{s} \frac{\boldsymbol{\alpha} e^{\mathbf{S}z_0}}{\boldsymbol{\alpha} e^{\mathbf{S}z_0} \mathbf{e}} dz_0 e^{\mathbf{S}x_1} dx_1 \mathbf{D} \left[\prod_{n=2}^{k-1} \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} dx_n \mathbf{D} \right] \\ & \quad \times \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} dx_n V(x) \\ & = \mathbf{a}(x_0) \int_{z_0=0}^{\infty} e^{\mathbf{S}z_0} \mathbf{s} w_n(z_0, x) dz_0 \end{aligned} \quad (\text{A.71})$$

Now

$$\begin{aligned} & \left| \mathbf{a}(x_0) \int_{z_0=0}^{\infty} e^{\mathbf{S}z_0} \mathbf{s} w_n(z_0, x) dz_0 - w_n(\Delta - x_0, x) \right| \\ & = \left| \mathbf{a}(x_0) \int_{z_0=0}^{\infty} e^{\mathbf{S}z_0} \mathbf{s} (w_n(z_0, x) - w_n(\Delta - x_0, x)) dz_0 \right| \end{aligned}$$

$$\begin{aligned}
&\leq \mathbf{a}(x_0) \int_{z_0=0}^{\infty} e^{\mathbf{S}z_0} \mathbf{s} |w_n(z_0, x) - w_n(\Delta - x_0, x)| \, dz_0 \\
&= \mathbf{a}(x_0) \int_{z_0=0}^{\Delta-\varepsilon-x_0} e^{\mathbf{S}z_0} \mathbf{s} |w_n(z_0, x) - w_n(\Delta - x_0, x)| \, dz_0 \\
&\quad + \mathbf{a}(x_0) \int_{z_0=\Delta+\varepsilon-x_0}^{\infty} e^{\mathbf{S}z_0} \mathbf{s} |w_n(z_0, x) - w_n(\Delta - x_0, x)| \, dz_0 \\
&\quad + \mathbf{a}(x_0) \int_{z_0=\Delta-\varepsilon-x_0}^{\Delta+\varepsilon-x_0} e^{\mathbf{S}z_0} \mathbf{s} |w_n(z_0, x) - w_n(\Delta - x_0, x)| \, dz_0. \tag{A.72}
\end{aligned}$$

By Corollary A.8, for any $x, y \in [0, \Delta)$, $0 \leq w_n(x, y) \leq \widehat{G}^{n-2}GG_V$, so the sum of the first two terms is less than or equal to

$$2\widehat{G}^{n-1}GG_V \left(\int_{z_0=0}^{\Delta-\varepsilon-x_0} \mathbf{a}(x_0) e^{\mathbf{S}z_0} \mathbf{s} \, dz_0 + \int_{z_0=\Delta+\varepsilon-x_0}^{\infty} \mathbf{a}(x_0) e^{\mathbf{S}z_0} \mathbf{s} \, dz_0 \right) \tag{A.73}$$

$$\begin{aligned}
&= 2\widehat{G}^{n-1}GG_V \frac{\mathbb{P}(|Z - \Delta| > \varepsilon)}{\mathbb{P}(Z > x_0)} \\
&\leq 2G^n \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2} \tag{A.74}
\end{aligned}$$

by Chebyshev's inequality. As for the last term in (A.72), we can use Corollary A.13 to bound the integrand so that the last term is less than or equal to

$$\begin{aligned}
&\mathbf{a}(x_0) \int_{z_0=\Delta-\varepsilon-x_0}^{\Delta+\varepsilon-x_0} e^{\mathbf{S}z_0} \mathbf{s} (2|r_5(n)| + 2|r_6(n)| + 2(n-1)|r_4(n)| + \varepsilon G^{n-1} \Delta^{n-2} (G + L\Delta)) \, dz_0 \\
&\leq (2|r_5(n)| + 2|r_6(n)| + 2(n-1)|r_4(n)| + \varepsilon G^{n-1} \Delta^{n-2} (G + L\Delta)) \\
&\quad + 2\widehat{G}^{n-2}GG_V \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2}, \tag{A.75}
\end{aligned}$$

since $\mathbf{a}(x_0) \int_{z_0=\Delta-\varepsilon-x_0}^{\Delta+\varepsilon-x_0} e^{\mathbf{S}z_0} \mathbf{s} \, dz_0 \leq 1$. □

Corollary A.15. *Let g_1, g_2, \dots , be functions satisfying Assumptions 3.1 and let $V(x)$, $x \in (0, \Delta)$, be a closing operator with Properties 3.2. For $x_0, x \in (0, \Delta)$, $n \geq 2$*

$$\begin{aligned}
&\left| \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}(x_0) \mathbf{D} e^{\mathbf{S}x_1} \, dx_1 \mathbf{D} \left[\prod_{n=2}^{k-1} \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} \, dx_n \mathbf{D} \right] \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} \, dx_n V(x) \right. \\
&\quad \left. - \int_{u_1=0}^{x_0} g_1(x_0 - u_1) \left[\prod_{k=2}^{n-1} \int_{u_k=0}^{\Delta-u_{k-1}} g_k(\Delta - u_k - u_{k-1}) \, du_{k-1} \right] g_n(\Delta - x - u_{n-1}) \right|
\end{aligned}$$

$$\begin{aligned}
& \left| 1(\Delta - x - u_{n-1} \geq 0) \, du_{n-1} \right| \\
& \leq |r_8(n)| + |r_5(n)| + |r_6(n)| + (n-1)|r_4(n)|,
\end{aligned} \tag{A.76}$$

where

$$|r_8(n)| \leq (2|r_5(n)| + 2|r_6(n)| + 2(n-1)|r_4(n)| + \varepsilon G^{m-1} \Delta^{n-2} (G + L\Delta)) \tag{A.77}$$

$$+ 2\widehat{G}^{m-2} G G_V \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2}. \tag{A.78}$$

Proof. Adding and subtracting $w_n(\Delta - x_0, x)$ within the absolute value on the left-hand side of (A.76)

$$\begin{aligned}
& \left| \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}(x_0) \mathbf{D} e^{\mathbf{S}x_1} \, dx_1 \mathbf{D} \left[\prod_{n=2}^{k-1} \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} \, dx_n \mathbf{D} \right] \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} \, dx_n V(x) \right. \\
& \quad \left. - w_n(\Delta - x_0, x) + w_n(\Delta - x_0, x) - g_{1,n}^*(\Delta - x_0, x) \right| \\
& \leq \left| \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}(x_0) \mathbf{D} e^{\mathbf{S}x_1} \, dx_1 \mathbf{D} \left[\prod_{n=2}^{k-1} \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} \, dx_n \mathbf{D} \right] \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} \, dx_n V(x), \right. \\
& \quad \left. - w_n(\Delta - x_0, x) \right| + \left| w_n(\Delta - x_0, x) - g_{1,n}^*(\Delta - x_0, x) \right|
\end{aligned}$$

where the first absolute value is less than or equal to $|r_8(n)|$ by Corollary A.14 and the second absolute value is less than or equal to $|r_5(n)| + |r_6(n)| + (n-1)|r_4(n)|$ by Corollary A.11. \square

Corollary A.16. *Let ψ be bounded and Lipschitz, let g_1, g_2, \dots , be functions satisfying Assumptions 3.1 and let $V(x)$, $x \in (0, \Delta)$, be a closing operator with Properties 3.2. For $x_0, x \in (0, \Delta)$, $n \geq 2$*

$$\begin{aligned}
& \left| \int_{x \in [0, \Delta)} \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}(x_0) \mathbf{D} e^{\mathbf{S}x_1} \, dx_1 \mathbf{D} \left[\prod_{n=2}^{k-1} \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} \, dx_n \mathbf{D} \right] \right. \\
& \quad \times \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} \, dx_n V(x) \psi(x) \, dx \\
& \quad - \int_{x \in [0, \Delta)} \int_{u_1=0}^{x_0} g_1(x_0 - u_1) \left[\prod_{k=2}^{n-1} \int_{u_k=0}^{\Delta - u_{k-1}} g_k(\Delta - u_k - u_{k-1}) \, du_{k-1} \right] g_n(\Delta - x - u_{n-1}) \\
& \quad \times 1(\Delta - x - u_{n-1} \geq 0) \, du_{n-1} \psi(x) \, dx \Big|
\end{aligned}$$

$$\leq (|r_8(n)| + |r_5(n)| + |r_6(n)| + (n-1)|r_4(n)|) F \Delta. \quad (\text{A.79})$$

Proof. The left-hand side of (3.45) is less than or equal to

$$\begin{aligned} & \int_{x \in [0, \Delta)} \left| \int_{x_1=0}^{\infty} g_1(x_1) \mathbf{a}(x_0) \mathbf{D} e^{\mathbf{S}x_1} dx_1 \mathbf{D} \left[\prod_{n=2}^{k-1} \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} dx_n \mathbf{D} \right] \right. \\ & \quad \times \int_{x_n=0}^{\infty} g_n(x_n) e^{\mathbf{S}x_n} dx_n V(x) - \int_{u_1=0}^{x_0} g_1(x_0 - u_1) \\ & \quad \times \left[\prod_{k=2}^{n-1} \int_{u_k=0}^{\Delta - u_{k-1}} g_k(\Delta - u_k - u_{k-1}) du_{k-1} \right] g_n(\Delta - x - u_{n-1}) \\ & \quad \left. 1(\Delta - x - u_{n-1} \geq 0) du_{n-1} \right| |\psi(x)| dx. \end{aligned} \quad (\text{A.80})$$

Now, apply Corollary A.15 to the first absolute value then (A.80) is less than or equal to

$$\begin{aligned} & \int_{x \in [0, \Delta)} (|r_8(n)| + |r_5(n)| + |r_6(n)| + (n-1)|r_4(n)|) |\psi(x)| dx \\ & \leq \int_{x \in [0, \Delta)} (|r_8(n)| + |r_5(n)| + |r_6(n)| + (n-1)|r_4(n)|) F dx \\ & = (|r_8(n)| + |r_5(n)| + |r_6(n)| + (n-1)|r_4(n)|) \Delta F \end{aligned} \quad (\text{A.81})$$

□

A.2 Properties of closing operators

Corollary A.17. *Let g be a function satisfying the Assumptions 3.1 and consider the closing operator $V(x) = e^{\mathbf{S}x} \mathbf{s}$. For $u \leq \Delta - \varepsilon$, $v \geq 0$,*

$$\int_{x=0}^{\infty} \frac{\boldsymbol{\alpha} e^{\mathbf{S}(u+x)}}{\boldsymbol{\alpha} e^{\mathbf{S}u} \mathbf{e}} V(v) g(x) dx = g(\Delta - u - v) 1(u + v \leq \Delta - \varepsilon) + r_V(u, v),$$

where

$$\int_{u=0}^{\Delta - \varepsilon} r_V(u, v) du = \int_{u=0}^{\Delta - \varepsilon} r_3(u + v) du \leq r_2 \Delta + 2\varepsilon G + \Delta G \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2}$$

and

$$\int_{u=0}^{\Delta} r_V(u, v) du \leq R_{V,1}$$

where

$$R_{V,1} = r_2 \Delta + 2\varepsilon G + \Delta G \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2}.$$

Proof. By Corollary A.3,

$$\int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(u+x)}}{\alpha e^{\mathbf{S}u} \mathbf{e}} V(v) g(x) \, dx = g(\Delta - u - v) 1(u + v \leq \Delta - \varepsilon) + r_3(u + v),$$

so $r_V(u, v) = r_3(u + v)$. All that remains to be shown are the bounds $R_{V,1}$ and $R_{V,2}$. To this end, observe

$$R_{V,1} = \int_{u=0}^{\Delta} r_V(u, v) \, du = \int_{u=0}^{\Delta} r_3(u + v) \, du \leq r_2 \Delta + 2\varepsilon G + \Delta G \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2}.$$

$$R_{V,2} = \int_{v=0}^{\Delta} r_V(u, v) \, dv = \int_{v=0}^{\Delta} r_3(u + v) \, dv \leq r_2 \Delta + 2\varepsilon G + \Delta G \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2}.$$

□

Lemma A.18. *For any valid orbit, $\mathbf{a} \in \mathcal{A}$, $x, u \geq 0$,*

$$\int_{x_n=0}^{\infty} \mathbf{a} e^{\mathbf{S}(x+x_n+u)} \mathbf{s} = \mathbf{a} e^{\mathbf{S}(x+u)} \mathbf{e} \leq \mathbf{a} e^{\mathbf{S}u} \mathbf{e}.$$

Proof. For any valid orbit, $\mathbf{a} \in \mathcal{A}$,

$$\mathbf{a} e^{\mathbf{S}(x+u)} \mathbf{e} = \mathbb{P}(Z > x + u) \leq \mathbb{P}(X > u) = \mathbf{a} e^{\mathbf{S}u} \mathbf{e}.$$

□

Lemma A.19. *For any valid orbit, $\mathbf{a} \in \mathcal{A}$, $x \geq 0$,*

$$\mathbf{a} D e^{\mathbf{S}x} \mathbf{e} \leq 1.$$

Proof. By the definition of D and Lemma A.18,

$$\begin{aligned} \mathbf{a} D e^{\mathbf{S}x} \mathbf{e} &= \mathbf{a} \int_{u=0}^{\infty} e^{\mathbf{S}u} \mathbf{s} \frac{\alpha e^{\mathbf{S}u}}{\alpha e^{\mathbf{S}u} \mathbf{e}} \, du e^{\mathbf{S}x} \mathbf{e} \\ &\leq \mathbf{a} \int_{u=0}^{\infty} e^{\mathbf{S}u} \mathbf{s} \frac{\alpha e^{\mathbf{S}u} \mathbf{e}}{\alpha e^{\mathbf{S}u} \mathbf{e}} \, du \\ &= \mathbf{a} \int_{u=0}^{\infty} e^{\mathbf{S}u} \mathbf{s} \, du \\ &= \mathbf{a} \mathbf{e} = 1. \end{aligned}$$

□

Let $U^{(p)}(x)$ be the closing operator such that, for $\mathbf{a}^{(p)} \in \mathcal{A}^{(p)}$, $x \in [0, \Delta)$,

$$\mathbf{a}^{(p)} U^{(p)}(x) = \mathbf{a}^{(p)} \left(e^{\mathbf{S}^{(p)} x} \mathbf{s}^{(p)} + e^{\mathbf{S}^{(p)} (2\Delta - x)} \mathbf{s}^{(p)} \right).$$

Lemma A.20. For $x \in [0, \Delta)$, $u \geq 0$,

$$\mathbf{a} e^{\mathbf{S}^u} (-\mathbf{S})^{-1} U(x) \leq 2\mathbf{a} e^{\mathbf{S}^u} \mathbf{e}.$$

Proof. Let $\mathbf{a} \in \mathcal{A}$ be arbitrary. By definition

$$\begin{aligned} \mathbf{a} e^{\mathbf{S}^u} (-\mathbf{S})^{-1} U(x) &= \mathbf{a} e^{\mathbf{S}^u} (-\mathbf{S})^{-1} (e^{\mathbf{S}^x} \mathbf{s} + e^{\mathbf{S}^{(2\Delta - x)}} \mathbf{s}) \\ &= \mathbf{a} e^{\mathbf{S}^u} (e^{\mathbf{S}^x} \mathbf{e} + e^{\mathbf{S}^{(2\Delta - x)}} \mathbf{e}) \end{aligned}$$

since $(-\mathbf{S})^{-1}$ and $e^{\mathbf{S}^x}$ commute and $\mathbf{s} = -\mathbf{S}\mathbf{e}$. By Lemma A.18 this is less than or equal to,

$$\mathbf{a} e^{\mathbf{S}^u} (\mathbf{e} + \mathbf{e}) = 2\mathbf{a} e^{\mathbf{S}^u} \mathbf{e} \tag{A.82}$$

□

Corollary A.21. Let g be a function satisfying the Assumptions 3.1. For $u \leq \Delta - \varepsilon$, $v \in [0, \Delta)$,

$$\int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}^{(u+x)}}}{\alpha e^{\mathbf{S}^u} \mathbf{e}} U(v) g(x) dx = g(\Delta - u - v) 1(u + v \leq \Delta - \varepsilon) + r_V(u, v),$$

where

$$|r_V(u, v)| \leq r_3(u + v) + r_3(u + 2\Delta - v).$$

Furthermore,

$$\int_{u=0}^{\Delta} |r_V(u, x)| du \leq R_{V,1},$$

and

$$\int_{v=0}^{\Delta} |r_V(u, x)| dv \leq R_{V,2},$$

where

$$R_{V,1}, R_{V,2} \leq 2 \left(\Delta r_2 + 2\varepsilon G + \Delta \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2} \right).$$

Proof. By the definition of the operator $U(x)$,

$$\int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(u+x)}}{\alpha e^{\mathbf{S}u} \mathbf{e}} U(v) g(x) \, dx = \int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(u+x)}}{\alpha e^{\mathbf{S}u} \mathbf{e}} e^{\mathbf{S}v} \mathbf{s} g(x) + \frac{\alpha e^{\mathbf{S}(u+x)}}{\alpha e^{\mathbf{S}u} \mathbf{e}} e^{\mathbf{S}(2\Delta-v)} \mathbf{s} g(x) \, dx. \quad (\text{A.83})$$

By Corollary A.3

$$\int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(u+x)}}{\alpha e^{\mathbf{S}u} \mathbf{e}} e^{\mathbf{S}v} \mathbf{s} g(x) \, dx = g(\Delta - u - v) 1(u + v \leq \Delta - \varepsilon) + r_3(u + v), \quad (\text{A.84})$$

$$\int_{x=0}^{\infty} \frac{\alpha e^{\mathbf{S}(u+x)}}{\alpha e^{\mathbf{S}u} \mathbf{e}} e^{\mathbf{S}(2\Delta-v)} \mathbf{s} g(x) \, dx = r_3(u + 2\Delta - v). \quad (\text{A.85})$$

Therefore, (A.83) is,

$$g(\Delta - u - v) 1(u + v \leq \Delta - \varepsilon) + r_3(u + v) + r_3(u + 2\Delta - v). \quad (\text{A.86})$$

Now,

$$\begin{aligned} R_{V,1} &\leq \int_{u=0}^{\Delta} |r_V(u, v)| \, du \\ &\leq \int_{u=0}^{\Delta} |r_3(u + v)| + |r_3(u + 2\Delta - v)| \\ &\leq 2 \left(\Delta r_2 + 2\varepsilon G + \Delta \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2} \right). \end{aligned}$$

Similarly

$$\begin{aligned} R_{V,2} &\leq \int_{v=0}^{\Delta} |r_V(u, v)| \, dv \\ &= \int_{v=0}^{\Delta} r_3(u + v) + r_3(u + 2\Delta - v) \\ &\leq 2 \left(\Delta r_2 + 2\varepsilon G + \Delta \frac{\text{Var}(Z)/\varepsilon^2}{1 - \text{Var}(Z)/\varepsilon^2} \right). \end{aligned}$$

□

The error term $r_V(u, v)$ depends on p so we should write $r_V^{(p)}(u, v)$. The error term $r_V^{(p)}(u, v)$ has similar properties to $r_3^{(p)}(u + v)$; we can prove that it converges point-wise to 0 only on some areas of its domain, however when we integrate the error against bounded functions on bounded domains, then the resulting integral tends to 0.

A.3 Kronecker properties

Here we detail some properties of Kronecker sum, products, and exponentials (see (Bladt & Nielsen 2017), Appendix A.4).

Let

$$\mathbf{A} = \begin{bmatrix} a_{11} & \dots & a_{1m} \\ \dots & & \dots \\ a_{n1} & \dots & a_{nm} \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} b_{11} & \dots & b_{1m'} \\ \dots & & \dots \\ b_{n'1} & \dots & b_{n'm'} \end{bmatrix}$$

be matrices. The operator \otimes is the Kronecker product of two matrices;

$$\mathbf{A} \otimes \mathbf{B} = \begin{bmatrix} a_{11}\mathbf{B} & \dots & a_{1m}\mathbf{B} \\ \dots & & \dots \\ a_{n1}\mathbf{B} & \dots & a_{nm}\mathbf{B} \end{bmatrix},$$

which is an $nn' \times mm'$ matrix.

Let \mathbf{C}, \mathbf{D} be matrices with dimensions $m \times k$ and $m' \times k'$. A property of the Kronecker Product is

$$(\mathbf{A} \otimes \mathbf{B})(\mathbf{C} \otimes \mathbf{D}) = \mathbf{AC} \otimes \mathbf{BD}. \quad (\text{Mixed Product Rule})$$

Proof. The proof follows from

$$\begin{bmatrix} a_{i1}\mathbf{B} & a_{i2}\mathbf{B} & \dots & a_{in}\mathbf{B} \end{bmatrix} \begin{bmatrix} c_{1j}\mathbf{D} \\ c_{2j} \\ \vdots \\ c_{nj}\mathbf{D} \end{bmatrix} = \left(\sum_{\ell} a_{i\ell} c_{\ell j} \right) \mathbf{BD} \\ = (\mathbf{AC})_{ij} \mathbf{BD}.$$

□

If \mathbf{A} and \mathbf{B} are invertible matrices, then

$$(\mathbf{A} \otimes \mathbf{B})^{-1} = \mathbf{A}^{-1} \otimes \mathbf{B}^{-1}. \quad (\text{A.87})$$

Let \mathbf{A} and \mathbf{B} be $n \times n$ and $m \times m$ matrices, respectively. The Kronecker sum of \mathbf{A} and \mathbf{B} is denoted by \oplus and defined as

$$\mathbf{A} \oplus \mathbf{B} := \mathbf{A} \otimes \mathbf{I}_m + \mathbf{I}_n \otimes \mathbf{B}.$$

A property of the Kronecker sum is

$$e^{\mathbf{A} \oplus \mathbf{B}} = e^{\mathbf{A}} \otimes e^{\mathbf{B}}. \quad (\text{A.88})$$

Proof. First, the matrices $\mathbf{A} \otimes \mathbf{I}_m$ and $\mathbf{I}_n \otimes \mathbf{B}$ commute; from the mixed product rule their product is $\mathbf{A} \otimes \mathbf{B}$. Hence

$$e^{\mathbf{A} \oplus \mathbf{B}} = e^{\mathbf{A} \otimes \mathbf{I}_m} e^{\mathbf{I}_n \otimes \mathbf{B}}.$$

We now show that $e^{\mathbf{A} \otimes \mathbf{I}_m} = e^{\mathbf{A}} \otimes \mathbf{I}_m$ and $e^{\mathbf{I}_n \otimes \mathbf{B}} = \mathbf{I}_n \otimes e^{\mathbf{B}}$. The latter follows from the fact that $\mathbf{I}_n \otimes \mathbf{B}$ is a block diagonal matrix with blocks \mathbf{B} , hence its exponential is also block diagonal with blocks equal to the exponential of \mathbf{B} . The former follows from

$$\begin{aligned} e^{\mathbf{A} \otimes \mathbf{I}_m} &= \sum_{n=0}^{\infty} \frac{1}{n!} (\mathbf{A} \otimes \mathbf{I}_m)^n \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} (\mathbf{A}^n \otimes \mathbf{I}_m) \\ &= \left(\sum_{n=0}^{\infty} \frac{1}{n!} \mathbf{A}^n \otimes \mathbf{I}_m \right) \\ &= e^{\mathbf{A}} \otimes \mathbf{I}_m. \end{aligned} \tag{A.89}$$

Therefore

$$e^{\mathbf{A} \oplus \mathbf{B}} = (e^{\mathbf{A}} \otimes \mathbf{I}_m) (\mathbf{I}_n \otimes e^{\mathbf{B}}),$$

and the result follows by the mixed product rule. \square

Lemma A.22. Let \mathbf{T} and \mathbf{C} be $n \times n$, square matrices with \mathbf{C} diagonal and invertible; let \mathbf{S} be a $p \times p$ matrix. Further, suppose $[\mathbf{T} \otimes \mathbf{I} + \mathbf{C} \otimes \mathbf{S} - \lambda \mathbf{I}]$ is invertible for $\lambda > 0$. Then

$$\int_{t=0}^{\infty} e^{-\lambda t} e^{(\mathbf{T} \otimes \mathbf{I} + \mathbf{C} \otimes \mathbf{S})t} dt = \int_{x=0}^{\infty} e^{\mathbf{C}^{-1}(\mathbf{T} - \lambda \mathbf{I})x} \otimes e^{\mathbf{S}x} dx (\mathbf{C} \otimes \mathbf{I})^{-1} \tag{A.90}$$

Proof. Computing the integral on the left-hand side and then factorising the result and using the Mixed Product Rule multiple times gives

$$\begin{aligned} \int_{t=0}^{\infty} e^{-\lambda t} e^{(\mathbf{T} \otimes \mathbf{I} + \mathbf{C} \otimes \mathbf{S})t} dt &= -[\mathbf{T} \otimes \mathbf{I} + \mathbf{C} \otimes \mathbf{S} - \lambda \mathbf{I}]^{-1} \\ &= -[\mathbf{T} \otimes \mathbf{I} + (\mathbf{C} \otimes \mathbf{I})(\mathbf{I} \otimes \mathbf{S}) - \lambda \mathbf{I}]^{-1} \\ &= -[(\mathbf{C} \otimes \mathbf{I})((\mathbf{C} \otimes \mathbf{I})^{-1}(\mathbf{T} \otimes \mathbf{I}) + \mathbf{I} \otimes \mathbf{S} - (\mathbf{C} \otimes \mathbf{I})^{-1} \lambda \mathbf{I})]^{-1}. \end{aligned} \tag{A.91}$$

By Equation (A.87) and since \mathbf{C} is invertible, (A.91) is equal to

$$-[(\mathbf{C} \otimes \mathbf{I})((\mathbf{C}^{-1} \otimes \mathbf{I})(\mathbf{T} \otimes \mathbf{I}) + \mathbf{I} \otimes \mathbf{S} - (\mathbf{C}^{-1} \otimes \mathbf{I}) \lambda \mathbf{I})]^{-1}. \tag{A.92}$$

Using the Mixed Product Rule and algebraic manipulation, (A.92) is equal to

$$\begin{aligned}
& - [(\mathbf{C} \otimes \mathbf{I}) ((\mathbf{C}^{-1}\mathbf{T}) \otimes \mathbf{I} + \mathbf{I} \otimes \mathbf{S} - (\mathbf{C}^{-1}\lambda\mathbf{I}) \otimes \mathbf{I})]^{-1} \\
& = - [(\mathbf{C} \otimes \mathbf{I}) ((\mathbf{C}^{-1}(\mathbf{T} - \lambda\mathbf{I})) \otimes \mathbf{I} + \mathbf{I} \otimes \mathbf{S})]^{-1} \\
& = - [(\mathbf{C}^{-1}(\mathbf{T} - \lambda\mathbf{I})) \otimes \mathbf{I} + \mathbf{I} \otimes \mathbf{S}]^{-1} (\mathbf{C} \otimes \mathbf{I})^{-1} \\
& = - [(\mathbf{C}^{-1}(\mathbf{T} - \lambda\mathbf{I})) \oplus \mathbf{S}]^{-1} (\mathbf{C} \otimes \mathbf{I})^{-1}, \tag{A.93}
\end{aligned}$$

by definition of the Kronecker sum.

Now, for an invertible matrix \mathbf{A} we can write $-\mathbf{A}^{-1} = \int_{x=0}^{\infty} e^{\mathbf{A}x} dx$. Therefore (A.93) is

$$- [(\mathbf{C}^{-1}(\mathbf{T} - \lambda\mathbf{I})) \oplus \mathbf{S}]^{-1} (\mathbf{C} \otimes \mathbf{I})^{-1} = \int_{x=0}^{\infty} e^{(\mathbf{C}^{-1}(\mathbf{T} - \lambda\mathbf{I})x) \oplus \mathbf{S}x} dx (\mathbf{C} \otimes \mathbf{I})^{-1}.$$

Using the rule in Equation (A.88) gives

$$\int_{x=0}^{\infty} e^{(\mathbf{C}^{-1}(\mathbf{T} - \lambda\mathbf{I})x) \otimes e^{\mathbf{S}x}} dx (\mathbf{C} \otimes \mathbf{I})^{-1},$$

which is the result. □

The exponential of a matrix \mathbf{B} is

$$e^{\mathbf{B}} := \sum_{n=0}^{\infty} \frac{1}{n!} \mathbf{B}^n.$$

Latouche & Nguyen (2015) show the following.

Lemma A.23. *Let \mathbf{B} be the block-partitioned matrix*

$$\mathbf{B} = \begin{bmatrix} \mathbf{B}_{11} & \mathbf{B}_{12} \\ \mathbf{B}_{21} & \mathbf{B}_{22} \end{bmatrix}$$

where \mathbf{B}_{11} and \mathbf{B}_{22} are matrices of order m_1 and m_2 , respectively. Denote by $\mathbf{H}_{11}(t)$ the top-left quadrant of order m_1 of $e^{\mathbf{B}t}$:

$$\mathbf{H}_{11}(t) = [\mathbf{I}_{m_1 \times m_1} \quad \mathbf{0}] e^{\mathbf{B}t} \begin{bmatrix} \mathbf{I}_{m_1 \times m_1} \\ \mathbf{0} \end{bmatrix}.$$

The matrix $\mathbf{H}_{11}(t)$ is the solution of

$$\mathbf{H}_{11}(t) = e^{\mathbf{B}_{11}t} + \int_{v=0}^t \int_{u=v}^t e^{\mathbf{B}_{11}(t-u)} \mathbf{B}_{12} e^{\mathbf{B}_{22}(u-v)} \mathbf{B}_{21} \mathbf{H}_{11}(v) du dv. \tag{A.94}$$

Let $\mathbf{H}_{12}(t)$ be the top-right quadrant of $e^{\mathbf{B}t}$ of size $m_1 \times m_2$, i.e.

$$\mathbf{H}_{12}(t) = \begin{bmatrix} \mathbf{I}_{m_1 \times m_1} & \mathbf{0} \end{bmatrix} e^{\mathbf{B}t} \begin{bmatrix} \mathbf{0} \\ \mathbf{I}_{m_2 \times m_2} \end{bmatrix}. \quad (\text{A.95})$$

Denote by $\widehat{\mathbf{H}}_{11}(\lambda) := \int_{t=0}^{\infty} e^{-\lambda t} \mathbf{H}_{11}(t) dt$ and by $\widehat{\mathbf{H}}_{12}(\lambda) := \int_{t=0}^{\infty} e^{-\lambda t} \mathbf{H}_{12}(t) dt$, the Laplace transforms of $\mathbf{H}_{11}(t)$ and $\mathbf{H}_{12}(t)$, respectively. Using Lemma A.23 we can show the following result.

Lemma A.24.

$$\widehat{\mathbf{H}}_{11}(\lambda) = \int_{x=0}^{\infty} e^{(\mathbf{B}_{11} - \lambda \mathbf{I}_{m_1 \times m_1} + \mathbf{B}_{12}(\lambda \mathbf{I}_{m_2 \times m_2} - \mathbf{B}_{22})^{-1} \mathbf{B}_{21})x} dx, \quad (\text{A.96})$$

$$\widehat{\mathbf{H}}_{12}(\lambda) = \int_{x=0}^{\infty} e^{(\mathbf{B}_{11} - \lambda \mathbf{I}_{m_1 \times m_1} + \mathbf{B}_{12}(\lambda \mathbf{I}_{m_2 \times m_2} - \mathbf{B}_{22})^{-1} \mathbf{B}_{21})x} \mathbf{B}_{12}(\lambda \mathbf{I}_{m_2 \times m_2} - \mathbf{B}_{22})^{-1} dx. \quad (\text{A.97})$$

Proof. First we show the result for $\widehat{\mathbf{H}}_{11}(\lambda)$. Taking the Laplace transform of (A.94) shows that $\widehat{\mathbf{H}}_{11}(\lambda)$ is equal to

$$\begin{aligned} & \int_{t=0}^{\infty} \int_{v=0}^t \int_{u=v}^t e^{-\lambda(t-u)} e^{\mathbf{B}_{11}(t-u)} \mathbf{B}_{12} e^{-\lambda(u-v)} e^{\mathbf{B}_{22}(u-v)} \mathbf{B}_{21} e^{-\lambda v} \mathbf{H}_{11}(v) du dv \\ & \quad + (\lambda \mathbf{I}_{m_1 \times m_1} - \mathbf{B}_{11})^{-1} \\ & = (\lambda \mathbf{I}_{m_1 \times m_1} - \mathbf{B}_{11})^{-1} + (\lambda \mathbf{I}_{m_1 \times m_1} - \mathbf{B}_{11})^{-1} \mathbf{B}_{12} (\lambda \mathbf{I}_{m_2 \times m_2} - \mathbf{B}_{22})^{-1} \mathbf{B}_{21} \widehat{\mathbf{H}}_{11}(\lambda), \end{aligned} \quad (\text{A.98})$$

by the convolution theorem for Laplace transforms. This implies

$$\begin{aligned} & [\mathbf{I}_{m_1 \times m_1} - (\lambda \mathbf{I}_{m_1 \times m_1} - \mathbf{B}_{11})^{-1} \mathbf{B}_{12} (\lambda \mathbf{I}_{m_2 \times m_2} - \mathbf{B}_{22})^{-1} \mathbf{B}_{21}] \widehat{\mathbf{H}}_{11}(\lambda) \\ & = (\lambda \mathbf{I}_{m_1 \times m_1} - \mathbf{B}_{11})^{-1}, \end{aligned}$$

and therefore

$$\begin{aligned} & \widehat{\mathbf{H}}_{11}(\lambda) \\ & = [\mathbf{I}_{m_1 \times m_1} - (\lambda \mathbf{I}_{m_1 \times m_1} - \mathbf{B}_{11})^{-1} \mathbf{B}_{12} (\lambda \mathbf{I}_{m_2 \times m_2} - \mathbf{B}_{22})^{-1} \mathbf{B}_{21}]^{-1} (\lambda \mathbf{I}_{m_1 \times m_1} - \mathbf{B}_{11})^{-1} \\ & = [(\lambda \mathbf{I}_{m_1 \times m_1} - \mathbf{B}_{11}) (\mathbf{I}_{m_1 \times m_1} - (\lambda \mathbf{I}_{m_1 \times m_1} - \mathbf{B}_{11})^{-1} \mathbf{B}_{12} (\lambda \mathbf{I}_{m_2 \times m_2} - \mathbf{B}_{22})^{-1} \mathbf{B}_{21})]^{-1} \\ & = [\lambda \mathbf{I}_{m_1 \times m_1} - \mathbf{B}_{11} - \mathbf{B}_{12} (\lambda \mathbf{I}_{m_2 \times m_2} - \mathbf{B}_{22})^{-1} \mathbf{B}_{21}]^{-1} \\ & = \int_{t=0}^{\infty} e^{(\mathbf{B}_{11} - \lambda \mathbf{I}_{m_1 \times m_1} + \mathbf{B}_{12}(\lambda \mathbf{I}_{m_2 \times m_2} - \mathbf{B}_{22})^{-1} \mathbf{B}_{21})t} dt, \end{aligned}$$

which is (A.96).

Now, to show (A.97), differentiate (A.95)

$$\begin{aligned} \frac{d}{dt} \mathbf{H}_{12}(t) &= [\mathbf{I}_{m_1 \times m_1} \quad \mathbf{0}] e^{\mathbf{B}t} \begin{bmatrix} \mathbf{B}_{11} & \mathbf{B}_{12} \\ \mathbf{B}_{21} & \mathbf{B}_{22} \end{bmatrix} \begin{bmatrix} \mathbf{0} \\ \mathbf{I}_{m_2 \times m_2} \end{bmatrix} \\ &= [\mathbf{I}_{m_1 \times m_1} \quad \mathbf{0}] e^{\mathbf{B}t} \begin{bmatrix} \mathbf{B}_{12} \\ \mathbf{B}_{22} \end{bmatrix} \\ &= \mathbf{H}_{11}(t) \mathbf{B}_{12} + \mathbf{H}_{12}(t) \mathbf{B}_{22}. \end{aligned} \quad (\text{A.99})$$

Now take the Laplace transform

$$\lambda \widehat{\mathbf{H}}_{12}(\lambda) - \mathbf{H}_{12}(0) = \widehat{\mathbf{H}}_{11}(\lambda) \mathbf{B}_{12} + \widehat{\mathbf{H}}_{12}(\lambda) \mathbf{B}_{22}. \quad (\text{A.100})$$

Since $\mathbf{H}_{12}(0) = \mathbf{0}$ and after rearranging we get

$$\widehat{\mathbf{H}}_{12}(\lambda) = \widehat{\mathbf{H}}_{11}(\lambda) \mathbf{B}_{12} (\lambda \mathbf{I}_{m_2 \times m_2} - \mathbf{B}_{22})^{-1}, \quad (\text{A.101})$$

which gives (A.97) upon substituting (A.96). \square

Corollary A.25. For $m \in \{+, -\}$ the top-left quadrant of size $m_1 \times m_1 = |\mathcal{S}_m| \cdot p \times |\mathcal{S}_m| \cdot p$ of $e^{\mathbf{B}_{mm}t}$,

$$[\mathbf{I} \quad \mathbf{0}] \int_{t=0}^{\infty} e^{-\lambda t} \exp \left\{ \begin{bmatrix} \mathbf{T}_{mm} \otimes \mathbf{I} + \mathbf{C}_m \otimes \mathbf{S} & \mathbf{T}_{m0} \otimes \mathbf{I} \\ \mathbf{T}_{0m} \otimes \mathbf{I} & \mathbf{T}_{00} \otimes \mathbf{I} \end{bmatrix} t \right\} dt \begin{bmatrix} \mathbf{I} \\ \mathbf{0} \end{bmatrix},$$

is given by

$$\int_{x=0}^{\infty} e^{\mathbf{Q}_{mm}(\lambda)x} \otimes e^{\mathbf{S}x} dx (\mathbf{C}_m^{-1} \otimes \mathbf{I}). \quad (\text{A.102})$$

For $m \in \{+, -\}$ the top-right quadrant of size $m_1 \times m_2 = |\mathcal{S}_m| \cdot p \times |\mathcal{S}_0| \cdot p$ of $e^{\mathbf{B}_{mm}t}$,

$$[\mathbf{I} \quad \mathbf{0}] \int_{t=0}^{\infty} e^{-\lambda t} \exp \left\{ \begin{bmatrix} \mathbf{T}_{mm} \otimes \mathbf{I} + \mathbf{C}_m \otimes \mathbf{S} & \mathbf{T}_{m0} \otimes \mathbf{I} \\ \mathbf{T}_{0m} \otimes \mathbf{I} & \mathbf{T}_{00} \otimes \mathbf{I} \end{bmatrix} t \right\} dt \begin{bmatrix} \mathbf{0} \\ \mathbf{I} \end{bmatrix},$$

is given by

$$\int_{x=0}^{\infty} e^{\mathbf{Q}_{mm}(\lambda)x} \otimes e^{\mathbf{S}x} dx ((\mathbf{C}_m^{-1} \mathbf{T}_{m0} (\lambda \mathbf{I} - \mathbf{T}_{00})^{-1}) \otimes \mathbf{I}). \quad (\text{A.103})$$

Also,

$$[\mathbf{I} \quad \mathbf{0}] \int_{t=0}^{\infty} e^{-\lambda t} e^{\mathbf{B}_{mm}t} dt \mathbf{B}_{mn} = \int_{x=0}^{\infty} \mathbf{H}^{mn}(\lambda, x) \otimes e^{\mathbf{S}x} \mathbf{D} dx ([\mathbf{I}_{pn} \quad \mathbf{0}_{pn \times p|\mathcal{S}_0|}]), \quad (\text{A.104})$$

for $m, n \in \{+, -\}$, $m \neq n$.

Proof. From Lemma A.23 the top-left quadrant of size $m_1 \times m_1 = |\mathcal{S}_m| \cdot p \times |\mathcal{S}_m| \cdot p$ of the integral with respect to t on the left-hand side of (A.102) is

$$\int_{t=0}^{\infty} e^{(\mathbf{T}_{mm} \otimes \mathbf{I} + \mathbf{C}_m \otimes \mathbf{S} - \lambda \mathbf{I} + (\mathbf{T}_{m0} \otimes \mathbf{I})(\lambda \mathbf{I} - \mathbf{T}_{00} \otimes \mathbf{I})^{-1}(\mathbf{T}_{0m} \otimes \mathbf{I}))t} dt. \quad (\text{A.105})$$

By Lemma A.22, (A.105) is equal to

$$\begin{aligned} & \int_{x=0}^{\infty} e^{\mathbf{C}_m^{-1}(\mathbf{T}_{mm} - \lambda \mathbf{I} + \mathbf{T}_{m0}(\lambda \mathbf{I} - \mathbf{T}_{00})^{-1}\mathbf{T}_{0m})x} \otimes e^{\mathbf{S}x} dx (\mathbf{C}_m \otimes \mathbf{I})^{-1} \\ &= \int_{x=0}^{\infty} e^{\mathbf{Q}_{mm}(\lambda)x} \otimes e^{\mathbf{S}x} dx (\mathbf{C}_m \otimes \mathbf{I})^{-1}, \end{aligned} \quad (\text{A.106})$$

from the definition of $\mathbf{Q}_{mm}(\lambda)$. This proves (A.102).

Now, from Lemma A.23 the top-right quadrant of size $m_1 \times m_2 = |\mathcal{S}_m| \cdot p \times |\mathcal{S}_0| \cdot p$ of the integral with respect to t on the left-hand side of (A.103) is

$$\begin{aligned} & \int_{t=0}^{\infty} e^{(\mathbf{T}_{mm} \otimes \mathbf{I} + \mathbf{C}_m \otimes \mathbf{S} - \lambda \mathbf{I} + (\mathbf{T}_{m0} \otimes \mathbf{I})(\lambda \mathbf{I} - \mathbf{T}_{00} \otimes \mathbf{I})^{-1}(\mathbf{T}_{0m} \otimes \mathbf{I}))t} (\mathbf{T}_{m0} \otimes \mathbf{I})(\lambda \mathbf{I} - \mathbf{T}_{00} \otimes \mathbf{I})^{-1} \\ & \quad \times (\mathbf{T}_{0m} \otimes \mathbf{I}) dt. \end{aligned} \quad (\text{A.107})$$

By Lemma A.22, (A.107) is equal to

$$\int_{x=0}^{\infty} e^{\mathbf{Q}_{mm}(\lambda)x} \otimes e^{\mathbf{S}x} dx (\mathbf{C}_m \otimes \mathbf{I})^{-1} (\mathbf{T}_{m0} \otimes \mathbf{I})(\lambda \mathbf{I} - \mathbf{T}_{00} \otimes \mathbf{I})^{-1} (\mathbf{T}_{0m} \otimes \mathbf{I}). \quad (\text{A.108})$$

Now,

$$\begin{aligned} (\lambda \mathbf{I} - \mathbf{T}_{00} \otimes \mathbf{I})^{-1} &= \int_{u=0}^{\infty} e^{-(\lambda \mathbf{I} - \mathbf{T}_{00} \otimes \mathbf{I})u} du \\ &= \int_{u=0}^{\infty} e^{-\lambda u} e^{\mathbf{T}_{00} \otimes \mathbf{I} u} du \\ &= \int_{u=0}^{\infty} e^{-\lambda u} e^{\mathbf{T}_{00} u} \otimes \mathbf{I} du, \end{aligned}$$

by (A.89). Using this and the Mixed Product Rule we can write

$$\begin{aligned} & (\mathbf{C}_m \otimes \mathbf{I})^{-1} (\mathbf{T}_{m0} \otimes \mathbf{I})(\lambda \mathbf{I} - \mathbf{T}_{00} \otimes \mathbf{I})^{-1} (\mathbf{T}_{0m} \otimes \mathbf{I}) \\ &= (\mathbf{C}_m^{-1} \otimes \mathbf{I})(\mathbf{T}_{m0} \otimes \mathbf{I}) \int_{u=0}^{\infty} e^{-\lambda u} e^{\mathbf{T}_{00} u} \otimes \mathbf{I} du (\mathbf{T}_{0m} \otimes \mathbf{I}) \\ &= (\mathbf{C}_m^{-1} \mathbf{T}_{m0} (\lambda \mathbf{I} - \mathbf{T}_{00})^{-1} \mathbf{T}_{0m}) \otimes \mathbf{I}. \end{aligned} \quad (\text{A.109})$$

Substituting (A.109) into (A.108) completes the proof of (A.103).

Now, using (A.102) and (A.103) we can write

$$\begin{aligned}
[\mathbf{I} \quad \mathbf{0}] \int_{t=0}^{\infty} e^{-\lambda t} e^{\mathbf{B}_{mm}t} dt \mathbf{B}_{mn} &= [\mathbf{I} \quad \mathbf{0}] \int_{t=0}^{\infty} e^{-\lambda t} e^{\mathbf{B}_{mm}t} dt \begin{bmatrix} \mathbf{T}_{mn} \otimes \mathbf{D} & \mathbf{0} \\ \mathbf{T}_{0n} \otimes \mathbf{D} & \mathbf{0} \end{bmatrix} \\
&= \int_{x=0}^{\infty} e^{\mathbf{Q}_{mm}(\lambda)x} \otimes e^{\mathbf{S}x} dx (\mathbf{C}_m^{-1}(\mathbf{T}_{mn} + \mathbf{T}_{m0}(\lambda\mathbf{I} - \mathbf{T}_{00})^{-1}\mathbf{T}_{0n}) [\mathbf{I}_n \quad \mathbf{0}_{n \times |S_0|}] \otimes \mathbf{D}) \\
&= \int_{x=0}^{\infty} e^{\mathbf{Q}_{mm}(\lambda)x} \otimes e^{\mathbf{S}x} dx ((\mathbf{Q}_{mn}(\lambda) [\mathbf{I}_n \quad \mathbf{0}_{n \times |S_0|}]) \otimes \mathbf{D}) \\
&= \int_{x=0}^{\infty} (\mathbf{H}^{mn}(\lambda, x) [\mathbf{I}_n \quad \mathbf{0}_{n \times |S_0|}]) \otimes e^{\mathbf{S}x} \mathbf{D} dx, \\
&= \int_{x=0}^{\infty} \mathbf{H}^{mn}(\lambda, x) \otimes e^{\mathbf{S}x} \mathbf{D} dx, [\mathbf{I}_{pn} \quad \mathbf{0}_{pn \times p|S_0|}] \tag{A.110}
\end{aligned}$$

for $m, n \in \{+, -\}$, $m \neq n$ which is (A.104), where the last line holds from the Mixed Product Rule. \square

Appendix B

second appendix

Bibliography

- Ahn, S., Jeon, J. & Ramaswami, V. (20035), ‘Steady state analysis of finite fluid flow models using finite QBDs’, *Queueing Syst.* **49**, 223–259.
- Ahn, S. & Ramaswami, V. (2003), ‘Fluid flow models and queues—a connection by stochastic coupling’, *Stochastic Models* **19**(3), 325–348.
URL: <https://doi.org/10.1081/STM-120023564>
- Ahn, S. & Ramaswami, V. (2004), ‘Transient analysis of fluid flow models via stochastic coupling to a queue’, *Stochastic Models* **20**(1), 71–101.
URL: <https://doi.org/10.1081/STM-120028392>
- Aldous, D. & Shepp, L. (1987), ‘The least variable phase type distribution is Erlang’, *Communications in Statistics. Stochastic Models* **3**(3), 467–473.
URL: <https://doi.org/10.1080/15326348708807067>
- Anick, D., Mitra, D. & Sondhi, M. (1982), ‘Stochastic theory of a data-handling system with multiple sources’, *The Bell System Technical Journal* **61**(8), 1871–1894.
- Asmussen, S. (2008), *Applied probability and queues*, Vol. 51, Springer Science & Business Media.
- Badescu, A., Breuer, L., Soares, A. D. S., Latouche, G., Remiche, M.-A. & Stanford, D. (2005), ‘Risk processes analyzed as fluid queues’, *Scandinavian Actuarial Journal* **2005**(2), 127–141.
- Bean, N. G. & Nielsen, B. F. (2010), ‘Quasi-birth-and-death processes with rational arrival process components’, *Stochastic Models* **26**(3), 309–334.
- Bean, N. G. & O’Reilly, M. M. (2013), ‘Spatially-coherent uniformization of a stochastic fluid model to a quasi-birth-and-death process’, *Performance Evaluation* **70**(9), 578 – 592.
- Bean, N. G. & O’Reilly, M. M. (2014), ‘The stochastic fluid-fluid model: A stochastic fluid model driven by an uncountable-state process, which is a stochastic fluid itself’, *Stochastic Processes and their Applications* **124**, 1741–1772.

- Bean, N. G., O'Reilly, M. M. & Sargison, J. (2010), 'A stochastic fluid flow model of the operation and maintenance of power generation systems', *IEEE Transactions on Power Systems* **25**(3), 1361–1374.
- Bean, N. G., O'Reilly, M. M. & Taylor, P. G. (2005a), 'Algorithms for return probabilities for stochastic fluid flows', *Stochastic Models* **21**(1), 149–184.
URL: <https://doi.org/10.1081/STM-200046511>
- Bean, N. G., O'Reilly, M. M. & Taylor, P. G. (2005b), 'Hitting probabilities and hitting times for stochastic fluid flows', *Stochastic Processes and their Applications* **115**(9), 1530–1556.
- Bean, N. G., O'Reilly, M. M. & Taylor, P. G. (2009a), 'Algorithms for the Laplace-Stieltjes transforms of first return times for stochastic fluid flows', *Methodology and Computing in Applied Probability* **10**, 381–408.
- Bean, N. G., O'Reilly, M. M. & Taylor, P. G. (2009b), 'Hitting probabilities and hitting times for stochastic fluid flows: the bounded model', *Probability in the Engineering and Informational Sciences* **23**(1), 121–147.
- Bean, N., Lewis, A., Nguyen, G. T., O'Reilly, M. M. & Sunkara, V. (2021), 'A discontinuous galerkin method for approximating the stationary distribution of stochastic fluid-fluid processes', *arXiv preprint arXiv:1901.10635*.
- Bladt, M. & Nielsen, B. (2017), *Matrix-Exponential Distributions in Applied Probability*, Springer.
- da Silva Soares, A. (2005), Fluid queues: building upon the analogy with QBD processes, PhD thesis, Université Libre de Bruxelles.
- Élteto, T., Rácz, S. & Telek, M. (2006), Minimal coefficient of variation of matrix exponential distributions, in '2nd Madrid Conference on Queueing Theory'.
- Feller, W. (1957), *An introduction to probability theory and its applications*, Wiley series in probability and mathematical statistics, 2nd ed. edn, Wiley, New York.
- Hautphenne, S., Massaro, M. & Taylor, P. (2017), 'How old is this bird? The age distribution under some phase sampling schemes', *Journal of mathematical biology* **75**(6), 1319–1347.
- Hesthaven, J. S. & Warburton, T. (2007), *Nodal discontinuous Galerkin methods: algorithms, analysis, and applications*, Springer Science & Business Media.
- Horváth, I., Sáfár, O., Telek, M. & Zambó, B. (2016), Concentrated matrix exponential distributions, in 'In proceedings of European Performance Evaluation Workshop'.

- Horváth, G., Horváth, I. & Telek, M. (2020), ‘High order concentrated matrix-exponential distributions’, *Stochastic Models* **36**(2), 176–192.
- Karandikar, R. L. & Kulkarni, V. G. (1995), ‘Second-order fluid flow models: Reflected Brownian motion in a random environment’, *Operations Research* **43**(1), 77–88.
- Koltai, P. (2011), *Efficient approximation methods for the global long-term behavior of dynamical systems: theory, algorithms and examples*, Logos Verlag Berlin GmbH.
- Latouche, G. & Nguyen, G. T. (2015), ‘The morphing of fluid queues into Markov-modulated Brownian motion’, *Stochastic Systems* **5**(1), 62–86.
- Latouche, G. & Nguyen, G. T. (2019), ‘Analysis of fluid flow models’, *arXiv preprint arXiv:1901.10635*.
- Mészáros, A. & Telek, M. (2021), ‘Concentrated matrix exponential distributions with real eigenvalues’, *Probability in the Engineering and Informational Sciences* p. 1–17.
- Wurm, M. (2020), Stochastic modelling of coral-algal symbiosis on the Great Barrier Reef, Master’s thesis, The University of Adelaide.