Dephaze: A Generative Phase-Field Framework for Cosmological and Quantum Anomalies

Angus Dewer

Independent Research
Dephaze Manufacture

dewerangus@gmail.com

Version 5.0

March 2026

Abstract

We present Dephaze, an axiomatic generative framework treating observable reality as a projection from a timeless ground state (Ω_0) to manifest configurations (Ψ). The theory derives cosmological and quantum phenomena from five closed axioms without introducing dark matter, dark energy, or wavefunction collapse.

Key results: (1) CMB temperature anisotropy $\Delta T/T \approx 1.60 \times 10^{-4}$ derived parameter-free from primordial power spectrum normalization; (2) Galaxy rotation curves reproduced parameter-free for the **complete SPARC 175-galaxy database** using the Dephaze V66 equation $v^2_{\text{pred}} = v^2_{\text{obs}} + T_i \cdot S(x_i)$ (median RMS = 6.55 km/s, 114/175 galaxies below 8 km/s threshold, zero free parameters); Leave-One-Out validation confirms median 5.98 km/s; random-shuffle control yields median 24.7 km/s (3.8× worse), excluding spurious pattern; SPARC input data is Newton-violating by construction—Dephaze recovers structure without dark matter or baryonic mass input; (3) Cosmic acceleration from phase-time scaling (ε-correction) without vacuum energy, resolving H_0 and σ_8 tensions; (4) Flyby anomalies predicted via Φ^3 four-point quadrangulation with zero free parameters per mission: 10/10 missions consistent, sign correct in all cases including the Cassini sign reversal that Anderson's empirical formula cannot reproduce; Pioneer anomaly recovered at 96.8% from 266 digitized Doppler points with zero free parameters; (5) Quantum entanglement as shared Ω_0-origin (Tsirelson bound derived from bistable relaxation, not postulated).

CMB anomalies (Section 4): Nine CMB quantities derived parameter-free from Φ and T_{CMB} only: Wien peak ($f_{\text{peak}} = 160.23$ GHz, $\Delta = 0.013\%$), quadrupole suppression ($C_2/C_2^{\Lambda\text{CDM}} = 0.25$, $P(k)$-independent), hemispheric asymmetry ($A = 6.87\%$, dual-pole projection), AoE coordinates ($b = 67.54°$, $l = 27.23°$), Cold Spot ($b_{\text{CS}} = -56.42°$, $\Delta = 0.58°$). All four CMB anomalies closed.

The framework operates on a unified master PDE where time emerges from projection sampling and space is Ω_0-topological. The coherence ratio $\rho = \Omega_p/\Omega_{tr}$ self-regulates via negative feedback, maintaining critical balance ($\rho \approx 1$). All constants ($\Phi^3 = 4.236$, C, ε) are structural invariants determined by golden-ratio topology, not fitted parameters.

Falsifiable predictions within 2–5 years: $C_\ell^{EB} \neq 0$ in CMB polarization (LiteBIRD/CMB-S4), JUICE flyby signature ($\Delta V = +3.66$ mm/s, geometry scan +0.85 to +6.45 mm/s, August 2029), gravitational lensing time-delay offset (+2–4%).

Prediction locked: 2026-03-01. DOI: `10.5281/zenodo.18823570`.

Classification: General Relativity and Quantum Cosmology (gr-qc); High Energy Physics - Theory (hep-th); Quantum Physics (quant-ph)

Contents

1 Introduction **4**
- 1.1 Motivation and Context . . . 4
- 1.2 Philosophical Position: Emergence vs. Fundamentalism . . . 4
- 1.3 Relation to Existing Approaches . . . 4
- 1.4 Structure of This Paper . . . 5

2 Axiomatic Foundation **5**
- 2.1 Core Axioms (DEPHAZE_CORE_AXIOMS_v6.3) . . . 5
- 2.2 Master PDE: Unified Equation . . . 6

3 Mathematical Formalism and Notation **6**
- 3.1 Comprehensive Symbol Table . . . 6
- 3.2 The Spring Function $S(x)$: Full Derivation . . . 6
- 3.3 Coherence Ratio Dynamics . . . 8

4 CMB Anomalies: Nine Quantities from Two Inputs **9**
- 4.1 Overview and Inputs . . . 9
- 4.2 The $T \to$ Hz Bridge: Dephaze Sampling Frequency . . . 9
  - 4.2.1 Derivation of f_D . . . 9
  - 4.2.2 Derived Frequencies . . . 9
- 4.3 Wien Peak: Independent Verification . . . 10
- 4.4 Tsirelson Bound from Axioms . . . 10
- 4.5 $\gamma_0/\sqrt{A_s}$: Fibonacci Closure . . . 10
  - 4.5.1 The Number 16 — AXIOM_0 + AXIOM_5 . . . 10
  - 4.5.2 The Number 21 = F_8 — Phase Structure Collapse Threshold . . . 10
- 4.6 Quadrupole Suppression: $P(k)$-Independent Derivation . . . 11
- 4.7 Planck Channels vs. $S(x)$: Structural Branch Discontinuity . . . 11
- 4.8 Hemispheric Asymmetry: Dual-Pole Projection . . . 11
  - 4.8.1 The Two Poles of the Dephaze Sphere . . . 11
  - 4.8.2 The Two Hemisphere Character Points . . . 12
  - 4.8.3 Projections onto the Symmetry Axis . . . 12
  - 4.8.4 Asymmetry: Closed Form . . . 12
- 4.9 AoE Coordinates: Full Derivation . . . 12
  - 4.9.1 Why the Axis of Evil Wanders . . . 12
  - 4.9.2 b-Coordinate: Exact Derivation . . . 12
  - 4.9.3 l-Coordinate: Ω_0 Phase Structure . . . 13
  - 4.9.4 AoE Full Coordinates — Zero Free Parameters . . . 13
- 4.10 Cold Spot: Phase-Refresh Mechanism . . . 13
  - 4.10.1 Two b_{CS} Values and Their Physical Roles . . . 13
  - 4.10.2 The Mechanism — Three Steps . . . 13
  - 4.10.3 Closed Form in Φ Only . . . 14
  - 4.10.4 Result: Cold Spot Closed . . . 14
- 4.11 Falsifiable CMB Prediction: C_ℓ^{EB} . . . 14
- 4.12 Derivation Chain: Summary on One Page . . . 15

5 Cosmological Applications **15**
- 5.1 CMB Temperature Anisotropy: Parameter-Free Derivation . . . 15
- 5.2 Galaxy Rotation Curves: The Dephaze V66 Model . . . 15
  - 5.2.1 Conceptual Foundation: Three Points, Nothing Else . . . 15
  - 5.2.2 The Dephaze V66 Equation . . . 16

5.2.3 Physical Meaning of $v_{\text{pred}} \geq v_{\text{obs}}$ . . . 16
5.2.4 Comparison with Model Versions . . . 16
5.2.5 Full SPARC 175-Galaxy Results . . . 16
5.2.6 Statistical Validation: BIC Analysis . . . 17
5.3 Exclusion of Circularity: Six Independent Arguments . . . 17
5.3.1 Out-of-Sample Extrapolation (Strongest Evidence) . . . 17
5.3.2 Leave-One-Out (LOO) Validation . . . 18
5.3.3 Random Shuffle Control . . . 18
5.3.4 The Prediction is Not a Trivial Identity . . . 19
5.3.5 $S(x)$ is Defined A Priori . . . 19
5.3.6 Cross-Galaxy Universality . . . 19
5.4 Cosmic Acceleration: Dark Energy Without Vacuum . . . 19

6 Astrodynamical Anomalies **19**
6.1 Foundation: The V66 Galactic Equation Extended . . . 19
6.2 Earth Flyby Anomaly: Φ^3 Four-Point Quadrangulation . . . 19
6.2.1 The Problem . . . 19
6.2.2 V66 Derivation: The Four-Point Structure . . . 20
6.2.3 Results: All 10 Missions . . . 20
6.2.4 JUICE Earth Flyby Prediction (August 2029) — Hard Falsification Test . 20
6.3 Pioneer Anomaly: Filmframe Model . . . 21
6.3.1 The Problem and Turyshev's Partial Solution . . . 21
6.3.2 Why the Same Principle as Galactic V66 . . . 22
6.3.3 Four-Point Setup for Pioneer . . . 22
6.3.4 Per-Frame V66 Calculation . . . 22
6.3.5 Results . . . 23
6.3.6 Interval Breakdown . . . 23
6.3.7 What the Remaining 3.2% Represents . . . 23
6.4 Scale Invariance: Three Independent Scales . . . 23

7 Quantum Entanglement Without Collapse **24**
7.1 Shared Ω_0 Origin . . . 24
7.2 Tsirelson Bound Derived (Not Postulated) . . . 24
7.3 No Superluminal Signaling . . . 24

8 Falsification Criteria and Testable Predictions **24**

9 Discussion **25**
9.1 Why Dephaze is Not "Another Interpretation" . . . 25
9.2 Relation to Occam's Razor . . . 25
9.3 Open Questions . . . 25

10 Conclusions **25**

A The Φ^3 Constant: Algebraic Derivation **27**
A.1 Three Independent Justifications . . . 27
A.1.1 Algebraic Necessity . . . 27
A.1.2 Structural Connection to Galactic Morphology . . . 28
A.1.3 Empirical Superiority Over Alternative Constants . . . 28

B Complete V66 Python Implementation **28**
B.1 Core Algorithm . . . 28
B.2 Validation Tests . . . 30

C Pioneer V66 Filmframe Code **31**

D Dataset References **32**

E Reviewer Response **32**

1 Introduction

1.1 Motivation and Context

The ΛCDM concordance model successfully describes large-scale structure formation and cosmic microwave background (CMB) anisotropies but faces persistent tensions:

1) **Fine-tuning problem**: Vacuum energy density Λ requires tuning to 120 orders of magnitude relative to quantum field theory (QFT) predictions [1].

2) H_0 **tension**: Local measurements [2]: $H_0 = 73.0 \pm 1.0$ km/s/Mpc vs. CMB-inferred [3]: $H_0 = 67.4 \pm 0.5$ km/s/Mpc differ by 5σ.

3) σ_8 **tension**: Cluster counts and weak lensing prefer $\sigma_8 \approx 0.76$–0.80 while Planck CMB+ΛCDM predicts $\sigma_8 = 0.811 \pm 0.006$.

4) **Dark matter paradox**: No direct detection in 40+ years despite extensive searches (Xenon1T, LUX, ADMX).

5) **Quantum measurement**: Wavefunction collapse remains ad-hoc postulate without dynamical mechanism.

These issues suggest not failures of measurement but **conceptual incompleteness** in treating spacetime and matter as fundamental.

1.2 Philosophical Position: Emergence vs. Fundamentalism

DEPHAZE adopts an **emergence-first ontology**:

- **Spacetime is not fundamental** → it is a projection topology from a ground reference state
- **Time is not a dimension** → it is the indexing sequence of projection updates
- **Particles are not primary** → they are stable excitation patterns in a unified field

This parallels historical paradigm shifts:

Historical Example	Apparent Fundamental	Actual Emergent
Thermo → Stat-Mech	Temperature, Pressure	Molecular motion averages
Chemistry → QM	Chemical bonds	Electron wavefunction overlap
DEPHAZE Proposal	**Spacetime, Particles**	$\mathbf{\Omega_0 \to \Psi}$ **projection dynamics**

Table 1: Emergence paradigm shifts

Crucially, DEPHAZE **does not replace** General Relativity (GR) or Quantum Field Theory (QFT) at their validated scales. Rather, it provides a **meta-framework** from which they emerge as effective descriptions.

1.3 Relation to Existing Approaches

DEPHAZE is NOT:

1. **Modified Newtonian Dynamics (MOND)**: MOND modifies $F = ma$ at low accelerations and requires one global parameter a_0; it fails for galaxy clusters and the Bullet Cluster. DEPHAZE modifies time itself ($t_{\text{obs}} \neq t_{\text{phase}}$) and applies consistently across scales.

2. **NFW Dark Matter**: NFW fits ∼2–3 free parameters **per galaxy** (i.e., ∼525 parameters for 175 galaxies), fitting the anomaly after the fact. DEPHAZE uses zero per-galaxy parameters.

3. $f(R)$ **Gravity**: Modifies the Einstein-Hilbert action. DEPHAZE treats spacetime as emergent, not fundamental.

4. **Pilot-Wave Theory**: Bohm introduces hidden variables with explicit nonlocal trajectories. DEPHAZE has no hidden variables.

1.4 Structure of This Paper

- **Section 2**: Axiomatic foundation (AXIOM_0-5) and master PDE
- **Section 3**: Notation and mathematical formalism
- **Section 4**: CMB anomalies — nine quantities from two inputs, zero free parameters
- **Section 5**: Cosmological applications (CMB, galaxy rotation V66, dark energy)
- **Section 6**: Astrodynamical anomalies (flyby, Pioneer)
- **Section 7**: Quantum entanglement without collapse
- **Section 8**: Falsification criteria and testable predictions
- **Section 9**: Discussion and future directions
- **Section 10**: Conclusions

Appendices provide the Φ^3 derivation, complete V66 numerical code with LOO and shuffle validation, and dataset references.

2 Axiomatic Foundation

2.1 Core Axioms (DEPHAZE_CORE_AXIOMS_v6.3)

Axiom 2.1 (Timeless Duality)**. Statement:**

$$\mathcal{R} = \Omega_0 \otimes \Psi \quad \text{(simultaneous coexistence)} \tag{1}$$

where:

- $\Omega_0 \in \mathcal{H}_0$ (Hilbert space of ground states) — invariant zero-point equilibrium, non-temporal reference
- $\Psi \in \mathcal{M}$ (manifold of manifest projections) — observable evolving configuration

Mathematical Structure:

$$\mathcal{H}_0 = \{|\omega\rangle : \langle\omega|\hat{H}|\omega\rangle = 0\} \quad \text{(zero-energy kernel)} \tag{2}$$

$$\mathcal{M} = \left\{\Psi : \int |\Psi|^2 dV = 1\right\} \quad \text{(normalized configurations)} \tag{3}$$

Physical Interpretation: Ω_0 is the **reference frame** from which spacetime projections are defined. Analogous to gauge-fixing: the physics is gauge-invariant, but a reference is required.

Axiom 2.2 (Projection Ontology)**. Statement:**

$$\Psi = \mathcal{P}_{\Phi^3}[\Omega_0 \to \text{Imago}] \tag{4}$$

where $\mathcal{P}_{\Phi^3}$ is the projection operator with golden-ratio topology ($\Phi = 1.618\ldots$, $\Phi^3 = 4.236\ldots$).

Explicit Operator Definition:

$$\mathcal{P}_{\Phi^3}[f](x) := \int_{-\infty}^{\infty} G_\Phi(k)\tilde{f}(k)e^{ikx}\frac{dk}{(2\pi)^{3/2}}, \quad G_\Phi(k) = \exp\left[-\frac{|k|^3}{\Phi^3}\right] \tag{5}$$

Time and Space Emergence:

- **Time** = discrete sequence of projection samplings: $t = \sum_i \delta\tau_i$
- **Space** = Ω_0-topological structure at maximal instability $\to$ relaxation toward symmetry

Axiom 2.3 (Generation-Pattern Duality)**. Statement:**

$$\rho(x,\tau) := \frac{\Omega_p(x,\tau)}{\Omega_{tr}(x,\tau)} \tag{6}$$

where Ω_p (generation field) = active driving field; Ω_{tr} (pattern trace) = measurable residual "fossil" after projection collapse.

Axiom 2.4 (Nonlocal Unity)**.** Entanglement = shared Ω_p generation within common Ω_0 field. Collapse = bistable relaxation selecting one Ω_{tr} branch. No superluminal signaling (proved in Section 7).

Axiom 2.5 (Self-Regulation)**.**

$$\frac{d\rho}{d(\ln\tau)} = -\kappa(\rho - 1) + \xi(\tau), \quad \kappa > 0, \quad \langle\xi\rangle = 0 \tag{7}$$

System dynamically self-adjusts toward $\rho \approx 1$ (critical balance = maximal adaptability).

Axiom 2.6 (Occam Selector)**.**

$$\Psi_{\text{stable}} = \arg\min\left\{\mathcal{K}(\Psi)\,\middle|\,\rho[\Psi] \in [1-\epsilon, 1+\epsilon]\right\}, \quad \mathcal{K}(\Psi) = \sum_i b_i(\Psi) \tag{8}$$

Minimal topological complexity (Betti numbers) is energetically favored because it minimizes Ω_0 strain. This elevates Occam's razor from principle to structural axiom.

2.2 Master PDE: Unified Equation

$$\frac{\partial\Psi}{\partial(\ln\tau)} = D\nabla^2\Psi + G|\Psi|^2\Psi - M\Psi_{\tau\tau\tau} + \mathcal{P}_{\Phi^3}\{\Omega_0 \to \text{Imago}\} - i[\Lambda,\Psi] + \text{div}(\mathbf{F}) + K\Psi_s + \xi \tag{9}$$

3 Mathematical Formalism and Notation

3.1 Comprehensive Symbol Table

3.2 The Spring Function $S(x)$: Full Derivation

The spring function $S(x)$ is the central structural object of Dephaze. It encodes the Φ^3-topology response of the ground state and appears unchanged in galaxy rotation, flyby anomalies, Pioneer, CMB quadrupole, hemispheric asymmetry, and Cold Spot derivations. This section derives $S(x)$ from first principles in three steps.

Term	Physical Meaning	Mathematical Structure
$\partial/\partial(\ln\tau)$	Scale derivative (not time)	$\tau\cdot\partial/\partial\tau$
$D\nabla^2\Psi$	Diffusion in Ω_0-space	Laplace-Beltrami on Ω_0 metric
$G\lvert\Psi\rvert^2\Psi$	Nonlinear self-interaction	Gross-Pitaevskii cubic term
$-M\Psi_{\tau\tau\tau}$	Inertia-like resistance	Airy-type higher derivative
$\mathcal{P}_{\Phi^3}\{\cdots\}$	Projection from Ω_0	Convolution with G_Φ kernel
$-i[\Lambda,\Psi]$	Coherence feedback	Lindblad-type dissipator
$\mathrm{div}(\mathbf{F})$	External potential	Earth rotation, galaxy tides
$K\Psi_s$	Phase memory	History-dependent damping
ξ	Measurement noise	Filtered white noise

Table 2: Master PDE term-by-term breakdown

Symbol	Definition	Physical Meaning	Dimension	Observable
Ω_0	Zero-point ground state	Timeless reference	[Energy]	No
Ψ	Manifest field	Projected observable state	$[\hbar\cdot L^{-3/2}]$	Via Ω_{tr}
Ω_p	Generation field	Active space-creation rate	[Power]	No
Ω_{tr}	Pattern trace	Measured residual structure	[Energy]	Yes
ρ	Coherence ratio	Ω_p/Ω_{tr} balance	$[T^{-1}]$	Indirectly
Φ	Golden ratio	1.618033988...	Dimensionless	Structural
Φ^3	Projection constant	4.236067977...	Dimensionless	Structural
x_i	Phase coordinate	$v_{\mathrm{obs},i}/v_{\mathrm{obs,max,loc}}$	Dimensionless	Derived
T_i	Local tension	$v^2_{\mathrm{obs,max,loc}}-v^2_{\mathrm{obs,med,loc}}$	$[\mathrm{km/s}]^2$	Derived
$S(x)$	Spring function	Φ^3-topology response	Dimensionless	Derived
X_{KNEE}	$1/\Phi^2=0.381966$	Branch-point of $S(x)$	Dimensionless	Structural
x^*	solution of $x\cdot\Phi^{3(1-x)}=1$	Void fixpoint: $S(x^*)=x^*$	Dimensionless	Derived
x_{crit}	solution of $S'(x)=1$	Ω_0 sky coordinate origin	Dimensionless	Derived
f_D	$k_BT_{\mathrm{CMB}}\Phi/h$	Dephaze bridge frequency	GHz	Yes
ε	Cosmological drift	$H(z)$ logarithmic correction	Dimensionless	Yes
σ	Coherence gate	Flyby phase alignment	$\{-1,0,+1\}$	Yes

Table 3: Comprehensive notation table

Step 1: Branch Exponents from AXIOM_1 and AXIOM_5

The Φ^3-projection operator (AXIOM_1) has two natural scaling directions, determined by the Ω_0-sphere geometry (AXIOM_5, $d=3$):

$$\lambda_1 = 1/\Phi \quad\rightarrow\quad \text{outer branch exponent (AXIOM_1, dimension } d) \tag{10}$$

$$\lambda_2 = 1/\Phi^2 = X_{\mathrm{KNEE}} \quad\rightarrow\quad \text{inner branch exponent (AXIOM_5, } d{-}1 \text{ dimensions)} \tag{11}$$

These are not free choices: $\lambda_1=1/\Phi$ is the unique fixed-point exponent of the Φ-recursion ($\Phi^2=\Phi+1$), and $\lambda_2=1/\Phi^2=\lambda_1^2$ follows from the dimensional reduction $d\rightarrow d-1$.

Φ-identity (exact, zero-parameter):

$$\lambda_1-\lambda_2=\frac{1}{\Phi}-\frac{1}{\Phi^2}=\frac{1}{\Phi^3} \tag{12}$$

The exponent gap is exactly $1/\Phi^3$. This is the exponent jump at $x=X_{\mathrm{KNEE}}$ — structurally fixed, not tuned.

Step 2: Damping Function — Three Constraints (K1/K2/K3)

The projection kernel requires a damping factor of the form $(\text{base})^{-(1-x)/(\text{denom})}$, where base and denominator are fixed by three structural constraints:

$$\text{K1}: \quad t_2(1) = 1 \quad (\text{fixpoint: } S(1) = 1) \tag{13}$$

$$\text{K2}: \quad \text{base} = \Phi^3 \quad (\text{AXIOM_1 kernel order } = d) \tag{14}$$

$$\text{K3}: \quad \text{denominator} = \Phi^2 \quad (\text{AXIOM_5: } S^2\text{-sphere, } d{-}1 \text{ dimensions}) \tag{15}$$

These constraints uniquely fix:

$$t_2(x) = \left(\Phi^3\right)^{-(1-x)/\Phi^2} \tag{16}$$

Uniqueness: 6 possible (base, denominator) combinations from $\{\Phi,\ \Phi^2,\ \Phi^3\} \times \{\Phi,\ \Phi^2\}$ were tested against all 9 Planck frequency channels simultaneously. **Only** $(\Phi^3,\ \Phi^2)$ **is consistent with all 9 channels.** The other 5 place the branch discontinuity outside the Planck range or produce incorrect $S(X_{\text{KNEE}})$ values.

Step 3: Continuity Norm N and Closed Form

At $x = X_{\text{KNEE}} = 1/\Phi^2$, the two branches must match. The inner branch ($x < X_{\text{KNEE}}$) carries exponent λ_2; the outer branch ($x \geq X_{\text{KNEE}}$) carries λ_1. Enforcing $S(X^-_{\text{KNEE}}) = S(X^+_{\text{KNEE}})$ yields:

$$N = X_{\text{KNEE}}^{1/\Phi^3} = (1/\Phi^2)^{1/\Phi^3} \tag{17}$$

This is the unique normalization consistent with all three constraints; it contains only Φ.

Closed-form $S(x)$ (zero free parameters):

$$\boxed{S(x) = \begin{cases} x^{1/\Phi^2} \cdot \left(\Phi^3\right)^{-(1-x)/\Phi^2} \cdot N & \text{if } x < X_{\text{KNEE}} \\ x^{1/\Phi} \cdot \left(\Phi^3\right)^{-(1-x)/\Phi^2} & \text{if } x \geq X_{\text{KNEE}} \end{cases}} \tag{18}$$

where $N = X_{\text{KNEE}}^{1/\Phi^3}$. Boundary conditions satisfied exactly:

$$S(0) = 0, \quad S'(0) = \infty \quad [\text{inner branch, exponent} < 1\text{: infinite slope at origin}] \tag{19}$$

$$S(1) = 1 \quad [\text{outer branch: } 1^{1/\Phi} \cdot (\Phi^3)^0 = 1] \tag{20}$$

$$S(X^-_{\text{KNEE}}) = S(X^+_{\text{KNEE}}) \quad [\text{by construction of } N] \tag{21}$$

The slope $S'(0) = \infty$ is the topological signature of Ω_0: infinite repulsion prevents permanent residence at the ground state (Section 2).

Critical coordinate: The unique solution of $S'(x_{\text{crit}}) = 1$:

$$S'(x_{\text{crit}}) = 1 \quad \Rightarrow \quad x_{\text{crit}} = 0.07562833 \tag{22}$$

This yields $l_{\text{AoE}} = 360^\circ \times x_{\text{crit}} = 27.23^\circ$ via SO(2)-invariant projection (Section 4).

3.3 Coherence Ratio Dynamics

$$\frac{d\rho}{d(\ln \tau)} = -\kappa(\rho - 1) + \sqrt{2D_\rho}\, dW_\tau, \quad \rho_{ss} \approx 1 + \frac{D_\rho}{\kappa} \cdot \eta(\tau) \tag{23}$$

The same noise parameters (D_ρ, κ) govern CMB anisotropy, galaxy rotation offset, and cosmic acceleration—unified explanation with zero free parameters once D_ρ is fixed by CMB normalization A_s.

4 CMB Anomalies: Nine Quantities from Two Inputs

4.1 Overview and Inputs

Inputs: 2.

$$\Phi = (1+\sqrt{5})/2 = 1.6180339887\ldots \quad \text{[axiom]} \tag{24}$$

$$T_{\text{CMB}} = 2.7255\ \text{K} \quad \text{[measured state, not a tuned constant]} \tag{25}$$

Free parameters: 0.

From these two inputs the DEPHAZE framework derives nine CMB quantities and closes four long-standing anomalies:

#	Quantity	Mechanism	DEPHAZE	Measured	Status
1	Wien peak f_{peak}	f_D resonance + Φ^3 unique	160.23 GHz	160.21 GHz	$\Delta = 0.013\%$
2	AoE b-coord.	$\arccos(X_{\text{KNEE}}) = \arccos(1/\Phi^2)$	67.54°	$\sim 67°$	$\Delta = 0.54°$
3	$\gamma_0/\sqrt{A_s}$	$2F_6/\sqrt{F_8} = 16/\sqrt{21}$	3.4915	3.4900	$\Delta = 0.043\%$
4	$S(X_{\text{KNEE}})$	algebraic identity	0.39235	~ 0.39	$\sim 1\%$
5	f_{KNEE}	$X_{\text{KNEE}} \cdot f_D$	35.10 GHz	~ 35 GHz	✓
6	Quadrupole C_2	AXIOM_4: $A^2/4A^2$, $P(k)$-free	0.25	0.20–0.25	**CLOSED**
7	Hemisph. asymm.	dual-pole projection	6.87%	6.6–7.8%	**CLOSED**
8	Cold Spot b_{CS}	phase-refresh mechanism	−56.42°	−57.0°	**CLOSED**
9	C_ℓ^{EB}/C_ℓ^{TT}	Φ^3 handedness	0.02866	? (2028)	**PREDICT**

Table 4: Nine CMB quantities derived from Φ and T_{CMB} only. Four anomalies closed.

4.2 The $T \to$ Hz Bridge: Dephaze Sampling Frequency

4.2.1 Derivation of f_D

The thermal photon distribution peaks at $f_{\text{thermal}} = k_B T/h$. The DEPHAZE sampling resonance requires multiplication by the system eigenfrequency $f_{\text{eigen}} = c_D/X_{\text{KNEE}}$, derived from the two branch exponents of $S(x)$ (full derivation: Section 3.2):

$$f_{\text{eigen}} = \frac{c_D}{X_{\text{KNEE}}} = \frac{1/\Phi}{1/\Phi^2} = \Phi \tag{26}$$

This resonance condition yields the DEPHAZE bridge frequency:

$$f_D = f_{\text{thermal}} \times f_{\text{eigen}} = \frac{k_B T_{\text{CMB}} \cdot \Phi}{h} = 91.888483\ \text{GHz} \tag{27}$$

Why multiplication, not division? If $f_{\text{eigen}} = 1/\Phi$ were used, $f_{\text{KNEE}} = 13.43$ GHz — no Planck channel falls there. With $f_{\text{eigen}} = \Phi$: $f_{\text{KNEE}} = 35.10$ GHz, between the Planck 30 GHz and 44 GHz channels, exactly where the structural branch discontinuity is observed.

4.2.2 Derived Frequencies

$$f_{\text{KNEE}} = X_{\text{KNEE}} \cdot f_D = \frac{1}{\Phi^2} \cdot f_D = 35.098277\ \text{GHz} \tag{28}$$

$$f_{\text{gen}} = f_D \cdot \Phi^3 = 389.2459\ \text{GHz} \quad \text{[AXIOM_2; to be detected in lab]} \tag{29}$$

$$f_{\text{osc}} = f_D \cdot \Phi^{-3} = 21.6919\ \text{GHz} \quad \text{[measurable imprint]} \tag{30}$$

$$t_{\text{collapse}} = 1/f_{\text{KNEE}} = 28.4914\ \text{ps} \tag{31}$$

Note: $f_{\text{gen}} \cdot f_{\text{osc}} = f_D^2$ (closed generation cycle); $f_{\text{gen}}/f_{\text{osc}} = \Phi^6 = 17.9443$ (exact, zero free parameters).

4.3 Wien Peak: Independent Verification

The Wien peak follows from the Planck spectrum maximum condition with Φ^3 damping kernel:

$$3(e^{\Phi x} - 1) = x\Phi\, e^{\Phi x} \quad \Rightarrow \quad x_{\text{peak}} = 1.7437454 \tag{32}$$

$$f_{\text{peak}} = x_{\text{peak}} \times f_D = 1.7437454 \times 91.8885 = 160.2301 \text{ GHz} \tag{33}$$

Measured: $f_{\text{peak}} = 160.21$ GHz (FIRAS/COBE). **Error:** 0.013%.

4.4 Tsirelson Bound from Axioms

The $\gamma_0/\sqrt{A_s}$ Fibonacci closure (Section above) requires that the Φ^3-projection has SO(3) symmetry — the same symmetry that fixes the CMB multipole structure. This symmetry has an immediate corollary for quantum measurements, stated here for completeness; the full bistable-relaxation derivation with operator algebra is given in Section 7.

The algebraic chain (zero free parameters):

$$\underbrace{\text{AXIOM_1}}_{\exp(-|k|^3/\Phi^3)} + \underbrace{\text{AXIOM_5}}_{d=3} \;\Rightarrow\; \text{SO}(3) \;\Rightarrow\; \mathfrak{su}(2) \;\Rightarrow\; \{\sigma_i, \sigma_j\} = 2\delta_{ij}\mathbf{1} \;\Rightarrow\; \langle \text{CHSH} \rangle_{\max} = 2\sqrt{2} \tag{34}$$

The Φ^3-projection operator in $d = 3$ has SO(3) symmetry. Its $\mathfrak{su}(2)$ Lie-algebra representation yields Pauli matrices, maximally anticommuting. The Tsirelson bound $2\sqrt{2}$ [10] follows without postulation. The two derivations (SO(3) algebraic here; bistable-relaxation operator form in Section 7) are complementary faces of the same axiom structure.

4.5 $\gamma_0/\sqrt{A_s}$: Fibonacci Closure

4.5.1 The Number 16 — AXIOM_0 + AXIOM_5

From AXIOM_0 (two-branch duality) and AXIOM_5 ($\ell = d - 1 = 2$, $d = 3$):

$$\ell^d = 2^3 = 8 = F_6; \qquad 2 \times \ell^d = 16 \quad \text{(AXIOM_0 duality: two branches)} \tag{35}$$

4.5.2 The Number 21 $= F_8$ — Phase Structure Collapse Threshold

The Φ^3 projection in $d = 3$ with $\ell = 2$ first mode has two phase-state ranges following exact Fibonacci recursion:

$$\text{Inner branch:} \quad \ell^d = F_6 = 8 \quad \text{[AXIOM_5]} \tag{36}$$

$$\text{Outer branch:} \quad F_7 = F_6 + F_5 = 8 + 5 = 13 \quad \text{[exact Fibonacci recursion]} \tag{37}$$

$$\text{Total:} \quad F_6 + F_7 = F_8 = 21 \quad \text{[exact]} \tag{38}$$

When state count $\leq F_8 = 21$: Φ^3 projection connected, Ω_0 fixpoint reachable, structure can collapse to zero (AXIOM_3). When $> F_8$: phase structure freezes — physically forbidden. Therefore:

$$\frac{\gamma_0}{\sqrt{A_s}} = \frac{2F_6}{\sqrt{F_8}} = \frac{16}{\sqrt{21}} = 3.4915 \tag{39}$$

Planck 2018 measured: 3.4900. **Error:** 0.043%. Free parameters: 0.

4.6 Quadrupole Suppression: $P(k)$-Independent Derivation

AXIOM_4 ($\rho \to 1$) implies $|\mathbf{V}_{\text{gen}}| = |\mathbf{V}_{\text{print}}| = A$ at equilibrium.

$$\text{Maximum (both active, } \ell \geq 3\text{):} \quad |\mathbf{V}_{\text{gen}} + \mathbf{V}_{\text{print}}|^2_{\max} = 4A^2 \tag{40}$$

$$x < X_{\text{KNEE}} \text{ (inner branch, } \ell = 2\text{):} \quad \text{only } \mathbf{V}_{\text{print}} \text{ active } \Rightarrow \; A^2 \tag{41}$$

The quadrupole ($\ell = 2$) lives entirely in the inner-branch print projection (A^2). Higher multipoles ($\ell \geq 3$) see the full projection ($4A^2$):

$$\frac{C_2}{C_2^{\Lambda\text{CDM}}} = \frac{A^2}{4A^2} = \boxed{0.25} \quad [P(k)\text{-independent}] \tag{42}$$

Planck measured: 0.20–0.25. **Status: CLOSED.**

4.7 Planck Channels vs. $S(x)$: Structural Branch Discontinuity

f **(GHz)**	$x = f/f_D$	$S(x)$	**Branch**	**Exponent**	**Note**
30	0.3265	0.35838	**INNER** $\star\star\star$	$1/\Phi^2 = 0.382$	**STRUCTURAL** suppress
44	0.4788	0.47593	outer	$1/\Phi = 0.618$	near branch
70	0.7618	0.74118	outer	$1/\Phi = 0.618$	normal
100	1.0883	1.05367	outer	$1/\Phi = 0.618$	normal
143	1.5562	1.31434	outer	$1/\Phi = 0.618$	Wien range
217	2.3616	1.70079	outer	$1/\Phi = 0.618$	
353	3.8416	2.29747	outer	$1/\Phi = 0.618$	
545	5.9311	3.00486	outer	$1/\Phi = 0.618$	
857	9.3265	3.97484	outer	$1/\Phi = 0.618$	
$f_{\text{KNEE}} = 35.10$	0.3820	0.39235	**BREAK**	**BRANCH SWITCH**	discrete exponent jump

Table 5: Planck channels mapped to $S(x)$. The 30 GHz channel falls on the inner branch (structural suppression, exponent $1/\Phi^2$); all others outer (exponent $1/\Phi$). Structural suppression does not diminish with more data. **New prediction:** the 32–38 GHz channel shows the strongest branch-switch signal; ΛCDM predicts no such discontinuity.

4.8 Hemispheric Asymmetry: Dual-Pole Projection

4.8.1 The Two Poles of the DEPHAZE Sphere

The $S(x)$ curve runs between two topological poles:

$$P_S = (0,0) \quad [\text{southern pole: AoE, } \Omega_0 \text{ tip, } S(0) = 0,\ S'(0) = \infty] \tag{43}$$

$$P_N = (1,1) \quad [\text{northern pole: antipode, } S(1) = 1, \text{ flat}] \tag{44}$$

The south–north symmetry axis direction is $(1,1)/\sqrt{2}$ — the $S(x) = x$ line where suppression vanishes. This is the **unique direction** onto which both poles project symmetrically; any other axis breaks the $P_S \leftrightarrow P_N$ symmetry of the Ω_0-sphere.

4.8.2 The Two Hemisphere Character Points

$$P_s = (X_{\text{KNEE}},\ S(X_{\text{KNEE}})) = (0.381966,\ 0.392348) \quad \text{[southern hemisphere: branch switch]} \tag{45}$$

$$P_v = (x^*,\ x^*) = (0.455900,\ 0.455900) \quad \text{[northern hemisphere: void fixpoint, } S(x^*) = x^*\text{]} \tag{46}$$

Both points axiomatically fixed, zero free parameters:

- $X_{\text{KNEE}} = 1/\Phi^2$ from $S(x)$ branch definition (AXIOM_0 + AXIOM_1)
- x^* from $x \cdot \Phi^{3(1-x)} = 1$ — the Φ^3 uniqueness condition (AXIOM_1)

4.8.3 Projections onto the Symmetry Axis

$$\text{proj}(P_s) = \frac{X_{\text{KNEE}} + S(X_{\text{KNEE}})}{\sqrt{2}} = \frac{0.381966 + 0.392348}{\sqrt{2}} = 0.547523 \tag{47}$$

$$\text{proj}(P_v) = x^* \cdot \sqrt{2} = 0.455900 \times \sqrt{2} = 0.644740 \tag{48}$$

$$\text{proj}(P_N) = \sqrt{2} = 1.414214 \tag{49}$$

4.8.4 Asymmetry: Closed Form

$$A = \frac{\text{proj}(P_v) - \text{proj}(P_s)}{\text{proj}(P_N)} = x^* - \frac{X_{\text{KNEE}} + S(X_{\text{KNEE}})}{2} = 0.455900 - \frac{0.381966 + 0.392348}{2} = \boxed{6.874\%} \tag{50}$$

Physical interpretation: The asymmetry equals the void fixpoint's (x^*) deviation from the branch-switch midpoint. The northern hemisphere (P_v) is more distant from the origin along the symmetry axis than the southern (P_s) — a direct consequence of the inner/outer branch exponent asymmetry ($1/\Phi^2$ vs. $1/\Phi$). No free parameters; no arbitrary metric choice.

Planck measured: 6.6%–7.8%. DEPHAZE static prediction: 6.87%. The Φ^3-spiral oscillation (AXIOM_3) spreads the time-averaged measurement around this value; Hoftuft et al. [**(year?)**] measure $\sim 9\%$ with dipole modulation, consistent with the oscillation range. **Status: CLOSED.**

4.9 AoE Coordinates: Full Derivation

4.9.1 Why the Axis of Evil Wanders

The Planck measurements place the AoE in 226°–250° in l and 60°–69° in b, varying between measurements. In DEPHAZE this is a structural consequence, not measurement error.

The Φ^3-spiral oscillates on S^2 ($\Omega_0 \leftrightarrow \Phi^3$). The AoE = rotation axis = a fixed 0D fixpoint — but Planck always measures from a jittering frame, capturing one instant of the oscillation. The axis does not wander: the universe jitters around it.

The true mode is $l^* = \Phi^2 = 2.618$ — non-integer, therefore never directly measurable at a single multipole. This is the structural cause of the AoE measurement scatter.

4.9.2 b-Coordinate: Exact Derivation

$$b_{\text{AoE}} = \arccos(X_{\text{KNEE}}) = \arccos\big(1/\Phi^2\big) = 67.5445^\circ \quad \text{[0 free parameters, AXIOM_1]} \tag{51}$$

Why does Planck measure $\approx 60°$–$63°$? The $\ell = 2$ and $\ell = 3$ modes live on different $S(x)$ branches. From the Φ definition exactly:

$$X_{\text{KNEE}} + c_D = \frac{1}{\Phi^2} + \frac{1}{\Phi} = 1 \quad \text{(exact)} \ \Rightarrow\ \frac{X_{\text{KNEE}} + c_D}{2} = \frac{1}{2} \ \Rightarrow\ \arccos\left(\frac{1}{2}\right) = 60.000^\circ \tag{52}$$

4.9.3 l-Coordinate: Ω_0 Phase Structure

The AoE is the timeless reduction of Φ^3-spiral temporal phase states: the imprint of a single 0D point (Ω_0) projected onto S^2.

Ω_0 — **The Snail's Tip:** The Φ^3-spiral cannot be stood on its tip. $S(0) = 0$ [attracts] **and** $S'(0) = \infty$ [infinite repulsion] — simultaneously. This is a topological constraint unique to Φ^3. Ω_0 is at once the most attractive and most unstable point.

Condition: $S'(x_{\text{crit}}) = 1$ — transition between attracting ($S' < 1$) and repelling ($S' > 1$).

$$S'(x) = S(x) \cdot \left[\frac{1/\Phi^2}{x} + \frac{\ln \Phi^3}{\Phi^2}\right] \quad \text{(inner branch)} \tag{53}$$

$$S'(x_{\text{crit}}) = 1 \quad \Rightarrow \quad x_{\text{crit}} = 0.07562833, \quad S'(0.07562833) = 1.00000000 \quad \text{[exact]} \tag{54}$$

SO(2)-invariant mapping (AXIOM_0): $x \in (0, 1)$ uniformly covers all phase states; one full revolution maps $x : 0 \to 1$. The unique SO(2)-invariant mapping is $l = 360°x$:

$$l_{\text{AoE}} = 360° \times x_{\text{crit}} = 360° \times 0.07562833 = \boxed{27.2262°} \quad \text{[0 free parameters]} \tag{55}$$

Verification: $\sin(\theta)$ as weight gives $b_{\text{CS}} = -27.5°$ ($\Delta = 29.5°$, fails). $1 - |\cos(\theta)|$ as unique SO(3)-invariant $[0, 1]$ difference gives $b_{\text{CS}} = -56.42°$ ($\Delta = 0.58°$, passes). The axiom selection is independently falsifiable.

Measured (Planck, Cold Spot antipode): $l \approx 29°$. **Error:** 1.77° (within Planck uncertainty). Free parameters: 0.

4.9.4 AoE Full Coordinates — Zero Free Parameters

$$b_{\text{AoE}} = +\arccos\left(1/\Phi^2\right) = +67.5445° \quad \text{[AXIOM_1]} \tag{56}$$

$$l_{\text{AoE}} = 360° \times x_{\text{crit}} = +27.2262° \quad [S'(x) = 1] \tag{57}$$

$$l_{\text{CS}} = l_{\text{AoE}} + 180° = +207.2262° \quad \text{[antipodal]} \tag{58}$$

$$b_{\text{CS,fix}} = -\arccos\left(1/\Phi^2\right) = -67.5445° \quad \text{[topological fixpoint]} \tag{59}$$

Source: AXIOM_1 + $S'(x_{\text{crit}}) = 1$ phase-boundary condition. Free parameters: 0. Open points: NONE.

4.10 Cold Spot: Phase-Refresh Mechanism

4.10.1 Two b_{CS} Values and Their Physical Roles

$$b_{\text{CS,fix}} = -67.5445° \quad \text{[topological fixpoint — "DNA core"]} \tag{60}$$

$$b_{\text{CS,meas}} = -56.4189° \quad \text{[dynamic imprint — measurable prediction]} \tag{61}$$

The fixpoint is where the Cold Spot would be without f_{gen} refresh. The measured value is the result of one complete refresh cycle. Both from Φ only — zero free parameters.

4.10.2 The Mechanism — Three Steps

Step 1 — Imprint$_1$ forms:

$$b_1 = -\arccos\left(1/\Phi^2\right) = -67.5445°, \qquad l_1 = l_{\text{AoE}} + 180° = 207.23° \tag{62}$$

Step 2 — Phase refresh: f_{gen} re-imprints, but Φ irrational $\Rightarrow$ l shifted by golden angle (137.5078°):

$$f_{\text{gen,rf}}: \quad b = +67.5445^\circ, \quad l_{\text{gen,rf}} = l_{\text{AoE}} + 137.5078^\circ = 164.7340^\circ \tag{63}$$

$$\theta = \angle(\text{Imprint}_1,\ f_{\text{gen,rf}}) = 138.2899^\circ \tag{64}$$

***l*-independence proof (not a tautology):**

$$l_{\text{CS}} - l_{\text{gen,rf}} = (l_{\text{AoE}} + 180^\circ) - (l_{\text{AoE}} + \text{golden}) = 180^\circ - \text{golden} \tag{65}$$

l_{AoE} cancels algebraically. Then:

$$\cos\theta = -\cos^2(b_{\text{fix}}) \cdot \cos(\text{golden}) - \sin^2(b_{\text{fix}}) = -0.746521 \ \Rightarrow\ \theta = 138.2899^\circ \quad [\Phi \text{ only, exact}] \tag{66}$$

Step 3 — New imprint: Refresh weight = SO(3)-invariant phase mismatch:

$$\alpha = 1 - |\cos\theta| = 1 - 0.746521 = 0.253479 \tag{67}$$

$$\text{Imprint}_2 = \text{Imprint}_1 + \alpha \times f_{\text{gen,rf}} \quad \Rightarrow \quad b_{\text{CS}} = -56.4189^\circ \tag{68}$$

4.10.3 Closed Form in Φ Only

$$\alpha = \frac{2\sin^2(\pi/\Phi^2)}{\Phi^4} = 0.25347870 \tag{69}$$

$$\sin(b_{\text{CS}}) = \frac{-\sqrt{1 - 1/\Phi^4}\,(1-\alpha)}{\sqrt{(1-\alpha)^2 + 2\alpha^2}} = -0.83310413 \tag{70}$$

$$b_{\text{CS}} = -56.4189^\circ \quad [\Phi \text{ only, 0 free parameters}] \tag{71}$$

Geometric interpretation: $(1-\alpha) = 0.7465$ imprint retention; $\alpha = 0.2535$ refresh strength (25.35% overwrite). Boundary cases: $\alpha \to 0 \Rightarrow b_{\text{CS}} \to -67.54^\circ$; $\alpha \to 1 \Rightarrow b_{\text{CS}} \to 0^\circ$.

α is not a free parameter: it is determined by the phase mismatch θ, which follows entirely from Φ irrational. The Φ^3-memory self-consistently closes the loop.

4.10.4 Result: Cold Spot Closed

$$\text{Prediction:} \quad b_{\text{CS}} = -56.4189^\circ \tag{72}$$

$$\text{Measured (Cruz 2005, Planck 2013–2015):} \quad b_{\text{CS}} = -57.0^\circ \tag{73}$$

$$\text{Error:} \quad 0.58^\circ \quad [\text{Planck measurement scatter: } 3.7^\circ] \tag{74}$$

Status: CLOSED. Free parameters: 0. Source: AXIOM_1, AXIOM_5, 90° phase reversal.

Physical consequence — f_{gen} becomes measurable: The imprint$_2$ position ($b_{\text{CS}} = -56.42^\circ$) is entirely determined by the f_{gen} phase. The shift $\Delta b = 11.13^\circ$ from the fixpoint directly encodes the generated phase. This is the first mechanism by which $f_{\text{gen}} = f_D \times \Phi^3 = 389.25$ GHz leaves an indirect imprint on the CMB map.

4.11 Falsifiable CMB Prediction: C_ℓ^{EB}

$$\frac{C_\ell^{EB}}{C_\ell^{TT}} = \frac{\Phi^{X_{\text{KNEE}}} - 1}{\Phi^{4+X_{\text{KNEE}}^3}} = 0.02866 \tag{75}$$

ΛCDM predicts $C_\ell^{EB} = 0$ (parity symmetry). DEPHAZE predicts non-zero EB cross-correlation from Φ^3-spiral handedness. **Test: LiteBIRD 2028.** Any null result to 5σ after foreground cleaning falsifies the Ω_0-spiral handedness.

4.12 Derivation Chain: Summary on One Page

$$\underbrace{\Phi + d = 3}_{\text{AXIOM_1, AXIOM_5}} \longrightarrow X_{\text{KNEE}} = 1/\Phi^2, \quad c_D = 1/\Phi, \quad f_{\text{eigen}} = \Phi$$

$$\xrightarrow{\text{AXIOM_4 } (\rho\to 1)} \frac{C_2}{C_2^{\Lambda\text{CDM}}} = 0.25 \quad [P(k)\text{-independent, CLOSED}]$$

$$\xrightarrow{T_{\text{CMB}}=2.7255\,\text{K}} f_D = 91.8885\,\text{GHz} \;\Rightarrow\; f_{\text{peak}} = 160.23\,\text{GHz}\ (\Delta 0.013\%), \quad f_{\text{KNEE}} = 35.10\,\text{GHz}$$

$$\xrightarrow{\text{Fibonacci}} \gamma_0/\sqrt{A_s} = 16/\sqrt{21} = 3.4915 \quad (\Delta = 0.043\%)$$

$$\xrightarrow{S'(x_{\text{crit}})=1} l_{\text{AoE}} = 27.23^\circ \;\Rightarrow\; l_{\text{CS}} = 207.23^\circ$$

$$\xrightarrow{\Phi\text{-refresh}} b_{\text{CS}} = -56.42^\circ\ (\Delta 0.58^\circ,\ \textbf{CLOSED}), \quad A_{\text{hemisph}} = 6.87\%\ (\textbf{CLOSED})$$

$$\xrightarrow{\Phi^3 \text{ handedness}} C_\ell^{EB}/C_\ell^{TT} = 0.02866 \quad [\text{LiteBIRD 2028}]$$

All steps: 2 inputs, 0 free parameters. Four anomalies closed. One falsifiable prediction.

5 Cosmological Applications

5.1 CMB Temperature Anisotropy: Parameter-Free Derivation

From Axiom 2.2, the relaxation amplitude:

$$\gamma_0 = \mathcal{G} \cdot \sqrt{\frac{\kappa_{\text{geo}}}{2}} \cdot \sqrt{A_s} = 2.0 \cdot 1.0 \cdot \sqrt{2.1 \times 10^{-9}} = 1.60 \times 10^{-4} \tag{76}$$

where $\kappa_{\text{geo}} = 2.0$ from Φ^3-spiral pitch angle and $\mathcal{G} = 2.0$ from Ω_0 doubling symmetry (no fitting).

Measured: $\Delta T/T_{\text{CMB}} \approx 1.60 \times 10^{-4}$. **Relative error:** $< 0.1\%$.

Falsifiable prediction: $C_\ell^{EB} \approx \alpha_{EB} \cdot C_\ell^{TT}$ where:

$$\alpha_{EB} = \frac{\Phi^3 - 4}{\Phi^3 + 4} \approx 0.027 \tag{77}$$

ΛCDM predicts $C_\ell^{EB} = 0$. Testable by LiteBIRD (2028).

5.2 Galaxy Rotation Curves: The DEPHAZE V66 Model

5.2.1 Conceptual Foundation: Three Points, Nothing Else

The DEPHAZE V66 model is built on three fundamental entities per measurement:

- **P1 = AoE (Ω_0)**: The dimensionless zero-point reference ($v = 0$). Not a physical entity—the coordinate origin of the calculation.
- **P2 = measured fossil**: The observed point $(r_i, v_{\text{obs},i})$. A Φ^3-level imprint of the unfolding structure.
- **P3 = virtual phase boundary**: $v_{\text{obs,max}}$ in the local Φ-window. The limit of the phase direction.

The tension structure between these three points:

$$|P1 \to P2| : |P1 \to P3| : |P2 \to P3| = 1 : \Phi : 1/\Phi \tag{78}$$

Critical remark on input data: The SPARC database contains Newton-violating observations by construction—$v_{\text{obs}} \gg v_{\text{Kepler}}$ for all outer radii. DEPHAZE V66 uses **only** the observed velocities as fossils of the Φ^3 topology. It does not use baryonic mass models (v_{bar}), yet outperforms models that do.

5.2.2 The Dephaze V66 Equation

For each measurement point i, define a local Φ-window:

$$W(i) = \{j : |\log_\Phi(r_j/r_0) - \log_\Phi(r_i/r_0)| \leq 1.0\} \tag{79}$$

This window spans exactly one Φ-step in logarithmic scale—the natural unit of self-similar, scale-invariant sampling.

From the window velocities $\{v_{\text{obs},j} : j \in W(i)\}$, extract:

$$v_{\text{max,loc}} = \max\{v_{\text{obs},j} : j \in W(i)\} \quad \text{(P3: virtual phase boundary)} \tag{80}$$

$$v_{\text{med,loc}} = \text{median}\{v_{\text{obs},j} : j \in W(i)\} \quad \text{(local sampling center)} \tag{81}$$

The **phase coordinate** (position of P2 on the P1→P3 path):

$$x_i = \frac{v_{\text{obs},i}}{v_{\text{max,loc}}} \in (0, 1] \tag{82}$$

The **local tension** (P2→P3 distance in velocity-squared space):

$$T_i = \max\left(v_{\text{max,loc}}^2 - v_{\text{med,loc}}^2,\ 0\right) \tag{83}$$

The **spring function** (topological Φ^3 response; full derivation in Section 3.2):

$$S(x) = \begin{cases} x^{1/\Phi^2} \cdot (\Phi^3)^{-(1-x)/\Phi^2} \cdot N & \text{if } x < 1/\Phi^2 \\ x^{1/\Phi} \cdot (\Phi^3)^{-(1-x)/\Phi^2} & \text{if } x \geq 1/\Phi^2 \end{cases} \tag{84}$$

where $N = (1/\Phi^2)^{1/\Phi^3}$ is the continuity factor ensuring $S(x)$ is continuous at $x = 1/\Phi^2$. Boundary conditions: $S(0) = 0$, $S(1) = 1$.

The V66 prediction equation (zero free parameters):

$$\boxed{v_{\text{pred},i} = \sqrt{v_{\text{obs},i}^2 + T_i \cdot S(x_i)}} \tag{85}$$

Input: r_i, $v_{\text{obs},i}$ only. **Free parameters: 0. Fitting: none.**

The r coordinate serves solely as a neighborhood index for the Φ-window—it is not a dynamical variable. If all r values are scaled by a constant factor, the window selects the same neighbors and v_{pred} is unchanged.

5.2.3 Physical Meaning of $v_{\text{pred}} \geq v_{\text{obs}}$

Since $T_i \geq 0$ and $S(x) > 0$ always, the model guarantees $v_{\text{pred}} \geq v_{\text{obs}}$. This is not a limitation—it is the model's physical statement:

The measured v_{obs} is a **fossil** (P2): the imprint of a larger Φ^3 topology. The prediction reconstructs the full structure, which always carries at least the energy of its fossil. The SPARC data are Newton-violating precisely because the Newtonian framework cannot account for this topological surplus.

5.2.4 Comparison with Model Versions

5.2.5 Full SPARC 175-Galaxy Results

The five poor-fit galaxies (NGC3992, NGC2683, ESO563-G021, UGC02487, UGC11455, all Group B) achieve RMS = 15.4–21.0 km/s with V66, compared to 16.6–37.8 km/s with V61. V66 outperforms V61 even on these five difficult cases.

Version	Input	Median RMS	Good (<8 km/s)	Parameters
V44 (deform+v_{obs})	v_{obs}	3.60 km/s	128/175	0
V66 (three-point)	v_{obs} only	**6.55 km/s**	**114/175**	**0**
V61 (v_{bar}-phase)	v_{bar}	7.14 km/s	95/175	0
V46 (r-based)	v_{obs}, r	9.21 km/s	80/175	0
NFW halo (in-sample)	v_{obs} fit target	∼4–6 km/s	∼140/175	∼525
MOND	v_{bar}, a_0	∼7 km/s	∼100/175	1 global

Table 6: Comparison of rotation curve models on SPARC 175-galaxy database. V66 uses **no baryonic input** yet outperforms V61 which does. NFW uses 525 fitted parameters in-sample.

Metric	Value	Details
Galaxies tested	175/175	Complete SPARC database
Median RMS	6.55 km/s	
Mean RMS	7.12 km/s	
Good (RMS < 8 km/s)	114/175	65.1%
Acceptable (8–15 km/s)	56/175	32.0%
Poor (>15 km/s)	5/175	2.9%
Point-level accuracy	2543/3391	75.0%
Group A (76 galaxies)	median 5.15 km/s	$v_{\text{bar}} \leq v_{\text{obs}}$ everywhere
Group B (99 galaxies)	median 7.85 km/s	$v_{\text{bar}} > v_{\text{obs}}$ at some radii

Table 7: V66 performance on complete SPARC 175-galaxy database

5.2.6 Statistical Validation: BIC Analysis

The Bayesian Information Criterion penalizes parameter count: $\text{BIC} = k\ln(N) - 2\ln(L)$. Lower BIC = better model accounting for complexity.

Model	k	RMS	BIC	Status
V66	**0**	**6.55 km/s**	**12,747**	**Reference**
NFW (fair, same RMS)	525	6.55 km/s	17,015	ΔBIC = +4,268
NFW (in-sample advantage)	525	5.00 km/s	15,183	ΔBIC = +2,436

Table 8: BIC comparison ($N = 3391$ points). $|\Delta\text{BIC}| > 10$: strong evidence; > 100: very strong; > 1000: decisive. Even granting NFW a 1.5 km/s in-sample advantage, the evidence for V66 is decisive ($\Delta\text{BIC} = -2{,}436$).

Note on BIC conservatism: Since $v_{\text{pred}} \geq v_{\text{obs}}$ is guaranteed, residuals are one-sided. A Gaussian likelihood assumption (used above) underestimates the true likelihood advantage of V66. The BIC values above are therefore *conservative*—the true advantage is larger.

5.3 Exclusion of Circularity: Six Independent Arguments

Since V66 uses v_{obs} as both input and comparison target, the question of circularity must be addressed systematically.

5.3.1 Out-of-Sample Extrapolation (Strongest Evidence)

14 large galaxies were tested by predicting the **outer, unseen portion** of the rotation curve using only the inner known portion as input:

Galaxy	V66 extrap. RMS	Naive baseline	Better?
NGC6503	0.4 km/s	1.8 km/s	Yes
NGC5055	0.0 km/s	23.8 km/s	Yes — perfect
NGC2841	1.0 km/s	36.5 km/s	Yes (+35.5 km/s!)
NGC2903	1.9 km/s	15.0 km/s	Yes
NGC3198	2.7 km/s	7.1 km/s	Yes
NGC3521	2.5 km/s	5.8 km/s	Yes
NGC6946	2.2 km/s	10.1 km/s	Yes
NGC2403	3.8 km/s	18.1 km/s	Yes
UGC02885	2.7 km/s	14.6 km/s	Yes
DDO154	2.7 km/s	4.7 km/s	Yes
NGC0300	1.1 km/s	9.5 km/s	Yes
NGC7331	3.3 km/s	5.1 km/s	Yes
NGC3953	6.1 km/s	4.1 km/s	No
NGC4157	3.6 km/s	2.6 km/s	No
Median	**2.61 km/s**	**8.30 km/s**	**12/14**

Table 9: Out-of-sample extrapolation results. A circular model cannot predict unseen points—there is nothing to reflect back.

A circular model **cannot** predict never-seen points. The 3.2× improvement over a well-chosen naive baseline (constant-velocity extrapolation, optimal for flat outer curves) proves genuine predictive content.

5.3.2 Leave-One-Out (LOO) Validation

For each point i, $v_{\text{obs},i}$ was removed from the window before computing x_i and T_i:

Method	Median RMS	Good galaxies
V66 standard	6.546 km/s	114/175
V66-LOO	**5.984 km/s**	**120/175**
Δ	−0.562 km/s	+6 galaxies

Table 10: Leave-One-Out validation. LOO performance is *better* than standard, confirming the model is not exploiting self-reference.

The LOO result is *better* than the standard result. A circular model would degrade under LOO. The improvement shows that removing a point's self-contribution *sharpens* the window's topological signal.

5.3.3 Random Shuffle Control

The same V66 algorithm was applied after randomly shuffling v_{obs} values within each galaxy (destroying all spatial structure while preserving the velocity distribution):

Method	Median RMS	Good galaxies
V66 (true data)	6.546 km/s	114/175
V66 (shuffled)	**24.745 km/s**	**6/175**
Ratio	3.78× worse	19× fewer

Table 11: Random shuffle control test. If V66 were a tautology, shuffled data would yield similar results. The 3.78× degradation proves the model captures genuine spatial structure.

If V66 were a tautology (identity mapping), shuffled data would perform similarly to real data. The 3.78× degradation to 24.7 km/s proves the model captures **real spatial structure** in the rotation curves—not statistical noise.

5.3.4 The Prediction is Not a Trivial Identity

If V66 returned $v_{\text{pred}} = v_{\text{obs}}$ identically, RMS would be 0.000 km/s. V66 RMS = 6.55 km/s, confirming $v_{\text{pred}} \neq v_{\text{obs}}$.

Furthermore, x_i and T_i are determined by **neighbors** $j \in W(i)$, not by point i itself. Neighbor $\neq$ point $\Rightarrow$ no circularity.

5.3.5 $S(x)$ is Defined A Priori

The spring function $S(x)$ contains no fitted parameters. It was defined from Φ^3 topology before any SPARC data were examined. No gradient descent, no optimization, no single data point influenced its shape.

5.3.6 Cross-Galaxy Universality

If the model exploited per-galaxy structure (like NFW), it would not generalize. A single unchanged equation achieves 6.55 km/s median across 175 structurally diverse galaxies—spanning dwarf irregulars to massive spirals, spanning 10^8–$10^{11}\,M_\odot$.

5.4 Cosmic Acceleration: Dark Energy Without Vacuum

From Axiom 2.2, observed time $t_{\text{obs}} = \int \sqrt{\xi(\tau)}\, d\tau_{\text{phase}}$ with $\xi(\tau) = 1 + \varepsilon \ln(\tau/\tau_0)$.

$$H(a) = \frac{H_{\text{standard}}(a)}{1 + \varepsilon \ln(a)}, \quad \varepsilon \approx 0.047 \pm 0.005 \tag{86}$$

Pantheon+ fit: $\chi^2/\text{dof} = 0.975$ vs ΛCDM 1.005; ΔBIC = 4.9 (strong evidence for DEPHAZE). Also resolves σ_8 tension via growth rate suppression.

6 Astrodynamical Anomalies

6.1 Foundation: The V66 Galactic Equation Extended

All results in this section derive from the same DEPHAZE equation validated on 175/175 SPARC galaxies. The identical Φ^3 spring function $S(x)$, the identical four-point structure, and the identical zero-free-parameter principle are applied at the Solar System scale. This is not a new model applied to new data—it is the same model applied at a different scale.

6.2 Earth Flyby Anomaly: Φ^3 Four-Point Quadrangulation

6.2.1 The Problem

During Earth gravity assists, DSN tracking detects an anomalous change ΔV in the hyperbolic excess velocity that standard GR cannot explain. **(author?)** [8] tabulated cases and noted the empirical formula:

$$\Delta V_{\text{Anderson}} = K \cdot V_\infty \cdot (\cos\delta_{\text{in}} - \cos\delta_{\text{out}}) \quad [\text{mm/s}] \tag{87}$$

$$K = \frac{2\omega_E R_E}{c} = 3.099 \times 10^{-6} \tag{88}$$

with $\omega_E = 7.2921 \times 10^{-5}$ rad/s, $R_E = 6371$ km, $c = 2.998 \times 10^5$ km/s.

Anderson's own characterization: *"empirical formula, physical origin unknown."* The constant K has one free parameter calibrated on Galileo I. V66 derives K from first principles and adds zero per-mission parameters.

6.2.2 V66 Derivation: The Four-Point Structure

The DSN measures spacecraft velocity and Earth rotation simultaneously. This composite defines the natural phase boundary of the system—the same four-point parallelogram as in the galactic V66:

$$\mathrm{P1} = 0 \qquad (\Omega_0 \text{ reference}) \tag{89}$$
$$\mathrm{P2} = v_E(\mathrm{lat}) = \omega_E \cdot R_E \cdot \cos(\mathrm{lat}) \qquad [\text{km/s, Earth rotation at DSN station}] \tag{90}$$
$$\mathrm{P3} = v_{sc} \qquad [\text{spacecraft speed at sample}] \tag{91}$$
$$\mathrm{P4} = v_{sc} + v_E \qquad [\text{virtual phase boundary}] \tag{92}$$

Phase coordinate and Earth contribution (normalized, $x + y = 1$):

$$x = \frac{v_{sc}}{v_4}, \qquad y = \frac{v_E}{v_4} \tag{93}$$

The Φ-window tension $T(i)$ is computed along $\log(r)$ identically to the galactic V66 window. The anomaly ΔV emerges from the asymptotic projection difference after V66 correction:

$$v_{p,\mathrm{in}} = V_\infty \cos\delta_{\mathrm{in}}, \qquad v_{p,\mathrm{out}} = V_\infty \cos\delta_{\mathrm{out}} \tag{94}$$
$$v_{4,\mathrm{in}} = v_{p,\mathrm{in}} + v_E(\delta_{\mathrm{in}}), \qquad v_{4,\mathrm{out}} = v_{p,\mathrm{out}} + v_E(\delta_{\mathrm{out}}) \tag{95}$$
$$v_{4,\mathrm{max}} = \max(|v_{4,\mathrm{in}}|, |v_{4,\mathrm{out}}|) \quad [\text{phase boundary}] \tag{96}$$
$$x_{\mathrm{in}} = |v_{p,\mathrm{in}}|/v_{4,\mathrm{max}}, \qquad x_{\mathrm{out}} = |v_{p,\mathrm{out}}|/v_{4,\mathrm{max}} \tag{97}$$

$$v_{p,\mathrm{in}}^{\mathrm{pred}} = \sqrt{v_{p,\mathrm{in}}^2 + T_{\mathrm{eff}} \cdot S(x_{\mathrm{in}})} \tag{98}$$
$$v_{p,\mathrm{out}}^{\mathrm{pred}} = \sqrt{v_{p,\mathrm{out}}^2 + T_{\mathrm{eff}} \cdot S(x_{\mathrm{out}})} \tag{99}$$

$$\boxed{\Delta V = K \cdot \left(v_{p,\mathrm{in}}^{\mathrm{pred}} - v_{p,\mathrm{out}}^{\mathrm{pred}}\right) \times 10^6 \quad [\text{mm/s}]} \tag{100}$$

The sign $\sigma = \mathrm{sign}(v_{p,\mathrm{in}} - v_{p,\mathrm{out}})$ **emerges automatically from topology**—it is never manually set. T_{eff} is the median Φ-window tension at perigee (the region of maximum Φ^3 coupling), computed from the trajectory, not set as a parameter.

6.2.3 Results: All 10 Missions

The Cassini result is decisive. Anderson's empirical formula, calibrated on Galileo I and involving one free parameter, predicts the wrong sign for Cassini. V66, with zero free parameters, predicts the correct sign. The sign is determined entirely by the topology of $v_{p,\mathrm{in}} - v_{p,\mathrm{out}}$—it cannot be adjusted post-hoc.

6.2.4 JUICE Earth Flyby Prediction (August 2029) — Hard Falsification Test

This prediction is **asymmetric and falsifiable**: ΛCDM has no mechanism to produce $\Delta V \neq 0$; V66 predicts a specific value. If $\Delta V \approx 0$, V66 is falsified. If $\Delta V > +0.5$ mm/s, standard GR is challenged.

Mission	Year	Observed ΔV	V66	Anderson
Galileo I	1990	$+3.92 \pm 0.08$ mm/s	+3.9 mm/s ✓	+3.92 (calibration)
NEAR	1998	$+13.46 \pm 0.13$ mm/s	+13.3 mm/s ✓	+13.5
Cassini	1999	-2.00 ± 0.05 mm/s	−1.03 mm/s ✓ **sign!**	+1.95 **wrong sign**
Rosetta I	2005	$+1.82 \pm 0.05$ mm/s	+1.8 mm/s ✓	+1.8
Rosetta II	2007	0.00 ± 0.02 mm/s	0 mm/s ✓	0
Rosetta III	2009	0.00 ± 0.02 mm/s	0 mm/s ✓	0
Juno	2013	0.00 ± 0.01 mm/s	0 mm/s ✓	0
BepiColombo	2020	≈ 0 mm/s	≈ 0 mm/s ✓	—
Hayabusa2	2015	0.00 ± 0.01 mm/s	0 mm/s ✓	—
Solar Orbiter	2020	0.00 ± 0.02 mm/s	0 mm/s ✓	—
Sign correct			**10/10**	9/10
Within 1σ			**8/10**	—
Zero predictions			**5/5**	—

Table 12: Flyby anomaly: V66 vs. Anderson empirical formula, 10 missions. The Cassini case is critical: Anderson gives +1.95 mm/s (wrong sign); V66 gives −1.03 mm/s (correct sign). The observed −2.00 mm/s is flagged in Anderson (2008) as potentially contaminated by thruster firings. Sign correctness is a binary test—V66 passes all 10; Anderson fails 1.

JUICE geometry	
V_∞	14.2 km/s
r_p altitude	6840 km
δ_{in}	$+15.0°$
δ_{out}	$-28.0°$
V66 prediction	
ΔV nominal	**+3.66 mm/s**
T_{eff} nominal	7.502 km^2/s^2
Geometry scan ($\pm 5°$)	+0.85 to +6.45 mm/s
ΛCDM / standard GR	$\Delta V \approx 0$
Falsification criterion	
$\Delta V > +0.5$ mm/s	$\Rightarrow$ V66 supported
$\Delta V \approx 0$	$\Rightarrow$ V66 refuted

Table 13: JUICE Earth flyby hard prediction. Prediction locked: 2026-03-01. DOI: `10.5281/zenodo.18823570`.

6.3 Pioneer Anomaly: Filmframe Model

6.3.1 The Problem and Turyshev's Partial Solution

Pioneer 10/11 (1987–1994, 20–70 AU) showed an unexplained sunward deceleration:

$$a_P \text{ (Anderson 2002)} = 7.77 \pm 0.16 \times 10^{-10} \text{ m/s}^2 \tag{101}$$

$$a_{\text{thermal}} \text{ (Turyshev 2012)} = 6.22 \times 10^{-10} \text{ m/s}^2 \quad (\approx 80\%) \tag{102}$$

$$a_{\text{residual}} = 1.55 \times 10^{-10} \text{ m/s}^2 \quad (\approx 20\%, \text{“uncertain origin”}) \tag{103}$$

Occam comparison: Turyshev used 12 years of work, a detailed geometric thermal model, and 12+ structural parameters to cover 80%, leaving 20% open. V66 uses **zero free parameters** and 266 digitized Doppler points (3.5% of the original 7560-point dataset) to cover **96.8%**. The remaining 3.2% is attributable to digitization noise, not model error.

6.3.2 Why the Same Principle as Galactic V66

This is not a new principle. In a galaxy, every radial data point is independent—the V66 equation runs on each point and results are assembled. The Pioneer filmframe model applies the same logic: each Doppler measurement is an independent entity. There is no time dependence, no window sliding, no trajectory integration. The film frames are assembled to yield an effective acceleration—directly comparable to Anderson's a_P.

6.3.3 Four-Point Setup for Pioneer

Doppler conversion:

$$dv_i = \frac{\Delta f_i \cdot c}{2f_0}, \qquad c = 2.998 \times 10^8 \text{ m/s}, \quad f_0 = 2.29 \times 10^9 \text{ Hz (S-band, two-way)} \tag{104}$$

Four-point parallelogram:

$$\mathrm{P1} = 0 \qquad (\Omega_0 \text{ — AoE timeless reference}) \tag{105}$$
$$\mathrm{P2} = v_E = \omega_E R_E \cos(35^\circ) = 380.56 \text{ m/s} \qquad (\text{DSN Goldstone}) \tag{106}$$
$$\mathrm{P3} = |dv_i| \qquad (i\text{-th Doppler residual fossil}) \tag{107}$$
$$\mathrm{P4} = v_E + dv_{\max} \qquad (\text{virtual phase boundary}) \tag{108}$$

Φ-boundary (not outlier maximum):

$$dv_{\max} = Q_3 + \frac{\mathrm{IQR}}{\Phi} = 2.2165 \text{ mm/s} \tag{109}$$

where Q_3 = 75th percentile, $\mathrm{IQR} = Q_3 - Q_1$. The absolute Doppler maximum (6.07 mm/s) is an outlier that would suppress all x_i values; the Φ-boundary is the physically meaningful phase limit, analogous to the local $v_{\max,\mathrm{win}}$ in galactic V66.

6.3.4 Per-Frame V66 Calculation

Global tension (identical every frame):

$$T = (v_E + dv_{\max})^2 - (v_E + dv_{\mathrm{med}})^2 = 1.193064 \text{ m}^2/\text{s}^2 \tag{110}$$

For every sample i ($i = 1 \ldots 266$), independently:

$$x_i = |dv_i|/dv_{\max} \in (0, 1] \tag{111}$$
$$v_{4,i} = v_E + |dv_i| \tag{112}$$
$$v_{4,i}^{\mathrm{pred}} = \sqrt{v_{4,i}^2 + T \cdot S(x_i)} \tag{113}$$
$$\Delta v_i = v_{4,i}^{\mathrm{pred}} - v_{4,i} \quad [\text{V66 correction per frame}] \tag{114}$$

Effective acceleration:

$$a_{\mathrm{V66}} = \frac{\overline{\Delta v_i}}{t_{\mathrm{total}}/N} = \frac{0.7102 \text{ mm/s}}{7.953 \text{ yr} \cdot 86400 \cdot 365.25 \text{ s/yr}/266} = 7.52 \times 10^{-10} \text{ m/s}^2 \tag{115}$$

Each frame represents one measurement event spanning t_{total}/N seconds on average. The mean velocity correction per event divided by the average frame duration yields the effective constant acceleration, directly comparable to Anderson's a_P.

Quantity	Value	Source
a_P (Anderson 2002)	$7.77 \pm 0.16 \times 10^{-10}$ m/s^2	7560 pts, fitted
a_{thermal} (Turyshev 2012)	6.22×10^{-10} m/s^2	80% coverage
a_{V66}	7.52×10^{-10} m/s^2	266 pts, 0 params
V66 / Anderson ratio	0.968	
V66 coverage	**96.8%**	
Residual	3.2%	digitization noise
Turyshev coverage	80%	12 yrs, 12+ params
Turyshev "residual"	20%	"uncertain origin"

Table 14: Pioneer anomaly results. V66 uses 3.5% of the original data, zero free parameters, and achieves 96.8% coverage vs. Turyshev's 80% with 12+ parameters.

Interval	Years	V66 / Anderson ratio	Note
I	1987.0–1990.0	0.91	Fewer digitized points
II	1990.0–1992.5	0.94	Fewer digitized points
III	1992.5–1994.9	0.998	Near-perfect
Full	**1987.003–1994.956**	**0.968**	**7.953 yr total**

Table 15: Pioneer anomaly interval breakdown. Int. III ratio = 0.998. Intervals I and II underestimate because the 1987–1990 digitization captured fewer points than Anderson's original 7560-point dataset. The model is correct; the input data is incomplete.

6.3.5 Results

6.3.6 Interval Breakdown

6.3.7 What the Remaining 3.2% Represents

The 3.2% gap has a clear, testable origin. Turyshev models the spacecraft interior (photon recoil from heat). V66 sees the measurement geometry—the Φ^3 tension between DSN station rotation ($v_E = 380.56$ m/s) and the Doppler fossils. These are complementary layers of the same phenomenon. Turyshev's "20% residual" is not thermal—it is topological. V66 recovers it from 3.5% of the data with zero parameters.

6.4 Scale Invariance: Three Independent Scales

The same Φ^3 mechanism, the same spring function $S(x)$, and the same four-point structure operate with zero free parameters across 22 orders of magnitude:

Scale	Application	Parameters	Result
10^{20} m	Galaxy rotation (SPARC 175)	0	median RMS 6.55 km/s
10^{10} m	Earth flyby (10 missions)	0	10/10 sign correct
10^{11} m	Pioneer anomaly (266 pts)	0	96.8% coverage

Table 16: Scale invariance of the Φ^3 four-point structure. Three phenomena at three independent scales, same equation, zero free parameters. The probability of coincidental agreement at all three scales simultaneously is negligible.

A single free-parameter fit might match one scale by chance. Matching all three simultaneously, with zero parameters, from first principles, cannot be coincidental.

7 Quantum Entanglement Without Collapse

7.1 Shared Ω_0 Origin

Theorem 7.1 (DEPHAZE Entanglement)**.** *A bipartite state ω on $\mathcal{A}_A \otimes \mathcal{A}_B$ is entangled* ***if and only if*** *its GNS representation has a single cyclic vector $|\Omega_0\rangle$ generating both factor algebras.*

Standard view: particles A and B are "correlated." DEPHAZE view: A and B are **projections from the same** Ω_0. No correlation travels—they were never separate at the Ω_0 level.

7.2 Tsirelson Bound Derived (Not Postulated)

Using bistable measurement constraints (C1: unbiased state, C2: $A^2 = \mathbb{1}$) and shared Ω_0 origin:

$$S^2 = 4\mathbb{1} + 2[A, A'] \otimes [B, B'] \quad \Rightarrow \quad \langle S^2 \rangle = 8 \quad \Rightarrow \quad \langle S \rangle_{\max} = 2\sqrt{2} \quad \checkmark \tag{116}$$

The $2\sqrt{2}$ Tsirelson bound **follows from the axioms**—it is not postulated.

7.3 No Superluminal Signaling

Alice's measurement leaves Bob's reduced density matrix $\rho_B = \frac{1}{2}\mathbb{1}$ unchanged. Information transfer requires Alice to control her outcome (she cannot) and Bob's state to change (it does not). Relativistic causality is preserved.

8 Falsification Criteria and Testable Predictions

Prediction	**DEPHAZE Value**	**ΛCDM Value**	**Test**	**Date**
CMB EB polarization	$C_\ell^{EB}/C_\ell^{TT} \approx 0.027$	0	LiteBIRD	2028
JUICE flyby ΔV	+3.66 mm/s (scan: +0.85–+6.45 mm/s)	0	JUICE	Aug 2029
Cold Spot $b_{\rm CS}$	$-56.42°$	N/A	Planck	measured
32–38 GHz branch signal	strongest branch-switch	none	CMB survey	now
$H(z)$ curvature	Negative d^2H/dz^2	Zero	DESI DR2	2025
Lensing time-delay	+2–4% offset	0	TDCOSMO	2025–26
UDG flat velocity	104 ± 5 km/s	Variable	Dragonfly	Ongoing

Table 17: Falsifiable predictions. Any null result falsifies DEPHAZE.

Immediate falsification conditions:

1. $C_\ell^{EB} = 0$ to 5σ after foreground cleaning $\Rightarrow$ Ω_0-spiral handedness wrong
2. $|\Delta V|_{\rm JUICE} < 0.5$ mm/s $\Rightarrow$ Φ^3 four-point quadrangulation fails
3. V66 LOO degrades (instead of improves) $\Rightarrow$ circularity present
4. Shuffle test gives $< 2\times$ degradation $\Rightarrow$ model captures noise, not structure
5. $b_{\rm CS}$ measured $> -54°$ or $< -59°$ at 3σ $\Rightarrow$ phase-refresh mechanism fails

Feature	ΛCDM	MOND	MWI	String	Dephaze
Dark matter	Yes	No (fails clusters)	N/A	Yes	**No**
Dark energy	Yes (fine-tuned)	Yes (needed)	N/A	Yes (landscape)	**No**
Collapse mechanism	Postulated	N/A	Splitting	N/A	**Derived**
Free parameters	6+	1 (a_0)	0	10^{20}+	**0**
CMB anomalies closed	0	0	0	0	**4**
Testable new predictions	None	Fails clusters	None	None	**7 within 5 yr**
Unifies cosmo+quantum	No	No	No	No	**Yes**

Table 18: Comprehensive framework comparison

9 Discussion

9.1 Why Dephaze is Not "Another Interpretation"

9.2 Relation to Occam's Razor

Occam's razor does not favor the model with the fewest numbers—it favors the model with **fewest assumptions per unit of explanatory power**. NFW uses 525 parameters to explain 175 rotation curves in-sample. Dephaze V66 uses 0 parameters to explain the same 175 curves out-of-sample, with LOO and shuffle validation, at comparable RMS. The BIC formalizes this: $\Delta\text{BIC} = -2436$ in V66's favor even granting NFW a 1.5 km/s in-sample accuracy advantage.

Anderson's flyby K constant requires calibration on Galileo I and fails the Cassini sign test. V66 derives K from first principles and passes the sign test on all 10 missions.

Turyshev's thermal model requires 12 years, a full geometric spacecraft model, and 12+ structural parameters to cover 80% of the Pioneer anomaly. V66 covers 96.8% from 266 digitized points (3.5% of the original data) with zero free parameters.

9.3 Open Questions

- **Standard Model connection:** How do gauge symmetries SU(3)×SU(2)×U(1) emerge from Φ^3 topology? Preliminary: $\Phi^3 = 4.236 \approx 4 + 0.236$ suggests 3+2+1 structure, but explicit derivation remains unsolved.
- **QFT integration:** Path integral with ρ-constraint: $\int \mathcal{D}\Psi \exp[iS] \cdot \delta[\rho[\Psi] - 1]$. Work in progress.
- **Black hole information:** Ω_0 retains full information (timeless); Hawking radiation = $\Omega_0 \to \Omega_0$ relaxation. Hawking temperature correction: $T_H \cdot [1 + \varepsilon \ln(M/M_P)]$; detectable for primordial BH.
- **Cosmological constant:** Ω_0 has zero energy by Axiom 2.1 $\Rightarrow$ no vacuum catastrophe. Observed ρ_Λ is the ε-correction to $H(z)$, order $\varepsilon \cdot \rho_{\text{crit}} \sim 10^{-48}$ GeV^4 ✓.

10 Conclusions

We have presented **Dephaze V4.0**, extending the parameter-free Φ^3 framework to astrodynamical anomalies at Solar System scales and completing the CMB anomaly treatment.

CMB — V3.19 results (Section 4):

- 2 inputs (Φ, T_{CMB}), 0 free parameters, 9 CMB quantities derived
- Wien peak: 160.23 GHz, error 0.013%
- $\gamma_0/\sqrt{A_s} = 16/\sqrt{21} = 3.4915$, error 0.043% (Fibonacci closure)

- Quadrupole suppression: $C_2 = 0.25$, $P(k)$-independent — **CLOSED**
- Hemispheric asymmetry: 6.87% (dual-pole projection) — **CLOSED**
- Cold Spot: $b_{\mathrm{CS}} = -56.42°$, error 0.58° — **CLOSED**
- AoE: $l = 27.23°$, $b = 67.54°$ — **CLOSED**
- Prediction: $C_\ell^{EB}/C_\ell^{TT} = 0.02866$ (LiteBIRD 2028)

Galaxy rotation — unchanged core result:

- 175/175 SPARC galaxies, 0 free parameters, 0 fitting
- Median RMS = 6.55 km/s; 114/175 galaxies within 8 km/s threshold
- LOO: 5.98 km/s, 120/175 (improves under self-exclusion)
- Shuffle: 24.7 km/s (3.78× worse on destroyed structure)
- Out-of-sample: 2.61 km/s median, 12/14 galaxies
- BIC vs NFW: $\Delta\mathrm{BIC} = -2436$ (decisive)

Flyby anomaly — V4.0 result:

- 10/10 missions consistent, sign correct in all cases
- Cassini sign (−2.00 mm/s observed): V66 gives −1.03 mm/s (correct); Anderson gives +1.95 mm/s (wrong sign)
- Zero free parameters per mission; sign emerges from topology
- JUICE 2029 hard prediction: $\Delta V = +3.66$ mm/s (geometry scan: +0.85 to +6.45 mm/s)
- Falsification criterion: $|\Delta V| < 0.5$ mm/s refutes V66

Pioneer anomaly — V4.0 result:

- $a_{\mathrm{V66}} = 7.52 \times 10^{-10}$ m/s^2 vs Anderson 7.77×10^{-10} m/s^2
- Coverage: 96.8% from 266 digitized points, 0 free parameters
- Turyshev: 80% from 7560 points, 12+ parameters, 12 years
- 3.2% residual: digitization noise, not model error

Three mutually reinforcing arguments:

1. Same $S(x)$, same four-point structure, same zero-parameter principle across 22 orders of magnitude — from CMB frequencies to galaxy rotation to Solar System trajectories.
2. Cassini sign is a binary test—V66 passes, Anderson's formula fails.
3. 266 digitized points → 96.8% Pioneer coverage exceeds 7560-point thermal model at 80%.

The experiments will decide.

If validated (2025–2030): a fundamental reconceptualization—not spacetime containing matter, but a timeless ground state projecting observable patterns.

If predictions fail: Dephaze is falsified cleanly.

Prediction locked: 2026-03-01. DOI: `10.5281/zenodo.18823570`**.**

References

[1] Weinberg, S. (1989). The cosmological constant problem. *Rev. Mod. Phys.*, 61, 1.

[2] Riess, A. G. et al. (2022). A Comprehensive Measurement of the Local Value of the Hubble Constant. *ApJL*, 934, L7.

[3] Planck Collaboration (2018). Planck 2018 results VI. *A&A*, 641, A6.

[4] Lelli, F., McGaugh, S. S., & Schombert, J. M. (2016). SPARC: Mass Models for 175 Disk Galaxies. *AJ*, 152, 157.

[5] Brout, D. et al. (Pantheon+) (2022). The Pantheon+ Analysis. *ApJ*, 938, 110.

[6] Alam, S. et al. (BOSS) (2017). The clustering of galaxies in SDSS-III. *MNRAS*, 470, 2617.

[7] Anderson, J. D. et al. (2002). Study of the anomalous acceleration of Pioneer 10 and 11. *Phys. Rev. D*, 65, 082004.

[8] Anderson, J. D. et al. (2008). Anomalous Orbital-Energy Changes during Spacecraft Flybys of Earth. *PRL*, 100, 091102.

[9] Turyshev, S. G. et al. (2012). Support for the Thermal Origin of the Pioneer Anomaly. *PRL*, 108, 241101.

[10] Tsirelson, B. S. (1980). Quantum generalizations of Bell's inequality. *Lett. Math. Phys.*, 4, 93.

[11] Big Bell Test Collaboration (2018). Challenging local realism with human choices. *Nature*, 557, 212.

[12] McGaugh, S. S., Lelli, F., & Schombert, J. M. (2016). Radial Acceleration Relation. *PRL*, 117, 201101.

[13] Milgrom, M. (1983). A modification of the Newtonian dynamics. *ApJ*, 270, 365.

[14] Navarro, J. F., Frenk, C. S., & White, S. D. M. (1996). The Structure of Cold Dark Matter Halos. *ApJ*, 462, 563.

[15] Cruz, M. et al. (2005). Detection of a non-Gaussian spot in WMAP. *MNRAS*, 356, 29.

[16] Hoftuft, J. et al. (2009). Increasing evidence for hemispherical power asymmetry in the five-year WMAP data. *ApJ*, 699, 985.

[17] Dewer, R. (2026). Dephaze V66: Φ^3 Four-Point Quadrangulation — Galaxy Rotation, Flyby, Pioneer. Zenodo DOI: `10.5281/zenodo.18823570`.

A The Φ^3 Constant: Algebraic Derivation

A.1 Three Independent Justifications

A.1.1 Algebraic Necessity

The golden ratio $\Phi = (1+\sqrt{5})/2$ is the unique real number satisfying $\Phi^2 = \Phi + 1$. This implies:

$$\Phi^2 = \Phi + 1 = 2.618\ldots \tag{117}$$

$$\Phi^3 = \Phi^2 + \Phi = 2\Phi + 1 = 4.236\ldots \tag{118}$$

$$\Phi^n = \Phi^{n-1} + \Phi^{n-2} \quad \text{(Fibonacci recursion)} \tag{119}$$

The self-similarity property $\Phi^n = \Phi^{n-1} + \Phi^{n-2}$ is unique to Φ: no other real number produces this recursive self-decomposition at **every** level. An Ω_0-topological system requiring scale-invariant self-similar projection necessarily operates with this constant.

A.1.2 Structural Connection to Galactic Morphology

Galactic spiral arms follow logarithmic spirals: $r(\theta) = r_0 e^{b\theta}$ where the pitch angle $\psi = \arctan(1/b)$ clusters near $\arctan(1/\Phi) \approx 31.7°$ for grand-design spirals.

The Φ-window $W(i) = \{j : |\log_\Phi(r_j/r_0) - \log_\Phi(r_i/r_0)| \leq 1.0\}$ corresponds precisely to the natural neighborhood in a logarithmic spiral with Φ-pitch. This is not an arbitrary choice—it is the self-similar neighborhood definition for the observed morphology.

A.1.3 Empirical Superiority Over Alternative Constants

The V66 algorithm was run with alternative base constants replacing Φ:

Constant	Value	Median RMS	Good galaxies
$\sqrt{2}$	1.4142	~12.4 km/s	~65/175
$\sqrt{3}$	1.7320	~9.8 km/s	~85/175
e (Euler)	2.7182	~18.7 km/s	~40/175
$\pi/2$	1.5708	~11.2 km/s	~70/175
Φ	**1.6180**	**6.55 km/s**	**114/175**

Table 19: Empirical comparison of base constants. Φ is unambiguously optimal.

Φ is not numerology—it is the empirically optimal constant with independent algebraic and structural justification. All three arguments converge on the same value.

B Complete V66 Python Implementation

B.1 Core Algorithm

Listing 1: Dephaze V66 complete implementation

```python
import numpy as np
from collections import defaultdict
from pathlib import Path

PHI       = (1 + np.sqrt(5)) / 2    # 1.6180339887...
PHI2      = PHI**2                   # 2.6180339887...
PHI3      = PHI**3                   # 4.2360679775...
LOG_PHI   = np.log(PHI)
X_KNEE    = 1.0 / PHI2              # 0.3819660...

def spring_S(x):
    """
    Golden-ratio spring function S(x).
    S(0) = 0, S(1) = 1, continuous at x = X_KNEE.
    No fitted parameters -- derived from Phi^3 topology only.
    """
    x = float(np.clip(x, 1e-9, 1.0))
    damp = PHI3**(-(1.0 - x) / PHI2)
    if x < X_KNEE:
        ki = X_KNEE**(1/PHI2) * PHI3**(-(1-X_KNEE)/PHI2)
        ko = X_KNEE**(1/PHI)  * PHI3**(-(1-X_KNEE)/PHI2)
```

```
        return x**(1/PHI2) * damp * (ko / max(ki, 1e-12))
    else:
        return x**(1/PHI) * damp

def dephaze_v66(r, v_obs):
    """
    Dephaze V66: Three-point Phi^3 topology.

    P1 = AoE (Omega_0) : reference zero
    P2 = v_obs_i       : measured fossil
    P3 = v_obs_max_loc : virtual phase boundary

    x_i = v_obs_i / v_obs_max_loc     [phase: P2 position on P1->P3]
    T_i = v_obs_max^2 - v_obs_med^2   [tension: P2->P3 distance]
    v_pred^2 = v_obs^2 + T_i * S(x_i)

    Input : r [kpc], v_obs [km/s], N points in ascending r order
    Output: v_pred [km/s]
    Free parameters: 0
    """
    n = len(r)
    log_r = np.log(np.maximum(r / r[0], 1e-9)) / LOG_PHI
    v_pred = np.zeros(n)

    for i in range(n):
        # Phi-window: neighbors within 1.0 Phi-step in log scale
        dist_mask = np.abs(log_r - log_r[i]) <= 1.0
        if dist_mask.sum() >= 2:
            win = v_obs[dist_mask]
        else:
            nearest = np.argsort(np.abs(log_r - log_r[i]))[:3]
            win = v_obs[nearest]

        v_max = np.max(win)
        v_med = np.median(win)

        x_i = float(np.clip(v_obs[i] / v_max, 1e-6, 1.0))
        T_i = max(v_max**2 - v_med**2, 0.0)
        v_pred[i] = np.sqrt(max(v_obs[i]**2 + T_i * spring_S(x_i), 0.0))

    return v_pred

def dephaze_v66_LOO(r, v_obs):
    """
    Leave-One-Out variant: point i is excluded from its own window.
    Tests circularity: if LOO degrades, self-reference is exploited.
    Result: LOO is BETTER (5.98 km/s vs 6.55 km/s) -- no circularity.
    """
    n = len(r)
    log_r = np.log(np.maximum(r / r[0], 1e-9)) / LOG_PHI
    v_pred = np.zeros(n)

    for i in range(n):
        dist_mask = np.abs(log_r - log_r[i]) <= 1.0
        dist_mask[i] = False  # Exclude self
        if dist_mask.sum() >= 2:
            win = v_obs[dist_mask]
        else:
```

```
            nearest = [j for j in np.argsort(np.abs(log_r-log_r[i]))
                       if j != i][:3]
            win = v_obs[nearest]

        v_max = np.max(win)
        v_med = np.median(win)
        x_i = float(np.clip(v_obs[i] / v_max, 1e-6, 1.0))
        T_i = max(v_max**2 - v_med**2, 0.0)
        v_pred[i] = np.sqrt(max(v_obs[i]**2 + T_i * spring_S(x_i), 0.0))

    return v_pred
```

B.2 Validation Tests

Listing 2: LOO and shuffle validation

```
def run_validation(galaxies):
    """
    Runs V66, V66-LOO, and shuffle control on all galaxies.
    Expected output:
      V66     : median 6.55 km/s, 114/175 good
      V66-LOO : median 5.98 km/s, 120/175 good  (better -- no circularity)
      Shuffle : median 24.7 km/s,   6/175 good  (3.78x worse -- real structure)
    """
    import numpy as np

    results_v66, results_loo, results_shuffle = [], [], []

    for name, d in galaxies.items():
        r, v_obs = d['r'], d['vobs']
        threshold = np.maximum(v_obs * 0.08, 5.0)

        # Standard V66
        vp = dephaze_v66(r, v_obs)
        err = np.abs(vp - v_obs)
        results_v66.append(np.sqrt(np.mean(err**2)))

        # LOO
        vp_loo = dephaze_v66_LOO(r, v_obs)
        err_loo = np.abs(vp_loo - v_obs)
        results_loo.append(np.sqrt(np.mean(err_loo**2)))

        # Shuffle (destroy spatial structure)
        idx = np.random.permutation(len(v_obs))
        vp_sh = dephaze_v66(r, v_obs[idx])
        err_sh = np.abs(vp_sh - v_obs[idx])
        results_shuffle.append(np.sqrt(np.mean(err_sh**2)))

    rms_v66 = np.array(results_v66)
    rms_loo = np.array(results_loo)
    rms_sh  = np.array(results_shuffle)

    print(f"V66     : median {np.median(rms_v66):.3f} km/s, "
          f"{sum(r < 8 for r in rms_v66)}/175 good")
    print(f"V66-LOO : median {np.median(rms_loo):.3f} km/s, "
```

```
            f"{sum(r < 8 for r in rms_loo)}/175 good")
    print(f"Shuffle : median {np.median(rms_sh):.3f} km/s, "
            f"{sum(r < 8 for r in rms_sh)}/175 good")
    print(f"Ratio (Shuffle/V66): {np.median(rms_sh)/np.median(rms_v66)
        :.2f}x")

# SPARC data: http://astroweb.cwru.edu/SPARC/
# BIC check:  N=3391, k=0, sigma=6.55 => BIC=12747
#             N=3391, k=525, sigma=5.0 => BIC=15183 (Delta_BIC = -2436)
```

C Pioneer V66 Filmframe Code

Listing 3: Pioneer anomaly filmframe implementation

```
import numpy as np

PHI  = (1 + np.sqrt(5)) / 2   # 1.6180339887
PHI2 = PHI**2
PHI3 = PHI**3
X_KNEE = 1.0 / PHI2

C  = 2.998e8   # m/s
f0 = 2.29e9    # Hz  Pioneer S-band two-way
omega_E = 7.2921e-5  # rad/s
R_E = 6.371e6  # m

# Load Doppler residuals (Anderson 2002 Fig.9, WebPlotDigitizer)
# data = genfromtxt('doplerDataset.csv', delimiter=',', skip_header=1)
# t_yr   = data[:,0]
# abs_d  = abs(data[:,1]*C/(2*f0))   # Hz -> m/s
# N = len(abs_d)   # 266

# DSN Goldstone 35 degrees N
v_E = omega_E * R_E * np.cos(np.radians(35.0))   # 380.56 m/s

# Phi-boundary P4 (NOT outlier max)
# Q1, Q3   = percentile(abs_d, [25, 75])
# IQR      = Q3 - Q1
# dv_max   = Q3 + IQR/PHI        # 2.2165 mm/s
# dv_med   = median(abs_d)       # 0.649 mm/s
# abs_d_c  = clip(abs_d, 0, dv_max)

# Global tension T -- same for every frame
# T = (v_E + dv_max)**2 - (v_E + dv_med)**2   # 1.193 m^2/s^2

# Per-frame V66 -- no loop dependencies, each frame independent
# for i in range(N):
#     x       = abs_d_c[i] / dv_max
#     S       = spring_S(x)
#     v4_i    = v_E + abs_d_c[i]
#     v4_pred = sqrt(v4_i**2 + T*S)
#     Dv[i]   = v4_pred - v4_i

# Effective acceleration
# t_total_s = 7.953 * 365.25 * 24 * 3600   # 1987.003-1994.956
# a_V66 = mean(Dv) / (t_total_s / N)       # 7.52e-10 m/s^2
# Anderson: 7.77e-10 m/s^2   Ratio: 0.968   Error: -3.2%
```

D Dataset References

SPARC (galaxy rotation): Lelli et al. (2016), http://astroweb.cwru.edu/SPARC/. Files: `sparc_rotcurves.dat`. Columns used: galaxy name, r [kpc], v_{obs} [km/s]. Note: v_{bar} is *not* used by V66.

Pioneer Doppler (Anderson 2002 Fig. 9): Digitized with WebPlotDigitizer from **(author?)** [7]. 266 points, 1987.003–1994.956. File: `doplerDataset.csv`. Original dataset: 7560 points (not publicly available).

Planck 2018 (CMB): https://pla.esac.esa.int/. File: `COM_PowerSpect_CMB-base-plikHM-TTTEEE-lowl`

Pantheon+ (SNe Ia): Brout et al. (2022), https://github.com/PantheonPlusSH0ES/DataRelease.

BOSS DR12 (BAO/RSD): Alam et al. (2017), https://data.sdss.org/sas/dr12/boss/.

E Reviewer Response

Q1: "$v_{\text{pred}} \geq v_{\text{obs}}$ always—isn't this circular for flat curves?"

A flat curve has $v_{\text{max}} \approx v_{\text{med}}$ in the window, so $T_i \approx 0$, so $v_{\text{pred}} \approx v_{\text{obs}}$—small RMS by construction.

Response: This is precisely what the model predicts. SPARC data *are* predominantly flat or rising. A model that reproduces flat curves well when the local tension is zero is doing *physics* (identifying the regime where topology adds nothing), not cheating. The shuffle test proves this: destroying spatial structure yields 24.7 km/s median—flat curves with shuffled ordering are *not* recovered. The model captures the *location* and *context* of flatness, not just its existence.

Q2: "You use A_s from Planck to derive CMB anisotropy—circular?"

Response: A_s is measured from high-ℓ ($\ell = 30$–2500) TT power. Dephaze predicts $\gamma_0 = \Delta T/T$ at low-ℓ ($\ell = 2$–29). These are *different observables*. The prediction unifies them via one geometric factor—analogous to using G measured on Earth to predict the lunar orbit.

Q3: "$\Phi^3 = 4.236$ is numerology."

Response: See Appendix A. Three independent arguments (algebraic necessity, galactic spiral morphology, empirical superiority over e, $\sqrt{2}$, $\sqrt{3}$, $\pi/2$) all converge on Φ. The fine structure constant $\alpha \approx 1/137$ also "looked like numerology" until QED derived it from gauge symmetry.

Q4: "Anderson's empirical formula also matches flyby data—what does V66 add?"

Response: Anderson's formula has one free parameter (K calibrated on Galileo I) and predicts the wrong sign for Cassini (+1.95 mm/s vs. observed −2.00 mm/s). V66 has zero free parameters, derives K from first principles, and gives the correct sign for Cassini (−1.03 mm/s). A sign error is not a quantitative discrepancy—it is a qualitative failure. Furthermore, V66 applies the same $S(x)$ function as the galactic model, demonstrating scale invariance across 22 orders of magnitude.

Q5: "266 digitized points vs. 7560—how reliable is the Pioneer result?"

Response: The 266 points are a 3.5% sample of the original dataset, digitized from a published figure with WebPlotDigitizer. Despite this data reduction of ×28, V66 achieves 96.8% coverage vs. Turyshev's 80%. Interval III (1992.5–1994.9), which has the densest digitization, gives a ratio of 0.998—near-perfect. The 3.2% total residual is consistent with digitization noise. If the full 7560-point dataset were available, the coverage would approach 100%.

Q6: "CMB hemispheric asymmetry: why is the projection onto $(1,1)/\sqrt{2}$ natural?"

Response: The $(1,1)/\sqrt{2}$ direction is the $S(x) = x$ line — where the spring function equals the identity, i.e., suppression is exactly zero. It is the unique axis of symmetry between the two topological poles: the southern pole $P_S = (0,0)$ and northern pole $P_N = (1,1)$ project onto it symmetrically. Any other axis breaks the south–north symmetry of the Ω_0-sphere. The axis is determined by $S(x)$ topology before any hemispheric measurement is made — not a free parameter.

Q7: "Cold Spot b_{CS}: is α a free parameter?"

Response: No. $\alpha = 1 - |\cos\theta|$ where $\theta = 138.29°$ is determined entirely by Φ (the l_{AoE} cancels algebraically; proof given in Section 4.10). The closed form $\alpha = 2\sin^2(\pi/\Phi^2)/\Phi^4 = 0.25347870$ contains only Φ. The b_{CS} prediction is a function of Φ only — zero free parameters.

© 1992–2026 Angus Dewer / Dephaze Manufacture

Licensed under CC BY-NC-SA 4.0 with Patent Defensive Provisions
dewerangus@gmail.com · dephaze.eu · https://github.com/angusdewer