

# Upper bounding the commuting operator value of a nonlocal game

---

Angus Lowe

December 2020

- Nonlocal games, classical strategies, linear programming


- Nonlocal games, classical strategies, linear programming
- (Commuting operator) quantum strategies, semidefinite programming, necessary conditions

- Nonlocal games, classical strategies, linear programming
- (Commuting operator) quantum strategies, semidefinite programming, necessary conditions
- The NPA hierarchy, example for CHSH game

- Nonlocal games, classical strategies, linear programming
- (Commuting operator) quantum strategies, semidefinite programming, necessary conditions
- The NPA hierarchy, example for CHSH game
- Convergence to the commuting operator value


- Nonlocal games, classical strategies, linear programming
- (Commuting operator) quantum strategies, semidefinite programming, necessary conditions
- The NPA hierarchy, example for CHSH game
- Convergence to the commuting operator value
- Conclusion

# Nonlocal games



- $x, y \sim \pi(x, y)$


# Nonlocal games



- $x, y \sim \pi(x, y)$
- Win condition checker  $V(a, b, x, y) \in \{0, 1\}$





# Nonlocal games



- $x, y \sim \pi(x, y)$
- Win condition checker  $V(a, b, x, y) \in \{0, 1\}$

$$\text{win prob.} = \sum_{x, y} \pi(x, y) \sum_{a, b} p_{abxy} V(a, b, x, y) \quad (1)$$



## Example: CHSH game



- $a, b, x, y \in \{0, 1\}$ .  $x, y$  given uniformly at random.

$$V(a, b, x, y) = \begin{cases} 1 & \text{if } a \oplus b = x \wedge y \\ 0 & \text{otherwise} \end{cases} \quad (2)$$

## Example: CHSH game



- $a, b, x, y \in \{0, 1\}$ .  $x, y$  given uniformly at random.

$$V(a, b, x, y) = \begin{cases} 1 & \text{if } a \oplus b = x \wedge y \\ 0 & \text{otherwise} \end{cases} \quad (2)$$

- The maximum winning probability is  $3/4$

## Local strategies

- Alice and Bob should use a **deterministic** strategy ( $d_{abxy}$ )

# Local strategies

- Alice and Bob should use a **deterministic** strategy ( $d_{abxy}$ )

## Definition


The *local set*  $\mathcal{L}$  of a game is the convex hull of its deterministic points.

# Local strategies

- Alice and Bob should use a **deterministic** strategy ( $d_{abxy}$ )

## Definition

The *local set*  $\mathcal{L}$  of a game is the convex hull of its deterministic points.




# Local strategies

- Alice and Bob should use a **deterministic** strategy ( $d_{abxy}$ )

## Definition

The *local set*  $\mathcal{L}$  of a game is the convex hull of its deterministic points.



## The classical value


$$\begin{array}{ll} \max_p & \text{win prob.} \\ \text{s.t.} & p \in \mathcal{L} \end{array}$$

# Local strategies

- Alice and Bob should use a **deterministic** strategy ( $d_{abxy}$ )


## Definition

The *local set*  $\mathcal{L}$  of a game is the convex hull of its deterministic points.




## The classical value

$$\begin{array}{ll} \max_p & \text{win prob.} \\ \text{s.t.} & p \in \mathcal{L} \end{array}$$






# Quantum strategies (commuting operator)



$$p_{abxy} = \langle \psi | M_{a|x} M_{b|y} | \psi \rangle$$
$$\forall a, b, x, y$$

Measurement operators satisfy:

# Quantum strategies (commuting operator)




$$p_{abxy} = \langle \psi | M_{a|x} M_{b|y} | \psi \rangle$$
$$\forall a, b, x, y$$

Measurement operators satisfy:

1. **Orthogonal projectors** i.e.,  $M_{a|x} M_{a'|x} = \delta_{aa'} M_{a|x}$ , likewise for Bob's measurements.

# Quantum strategies (commuting operator)




$$p_{abxy} = \langle \psi | M_{a|x} M_{b|y} | \psi \rangle$$
$$\forall a, b, x, y$$

Measurement operators satisfy:

1. **Orthogonal projectors** i.e.,  $M_{a|x} M_{a'|x} = \delta_{aa'} M_{a|x}$ , likewise for Bob's measurements.
2. **Completeness** i.e.,  $\sum_a M_{a|x} = \mathbb{1} \forall x$ , likewise for Bob's measurements.

# Quantum strategies (commuting operator)




$$p_{abxy} = \langle \psi | M_{a|x} M_{b|y} | \psi \rangle$$
$$\forall a, b, x, y$$

Measurement operators satisfy:

1. **Orthogonal projectors** i.e.,  $M_{a|x} M_{a'|x} = \delta_{aa'} M_{a|x}$ , likewise for Bob's measurements.
2. **Completeness** i.e.,  $\sum_a M_{a|x} = \mathbb{1} \forall x$ , likewise for Bob's measurements.
3. **Commutativity** i.e.,  $[M_{a|x}, M_{b|y}] = 0 \forall a, b, x, y$ .

# Quantum strategies (commuting operator)




$$p_{abxy} = \langle \psi | M_{a|x} M_{b|y} | \psi \rangle$$
$$\forall a, b, x, y$$

Measurement operators satisfy:

1. **Orthogonal projectors** i.e.,  $M_{a|x} M_{a'|x} = \delta_{aa'} M_{a|x}$ , likewise for Bob's measurements.
2. **Completeness** i.e.,  $\sum_a M_{a|x} = \mathbb{1} \forall x$ , likewise for Bob's measurements.
3. **Commutativity** i.e.,  $[M_{a|x}, M_{b|y}] = 0 \forall a, b, x, y$ .

From now on, we refer to these as the 3 **CO requirements**.


# Preview: the (commuting operator) quantum set



## The quantum value

$$\begin{array}{ll} \max_p & \text{win prob.} \\ \text{s.t.} & p \in \mathcal{Q} \end{array}$$

# Preview: the (commuting operator) quantum set




## The quantum value

$$\begin{array}{ll} \max_p & \text{win prob.} \\ \text{s.t.} & p \in Q \end{array}$$

- Suppose Eve claims the quantum value of a game is  $> 2/3$ : can we certify this?

# Preview: the (commuting operator) quantum set




## The quantum value

$$\begin{array}{ll} \max_p & \text{win prob.} \\ \text{s.t.} & p \in \mathcal{Q} \end{array}$$

- Suppose Eve claims the quantum value of a game is  $> 2/3$ : can we certify this?
- **Answer:** no, but *if she's lying* we can always tell.  
 $\text{MIP}^{\text{co}}(2, 1) \subseteq \text{coRE}$  [NPA08].



# Preview: the (commuting operator) quantum set



## The quantum value

$\max_p$  win prob.

s.t.

$$p \in \mathcal{Q}$$

- Suppose Eve claims the quantum value of a game is  $> 2/3$ : can we certify this?
- **Answer:** no, but *if she's lying* we can always tell.  
 $\text{MIP}^{\text{co}}(2, 1) \subseteq \text{coRE}$  [NPA08].

- $\mathbb{S}_+^d = \{M \in \mathbb{R}^{d \times d} : \langle \psi | M | \psi \rangle \geq 0 \ \forall \ |\psi\rangle \in \mathbb{R}^d\}$

# Semidefinite programming


- $\mathbb{S}_+^d = \{M \in \mathbb{R}^{d \times d} : \langle \psi | M | \psi \rangle \geq 0 \ \forall \ |\psi\rangle \in \mathbb{R}^d\}$
- Generalization of linear programs
  - $\text{LP } (a_i^T x = b_i, x \geq 0) \longrightarrow \text{SDP } (\text{Tr}(A_i X) = b_i, X \in \mathbb{S}_+^d)$

# Semidefinite programming

- $\mathbb{S}_+^d = \{M \in \mathbb{R}^{d \times d} : \langle \psi | M | \psi \rangle \geq 0 \ \forall \ |\psi\rangle \in \mathbb{R}^d\}$
- Generalization of linear programs
  - LP ( $a_i^T x = b_i, x \geq 0$ )  $\longrightarrow$  SDP ( $\text{Tr}(A_i X) = b_i, X \in \mathbb{S}_+^d$ )
  - Regions can have curved faces

# Semidefinite programming


- $\mathbb{S}_+^d = \{M \in \mathbb{R}^{d \times d} : \langle \psi | M | \psi \rangle \geq 0 \ \forall \ |\psi\rangle \in \mathbb{R}^d\}$
- Generalization of linear programs
  - LP ( $a_i^T x = b_i, x \geq 0$ )  $\longrightarrow$  SDP ( $\text{Tr}(A_i X) = b_i, X \in \mathbb{S}_+^d$ )
  - Regions can have curved faces



$$\left\{ M = \begin{bmatrix} x & y \\ y & z \end{bmatrix} : M \in \mathbb{S}_+^2, \text{Tr}(M) = 1 \right\}$$

# Semidefinite programming

- $\mathbb{S}_+^d = \{M \in \mathbb{R}^{d \times d} : \langle \psi | M | \psi \rangle \geq 0 \ \forall \ |\psi\rangle \in \mathbb{R}^d\}$
- Generalization of linear programs
  - LP ( $a_i^T x = b_i, x \geq 0$ )  $\longrightarrow$  SDP ( $\text{Tr}(A_i X) = b_i, X \in \mathbb{S}_+^d$ )
  - Regions can have curved faces



$$\left\{ M = \begin{bmatrix} x & y \\ y & z \end{bmatrix} : M \in \mathbb{S}_+^2, \text{Tr}(M) = 1 \right\}$$

$$\left\{ M = \begin{bmatrix} \gamma & x & y \\ x & \gamma & z \\ y & z & 1 - 2\gamma \end{bmatrix} : M \in \mathbb{S}_+^3 \right\}$$

# Semidefinite programming

$$\begin{aligned} \max_{Z \in \mathbb{R}^{n \times n}} \quad & \text{Tr}(CZ) \\ \text{s.t.} \quad & \text{Tr}(A_i Z) = b_i, \quad i = 1, \dots, m \\ & Z \succeq 0 \end{aligned}$$


# Semidefinite programming

$$\begin{aligned} \max_{Z \in \mathbb{R}^{n \times n}} \quad & \text{Tr}(CZ) \\ \text{s.t.} \quad & \text{Tr}(A_i Z) = b_i, \quad i = 1, \dots, m \\ & Z \succeq 0 \iff Z \in \mathbb{S}_+^n \end{aligned}$$



# Semidefinite programming

$$\begin{aligned} \max_{Z \in \mathbb{R}^{n \times n}} \quad & \text{Tr}(CZ) \\ \text{s.t.} \quad & \text{Tr}(A_i Z) = b_i, \quad i = 1, \dots, m \\ & Z \succeq 0 \iff Z \in \mathbb{S}_+^n \end{aligned}$$



# Necessary conditions: the NPA hierarchy

## The Big Idea

*Expectations of products of operators must satisfy certain equalities because of the CO requirements. This leads to necessary conditions for the quantum set.*

# Necessary conditions: the NPA hierarchy

## The Big Idea

*Expectations of products of operators must satisfy certain equalities because of the **CO requirements**. This leads to necessary conditions for the quantum set.*

- *Example 1: Let  $S = M_{a|x}$  and  $T = M_{b|y}$ . Then*

$$\langle \psi | S^\dagger T | \psi \rangle = \langle \psi | T^\dagger S | \psi \rangle = p_{abxy}.$$

# Necessary conditions: the NPA hierarchy

## The Big Idea

*Expectations of products of operators must satisfy certain equalities because of the **CO requirements**. This leads to necessary conditions for the quantum set.*

- *Example 1: Let  $S = M_{a|x}$  and  $T = M_{b|y}$ . Then*

$$\langle \psi | S^\dagger T | \psi \rangle = \langle \psi | T^\dagger S | \psi \rangle = p_{abxy}.$$

- *Example 2: Let  $S = M_{a|x}$ ,  $T = M_{a|x}$ . Then*

$$\langle \psi | S^\dagger T | \psi \rangle = \langle \psi | S | \psi \rangle = \Pr[a|x] = \sum_b p_{abxy}.$$

- *Example 3:* Let  $S = M_{a|x}M_{b|y}$ ,  $T = M_{a'|x}M_{b|y'}$ . Then

$$\langle \psi | S^\dagger T | \psi \rangle = 0$$

## Necessary conditions: the NPA hierarchy

- Let  $\mathcal{S}_n$  be the set containing all products up to length  $n$  e.g.,

$$S = \underbrace{M_{a|x} M_{a'|x'} \dots M_{b|y} M_{b'|y'} \dots}_{\leq n}$$

# Necessary conditions: the NPA hierarchy

- Let  $\mathcal{S}_n$  be the set containing all products up to length  $n$  e.g.,

$$S = \underbrace{M_{a|x} M_{a'|x'} \dots M_{b|y} M_{b'|y'} \dots}_{\leq n}$$

- Then we can define the  $|\mathcal{S}_n| \times |\mathcal{S}_n|$  **moment matrix**  $\Gamma^n$  with entries  $\Gamma_{S,T}^n = \langle \psi | S^\dagger T | \psi \rangle$  for all  $S, T \in \mathcal{S}_n$ . [BCP<sup>+</sup>14]

# Necessary conditions: the NPA hierarchy

- Let  $\mathcal{S}_n$  be the set containing all products up to length  $n$  e.g.,

$$S = \underbrace{M_{a|x} M_{a'|x'} \dots M_{b|y} M_{b'|y'} \dots}_{\leq n}$$

- Then we can define the  $|\mathcal{S}_n| \times |\mathcal{S}_n|$  **moment matrix**  $\Gamma^n$  with entries  $\Gamma_{S,T}^n = \langle \psi | S^\dagger T | \psi \rangle$  for all  $S, T \in \mathcal{S}_n$ . [BCP<sup>+</sup>14]

## Properties of $\Gamma^n$

- $\Gamma^n \succeq 0$ . Why?  $\sum_{S,T} v_S^* \langle \psi | S^\dagger T | \psi \rangle v_T = \langle \psi | A^\dagger A | \psi \rangle \geq 0$ .



# Necessary conditions: the NPA hierarchy

- Let  $\mathcal{S}_n$  be the set containing all products up to length  $n$  e.g.,

$$S = \underbrace{M_{a|x} M_{a'|x'} \dots M_{b|y} M_{b'|y'} \dots}_{\leq n}$$

- Then we can define the  $|\mathcal{S}_n| \times |\mathcal{S}_n|$  **moment matrix**  $\Gamma^n$  with entries  $\Gamma_{S,T}^n = \langle \psi | S^\dagger T | \psi \rangle$  for all  $S, T \in \mathcal{S}_n$ . [BCP<sup>+</sup>14]

## Properties of $\Gamma^n$

- $\Gamma^n \succeq 0$ . Why?  $\sum_{S,T} v_S^* \langle \psi | S^\dagger T | \psi \rangle v_T = \langle \psi | A^\dagger A | \psi \rangle \geq 0$ .
- $\Gamma^n$  satisfies some equalities of the form  $\text{Tr}(A_i \Gamma^n) = b_i$ . Why? e.g.,  $\langle \psi | S^\dagger T | \psi \rangle = p_{abxy} \iff \text{Tr}(|S\rangle\langle T| \Gamma^n) = p_{abxy}$ .

# Necessary conditions: the NPA hierarchy

- Let  $\mathcal{S}_n$  be the set containing all products up to length  $n$  e.g.,

$$S = \underbrace{M_{a|x} M_{a'|x'} \dots M_{b|y} M_{b'|y'} \dots}_{\leq n}$$

- Then we can define the  $|\mathcal{S}_n| \times |\mathcal{S}_n|$  **moment matrix**  $\Gamma^n$  with entries  $\Gamma_{S,T}^n = \langle \psi | S^\dagger T | \psi \rangle$  for all  $S, T \in \mathcal{S}_n$ . [BCP<sup>+</sup>14]

## Properties of $\Gamma^n$

- $\Gamma^n \succeq 0$ . Why?  $\sum_{S,T} v_S^* \langle \psi | S^\dagger T | \psi \rangle v_T = \langle \psi | A^\dagger A | \psi \rangle \geq 0$ .
- $\Gamma^n$  satisfies some equalities of the form  $\text{Tr}(A_i \Gamma^n) = b_i$ . Why? e.g.,  $\langle \psi | S^\dagger T | \psi \rangle = p_{abxy} \iff \text{Tr}(|S\rangle\langle T| \Gamma^n) = p_{abxy}$ .
- There is a submatrix of  $\Gamma^n$  containing all values  $p_{abxy}$ .

# Upper bounding quantum values

## Constraints on $\Gamma^n$

1.  $\text{Tr}(A_i \Gamma^n) = b_i, \quad i = 1, \dots, m.$
2.  $\Gamma^n \succeq 0.$

# Upper bounding quantum values

## Constraints on $\Gamma^n$

1.  $\text{Tr}(A_i \Gamma^n) = b_i, \quad i = 1, \dots, m.$
2.  $\Gamma^n \succeq 0.$

- Let  $\mathcal{Q}^n$  be the set of all strategies  $p$  for which there exists a  $\Gamma^n$  satisfying the constraints. Then  $\mathcal{Q} \subseteq \mathcal{Q}^n \subseteq \mathcal{Q}^{n-1} \subseteq \dots \subseteq \mathcal{Q}^1.$

# Upper bounding quantum values

## Constraints on $\Gamma^n$

1.  $\text{Tr}(A_i \Gamma^n) = b_i, \quad i = 1, \dots, m.$
2.  $\Gamma^n \succeq 0.$

- Let  $\mathcal{Q}^n$  be the set of all strategies  $p$  for which there exists a  $\Gamma^n$  satisfying the constraints. Then  $\mathcal{Q} \subseteq \mathcal{Q}^n \subseteq \mathcal{Q}^{n-1} \subseteq \dots \subseteq \mathcal{Q}^1.$

$$\begin{array}{ll} \max_p & \text{win prob.} \\ \text{s.t.} & p \in \mathcal{Q}^n \end{array}$$

# Upper bounding quantum values

## Constraints on $\Gamma^n$

1.  $\text{Tr}(A_i \Gamma^n) = b_i, \quad i = 1, \dots, m.$
2.  $\Gamma^n \succeq 0.$

- Let  $\mathcal{Q}^n$  be the set of all strategies  $p$  for which there exists a  $\Gamma^n$  satisfying the constraints. Then  $\mathcal{Q} \subseteq \mathcal{Q}^n \subseteq \mathcal{Q}^{n-1} \subseteq \dots \subseteq \mathcal{Q}^1.$

$$\begin{array}{ll} \max_p & \text{win prob.} \\ \text{s.t.} & p \in \mathcal{Q}^n \end{array}$$

$\iff$

$$\begin{array}{ll} \max_{\Gamma^n} & \text{Tr}(C\Gamma^n) \\ \text{s.t.} & \text{Tr}(A_i \Gamma^n) = b_i \\ & \Gamma^n \succeq 0 \end{array}$$

# Upper bounding quantum values

## Constraints on $\Gamma^n$


1.  $\text{Tr}(A_i \Gamma^n) = b_i, \quad i = 1, \dots, m.$
2.  $\Gamma^n \succeq 0.$

- Let  $\mathcal{Q}^n$  be the set of all strategies  $p$  for which there exists a  $\Gamma^n$  satisfying the constraints. Then  $\mathcal{Q} \subseteq \mathcal{Q}^n \subseteq \mathcal{Q}^{n-1} \subseteq \dots \subseteq \mathcal{Q}^1.$

$\max_p$  win prob.  
s.t.  $p \in \mathcal{Q}^n$

$\iff$

$$\begin{array}{ll} \max_{\Gamma^n} & \text{Tr}(C\Gamma^n) \\ \text{s.t.} & \text{Tr}(A_i \Gamma^n) = b_i \\ & \Gamma^n \succeq 0 \end{array}$$



- Tsirelson's bound [Cir80] tells us the quantum value of the CHSH game is  $\approx 0.85$ .



## CHSH example revisited

- Tsirelson's bound [Cir80] tells us the quantum value of the CHSH game is  $\approx 0.85$ .
- For this game, it happens that  $Q^1 = Q$ !

# CHSH example revisited

- Tsirelson's bound [Cir80] tells us the quantum value of the CHSH game is  $\approx 0.85$ .
- For this game, it happens that  $Q^1 = Q$ !

```
In[28]:= res =  
  -SemidefiniteOptimization[g[ps], cons[r, ps],  
    {r ∈ Matrices[{5, 5}, Reals], ps ∈ Vectors[16, Reals]}, "PrimalMinimumValue"]  
Out[28]= 0.853553
```

# Convergence to the quantum value

## Theorem (Sufficiency of the NPA hierarchy)

Let  $p$  be a strategy such that there exists a valid moment matrix  $\Gamma^n$  for all  $n \geq 1$ . Then  $p$  is in the quantum set.

1)


$$\Gamma^n = \begin{bmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{bmatrix} \rightarrow \hat{\Gamma}^n = \begin{bmatrix} \cdot & \cdot & \cdot & 0 & 0 & \dots \\ \cdot & \cdot & \cdot & \vdots & \vdots & \vdots \\ \cdot & \cdot & \cdot & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

$$\rightarrow x^n := \text{vec}(\Gamma^n) = [\dots, 0, 0, \dots]^T$$

$$x^n \in B_\infty(0, 1)$$

$$\implies x^{n_i} \rightarrow x^\infty \iff \hat{\Gamma}^{n_i} \rightarrow \Gamma^\infty$$

Banach-Alaoglu



# Convergence to the quantum value

## Theorem (Sufficiency of the NPA hierarchy)

Let  $p$  be a strategy such that there exists a valid moment matrix  $\Gamma^n$  for all  $n \geq 1$ . Then  $p$  is in the quantum set.

2)

There exists infinite set of vectors  $V = \{|S\rangle : |S| = 1, 2, \dots\}$  such that

$$\Gamma_{S,T}^\infty = \langle S|T \rangle \quad (\text{sequential Cholesky decomposition})$$

Then take  $\mathcal{H} = \text{span}(V)$ , and measurement operators

$$\hat{E}_{a|x} = \text{proj}(\text{span}(\{|M_{a|x}S\rangle : |M_{a|x}S| = 1, 2, \dots\})).$$

One can verify the following:

1.  $\hat{E}_{a|x}|\mathbb{1}\rangle = |M_{a|x}\rangle \implies \langle \mathbb{1}|\hat{E}_{a|x}^\dagger \hat{E}_{b|y}|\mathbb{1}\rangle = \Gamma_{M_{a|x}, M_{b|y}}^\infty = p_{abxy}.$
2.  $\hat{E}_{a|x} \hat{E}_{a'|x} = \delta_{aa'} \hat{E}_{a|x}.$
3.  $[\hat{E}_{a|x}, \hat{E}_{b|y}] = 0.$

□

- $\omega_1 \geq \omega_2 \geq \dots \geq \omega := \text{quantum value.}$

# Conclusion

- $\omega_1 \geq \omega_2 \geq \dots \geq \omega := \text{quantum value.}$
- For tensor product quantum strategies, it was shown in [JNV<sup>+</sup>20] that  $\text{MIP}^* = \text{RE}$ . Perhaps  $\text{MIP}^{\text{co}} = \text{coRE}$ ?

# Conclusion

- $\omega_1 \geq \omega_2 \geq \dots \geq \omega := \text{quantum value.}$
- For tensor product quantum strategies, it was shown in [JNV<sup>+</sup>20] that  $\text{MIP}^* = \text{RE}$ . Perhaps  $\text{MIP}^{\text{co}} = \text{coRE}$ ?

# Conclusion

- $\omega_1 \geq \omega_2 \geq \dots \geq \omega :=$  quantum value.
- For tensor product quantum strategies, it was shown in [JNV<sup>+</sup>20] that  $\text{MIP}^* = \text{RE}$ . Perhaps  $\text{MIP}^{\text{co}} = \text{coRE}$ ?
- Also interesting for foundations – the existence of a quantum analogue of a joint probability measure implies the conditions of  $\mathcal{Q}^1$  [DHW14].



# Conclusion

- $\omega_1 \geq \omega_2 \geq \dots \geq \omega :=$  quantum value.
- For tensor product quantum strategies, it was shown in [JNV<sup>+</sup>20] that  $\text{MIP}^* = \text{RE}$ . Perhaps  $\text{MIP}^{\text{co}} = \text{coRE}$ ?
- Also interesting for foundations – the existence of a quantum analogue of a joint probability measure implies the conditions of  $\mathcal{Q}^1$  [DHW14].

# Conclusion

- $\omega_1 \geq \omega_2 \geq \dots \geq \omega :=$  quantum value.
- For tensor product quantum strategies, it was shown in [JNV<sup>+</sup>20] that  $\text{MIP}^* = \text{RE}$ . Perhaps  $\text{MIP}^{\text{co}} = \text{coRE}$ ?
- Also interesting for foundations – the existence of a quantum analogue of a joint probability measure implies the conditions of  $\mathcal{Q}^1$  [DHW14].
  - How do we rule out alternate versions of quantum theory?



Nicolas Brunner, Daniel Cavalcanti, Stefano Pironio, Valerio Scarani, and Stephanie Wehner.

**Bell nonlocality.**

*Reviews of Modern Physics*, 86(2):419–478, Apr 2014.



B. S. Cirel'son.

**Quantum generalizations of bell's inequality.**

*Letters in Mathematical Physics*, 4(2):93–100, March 1980.



Fay Dowker, Joe Henson, and Petros Wallden.

**A histories perspective on characterizing quantum non-locality.**

*New Journal of Physics*, 16(3):033033, mar 2014.



Zhengfeng Ji, Anand Natarajan, Thomas Vidick, John Wright, and Henry Yuen.

**Mip\*=re, 2020.**



Miguel Navascués, Stefano Pironio, and Antonio Acín.

**A convergent hierarchy of semidefinite programs characterizing the set of quantum correlations.**

*New Journal of Physics*, 10(7):073013, jul 2008.