

1 Abelian categories

1.1 Basics

In this chapter we recall some basic results.

1.1.1 Building blocks

Additive categories

Definition 1 (*Ab-enriched category*). A category \mathcal{C} is called *Ab-enriched* if it is enriched over the symmetric monoidal category $(\mathbf{Ab}, \otimes_{\mathbb{Z}})$.

Definition 2 (*additive category*). A category \mathcal{C} is additive if it has products and is **Ab**-enriched.

Proposition 3. Let \mathcal{C} be an additive category. Then the product $A \times B$ satisfies the universal property for a coproduct $A \amalg B$.

Proof. By the universal property for products, we get a unique map $i_A: A \rightarrow A \times B$ making the following diagram commute.

$$\begin{array}{ccccc} & & A & & \\ & \swarrow \text{id}_A & \downarrow \exists! i_A & \searrow 0 & \\ A & \xleftarrow{\pi_A} & A \times B & \xrightarrow{\pi_B} & B \end{array}$$

Similarly, there exists a unique map $i_B: B \rightarrow A \times B$ making the diagram

$$\begin{array}{ccccc} & & B & & \\ & \swarrow 0 & \downarrow \exists! i_B & \searrow \text{id}_B & \\ A & \xleftarrow{\pi_A} & A \times B & \xrightarrow{\pi_B} & B \end{array}$$

commute.

1 Abelian categories

Consider the following diagram.

$$\begin{array}{ccccc}
 & & A \times B & & \\
 & \swarrow \pi_A & \downarrow \exists! & \searrow \pi_B & \\
 A & \xleftarrow{\pi_A} & A \times B & \xrightarrow{\pi_B} & B
 \end{array}$$

Obviously, the identity map $\text{id}_{A \times B}$ makes this diagram commute. However

$$\pi_A \circ (i_A \circ \pi_A + i_B \circ \pi_B) = \pi_A$$

and

$$\pi_B \circ (i_A \circ \pi_A + i_B \circ \pi_B) = \pi_B,$$

so the map $i_A \circ \pi_A + i_B \circ \pi_B$ also does. Thus,

$$\pi_A \circ i_A + \pi_B \circ i_B = \text{id}_{A \times B}.$$

We now claim that $A \times B$, together with the maps i_A and i_B , satisfy the universal property for coproducts. To see this, let X be an object and $f: A \rightarrow X$ and $g: B \rightarrow X$ be morphisms.

$$\begin{array}{ccccc}
 A & \xrightarrow{i_A} & A \times B & \xleftarrow{i_B} & B \\
 & \searrow f & \downarrow \phi & \swarrow g & \\
 & & X & &
 \end{array}$$

We need to show that there exists a unique morphism ϕ making the diagram commute. Note that, by logic identical to that above, taking

$$\phi = f \circ \pi_A + g \circ \pi_B$$

makes the above diagram commute. Now suppose that there exists another $\tilde{\phi}$ making the diagram commute. Then

$$\begin{aligned}
 \tilde{\phi} &= \tilde{\phi} \circ \text{id}_{A \times B} \\
 &= \tilde{\phi} \circ (i_A \circ \pi_A + i_B \circ \pi_B) \\
 &= f \circ \pi_A + g \circ \pi_B \\
 &= \phi.
 \end{aligned}$$

□

Proposition 3 tells us that, in an additive category with products, we also have coproducts, and that these agree with the products. Dually, we could have defined an additive category to be a category with coproducts; the dual to **Proposition 3** would then have shown us that these agreed with products. Rather than preferencing either the notation

for products or the notation for coproducts, we will use a new symbol.


Definition 4 (direct sum). Let \mathcal{C} be a category with products and coproducts which agree. Then we say that \mathcal{C} has direct sums, and denote the product/coproduct of A and B by $A \oplus B$.

Definition 5 (additive functor). Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor between additive categories. We say that F is additive if for each $X, Y \in \text{Obj}(\mathcal{C})$ the map

$$\text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{\mathcal{D}}(F(X), F(Y))$$

is a homomorphism of abelian groups.

Lemma 6. Let A and B be objects in an additive category \mathcal{A} , and let X be an object in \mathcal{A} equipped with morphisms as follows,



such that the following equations hold.

$$i_A \circ \pi_A + i_B \circ \pi_B = \text{id}_X$$


$$\pi_A \circ i_B = 0 = \pi_B \circ i_A$$

$$\pi_A \circ i_A = \text{id}_A$$

$$\pi_B \circ i_B = \text{id}_B$$

Then the π s and i s exhibit $X \cong A \oplus B$.

Proof. It suffices to show that X is a product of A and B . Let $f: Z \rightarrow A$ and $g: Z \rightarrow B$. We need to show that there exists a unique $\phi: Z \rightarrow X$ making the following diagram commute.



That is, we need

$$\pi_A \circ \phi = f, \quad \pi_B \circ \phi = g.$$

1 Abelian categories

Thus, we need

$$\begin{aligned} i_A \circ f + \pi_B \circ g &= i_A \circ \pi_A \circ \phi + i_B \circ \pi_B \circ \phi \\ &= \phi. \end{aligned}$$

It is easy to check that ϕ defined in this way *does* make the diagram commute, so ϕ exists and is unique. \square

Corollary 7. Let $F: \mathcal{A} \rightarrow \mathcal{B}$ be an additive functor between additive categories. Then F preserves direct sums in the sense that there is a natural isomorphism

$$F(A \oplus B) \simeq F(A) \oplus F(B).$$

Conversely, if F preserves products,

Proof. Apply F to the data of [Lemma 6](#). \square

Pre-abelian categories

Definition 8 (kernel, cokernel). Let $f: A \rightarrow B$ be a morphism in an abelian category.

- A pair (K, ι) , where $\iota: K \rightarrow A$ is a morphism, is a kernel of f if $f \circ \iota = 0$ and for every object X and morphism $g: X \rightarrow A$ such that $f \circ g = 0$, there exists a unique morphism $\alpha: X \rightarrow K$ such that $f \circ \alpha = 0$.

$$\begin{array}{ccccc} & & X & & \\ & \swarrow \exists! & \downarrow g & \searrow 0 & \\ K & \xrightarrow{\iota} & A & \xrightarrow{f} & B \\ & \searrow & \downarrow 0 & \swarrow & \\ & & 0 & & \end{array}$$

- A pair (C, π) , where $\pi: B \rightarrow C$ is a morphism, is a cokernel of f if $\pi \circ f = 0$ and for every object Y and morphism $g: B \rightarrow Y$ such that $g \circ f = 0$, there exists a unique morphism $\beta: C \rightarrow Y$ such that $\beta \circ \pi = 0$.

$$\begin{array}{ccccc} & & 0 & & \\ & \searrow & \downarrow 0 & \swarrow & \\ A & \xrightarrow{f} & B & \xrightarrow{\pi} & C \\ & \swarrow 0 & \downarrow g & \searrow \exists! & \\ & & Y & & \end{array}$$

Here are some equivalent ways of defining the kernel of a morphism $f: A \rightarrow B$.

- The equalizer of f with the zero map.

$$\ker f = \text{eq} \longrightarrow A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{0} \end{array} B$$

- The pullback along the zero morphism

$$\begin{array}{ccc} \ker f & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ A & \xrightarrow{f} & B \end{array}$$

Definition 9 (pre-abelian category). A pre-abelian category is an additive category such that every morphism has a kernel and a cokernel.

Lemma 10. Pre-abelian categories have all finite limits and colimits.

Proof. In a pre-abelian category, the equalizer of f and g is the kernel of $f - g$, and the coequalizer is the cokernel of $f - g$. Thus, pre-abelian categories have finite products and equalizers, and finite coproducts and coequalizers. \square

Proposition 11. In a pre-abelian category, kernels are monic and cokernels are epic.

Proof. The kernel of $f: A \rightarrow B$ is the pullback of the morphism $0 \rightarrow B$, which is monic. The pullback of a monic is a monic, which gives the result. \square

Proposition 12. Let \mathcal{C} be a pre-abelian category, and let $f: A \rightarrow B$ be a morphism.

1. The morphism f is a monomorphism if and only if $\ker f = 0$.
2. The morphism f is an epimorphism if and only if $\text{coker } f = 0$.

Proof.

1. Suppose f is a monomorphism, and consider a morphism g making the following diagram commute.

$$\begin{array}{ccccc} & & Z & & \\ & & \downarrow g & \searrow 0 & \\ \ker f & \xrightarrow{\iota} & A & \xrightarrow{f} & B \end{array}$$

Because f is a monomorphism, we have $g = 0$. By the universal property for kernels, there exists a unique map $Z \rightarrow \ker f$ making the above diagram commute.

$$\begin{array}{ccccc} & & Z & & \\ & \swarrow \exists! & \downarrow g & \searrow 0 & \\ \ker f & \xrightarrow{\iota} & A & \xrightarrow{f} & B \end{array}$$

1 Abelian categories

But this remains true if we replace $\ker f$ by 0, so by definition we must have $\ker f = 0$.

Now suppose that $\ker f = 0$, and let $g: X \rightarrow A$ such that $f \circ g = 0$. The universal property guarantees us a morphism $\alpha: X \rightarrow K$ making the left-hand triangle below commute.

$$\begin{array}{ccccc} & & Z & & \\ & \swarrow \exists! & \downarrow g & \searrow 0 & \\ \ker f = 0 & \xrightarrow{\iota} & A & \xrightarrow{f} & B \end{array}$$

However, the only way the left-hand triangle can commute is if $g = 0$. Thus, f is mono.

2. Dual.

□

1.1.2 Abelian categories

We have seen that in a pre-abelian category, kernels and cokernels more or less behave as they do in the category of modules over some ring R . However, the first isomorphism theorem tells us that every submodule is the kernel of some module homomorphism (namely the cokernel), and every quotient module of some module homomorphism (namely the kernel).

Definition 13 (abelian category). A pre-abelian category is said to be abelian if every monomorphism is the kernel of its cokernel, and every epimorphism is the cokernel of its kernel.

Proposition 14. Let \mathcal{A} be an abelian category, and let $f: A \rightarrow B$ be a morphism in \mathcal{A} . Then f is an isomorphism if and only if $\ker f = 0$ and $\operatorname{coker} f = 0$.

Proof. If f is an isomorphism, then it is mono and epi, and we are done by [Proposition 12](#).

Conversely, if $\ker f = 0$ and $\operatorname{coker} f = 0$, then f is mono and epi. We need only to show that any morphism in an abelian category which is monic and epic is an isomorphism.

Let f be monic and epic. Then f is the kernel of its cokernel and the cokernel of its kernel, and both of these are zero.

$$\begin{array}{ccccccc} & & & B & & & \\ & & \swarrow \exists! f_R^{-1} & \parallel & \searrow 0 & & \\ \ker f = 0 & \longrightarrow & A & \xrightarrow{f} & B & \longrightarrow & 0 = \operatorname{coker} f \\ & \nwarrow 0 & \parallel & \nearrow \exists! f_L^{-1} & & & \\ & & A & & & & \end{array}$$

This gives us morphisms f_L^{-1} and f_R^{-1} such that

$$f \circ f_R^{-1} = \text{id}_A, \quad f_L^{-1} \circ f = \text{id}_B.$$

The usual trick shows that $f_L^{-1} = f_R^{-1}$. □

As the following proposition shows, additive functors from additive categories preserve direct sums.

Example 15. Once it is known that a category \mathcal{A} is abelian, a number of other categories are immediately known to be abelian.

- The category $\text{Ch}(\mathcal{A})$ of chain complexes in \mathcal{A} is abelian, as we will see in
- For any small category I , the category $\text{Fun}(I, \mathcal{A})$ of I -diagrams in \mathcal{A} is abelian.

1.1.3 Abelian-ness is a property, not a structure

Our definition of an abelian category, namely an **Ab**-enriched category with products, kernels, and cokernels, such that every monomorphism is the kernel of its cokernel and every epimorphism is the cokernel of its kernel, makes it sound like ordinary categories could never aspire to be abelian categories. After all, in an ordinary category, one has to provide extra structure, namely an addition law on morphisms, to get an **Ab**-enriched category.

Not so! It turns out that demanding that finite products and coproducts exist and agree is sufficient to specify the addition law.

Proposition 16. Let \mathcal{C} be a category with finite direct sums¹ (Definition 4). Then \mathcal{C} has a canonical **Ab**-enrichment.

Proof. Let $A \in \mathcal{C}$ be an object. Denote the diagonal map $A \rightarrow A \oplus A$ by Δ , and the codiagonal map $A \oplus A \rightarrow A$ by ∇ . Let $f, g: A \rightarrow B$. We define

$$f + g = \nabla \circ (f \oplus g) \circ \Delta.$$

To see that this is an **Ab**-enrichment, note the following.

1. With 0 the zero morphism, we have

$$f + 0$$

□

¹Including 0-ary direct sums, i.e. a zero object.

In fact, in an abelian category, the **Ab**-enrichment is completely determined by the direct sum structure. Thus, we have the following.

Corollary 17. Let $F: \mathcal{A} \rightarrow \mathcal{B}$ be a functor between abelian categories. Then F is additive if and only if it preserves direct sums.

1.2 Embedding theorems

In ordinary category theory, when manipulating a locally small category it is often helpful to pass through the Yoneda embedding, which gives a (fully) faithful rendition of the category under consideration in the category **Set**. One of the reasons that this is so useful is that the category **Set** has a lot of structure which can use to prove things about the subcategory of **Set** in which one lands. Having done this, one can then use the fully faithfulness to translate results to the category under consideration.

This sort of procedure, namely embedding a category which is difficult to work with into one with more desirable properties and then translating results back and forth, is very powerful. In the context of (small) Abelian categories one has essentially the best possible such embedding, known as the Freyd-Mitchell embedding theorem, which we will revisit at the end of this section. However, for now we will content ourselves with a simpler categorical embedding, one which we can work with easily.

In abelian categories, one can define kernels and cokernels slickly, as for example the equalizer along the zero morphism. However, in full generality, we can define both kernels and cokernels in any category with a zero object: f is a kernel for g if and only if the diagram below is a pullback, and a g is a cokernel of f if and only if the diagram below is a pullback.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow & & \downarrow g \\ 0 & \longrightarrow & C \end{array}$$

This means that kernels and cokernels are very general, as categories with zero objects are a dime a dozen. For example, the category **Set**_{*} of pointed sets has the singleton $\{*\}$ as a zero object. This means that one can speak of the kernel or the cokernel of a map between pointed sets, and talk about an exact sequence of such maps.

Given an abelian category \mathcal{A} , suppose one could find an embedding $\mathcal{H}: \mathcal{A} \rightarrow \mathbf{Set}_*$ which reflected exactness in the sense that if one started with a sequence $A \rightarrow B \rightarrow C$ in \mathcal{A} , mapped it into **Set**_{*} finding a sequence $\mathcal{H}(A) \rightarrow \mathcal{H}(B) \rightarrow \mathcal{H}(C)$, and found that this sequence was exact, one could be sure that $A \rightarrow B \rightarrow C$ had been exact to begin with. Then every time one wanted to check the exactness of a sequence, one could embed that sequence in **Set**_{*} using \mathcal{H} and check exactness there. Effectively, one could check exactness of a sequence in \mathcal{A} by manipulating the objects making up the sequence as if

they had elements.

Or, suppose one could find an embedding as above which sent only zero morphisms to zero morphisms. Then one could check that a diagram commutes (equivalently, that the difference between any two different ways of getting between objects is equal to 0) by checking that the image of the diagram under our functor \mathcal{H} commutes.

In fact, for \mathcal{A} small, we will find a functor which satisfies both of these and more. Then, just as we are feeling pretty good about ourselves, we will state the Freyd-Mitchell embedding theorem, which blows our pitiful result out of the water.

We now construct our functor to \mathbf{Set}_* .

Definition 18 (category of contravariant epimorphisms). Let \mathcal{A} be a small abelian category. Define a category \mathcal{A}_\leftarrow with $\text{Obj}(\mathcal{A}_\leftarrow) = \text{Obj}(\mathcal{A})$, and whose morphisms are defined by

$$\text{Hom}_{\mathcal{A}_\leftarrow}(X, Y) = \{f: Y \rightarrow X \text{ in } \mathcal{A} \mid f \text{ epimorphism}\}.$$

Note that this is indeed a category since the identity morphism is an epimorphism and epimorphisms are closed under composition.

Now for each $A \in \mathcal{A}$, we define a functor

$$\mathcal{H}_A: \mathcal{A}_\leftarrow \rightarrow \mathbf{Set}_*; \quad Z \mapsto \text{Hom}_{\mathcal{A}}(Z, A)$$

and sends a morphism $f: Z_1 \rightarrow Z_2$ in \mathcal{A}_\leftarrow (which is to say, an epimorphism $\tilde{f}: Z_2 \twoheadrightarrow Z_1$ in \mathcal{A}) to the map

$$\mathcal{H}_A(f): \text{Hom}_{\mathcal{A}}(Z_1, A) \rightarrow \text{Hom}_{\mathcal{A}}(Z_2, A); \quad (\alpha: Z_1 \rightarrow A) \mapsto (\alpha \circ \tilde{f}: Z_2 \rightarrow A).$$

Note that the distinguished point in the hom sets above is given by the zero morphism.

Definition 19 (member functor). Let \mathcal{A} be a small abelian category. We define a functor $\mathcal{M}: \mathcal{A} \rightarrow \mathbf{Set}_*$ on objects by

$$A \mapsto \mathcal{M}(A) = \text{colim } \mathcal{H}_A.$$

On morphisms, functoriality comes from the functoriality of the colimit and the co-Yoneda embedding.

Strictly speaking, we have finished our construction, but it doesn't do us much good as stated. It turns out that the sets $\mathcal{M}(\mathcal{A})$ have a much simpler interpretation.

Proposition 20. for any $A \in \mathcal{A}$, the value of the member functor $\mathcal{M}(A)$ is

$$\mathcal{M}(A) = \coprod_{X \in \mathcal{A}} \text{Hom}(X, A) / \sim,$$

1 Abelian categories

where $g \sim g'$ if there exist epimorphisms f and f' making the below diagram commute.

$$\begin{array}{ccc} Z & \xrightarrow{f} \twoheadrightarrow & X \\ f' \downarrow & & \downarrow g \\ X' & \xrightarrow{g'} & A \end{array}$$

Proof. The colimit can be computed using the following coequalizer.

$$\coprod_{\substack{f \in \text{Morph}(\mathcal{A}_{\leftarrow}) \\ \tilde{f}: Y \rightarrow X}} \text{Hom}_{\mathcal{A}}(X, A) \xrightarrow[\sim]{\text{id}} \coprod_{Z \in \text{Obj}(\mathcal{A}_{\leftarrow})} \text{Hom}(Z, A) \xrightarrow{\text{coeq}} \mathcal{M}(A)$$

On elements, we have the following.

$$\begin{array}{ccc} & \xrightarrow{\text{id}} & (g: X \rightarrow A) \\ (g: X \rightarrow A) & \searrow \scriptstyle - \circ \tilde{f} & \parallel \\ & \searrow \scriptstyle - \circ \tilde{f} & (g \circ \tilde{f}: Y \rightarrow A) \end{array}$$

Thus,

$$\mathcal{M}(A) = \coprod_{Z \in \text{Obj}(\mathcal{A}_{\leftarrow})} \text{Hom}(Z, A) / \sim,$$

where \sim is the equivalence relation generated by the relation

$$(g: X \rightarrow A) R (g': X' \rightarrow A) \iff \exists f: X' \twoheadrightarrow X \text{ such that } g' = g \circ f.$$

The above relation is reflexive and transitive, but not symmetric. The smallest equivalence relation containing it is the following.

$$(g: X \rightarrow A) \sim (g': X' \rightarrow A) \iff \exists f: Z \twoheadrightarrow X, f': Z \twoheadrightarrow X' \text{ such that } g' \circ f' = g \circ f.$$

That is, $g \sim g'$ if there exist epimorphisms making the below diagram commute.

$$\begin{array}{ccc} Z & \xrightarrow{f} \twoheadrightarrow & X \\ f' \downarrow & & \downarrow g \\ X' & \xrightarrow{g'} & A \end{array}$$

□

In summary, the elements of $\mathcal{M}(A)$ are equivalence classes of morphisms into A modulo the above relation, and for a morphism $f: A \rightarrow B$ in \mathcal{A} , $\mathcal{M}(f)$ acts on an equivalence

class $[g]$ by

$$\mathcal{M}(f): [g] \mapsto [g \circ f].$$

Lemma 21. Let $f: A \rightarrow B$ be a morphism in a small abelian category \mathcal{A} such that $f \sim 0$. Then $f = 0$.

Proof. We have that $f \sim 0$ if and only if there exists an object Z and epimorphisms making the following diagram commute.

$$\begin{array}{ccc} Z & \xrightarrow{g} & A \\ \downarrow & & \downarrow f \\ 0 & \longrightarrow & B \end{array}$$

But by the universal property for epimorphisms, $f \circ g = 0$ implies $f = 0$. \square

Now we introduce some load-lightening notation: we write $\hat{(-)} = \mathcal{M}(-)$.

Lemma 22. Let $f: A \rightarrow B$ be a morphism in a small abelian category \mathcal{A} . Then $f = 0$ if and only if $\hat{f}: \hat{A} \rightarrow \hat{B} = 0$.

Proof. Suppose that $f = 0$. Then for any $[g] \in \hat{A}$

$$\hat{f}([g]) = [g \circ 0] = [0].$$

Thus, $\hat{f}([g]) = [0]$ for all g , so $\hat{f} = 0$.

Conversely, suppose that $\hat{f} = 0$. Then in particular $\hat{f}([\text{id}_A]) = [0]$. But

$$\hat{f}([\text{id}_A]) = [\text{id}_A \circ f] = [f].$$

By Lemma 21, $[f] = [0]$ implies $f = 0$. \square

Corollary 23. A diagram commutes in \mathcal{A} if and only if its image in Set_* under \mathcal{M} commutes.

Proof. A diagram commutes in \mathcal{A} if and only if any two ways of going from one object to another agree, i.e. if the difference of any two

I actually don't see this right now. \square

Lemma 24. Let $f: A \rightarrow B$ be a morphism in a small abelian category.

- The morphism f is a monomorphism if and only if $\mathcal{M}(f)$ is an