

CHAPTER 11

Gene Therapy Clinical Trials: Past, Present and Future

Francesca Capone¹, Filomena Nappi², Maria Cristina Galli²

¹Research Coordination and Support Service, Istituto Superiore di Sanità, Roma, Italy; ²National Centre for Evaluation and Control of Medicines, Istituto Superiore di Sanità, Roma, Italy

INTRODUCTION

Gene therapy is a medical technique that involves inserting one or more corrective genes that have been designed in the laboratory into patient cells to treat a disease. To do this, medicinal products are used that are defined as gene therapy medicinal products.

On the wave of recombinant DNA pioneering studies and related ethical discussions of early 1970s, in 1972, [Friedmann and Roblin \(1972\)](#) published a paper in *Science* which cited Stanfield Roger's proposal in 1970 that 'good DNA' could be used to replace defective DNA in people with genetic disorders. They also proposed that development and clinical application of gene therapy techniques should be guided by ethical and scientific criteria so as to ensure that gene therapy is used without prejudice to patient's safety and to progress of science.

Those pioneering thoughts have evolved through the decades into regulatory science applied to clinical development of advanced therapies. Regulatory authorities all around the world have issued requirements and guidance to control the development and use of medicinal products for gene therapy approaches. An overview of the regulations that have so far been applied to clinical development and market approval of gene therapy is out the scope of this chapter; a comprehensive discussion can be found in 'Regulatory Aspects Of Gene Therapy And Cell Therapy Products: A Global Perspective' ([Galli, 2015](#)).

It took several years to move from those initial thoughts to a practical application. The advent of improved gamma-retroviral vectors (RVs) with enhanced gene transfer efficiency and stable integration into the chromosome ([Armentano et al., 1990](#); [Gilboa, 1986](#); [Miller and Buttimore, 1986](#); [Miller and Rosman, 1989](#)) and the demonstration that this procedure of gene transfer could be effectively and safely used in humans ([Kasid et al., 1990](#); [Rosenberg et al., 1990](#); [Anderson, 1992](#)) allowed the development of gene therapy clinical trials in the United States and Europe ([Wolff and Lederberg, 1994](#)).

THE MAIN TYPES OF GENE THERAPY VECTORS AND THEIR MODES OF DELIVERY

A variety of different vectors have been applied in gene therapy trials, among which two main categories can be identified: viral and nonviral vectors. Treatment strategies are based on *in vivo* administration (i.e., injecting the vector directly into a patient's body) or on *ex vivo* manipulation of human cells that are firstly genetically modified *in vitro* with the vector and then administered to the patient.

All vectors, independently from their origins or from the *in vivo* or *ex vivo* use, are constructed to contain a backbone with the function of delivering to target cells and an expression cassette with the function of coding for the transgene with the desired clinical efficacy.

Nonviral approaches include modified or naked DNA, or mRNA which can be delivered to the target cell by physical or chemical methods. Physical gene delivery strategies include microinjection, needle injection, jet injection, gene gun/DNA injection/DNA-coated particle bombardment, electroporation, sonoporation, magnetofection (Jin et al., 2014; Wang et al., 2013; Manjila et al., 2013; Nayerossadat et al., 2012).

Chemical gene delivery methods are based on calcium phosphate precipitation or on the ability of polycations, cationic lipids or lipopolyplexes to complex nucleic acids via charge interaction, leading to the formation of particles comprising the DNA/RNA (Jin et al., 2014; Wang et al., 2013; Medina-Kauwe et al., 2005).

A number of advantages can be considered by using nonviral vectors such as low immunogenicity, easy scale-up production and the ability to deliver large molecular size genes. However, delivery efficiency can be affected by an inefficient uptake as nonviral vectors can be easily degraded intracellularly or extracellularly. Other disadvantages are represented by low gene expression and low efficacy, due to gradual dilution of episomal DNA in dividing target cells (Wang et al., 2013). This approach has been used for gene therapy of cardiovascular diseases (Taniyama et al., 2012) or in solid tumours (Thaker et al., 2017; Spanggaard et al., 2017); however, results attained there have not been encouraging. Generally, nonviral vectors are not able to integrate into the cellular genome, thus persisting into the cells as episomal entities (Hitt et al., 1995). However, they can be engineered as to contain transposons or other mobile DNA elements that stably integrate the therapeutic gene into the target cell chromosomes (VandenDriessche et al., 2009; Tipanee et al., 2017). The use of nonviral vectors containing the latest generation of transposons has been proposed for clinical application mainly in cancer immunotherapy (Hudecek et al., 2017).

Viral vector delivery technologies use a modified virus as a carrier to deliver the target gene. Different viral vector families have been investigated such as Adenovirus (AdV), Adeno-Associated Virus (AAV), Herpes Simplex Virus (HSV), Retrovirus (RV) (Gammavirus and Lentivirus (LV)), Vaccinia and Baculovirus (Bessis et al., 2004; Choudhury et al., 2017; Verlengia et al., 2017).

High cellular uptake and transduction efficiency are the main advantages of this type of vectors, but there are safety concerns associated with their use. Due to their biological characteristics, viral vectors can elicit inflammatory or immune responses that may impair long-term gene expression and raise safety concerns (Manno et al., 2006) or when integrating into cell genome can result into insertional oncogenesis (Hacein-Bey-Abina et al., 2008, 2003a).

To overcome those inherent defects, viral vector design has been progressively improved so that most of the viral vectors currently in clinical use are chimeras that share very few sequences with the parental viruses. For instance, delivery efficiency or selectivity can be improved by pseudotyping viral capsids as for AAV (Hastie and Samulski, 2015), or vector-related risks can be mitigated by deleting most of unnecessary sequences as for Lentiviral Vector (LVV) (Naldini, 2011). Those vector-engineering methods aim at enhancing vector safety while retaining the desired biological characteristics for clinical efficacy.

The vast majority (81.5%) of gene therapy clinical trials to date have addressed cancer, cardiovascular disease and inherited monogenic diseases (Ginn et al., 2013): the first two because of their prevalence, impact and potentially fatal outcomes, the latter because the concept of replacing a well-defined defective gene with its correctly functioning counterpart with the aim to ensure permanent correction has an enormous appeal.

THE EARLY DAYS

The gene therapy trial history starts with two unapproved trials in the early 70s and 80s (Wolff and Lederberg, 1994). In the first trial, two young girls with arginase deficiency syndrome were treated in vivo with a wild-type Shope papilloma virus to replace the missing viral arginase enzyme. In the second trial, marrow cells from two patients with β -thalassaemia were treated ex vivo with a β -globin-containing plasmid before being transplanted again into the patients by a bone marrow (BM) transplant. Both trials were stopped. In neither case was there any real follow-up reported.

The first human gene therapy protocol for treating severe combined immunodeficiency disease (SCID) due to adenosine deaminase (ADA) deficiency with autologous lymphocytes transduced with a human ADA gene was submitted for approval in 1990 (Blaese and Anderson, 1990) and published in 1995 (Blaese et al., 1995). The trial involved ex vivo gene transfer to umbilical cord blood cells or autologous T lymphocytes of children with SCID due to mutations in the ADA-coding gene (Blaese et al., 1995; Bordignon et al., 1995; Kohn et al., 1995). In those early studies, gene transfer protocols based on a murine RV were inefficient to transduce haematopoietic stem cells (HSCs), thereby limiting extent and duration of gene transfer.

However, those ex vivo studies demonstrated for the first time the feasibility of using RV vectors for the transduction of human lymphocytes.

In the following years, clinical gene therapy approaches tried to establish what were the best vectors to use and to explore their safety, while looking for clinical experience in genetic diseases, such as cystic fibrosis, or in cancers where an effective therapy is still missing, such as glioblastoma. The most used vectors were RVs and AdVs, and most applications were in oncological patients or genetic diseases (DeWeese et al., 2001; Kirn, 2001; Thomas et al., 2003; Greco and Dachs, 2001; Kochanek, 1999; Maatta et al., 2009), but plasmids (so called DNA vaccines) were also used (Alton et al., 1999; Mor et al., 1997; Tighe et al., 1998). As stated above, viral and nonviral vectors used were only able to add a genetic sequence into target cells that bear the defective gene. Apart from genetic diseases where the gene to be targeted is well known, approaches for gene therapy of cancer mostly targeted mutated proteins that were very popular at that time, such as p53, or used a suicide strategy aiming at selectively killing cancer cells. None of those approaches proved to be effective at the clinical level, and none of the investigational medicines involved in those trials were further developed and later approved by Food and Drug Administration (FDA) or European Medicines Agency (EMA) (News, 2010). A remarkable exception is Gendicine (Editorial, 2005; Peng, 2005), an AdV vector derived from Adenovirus type 5 (AdV5) and engineered to express p53, that was market approved by China FDA for the treatment of head and neck squamous cell carcinoma. This product was totally unknown in the western world as the data were all published in Chinese journals. Recently, an update on the clinical experience of Gendicine used in association with oncological standard therapy has been published (Zhang et al., 2018).

While clinical experience with gene therapy progressed, immune responses and insertional mutagenesis emerged as main concerns, particularly in light of the poor clinical efficacy found.

In 1999, a fatal event occurred when an 18-year-old boy suffering from ornithine transcarbamylase (OTC) deficiency, an X-linked genetic disease of the liver, died when treated with gene therapy (Somia and Verma, 2000; Raper et al., 2003). The investigational medicine was a replication defective AdV vector based on AdV5 expressing human OTC. The patient was included in the very high-dose treatment group via hepatic artery. The death was attributed to unexpected and devastating inflammatory reactions to the AdV vector. In January 2000, the US FDA put that trial on hold and several other trials were also halted (Sibbald, 2001).

Meanwhile, a very promising approach to gene therapy was coming from the clinical experience with BM transplantation: ex vivo transduction of BM cells.

A clinical trial started in France, where children suffering from a rare form of X-linked severe combined immunodeficiency (SCID-X1) were successfully treated with autologous BM cells transduced with an RV vector carrying the correct gene (Cavazzana-Calvo et al., 2000).

However, in 2002, a series of severe adverse events occurred as leukaemia developed late after gene therapy (at 23–68 months posttreatment; (Cavazzana, 2014)) in 4 out of

10 children in the French trial and 1 out of 10 children in a similar UK trial. The leukaemia was fatal in one patient but was cured in others. It was later found out to be caused by RV insertional oncogenesis (Hacein-Bey-Abina et al., 2008, 2003b; Howe et al., 2008). Nevertheless, efficacy of the gene therapy treatment in those SCID-X1 children was evident, as they achieved a stable immunological T cell compartment (Cavazzana, 2014).

Thus, the first SCID-X1 gene therapy trial not only provided proof of principle on the clinical feasibility of gene therapy but also highlighted the significant toxicity associated with the Long Terminal Repeat (LTR) strong enhancer effect on the RV vector used to transduce BM cells. Leukaemic T cell clones showed vector integration near the LMO2 proto-oncogene promoter leading to its aberrant transcription and expression (Hacein-Bey-Abina et al., 2008, 2003b). A similar proto-oncogenes activation caused by Murine Leukemia Virus (MLV)-derived RV integration was also later reported for Wiskott–Aldrich and chronic granulomatous disease patients treated with a similar gene therapy approach (Stein et al., 2010; Avedillo Díez et al., 2011).

It should be noted that integration-related toxicity is also linked to other characteristics of the gene therapy medicinal product (such as target cell nature and patient genetic background), as it was shown in different Italian clinical trials: in ADA-SCID patients treated with a similar gene therapy approach (but a different transgene) (Aiuti et al., 2007, 2009; Cicalese and Aiuti, 2015; Ferrua and Aiuti, 2017), in patients with high-risk haematological diseases treated with transduced T lymphocytes (Lupo-Stanghellini et al., 2010; Cieri et al., 2014) and in autologous genetically modified cultured epidermal stem cells for gene therapy of junctional epidermolysis bullosa (De Rosa et al., 2014). Those clinical trials were carried out in the same years as the French trial; none of those patients have experienced vector-related genotoxicity.

Due to the leukaemia events, the French SCID-X1 clinical trial was halted like many others in the European Union and in the United States in early 2000s, while the cause was investigated. An in-depth analysis of RV insertion sites was initiated, resulting into a wealth of research on molecular methods to track integration and study its effects (Schmidt et al., 2001; Schröder et al., 2002; Montini et al., 2009; Biasco et al., 2011; von Kalle et al., 2014). It was shown that the RV insertion pattern is not random, and MLV-derived vectors integrate preferentially in transcriptionally active promoters and regulatory regions (Laufs et al., 2004).

Those serious adverse events had the effect to slow down gene therapy clinical development, but being the proof of principle established, they also led to a worldwide effort to improve integrating vector safety. Regulatory authorities too reacted to those events, issuing guidance documents such as the EMA

Reflection paper on management of clinical risks deriving from insertional mutagenesis (EMA, 2012; Aiuti et al., 2013).

TIME FOR SUCCESS

While the experience with SCID-X1 was also showing limited clinical efficacy because of the small proportion of engrafted genetically corrected HSCs, indeed a breakthrough was achieved when successful gene therapy pilot studies conducted in Italy at San Raffaele Telethon Institute for Gene Therapy (SR-Tiget) introduced an improved gene transfer protocol for BM haematopoietic stem and progenitor cells and the use of non-myeloablative chemotherapy regimen prior to gene therapy to prepare space for the transduced progenitors in the BM (Aiuti et al., 2002). As none of the patients received concurrent enzyme replacement therapy (ERT), the efficacy of gene therapy as single treatment could be fully assessed, exploiting at the same time the growth advantage for ADA-transduced cells. Following those initial studies, results were confirmed and extended in another clinical trial conducted at SR-Tiget. Gene therapy resulted in sustained lymphoid reconstitution with gene-corrected T cells, improvement of immune functions and effective metabolic detoxification in the absence of adverse events related to gene therapy (Aiuti et al., 2009; Cicalese et al., 2016), and most importantly, as described above, there were no severe adverse events linked to target cell transduction. Similar results were obtained in another set of patients (Gaspar et al., 2011).

The same success was met by the Italian clinical trials with the suicide gene therapy in allogeneic haematopoietic stem cell transplantation (HSCT) (Lupo-Stanghellini et al., 2010; Cieri et al., 2014).

In 2016, the EMA granted marketing approval to the medicinal products used in those pioneering clinical trials, thus recognising their efficacy and safety: Strimvelis received EU market approval for patients affected by ADA-SCID without a suitable human donor, and Zalmoxis received EU market approval for patients affected by leukaemia or lymphoma who have received a haploidentical HSCT (EMA, 2012; Aiuti et al., 2017). Up to 2013, RV and AdV vectors were the most popular gene therapy approaches, having been used in approximately two-thirds of trials performed (Ginn et al., 2013).

After the serious adverse events in the French SCID trial, there was a decrease in the proportion of trials using RV: in 2012, only 19.7% of trials compared with 22.8% in 2007 and 28% in 2004 (Ginn et al., 2013).

RV preference for integration into the first introns of genes and transcriptional start sites still is a cause of concern, if used for transducing cells with proliferation potential.

Therefore, efforts have been shifted towards the development of new vectors where safety, in terms of low genotoxicity and immunogenicity, would combine with a high efficiency of delivery. Although AdV vectors continued to be developed for anticancer treatments, LV and AAV soon emerged as promising technological tools.

LV vectors were developed to overcome the inability of RV vectors to infect nondividing cells (Naldini et al., 1996a). LV such as Human Immunodeficiency Virus (HIV) do not require cell division for gene transfer and are capable of integrative transfection of several quiescent and postmitotic cell types as well as proliferating cells (Naldini et al.,

1996a,b). Nevertheless, due to the genotoxicity risk posed by their integration ability, the use of LV vectors for gene therapy has been so far mainly limited to ex vivo transduction of haematopoietic cells, where they have shown high levels of gene transfer and transgene expression in haematopoietic progenitors (Naldini, 2009; Sessa et al., 2016; Biffi, 2017; Piguet et al., 2017). With this approach, several trials have been initiated in X-linked adrenoleukodystrophy (Cartier et al., 2009), in beta thalassaemia (Cavazzana-Calvo et al., 2010; Ferrari et al., 2017; Lidonnici and Ferrari, 2018), in Wiskott–Aldrich syndrome (Morris et al., 2017), in lysosomal storage diseases (Penati et al., 2017) showing significant clinical benefits without so far any major adverse event, although a case of major clonal expansion has been reported (Cavazzana-Calvo et al., 2010).

A very important application of gene vector technologies to ex vivo cell transduction is the development of Chimeric Antigen Receptor (CAR) T cells for cancer therapy (Hartmann et al., 2017). A comprehensive discussion of the clinical trials with CAR-T cells can be found elsewhere in this book.

Up to 2012, AdVs were the most commonly used vectors (23.3%) (Ginn et al., 2013). AdVs can carry a larger DNA load than RV, but their capacity is still too small to accommodate the genes required for certain clinical applications. The main advantages of AdV vectors are their high transduction efficiency, high gene expression level and ability to infect nondividing cells. As they remain episomal (i.e., they do not integrate into the cell genome, therefore are devoid of the genotoxicity risk), the transgene expression declines fairly rapidly in dividing cells, so that successive doses are needed to obtain a clinical effect. However, as most people are seropositive to the most common AdVs, preexisting immunity may impair clinical efficacy (Verma and Somia, 1997). Another important safety issue is the possibility of provoking severe immune and inflammatory responses as it was tragically exemplified in the case described above (Jonsson and Kreppel, 2017).

The risk of unwanted immunological responses can be reduced by choosing viruses with low immunogenicity such as AAV. In addition, nonintegrating AAV can reduce the risk of insertional oncogenicity associated with integrating viruses. Target specificity can be increased by engineering the viral capsids with different peptides to select the target cell receptor. High efficiency of in vivo gene transfers to numerous cell types, and the development of a large arsenal of viral capsids with different tropism and low innate/inflammatory responses to viral particles have made these vectors very attractive (Hastie and Samulski, 2015).

AAV offers significant advantages over AdV and RV as a platform for therapeutic gene transfer, particularly with regard to safety considerations (Dismuke et al., 2013). Within the past several years, evidence of clinical efficacy for AAV vectors in the treatment of monogenic disorders has emerged (Hastie and Samulski, 2015). Notable examples include the use of AAV for the treatment of factor IX hemophilia (Nathwani et al., 2011; Nathwani et al., 2017; George et al., 2017), cystic fibrosis (Griesenbach et al., 2015), inherited diseases of liver metabolism (Castello et al., 2016) and retinal disorders (Maguire et al., 2008; Le Meur et al., 2018; Russell et al., 2017; Trapani et al., 2015; Kumaran et al., 2018). AAV have been also used in gene therapy approaches to heart disease (Penny and Hammond, 2017).

AAV-based gene therapy medicinal products have proven so far the most successful among vectors for in vivo use, as there are two market-approved medicines in EU and in United States based on AAV.

Glybera (alipogene tiparvovec, an AAV vector expressing lipoprotein lipase) was the first gene therapy product approved by EMA in 2012 for the treatment of a very rare monogenic disease (lipoprotein lipase deficiency) (Bryant et al., 2013).

Very recently, FDA has approved LUXTURN A (voretigene neparvovec-rzyl), an AAV vector indicated for the treatment of patients with confirmed biallelic *RPE65* mutation-associated retinal dystrophy (Maguire et al., 2008; FDA).

In the vast majority of clinical trials described so far, replication defective vectors have been used to mitigate the risk of dissemination to nontarget tissues (including the germ line) or horizontal transmission. However, replication competent vectors have been used in clinical trials exploiting their oncolytic effect (Duffy et al., 2017; Hamid et al., 2017; de Graaf et al., 2018). One of them has successfully obtained a market authorisation: Imlygic (talimogene laherparepvec) was approved in the EU and United States for the treatment of unresectable melanoma. It is a HSV type 1–derived oncolytic immunotherapy designed to selectively replicate within tumours and produce granulocyte macrophage colony–stimulating factor to enhance systemic antitumour immune responses (Andtbacka et al., 2015).

The milestones in gene therapy development are summarised in Fig. 11.1.

THE SECOND INNOVATION WAVE

All vectors (viral and nonviral) used so far for gene therapy clinical approaches are only able to add a genetic sequence into target cells that bear the defective gene.

A recent significant advance is targeted genome editing (Mussolino et al., 2017). This technology allows to correct a specific sequence within a genomic DNA region. Gene editing has a great potential for gene therapy, although evidence is still accumulating on its precision (i.e., off-target mutations) and ethical concerns exists due to potential genetic changes in germline cells.

Three important strategies have been developed based on zinc-finger nuclease (ZFN), transcription activator–like effector nucleases (TALENs) and clustered regularly interspaced short palindromic repeat (CRISPR)/CRISPR-associated protein 9 (CRISPR/Cas9) (Sander and Joung, 2014). A more detailed discussion on those technologies can be found elsewhere in this book.

The first clinical trial using a nuclease for targeted gene editing was conducted in HIV patients with autologous CD4–enriched T cells modified at the CCR5 gene locus by ZFN (Tebas et al., 2014). This strategy was also used for the treatment of cervical cancer, hemophilia B and mucopolysaccharidoses (Cornu et al., 2017).

The TALEN technology was used for the treatment of an 11-month-old baby with B acute lymphoblastic leukaemia. This was the first patient treated with TALEN engineered




Figure 11.1 Overview of the gene therapy medicinal products that have been granted market authorisation.

CAR19T Cells (Qasim et al., 2015). TALENs technology has also been used for the correction of COL7A1 gene involved in epidermolysis bullosa (Osborn et al., 2013).

Many potential applications of gene editing with CRISPR/Cas9 are currently under development. A first clinical trial using CRISPR/Cas9 technology was authorised in China in lung cancer patients and others similar in prostate, bladder and kidney cancer were announced (Cyranoski, 2016). Other clinical trials have been started in other regions of the world (Ginn et al., 2018).

PERSPECTIVES

Gene therapy for ADA-SCID, AAV gene therapy for eye disease and LV vector platform is the most significant success stories to date, but progress in many other areas has been significant and will hopefully result in additional approvals in the near future.

After decades of clinical trials, after the first frustrating unsuccessful trials and after many clinical successes over the past recent years, gene therapy has been proven to provide long-lasting benefits to patients, thus resulting into an important tool for patient treatment.

Due to the novelty of development approach as compared with traditional pharmaceuticals and the challenges thus posed to all actors in the field (scientists, industry, regulators) (CAT, 2010), gene therapy has also been pivotal in establishing a new paradigm for collaboration between academia, industry and regulators that has been instrumental in driving the field ahead (Maciulaitis et al., 2012).

Nowadays some gene therapy medicinal products are approved for the market (Table 11.1): Glybera, Strimvelis, Imlygic, Zalmoxis, YESCARTA, Kymriah, LUXTURNAL,

Table 11.1 Approved Gene Therapy Medicinal Products.

GTMP Commercial Name (INN)	Type of GTMP	MA Holder	Clinical Indication	MA Date	MA Authority	Note
Gendicine	Ad-p53	Shenzhen Sibiono GeneTech	Head and neck squamous cell carcinoma	2003	China FDA	
Glybera (alipogene tiparvovec)	AAV LPLS447X	UniQure	Lipoprotein lipase deficiency	2012	EMA	Withdrawn from EU market
Imlygic (talimogene laherparepvec)	OncolyticGM-CSF-HSV-1	Amgen	Metastatic melanoma	2015 2015	EMA US FDA	
Strimvelis	Autologous CD34+ cells genetically modified with RV-ADA	GSK	ADA-SCID	2016	EMA	
Zalmoxis	Allogeneic T cells genetically modified with RV-ΔLNGFR+ HSV-TK Mut2	MolMed		2016	EMA	
KYMRIAH (tisagenlecleucel)	Autologous CD19-CAR-T cells	Novartis	Relapsed/refractory large B-cell lymphomaRelapsed/refractory B-cell precursor acute lymphoblastic leukaemia	2017 2018	US FDA EMA	
LUXTURN A (voretigene neparvovec-rzyl)	Adeno-associated viral vector-hRPE65	Spark Therapeutics	Biallelic RPE65 mutation-associated retinal dystrophy	2017 2018	US FDA EMA	
YESCARTA (axicabtagene ciloleucel)	Autologous CD19-CAR-T cells	Kite Pharma	Relapsed or refractory large B-cell lymphoma	2017 2018	US FDA EMA	

AAV, adeno-associated virus; ADA, adenosine deaminase; EMA, European Medicines Agency; EU, European Union; FDA, Food and Drug Administration; GTMP, Gene Therapy Medicinal Product; MA, Marketing Authorisation; SCID, severe combined immunodeficiency disease.

Zynteglo/ *Autologous CD34+ cells encoding βA-T87Q-globin gene*/Bluebird bio/beta-Thalassemia/2019/EMA

ZOLGENSMA (onasemnogene abeparvovec-xioi)/Adeno-associated viral vector/AveXis, Inc/spinal muscular atrophy (SMA)/2019/FDA

Zynteglo in EU (EMA); Imlygic, YESCARTA, Kymriah, LUXTURN A, Zolgensma in the United States (FDA) (Table 11.1).

Since the first clinical trials of the 1990s, gene therapy has moved a long way from the investigational stage to being a part of evidence-based medicine, fulfilling the promises that were raised at its dawn.

REFERENCES

- Aiuti A, Slavin S, Aker M, Ficara F, Deola S, Mortellaro A, Morecki S, Andolfi G, Tabucchi A, Carlucci F, Marinello E, Cattaneo F, Vai S, Servida P, Miniero R, Roncarolo MG, Bordignon C. Correction of ADA SCID by stem cell gene therapy combined with nonmyeloablative conditioning. *Science* 2002;296(5677):2410–3.
- Aiuti A, Cassani B, Andolfi G, Mirolo M, Biasco L, Recchia A, Urbinati F, Valacca C, Scaramuzza S, Aker M, Slavin S, Cazzola M, Sartori D, Ambrosi A, Di Serio C, Roncarolo MG, Mavilio F, Bordignon C. Multilineage hematopoietic reconstitution without clonal selection in ADA-SCID patients treated with stem cell gene therapy. *J Clin Invest* 2007;117(8):2233–40.
- Aiuti A, Cattaneo F, Galimberti S, Benninghoff U, Cassani B, Callegaro L, Scaramuzza S, Andolfi G, Mirolo M, Brigida I, Tabucchi A, Carlucci F, Eibl M, Aker M, Slavin S, Al-Mousa H, Al Ghonaium A, Ferster A, Duppenhaler A, Notarangelo L, Wintergerst U, Buckley RH, Bregni M, Marktel S, Valsecchi MG, Rossi P, Ciceri F, Miniero R, Bordignon C, Roncarolo MG. Gene therapy for immunodeficiency due to adenosine deaminase deficiency. *N Engl J Med* 2009;360(5):447–58.
- Aiuti A, Cossu G, de Felipe P, Galli MC, Narayanan G, Renner M, Stahlbom A, Schneider CK, Voltz-Girolt C. The committee for advanced therapies' of the European Medicines Agency reflection paper on management of clinical risks deriving from insertional mutagenesis. *Hum Gene Ther Clin Dev* 2013;24(2):47–54.
- Aiuti A, Roncarolo MG, Naldini L. Gene therapy for ADA-SCID, the first marketing approval of an ex vivo gene therapy in Europe: paving the road for the next generation of advanced therapy medicinal products. *EMBO Mol Med* 2017;9(6):737–40.
- Alton EW, Stern M, Farley R, Jaffe A, Chadwick SL, Phillips J, Davies J, Smith SN, Browning J, Davies MG, Hodson ME, Durham SR, Li D, Jeffery PK, Scallan M, Balfour R, Eastman SJ, Cheng SH, Smith AE, Meeker D, Geddes DM. Cationic lipid-mediated CFTR gene transfer to the lungs and nose of patients with cystic fibrosis: a double-blind placebo-controlled trial. *Lancet* 1999;353:947–54.
- Anderson WF. Human gene therapy. *Science* May 8, 1992;256(5058):808–13.
- Andtbacka RH, Kaufman HL, Collichio F, Amatruda T, Senzer N, Chesney J, Delman KA, Spitler LE, Puzanov I, Agarwala SS, Milhem M, Cranmer L, Curti B, Lewis K, Ross M, Guthrie T, Linette GP, Daniels GA, Harrington K, Middleton MR, Miller Jr WH, Zager JS, Ye Y, Yao B, Li A, Doleman S, VanderWalde A, Gansert J, Coffin RS. Talimogene laherparepvec improves durable response rate in patients with advanced melanoma. *J Clin Oncol* 2015;33:2780–8.
- Armentano D, Thompson AR, Darlington G, Woo SL. Expression of human factor IX in rabbit hepatocytes by retrovirus-mediated gene transfer: potential for gene therapy of hemophilia B. *Proc Natl Acad Sci USA* 1990;87(16):6141–5.
- Avedillo Díez I, Zychlinski D, Coci EG, Galla M, Modlich U, Dewey RA, Schwarzer A, Maetzig T, Mpofo N, Jaekel E, Boztug K, Baum C, Klein C, Schambach A. Development of novel efficient SIN vectors with improved safety features for Wiskott-Aldrich syndrome stem cell based gene therapy. *Mol Pharm* 2011;8(5):1525–37.
- Bessis N, GarciaCozar FJ, Boissier M-C. Immune responses to gene therapy vectors: influence on vector function and effector mechanisms. *Gene Ther* 2004;11(Suppl. 1):10–7.
- Biasco L, Ambrosi A, Pellin D, Bartholomae C, Brigida I, Roncarolo MG, Di Serio C, von Kalle C, Schmidt M, Aiuti A. Integration profile of retroviral vector in gene therapy treated patients is cell-specific according to gene expression and chromatin conformation of target cell. *EMBO Mol Med* 2011;3(2):89–101.

- Biffi A. Hematopoietic gene therapies for metabolic and neurologic diseases. *Hematol Oncol Clin N Am* October 2017;31(5):869–81.
- Blaese R, Anderson F. The ADA human gene therapy clinical protocol. *Hum Gene Ther* 1990;1:327–9.
- Blaese RM, Culver KW, Miller AD, Carter CS, Fleisher T, Clerici M, Shearer G, Chang L, Chiang Y, Tolstoshev P, Greenblatt JJ, Rosenberg SA, Klein H, Berger M, Mullen CA, Ramsey WJ, Muul L, Morgan RA, Anderson WF. T lymphocyte directed gene therapy for ADA-SCID: initial trial results after 4 years. *Science* 1995;270:475–80.
- Bordignon C, Notarangelo LD, Nobili N, Ferrari G, Casorati G, Panina P, Mazzolari E, Maggioni D, Rossi C, Servida P, Ugazio AG, Mavilio F. Gene therapy in peripheral blood lymphocytes and bone marrow for ADA-immunodeficient patients. *Science* 1995;270:470–5.
- Bryant LM, Christopher DM, Giles AR, Hinderer C, Rodriguez JL, Smith JB, Traxler EA, Tycko J, Wojno AP, Wilson JM. Lessons learned from the clinical development and market authorization of Glybera. *Hum Gene Ther Clin Dev* 2013;24(2):55–64.
- Cartier N, Hacein-Bey-Abina S, Bartholomae CC, Veres G, Schmidt M, Kutschera I, Vidaud M, Abel U, Dal-Cortivo L, Caccavelli L, Mahlaoui N, Kiermer V, Mittelstaedt D, Bellesme C, Lahlou N, Lefrère F, Blanco S, Audit M, Payen E, Leboulch P, l'Homme B, Bougnères P, Von Kalle C, Fischer A, Cavazzana-Calvo M, Aubourg P. Hematopoietic stem cell gene therapy with a lentiviral vector in X-linked adrenoleukodystrophy. *Science* 2009;326(5954):818–23.
- Castello R, Borzone R, D'Aria S, Annunziata P, Piccolo P, Brunetti-Pierri N. Helper-dependent adenoviral vectors for liver-directed gene therapy of primary hyperoxaluria type 1. *Gene Ther* 2016;23(2):129–34.
- Committee for Advanced Therapy (CAT), CAT Scientific Secretariat, Schneider CK, Salmikangas P, Jilma B, Flamion B, Todorova LR, Paphitou A, et al. Challenges with advanced therapy medicinal products and how to meet them. *Nat Rev Drug Discov* 2010;9:195–201.
- Cavazzana-Calvo M, Payen E, Negre O, Wang G, Hehir K, Fusil F, Down J, Denaro M, Brady T, Westerman K, Cavallese R, Gillet-Legrand B, Caccavelli L, Sgarra R, Maouche-Chrétien L, Bernaudin F, Girot R, Dorazio R, Mulder GJ, Polack A, Bank A, Soulier J, Larghero J, Kabbara N, Dalle B, Gourmel B, Socie G, Chrétien S, Cartier N, Aubourg P, Fischer A, Cornetta K, Galacteros F, Beuzard Y, Gluckman E, Bushman F, Hacein-Bey-Abina S, Leboulch P. Transfusion independence and HMGA2 activation after gene therapy of human beta-thalassemia. *Nature* 2010;467:318–22.
- Cavazzana M. Hematopoietic stem cell gene therapy: progress on the clinical front. *Hum Gene Ther* 2014;25(3):165–70.
- Cavazzana-Calvo M, Hacein-Bey S, de Saint Basile G, Gross F, Yvon E, Nusbaum P, Selz F, Hue C, Certain S, Casanova JL, Bousso P, Deist FL, Fischer A. Gene therapy of human severe combined immunodeficiency (SCID)-X1 disease. *Science* 2000;288(5466):669–72.
- Choudhury SR, Hudry E, Maguire CA, Sena-Esteves M, Breakefield XO, Grandi P. Viral vectors for therapy of neurologic diseases. *Neuropharmacology* 2017;120:63–80.
- Cicalese MP, Aiuti A. *Hum Gene Ther* 2015;26(4):210–9.
- Cicalese MP, Ferrua F, Castagnaro L, Pajno R, Barzaghi F, Giannelli S, Dionisio F, Brigida I, Bonopane M, Casiraghi M, Tabucchi A, Carlucci F, Grunebaum E, Adeli M, Bredius RG, Puck JM, Stepensky P, Tezcan I, Rolfe K, De Boever E, Reinhardt RR, Appleby J, Ciceri F, Roncarolo MG, Aiuti A. Update on the safety and efficacy of retroviral gene therapy for immunodeficiency due to adenosine deaminase deficiency. *Blood* July 7, 2016;128(1):45–54.
- Cieri N, Mastaglio S, Oliveira G, Casucci M, Bondanza A, Bonini C. Adoptive immunotherapy with genetically modified lymphocytes in allogeneic stem cell transplantation. *Immunol Rev* 2014;257(1):165–80.
- Cornu T, Mussolino C, Cathomen T. Refining strategies to translate genome editing to the clinic. *Nat Med* April 3, 2017;23(4):415–23.
- Cyranoski D. Chinese scientists to pioneer first human CRISPR trial. *Nature* 2016;535(7613):476–7.
- de Graaf JF, de Vor L, Fouchier RAM, van den Hoogen BG. Armed oncolytic viruses: a kick-start for anti-tumor immunity. *Cytokine Growth Factor Rev* 2018. <https://doi.org/10.1016/j.cytogfr.2018.03.006>.
- De Rosa L, Carulli S, Cocchiarella F, Quaglino D, Enzo E, Franchini E, Giannetti A, De Santis G, Recchia A, Pellegrini G, De Luca M. Long-term stability and safety of transgenic cultured epidermal stem cells in gene therapy of junctional epidermolysis bullosa. *Stem Cell Rep* 2014;2(1):1–8.

- DeWeese TL, van der Poel H, Li S, Mikhak B, Drew R, Goemann M, Hamper U, DeJong R, Detorie N, Rodriguez R, Haulk T, DeMarzo AM, Piantadosi S, Yu DC, Chen Y, Henderson DR, Carducci MA, Nelson WG, Simons JW. A phase I trial of CV706, a replication-competent, PSA selective oncolytic adenovirus, for the treatment of locally recurrent prostate cancer following radiation therapy. *Cancer Res* 2001;61:7464–72.
- Dismuke DJ, Tenenbaum L, Samulski RJ. Biosafety of recombinant adeno-associated virus vectors. *Curr Gene Ther* 2013;13:434–52.
- Duffy MR, Fisher KD, Seymour LW. Making oncolytic virotherapy a clinical reality: the European contribution. *Hum Gene Ther* 2017;28:1033–46.
- Editorial. *Hum Gene Ther* 2005;14:1014.
- EMA/CAT/190186/2012, Committee for Advanced Therapies, EMA website.
- FDA website, Approved cellular and gene therapy products.
- Ferrari G, Cavazzana M, Mavilio F. Gene therapy approaches to hemoglobinopathies. *Hematol Oncol Clin N Am* 2017;31(5):835–52.
- Ferrua F, Aiuti A. Twenty-five years of gene therapy for ADA-SCID: from bubble babies to an approved drug. *Hum Gene Ther* 2017;28(11):972–81.
- Friedmann T, Roblin R. Gene therapy for human genetic disease? *Science* 1972;175(4025):949–55.
- Regulatory aspects of gene therapy and cell therapy products. A global perspective. In: Galli MC, Serabian M, editors. *Advances in experimental medicine and Biology*, vol. 871. Springer; 2015.
- Gaspar HB, Cooray S, Gilmour KC, Parsley KL, Zhang F, Adams S, Bjorkegren E, Bayford J, Brown L, Davies EG, Veys P, Fairbanks L, Bordon V, Petropoulou T, Kinnon C, Thrasher AJ. Hematopoietic stem cell gene therapy for adenosine deaminase deficient severe combined immunodeficiency leads to long term immunological recovery and metabolic correction. *Sci Transl Med* 2011;3:97ra80.
- George LA, Sullivan SK, Giermasz A, Rasko JEJ, Samelson-Jones BJ, Ducore J, Cuker A, Sullivan LM, Majumdar S, Teitel J, McGuinn CE, Ragni MV, Luk AY, Hui D, Wright JF, Chen Y, Liu Y, Wachtel K, Winters A, Tiefenbacher S, Arruda VR, van der Loo JCM, Zelenia O, Takefman D, Carr ME, Couto LB, Anguela XM, High KA. Hemophilia B gene therapy with a high-specific-activity factor IX variant. *N Engl J Med* 2017;377(23):2215–27.
- Gilboa E. Retrovirus vectors and their uses in molecular biology. *Bioessays* 1986;5(6):252–7.
- Ginn SL, Alexander IE, Edelstein ML, Abedi MR, Wixon J. Gene therapy clinical trials worldwide to 2012—an update. *J Gene Med* 2013;15(2):65–77.
- Ginn SL, Amaya AK, Alexander IE, Edelstein M, Abedi MR. Gene therapy clinical trials worldwide to 2017—an update. *J Gene Med* March 25, 2018:e3015. [Epub ahead of print] <https://doi.org/10.1002/jgm.3015>.
- Greco O, Dachs GU. Gene directed enzyme/prodrug therapy of cancer: historical appraisal and future perspectives. *J Cell Physiol* 2001;187:22–35.
- Griesenbach U, Pytel KM, Alton EFWF. Cystic fibrosis gene therapy in the UK and elsewhere. *Hum Gene Ther* 2015;26:266–75.
- Hacein-Bey-Abina S, von Kalle C, Schmidt M, Le Deist F, Wulffraat N, McIntyre E, Radford I, Villeval JL, Fraser CC, Cavazzana-Calvo M, Fischer A. A serious adverse event after successful gene therapy for X-linked severe combined immunodeficiency. *N Engl J Med* 2003;348(3):255–6.
- Hacein-Bey-Abina S, Von Kalle C, Schmidt M, McCormack MP, Wulffraat N, Leboulch P, Lim A, Osborne CS, Pawliuk R, Morillon E, Sorensen R, Forster A, Fraser P, Cohen JL, de Saint Basile G, Alexander I, Wintergerst U, Frebourg T, Aurias A, Stoppa-Lyonnet D, Romana S, Radford-Weiss I, Gross F, Valensi F, Delabesse E, Macintyre E, Sigaux F, Soulier J, Leiva LE, Wissler M, Prinz C, Rabbitts TH, Le Deist F, Fischer A, Cavazzana-Calvo M. LMO2-associated clonal T cell proliferation in two patients after gene therapy for SCID-X1. *Science* 2003;302(5644):415–9.
- Hacein-Bey-Abina S, Garrigue A, Wang GP, Soulier J, Lim A, Morillon E, Clappier E, Caccavelli L, Delabesse E, Beldjord K, Asnafi V, MacIntyre E, Dal Cortivo L, Radford I, Brousse N, Sigaux F, Moshous D, Hauer J, Borkhardt A, Belohradsky BH, Wintergerst U, Velez MC, Leiva L, Sorensen R, Wulffraat N, Blanche S, Bushman FD, Fischer A, Cavazzana-Calvo M. Insertional oncogenesis in 4 patients after retrovirus-mediated gene therapy of SCID-X1. *J Clin Invest* 2008;118(9):3132–42.
- Hamid O, Hoffner B, Gasal E, Hong J, Carvajal RD. Oncolytic immunotherapy: unlocking the potential of viruses to help target cancer. *Cancer Immunol Immunother* 2017;66(10):1249–64.

- Hartmann J, Schübler-Lenz M, Bondanza A, Buchholz CJ. Clinical development of CAR T cells—challenges and opportunities in translating innovative treatment concepts. *EMBO Mol Med* 2017;9:1183–97.
- Hastie E, Samulski RJ. Adeno-associated virus at 50: a golden anniversary of discovery, research, and gene therapy success—a personal perspective. *Hum Gene Ther* 2015;26:257–65.
- Hitt M, Bett AJ, Addison CL, Prevec L, Graham FL. Techniques for human adenovirus vector construction and characterization. In: *Methods in molecular genetics*, vol. 7. Academic Press; 1995.
- Howe SJ, Mansour MR, Schwarzwaelder K, Bartholomae C, Hubank M, Kempinski H, Brugman MH, Pike-Overzet K, Chatters SJ, de Ridder D, Gilmour KC, Adams S, Thornhill SI, Parsley KL, Staal FJ, Gale RE, Linch DC, Bayford J, Brown L, Quaye M, Kinnon C, Ancliff P, Webb DK, Schmidt M, von Kalle C, Gaspar HB, Thrasher AJ. Insertional mutagenesis combined with acquired somatic mutations causes leukemogenesis following gene therapy of SCID-X1 patients. *J Clin Invest* 2008;118(9):3143–50.
- Hudecek M, Izsvák Z, Johnen S, Renner M, Thumann G, Ivics Z. Going non-viral: the Sleeping Beauty transposon system breaks on through to the clinical side. *Crit Rev Biochem Mol Biol* 2017;52(4):355–80.
- Jin L, Zeng X, Liu M, Deng Y, He N. Current progress in gene delivery technology based on chemical methods and nanocarriers. *Theranostics* 2014;4(3):240–55.
- Jonsson F, Kreppel F. Barriers to systemic application of virus-based vectors in gene therapy: lessons from adenovirus type 5. *Virus Genes* 2017;53(5):692–9.
- Kasid A, Morecki S, Aebersold P, Cornetta K, Culver K, Freeman S, Director E, Lotze MT, Blaese RM, Anderson WF, et al. Human gene transfer: characterization of human tumor-infiltrating lymphocytes as vehicles for retroviral-mediated gene transfer in man. *Proc Natl Acad Sci USA* 1990;87(1):473–7.
- Kirn D. Clinical research results with dl1520 (Onyx-015), a replication-selective adenovirus for the treatment of cancer: what have we learned? *Gene Ther* 2001;8:89–98.
- Kochanek S. High-capacity adenoviral vectors for gene transfer and somatic gene therapy. *Hum Gene Ther* 1999;10(15):2451–9.
- Kohn DB, Weinberg KI, Nolta JA, et al. Engraftment of gene-modified umbilical cord blood cells in neonates with adenosine deaminase deficiency. *Nat Med* 1995;1:1017–23.
- Kumaran N, Michaelides M, Smith AJ, Ali RR, Bainbridge JWB. Retinal gene therapy. *Br Med Bull* March 1, 2018. [Epub ahead of print] <https://doi.org/10.1093/bmb/ldy005>.
- Laufs S, Nagy KZ, Giordano FA, Hotz-Wagenblatt A, Zeller WJ, Fruehauf S. *Mol Ther* 2004;10(5):874–81.
- Le Meur C, Lebranchu P, Billaud F, Adjali O, Schmitt S, Béziau S, Péréon Y, Valabregue R, Ivan C, Darmon C, Moullier P, Rolling F, Weber M. Safety and long-term efficacy of AAV4 gene therapy in patients with RPE65 Leber congenital amaurosis. *Mol Ther* 2018;26(1):256–68.
- Lidonnici MR, Ferrari G. Gene therapy and gene editing strategies for hemoglobinopathies. *Blood Cells Mol Dis* January 3, 2018;(17). 30429–1. [Epub ahead of print] <https://doi.org/10.1016/j.bcmd.2017.12.001>. pii:S1079–9796.
- Lupo-Stanghellini MT, Provasi E, Bondanza A, Ciceri F, Bordignon C, Bonini C. Clinical impact of suicide gene therapy in allogeneic hematopoietic stem cell transplantation. *Hum Gene Ther* 2010;21(3):241–50.
- Maatta AM, Samaranyake H, Pikkarainen J, Wirth T, Ylä-Herttuala S. Adenovirus mediated herpes simplex virus-thymidine kinase/ganciclovir gene therapy for resectable malignant glioma. *Curr Gene Ther* 2009;9(5):356–67.
- Maciulaitis R, D'Apote L, Buchanan A, Pioppo L, Schneider CK. Clinical development of advanced therapy medicinal products in Europe: evidence that regulators must be proactive. *Mol Ther* 2012;20(3):479–82.
- Maguire AM, Simonelli F, Pierce EA, Pugh Jr EN, Mingozzi F, Bennicelli J, Banfi S, Marshall KA, Testa F, Surace EM, Rossi S, Lyubarsky A, Arruda VR, Konkle B, Stone E, Sun J, Jacobs J, Dell'Osso L, Hertle R, Ma JX, Redmond TM, Zhu X, Hauck B, Zelenia O, Shindler KS, Maguire MG, Wright JF, Volpe NJ, McDonnell JW, Auricchio A, High KA, Bennett J. Safety and efficacy of gene transfer for Leber's congenital amaurosis. *N Engl J Med* 2008;358:2240–8.
- Manjila SB, Baby JN, Bijin EN, Constantine I, Pramod K, Valsalakumari J. Novel gene delivery systems. *Int J Pharm Investig* 2013;3(1):1–7.

- Manno CS, Pierce GF, Arruda VR, Glader B, Ragni M, Rasko JJ, Ozelo MC, Hoots K, Blatt P, Konkle B, Dake M, Kaye R, Razavi M, Zajko A, Zehnder J, Rustagi PK, Nakai H, Chew A, Leonard D, Wright JF, Lessard RR, Sommer JM, Tigges M, Sabatino D, Luk A, Jiang H, Mingozzi F, Couto L, Ertl HC, High KA, Kay MA. Successful transduction of liver in hemophilia by AAV-Factor IX and limitations imposed by the host immune response. *Nat Med* 2006;12:342–7.
- Medina-Kauwe LK, Xie J, Hamm-Alvarez S. Intracellular trafficking of nonviral vectors. *Gene Ther* 2005;12:1734–51.
- Miller AD, Buttimore C. Redesign of retrovirus packaging cell lines to avoid recombination leading to helper virus production. *Mol Cell Biol* 1986;6(8):2895–902.
- Miller AD, Rosman GJ. Improved retroviral vectors for gene transfer and expression. *BioTechniques* 1989;7:980–90.
- Montini E, Cesana D, Schmidt M, Sanvito F, Bartholomae CC, Ranzani M, Benedicenti F, Sergi LS, Ambrosi A, Ponzoni M, Doglioni C, Di Serio C, von Kalle C, Naldini L. The genotoxic potential of retroviral vectors is strongly modulated by vector design and integration site selection in a mouse model of HSC gene therapy. *J Clin Invest* 2009;119:964–75.
- Mor G, Singla M, Steinberg AD, Hoffman SL, Okuda K, Klinman DM. Do DNA vaccines induce autoimmune disease? *Hum Gene Ther* 1997;8:293–300.
- Morris EC, Fox T, Chakraverty R, Tendeiro R, Snell K, Rivat C, Grace S, Gilmour K, Workman S, Buckland K, Butler K, Chee R, Salama AD, Ibrahim H, Hara H, Duret C, Mavilio F, Male F, Bushman FD, Galy A, Burns SO, Gaspar HB, Thrasher AJ. Gene therapy for Wiskott–Aldrich syndrome in a severely affected adult. *Blood* 2017;130(11):1327–35.
- Mussolino C, Alzubi J, Pennucci V, Turchiano G, Cathomen T. Genome and epigenome editing to treat disorders of the hematopoietic system. *Hum Gene Ther* 2017;28(11):1105–15.
- Naldini L, Blömer U, Gally P, Ory D, Mulligan R, Gage FH, Verma IM, Trono D. In vivo gene delivery and stable transduction of nondividing cells by a lentiviral vector. *Science* 1996;272(5259):263–7.
- Naldini L, Blömer U, Gage FH, Trono D, Verma IM. Efficient transfer, integration, and sustained long-term expression of the transgene in adult rat brains injected with a lentiviral vector. *Proc Natl Acad Sci USA* 1996;93(21):11382–8.
- Naldini L. A comeback for gene therapy. *Science* 2009;326(6).
- Naldini L. Ex vivo gene transfer and correction for cell-based therapies. *Nat Rev Genet* 2011;12:301–15.
- Nathwani AC, Tuddenham EG, Rangarajan S, Rosales C, McIntosh J, Linch DC, Chowdary P, Riddell A, Pie AJ, Harrington C, O’Beirne J, Smith K, Pasi J, Glader B, Rustagi P, Ng CY, Kay MA, Zhou J, Spence Y, Morton CL, Allay J, Coleman J, Sleep S, Cunningham JM, Srivastava D, Basner-Tschakarjan E, Mingozzi F, High KA, Gray JT, Reiss UM, Nienhuis AW, Davidoff AM. Adenovirus-associated virus vector mediated gene transfer in hemophilia B. *N Engl J Med* 2011;365:2357–65.
- Nathwani AC, Davidoff AM, Tuddenham EGD. Advances in gene therapy for hemophilia. *Hum Gene Ther* 2017;28(11).
- Nayerossadat N, Maedeh T, Ali PA. Viral and nonviral delivery systems for gene delivery. *Adv Biomed Res* 2012;1:27.
- News. *Nat Biotechnol* 2010;28(3):183.
- Osborn MJ, Starker CG, McElroy AN, Webber BR, Riddle MJ, Xia L, DeFeo AP, Gabriel R, Schmidt M, von Kalle C, Carlson DF, Maeder ML, Joung JK, Wagner JE, Voytas DF, Blazar BR, Tolar J. TALEN-based gene correction for epidermolysis bullosa. *Mol Ther* 2013;21(6):1151–9.
- Penati R, Fumagalli F, Calbi V, Bernardo ME, Aiuti A. Gene therapy for lysosomal storage disorders: recent advances for metachromatic leukodystrophy and mucopolysaccharidosis I. *J Inherit Metab Dis* July 2017;40(4):543–54.
- Peng Z. *Hum Gene Ther* 2005;16:1016–27.
- Penny WF, Hammond HK. Randomized clinical trials of gene transfer for heart failure with reduced ejection fraction. *Hum Gene Ther* 2017;28(5):378–84.
- Piguet F, Alves S, Cartier N. Clinical gene therapy for neurodegenerative diseases: past, present, and future. *Hum Gene Ther* 2017;28(11):988–1003.
- Qasim W, Amrolia PJ, Samarasinghe S, Ghorashian S, Zhan H, Stafford S, et al. First clinical application of talen engineered universal CAR19 T cells in B-ALL. *Blood* 2015;126:2046.

- Raper SE, Chirmule N, Lee FS, Wivel NA, Bagg A, Gao GP, Wilson JM, Batshaw ML. Fatal systemic inflammatory response syndrome in a ornithine transcarbamylase deficient patient following adenoviral gene transfer. *Mol Genet Metab* 2003;80(1–2):148–58.
- Rosenberg SA, Blaese RM, Anderson WF. The N2–TIL human gene transfer clinical protocol. *Hum Gene Ther* 1990;1(1):73–92.
- Russell S, Bennett J, Wellman JA, Chung DC, Yu ZF, Tillman A, Wittes J, Pappas J, Elci O, McCague S, Cross D, Marshall KA, Walshire J, Kehoe TL, Reichert H, Davis M, Raffini L, George LA, Hudson FP, Dingfield L, Zhu X, Haller JA, Sohn EH, Mahajan VB, Pfeifer W, Weckmann M, Johnson C, Gewaily D, Drack A, Stone E, Wachtel K, Simonelli F, Leroy BP, Wright JF, High KA, Maguire AM. Efficacy and safety of voretigene neparvovec (AAV2-hRPE65v2) in patients with RPE65-mediated inherited retinal dystrophy: a randomised, controlled, open-label, phase 3 trial. *Lancet* 2017;390(10097):849–60.
- Sander J, Joung J. CRISPR–Cas systems for editing, regulating and targeting genomes. *Nat Biotechnol* 2014;32(4):347–55.
- Schmidt M, Hoffmann G, Wissler M, Lemke N, Müssig A, Glimm H, Williams DA, Ragg S, Hesemann CU, von Kalle C. Detection and direct genomic sequencing of multiple rare unknown flanking DNA in highly complex samples. *Hum Gene Ther* 2001;12:743–9.
- Schröder AR, Shinn P, Chen H, Berry C, Ecker JR, Bushman F. HIV-1 integration in the human genome favors active genes and local hotspots. *Cell* 2002;110(4):521–9.
- Sessa M, Liorioli L, Fumagalli F, Acquati S, Redaelli D, Baldoli C, Canale S, Lopez ID, Morena F, Calabria A, Fiori R, Silvani P, Rancoita PM, Gabaldo M, Benedicenti F, Antonioli G, Assanelli A, Cicalese MP, Del Carro U, Sora MG, Martino S, Quattrini A, Montini E, Di Serio C, Ciceri F, Roncarolo MG, Aiuti A, Naldini L, Biffi A. Lentiviral haemopoietic stem-cell gene therapy in early-onset metachromatic leukodystrophy: an ad-hoc analysis of a non-randomised, open-label, phase 1/2 trial. *Lancet* 2016;388(10043):476–87.
- Sibbald B. Death but one unintended consequence of gene-therapy trial. *CMAJ* May 29, 2001;164(11):1612.
- Somia N, Verma IM. Gene therapy: trials and tribulations. *Nat Rev Genet* 2000;1(2):91–9.
- Spanggaard I, Dahlstroem K, Laessoe L, Hansen RH, Johannesen HH, Hendel HW, Bouquet C, Attali P, Gehl J. Gene therapy for patients with advanced solid tumors: a phase I study using gene electrotransfer to muscle with the integrin inhibitor plasmid AMEP. *Acta Oncol* 2017;56(7):909–16.
- Stein S, Ott MG, Schultze-Strasser S, Jauch A, Burwinkel B, Kinner A, Schmidt M, Krämer A, Schwäble J, Glimm H, Koehl U, Preiss C, Ball C, Martin H, Göhring G, Schwarzwaelder K, Hofmann WK, Karakaya K, Tchatchou S, Yang R, Reinecke P, Kühlcke K, Schlegelberger B, Thrasher AJ, Hoelzer D, Seger R, von Kalle C, Grez M. Genomic instability and myelodysplasia with monosomy 7 consequent to EVI1 activation after gene therapy for chronic granulomatous disease. *Nat Med* 2010;16:198–204.
- Taniyama Y, Azuma J, Kunugiza Y, Iekushi K, Rakugi H, Morishita R. Therapeutic option of plasmid-DNA based gene transfer. *Curr Top Med Chem* 2012;12(15):1630–7.
- Tebas P, Stein D, Tang WW, Frank I, Wang SQ, Lee G, Spratt SK, Surosky RT, Giedlin MA, Nichol G, Holmes MC, Gregory PD, Ando DG, Kalos M, Collman RG, Binder-Scholl G, Plesa G, Hwang WT, Levine BL, June CH. Gene editing of CCR5 in autologous CD4 T cells of persons infected with HIV. *N Engl J Med* 2014;370(10):901–10.
- Thaker PH, Brady WE, Lankes HA, Odunsi K, Bradley WH, Moore KN, Muller CY, Anwer K, Schilder RJ, Alvarez RD, Fracasso PM. A phase I trial of intraperitoneal GEN-1, an IL-12 plasmid formulated with PEG-PEI-cholesterol lipopolymer, administered with pegylated liposomal doxorubicin in patients with recurrent or persistent epithelial ovarian, fallopian tube or primary peritoneal cancers: an NRG Oncology/Gynecologic Oncology Group study. *Gynecol Oncol* 2017;147(2):283–90.
- Thomas CE, Ehrhardt A, Kay MA. Progress and problems with the use of viral vectors for gene therapy. *Nat Rev Genet* 2003;4:346–58.
- Tighe H, Corr M, Roman M, Raz E. Gene vaccination: plasmid DNA is more than just a blueprint. *Immunol Today* 1998;19:89–97.
- Tipanee J, VandenDriessche T, Chuah MK. Transposons: moving forward from preclinical studies to clinical trials. *Hum Gene Ther* 2017;28:1087–104.
- Trapani I, Banfi S, Simonelli F, Surace EM, Auricchio A. Gene therapy of inherited retinal degenerations: prospects and challenges. *Hum Gene Ther* 2015;26:193–200.

- VandenDriessche T, Ivics Z, Izsvák Z, Chuah MK. Emerging potential of transposons for gene therapy and generation of induced pluripotent stem cells. *Blood* 2009;114:1461–8.
- Verlengia G, Miyagawa Y, Ingusci S, Cohen JB, Simonato M, Glorioso JC. Engineered HSV vector achieves safe long-term transgene expression in the central nervous system. *Sci Rep* 2017;7:1507.
- Verma IM, Somia N. Gene therapy-promises, problems, and prospects. *Nature* September 18, 1997;389(6648):239–42.
- von Kalle C, Deichmann A, Schmidt M. Vector integration and tumorigenesis. *Hum Gene Ther* 2014;25(6):475–81.
- Wang W, Li W, Ma N, Steinhoff G. Non-viral gene delivery methods. *Curr Pharm Biotechnol* 2013;14(1):46–60.
- Wolff JA, Lederberg J. An early history of gene transfer and therapy. *Hum Gene Ther* 1994;5:469–80.
- Zhang W-W, Li L, Li D, Liu J, Li X, Li W, Xu X, Zhang MJ, Chandler LA, Lin H, Hu A, Xu W, Lam DMK. The first approved gene therapy product for cancer Ad-p53 (Gendicine): 12 years in the clinic. *Hum Gene Ther* 2018;29(2):160–79.