

Chapter 4. Probability models

- A **probability model** is an assignment of probabilities to possible outcomes.
- We don't observe these probabilities. We observe a particular dataset.
- If we treat the dataset as an outcome of a probability model, we can answer questions such as,

"If there really is no association between unemployment and life expectancy, what is the probability we would see a least squares linear model coefficient as large as the one we actually observed, due to random fluctuations in the data?"

- Here, we are particularly interested in developing a probability model for the linear model.
- First, we need some basic tools for probability models: random variables, the normal distribution, mean, variance and standard deviation.

Random variables and events

- A **random variable** X is a random number with probabilities assigned to outcomes.

Example: Let X be a roll of a fair die. A natural probability model is to assign probability of $1/6$ to each of the possible outcomes $1, 2, 3, 4, 5, 6$.

- An **event** is a set of possible outcomes.

Example: For a die, $E = \{X \geq 4\} = \{4, 5, 6\}$ is the event that the die shows 4 or more.

- We can assign probabilities to events just like to outcomes.

Example: For a die, $P(E) = P(X \geq 4) = 3/6 = 1/2$.

Question 4.1. If an experiment can be repeated many times (like rolling a die) how can you check whether the probability model is correct?

Notation for combining events

- $\{E \text{ or } F\}$ is the event that either E or F or both happens.
- Since E and F are sets, we can write this as a union, $\{E \text{ or } F\} = E \cup F$
- $\{E \text{ and } F\}$ is the event that both E and F happen.
- We can write this as an intersection,

$$\{E \text{ and } F\} = E \cap F$$

- Usually, we prefer “and/or” to “intersection/union”.

Question 4.2. When does this formal use of “and” and “or” agree with usual English usage? When does it disagree?

The basic rules of probability

- ① Probabilities are numbers between 0 (impossible) and 1 (certain).
- ② Let \mathcal{S} be the set of all possible outcomes. Then, $P(\mathcal{S}) = 1$.
Example: For a die, $P(X \in \{1, 2, 3, 4, 5, 6\}) = 1$.
- ③ Events E and F are called **mutually exclusive** if they cannot happen at the same time. In other words, their intersection is the empty set. In this case,

$$P(E \text{ or } F) = P(E) + P(F).$$

Question 4.3. You roll a red die and a blue die. Let

$E = \{\text{red die shows 1}\}$, $F = \{\text{blue die shows 1}\}$, $G = \{\text{red die shows 6}\}$.

(a) Are E and F mutually exclusive? (b) How about E and G ? (c) How about F and G ?

Discrete random variables

- A **discrete random variable** is one where we can list all possible outcomes. Let's call them x_1, x_2, \dots
- A discrete random variable is specified by probability that the random variable takes each possible outcome,

$$p_i = P[X = x_i], \text{ for } i = 1, 2, 3, \dots$$

- It can be helpful to plot a graph of p_i against x_i .
- This graph is called the **probability mass function**.


Question 4.4. Sketch the probability mass function for a fair die.

Simulating the law of large numbers


- The “law of large numbers” says that the proportion of each outcome i in a large number of draws of a discrete random variable approaches p_i .
- We can test this by simulation, using the `replicate()` command.

Worked example 4.1. In R, a random draw with replacement from $\{1, 2, 3, 4, 5, 6\}$ can be obtained by `sample(1:6, size=1)`. This is equivalent to one roll of a fair die.

```
hist(replicate(n=100, sample(1:6, size=1) ),  
     main="", prob=TRUE, breaks=0.5:6.5, xlab="n=100", ylim=c(0, 0.21))
```



n=100



n=10000

Continuous random variables: the normal distribution

- A **continuous random variable** is one which can take any value in an interval of the real numbers.

Example: physical quantities such as time and speed are not limited to a discrete set of possible values.

- We will often see the **normal distribution**.
- Let's look at **normal random variables** simulated by R using `rnorm()`.


```
rnorm(n=10, mean=20, sd=5)
```

```
## [1] 14.01141 26.18597 17.18972 21.12226 15.20211 30.34660  
## [7] 19.94277 22.05179 27.73804 25.94906
```


- The arguments `mean=20, sd=5` of `rnorm()` are the **parameters** of the normal distribution.
- A normal random variable can take any numeric value: it is continuous.
- Values are centered on the mean and are usually less than twice the standard deviation (`sd`) from the mean.

A histogram of normal distribution simulations

```
hist(rnorm(n=100,mean=20,sd=5),  
     main="",xlab="n=100")
```



n=100



n=10000

- Large samples from the normal distribution follow a **bell curve** histogram.
- From smaller samples, this is harder to see.

Finding probabilities for a continuous random variable

- A continuous random variable X has a **probability density function** $f(x)$ with

$$P(a < X < b) = \int_a^b f(x) dx$$

- Write $X \sim \text{normal}(\mu, \sigma)$ to mean X is a normal random variable with mean μ and sd σ . The probability density function of X is

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2/2\sigma^2}$$


and so

$$P(a < X < b) = \int_a^b \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2/2\sigma^2} dx$$

- This integral has no closed form solution.
- Fortunately, R provides `pnorm()` and `qnorm()` that let us work with probabilities for the normal distribution numerically.

Calculating probabilities for the normal distribution

- `pnorm()` finds the **left tail** of the normal distribution.



Example: `pnorm(25, mean=20, sd=5)` computes the shaded area above.

- We don't have to do calculus with the normal integral, but we do use the relationship between the curve, area under the curve, and probability. And we must know how to compute these things in R.
- For $X \sim \text{normal}(\mu, \sigma)$,

$$\text{pnorm}(x, \text{mu}, \text{sigma}) = P(X \leq x) = \int_{-\infty}^x \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(y-\mu)^2/2\sigma^2} dy$$

Finding probabilities that are not a left tail


Question 4.5. Let $X \sim \text{normal}(10, 10)$. Sketch a shaded area under a curve giving $P(0 \leq X \leq 10)$. Write this probability as an integral and as R code.

Hint: To use `pnorm()`, think of the shaded area as the difference of two left tails.


Other continuous distributions

- There are many continuous random variables that are not normally distributed.
- For example, we could explore the **uniform** or **exponential** distributions.

```
hist(runif(100))
```



```
hist(rexp(100))
```



- Normal random variables are the only ones we will need in this class.
- Let's investigate why the normal distribution is so important.

Sums and averages follow the normal distribution

- When we sum many random quantities, the sum often follows a normal distribution even if each term in the sum is not normally distributed.
- This property is called **the central limit theorem**.
- It is an empirical fact as well as a mathematical theorem!
- Averaging is multiplying the sum by a constant ($1/n$). A bell curve is still a bell curve when we rescale by multiplication.

Question 4.6. Would you expect a histogram of student heights to follow a normal curve? Why? Why not?

Question 4.7. Would you expect a histogram of the daily change in the Dow Jones stock market index to follow a normal curve? Why? Why not?

Demonstrating the central limit theorem with dice


- `sample(1:6,2,replace=TRUE)` simulates the outcome of rolling two dice.
- `sum(sample(1:6,2,replace=TRUE))` simulates the sum of two dice.
- `replicate()` lets us see what happens if we do this many times

```
dice2 <- replicate(50000,sum(sample(1:6,2,replace=TRUE)))  
dice3 <- replicate(50000,sum(sample(1:6,3,replace=TRUE)))  
dice10 <- replicate(50000,sum(sample(1:6,10,replace=TRUE)))  
dice20 <- replicate(50000,sum(sample(1:6,20,replace=TRUE)))
```


- Guess how many dice you have to add up before the histogram looks normal?

```
hist(dice2,prob=TRUE,breaks=(min(dice2)-0.5):(max(dice2)+0.5))
normal.x <- seq(from=min(dice2),to=max(dice2),length=100)
normal.y <- dnorm(normal.x,mean=mean(dice2),sd=sd(dice2))
lines(normal.x,normal.y,col="blue")
```


2 dice




3 dice



10 dice



20 dice



Question 4.8. Why do we use `prob` and `breaks` arguments to `hist()`?

More normal approximation situations

Question 4.9. Would you expect detrended data on life expectancy at birth to follow a normal distribution? Explain.

Question 4.10. Consider the mice weight data for HW1 with mice randomized to two treatments: a high fat diet and a usual lab diet. (a) Would you expect the weights of mice in each treatment group to follow a normal distribution? (b) If the experiment were replicated ten times, and an average weight calculated for each of these ten replications, would you expect the ten averages to follow a normal distribution? Are your answers different for (a) and (b)?

The sample mean and the expectation of a random variable

- The **sample mean** or **average** of $\mathbf{y} = (y_1, \dots, y_n)$ is

$$\bar{\mathbf{y}} = \frac{1}{n} \sum_{i=1}^n y_i$$

- The **expected value** of a random variable X taking possible values x_1, x_2, \dots with probabilities p_1, p_2, \dots is

$$\mathbb{E}[X] = \sum_{i=1}^{\infty} x_i p_i$$

- If we have many draws of X , the sample proportion taking value x_i becomes close to p_i and so the sample mean becomes close to the expected value.
- If X is a continuous random variable with density $f(x)$ the sum for expected value becomes an integral,

$$\mathbb{E}[X] = \int_{-\infty}^{\infty} x f(x) dx$$

The sample mean and expectation using R

- We've already seen the `mean()` function which computes the sample mean of a numeric vector.
- One way to compute the expected value of a random variable is to take the sample mean of many realizations of the random variable

```
x <- rnorm(10000,mean=20,sd=5)
mean(x)
## [1] 20.04557
```

- The expectation of a random variable is sometimes called its mean.

The sample variance and the variance of a random variable

- The **sample variance** of $\mathbf{y} = (y_1, \dots, y_n)$ is

$$\text{Var}(\mathbf{y}) = \frac{1}{n} \sum_{i=1}^n (y_i - \bar{\mathbf{y}})^2$$

- The **variance** of a random variable X is defined in terms of the expected value as

$$\text{Var}(X) = \text{E}[(X - \text{E}(X))^2]$$