


MCMC from Diffusions and Geodesics



- ▶ Riemann manifold Langevin and Hamiltonian Monte Carlo Methods
Girolami, M. & Calderhead, B.
J.R.Statist. Soc. B, with discussion, (2011), 73, 2, 123 - 214.

Talk Outline

- ▶ Motivation to improve MCMC capability for challenging problems

Talk Outline

- ▶ Motivation to improve MCMC capability for challenging problems
- ▶ Exploring geometric concepts in MCMC methodology

Talk Outline

- ▶ Motivation to improve MCMC capability for challenging problems
- ▶ Exploring geometric concepts in MCMC methodology
- ▶ Diffusions across Riemann manifold as basis for MCMC

Talk Outline

- ▶ Motivation to improve MCMC capability for challenging problems
- ▶ Exploring geometric concepts in MCMC methodology
- ▶ Diffusions across Riemann manifold as basis for MCMC
- ▶ Geodesic flows on manifold form basis of MCMC methods

Talk Outline

- ▶ Motivation to improve MCMC capability for challenging problems
- ▶ Exploring geometric concepts in MCMC methodology
- ▶ Diffusions across Riemann manifold as basis for MCMC
- ▶ Geodesic flows on manifold form basis of MCMC methods
- ▶ Illustrative examples and Hilbert manifolds

Talk Outline

- ▶ Motivation to improve MCMC capability for challenging problems
- ▶ Exploring geometric concepts in MCMC methodology
- ▶ Diffusions across Riemann manifold as basis for MCMC
- ▶ Geodesic flows on manifold form basis of MCMC methods
- ▶ Illustrative examples and Hilbert manifolds
- ▶ Further Work and Conclusions

Motivation Simulation Based Inference

- ▶ Monte Carlo method employs samples from $\pi(\theta)$ to obtain estimate

$$\int \phi(\theta) \pi(\theta) d\theta = \frac{1}{N} \sum_n \phi(\theta^n) + \mathcal{O}(N^{-\frac{1}{2}})$$

Motivation Simulation Based Inference

- ▶ Monte Carlo method employs samples from $\pi(\theta)$ to obtain estimate

$$\int \phi(\theta) \pi(\theta) d\theta = \frac{1}{N} \sum_n \phi(\theta^n) + \mathcal{O}(N^{-\frac{1}{2}})$$

- ▶ Draw θ^n from ergodic Markov process with stationary distribution $\pi(\theta)$

Motivation Simulation Based Inference

- ▶ Monte Carlo method employs samples from $\pi(\theta)$ to obtain estimate

$$\int \phi(\theta) \pi(\theta) d\theta = \frac{1}{N} \sum_n \phi(\theta^n) + \mathcal{O}(N^{-\frac{1}{2}})$$

- ▶ Draw θ^n from ergodic Markov process with stationary distribution $\pi(\theta)$
- ▶ Construct transition kernel as product of two components

Motivation Simulation Based Inference

- ▶ Monte Carlo method employs samples from $\pi(\theta)$ to obtain estimate

$$\int \phi(\theta) \pi(\theta) d\theta = \frac{1}{N} \sum_n \phi(\theta^n) + \mathcal{O}(N^{-\frac{1}{2}})$$

- ▶ Draw θ^n from ergodic Markov process with stationary distribution $\pi(\theta)$
- ▶ Construct transition kernel as product of two components
 - ▶ Propose a move $\theta \rightarrow \theta'$ with probability $p_p(\theta'|\theta)$

Motivation Simulation Based Inference

- ▶ Monte Carlo method employs samples from $\pi(\theta)$ to obtain estimate

$$\int \phi(\theta) \pi(\theta) d\theta = \frac{1}{N} \sum_n \phi(\theta^n) + \mathcal{O}(N^{-\frac{1}{2}})$$

- ▶ Draw θ^n from ergodic Markov process with stationary distribution $\pi(\theta)$
- ▶ Construct transition kernel as product of two components
 - ▶ Propose a move $\theta \rightarrow \theta'$ with probability $p_p(\theta'|\theta)$
 - ▶ accept or reject proposal with probability

$$p_a(\theta'|\theta) = \min \left[1, \frac{\pi(\theta') p_p(\theta|\theta')}{\pi(\theta) p_p(\theta'|\theta)} \right]$$

Motivation Simulation Based Inference

- ▶ Monte Carlo method employs samples from $\pi(\theta)$ to obtain estimate

$$\int \phi(\theta) \pi(\theta) d\theta = \frac{1}{N} \sum_n \phi(\theta^n) + \mathcal{O}(N^{-\frac{1}{2}})$$

- ▶ Draw θ^n from ergodic Markov process with stationary distribution $\pi(\theta)$
- ▶ Construct transition kernel as product of two components
 - ▶ Propose a move $\theta \rightarrow \theta'$ with probability $p_p(\theta'|\theta)$
 - ▶ accept or reject proposal with probability

$$p_a(\theta'|\theta) = \min \left[1, \frac{\pi(\theta') p_p(\theta|\theta')}{\pi(\theta) p_p(\theta'|\theta)} \right]$$

- ▶ Convergence rate and asymptotic variance dependent on $p_p(\theta'|\theta)$

Motivation Simulation Based Inference

- ▶ Monte Carlo method employs samples from $\pi(\theta)$ to obtain estimate

$$\int \phi(\theta) \pi(\theta) d\theta = \frac{1}{N} \sum_n \phi(\theta^n) + \mathcal{O}(N^{-\frac{1}{2}})$$


- ▶ Draw θ^n from ergodic Markov process with stationary distribution $\pi(\theta)$
- ▶ Construct transition kernel as product of two components
 - ▶ Propose a move $\theta \rightarrow \theta'$ with probability $p_p(\theta'|\theta)$
 - ▶ accept or reject proposal with probability

$$p_a(\theta'|\theta) = \min \left[1, \frac{\pi(\theta') p_p(\theta|\theta')}{\pi(\theta) p_p(\theta'|\theta)} \right]$$


- ▶ Convergence rate and asymptotic variance dependent on $p_p(\theta'|\theta)$
- ▶ Success of MCMC reliant upon appropriate proposal design

Simple Dynamics


$$\begin{aligned}\frac{dx}{dt} &= \theta_1 yz \\ \frac{dy}{dt} &= -xz \\ \frac{dz}{dt} &= -\theta_2 xy\end{aligned}$$




Simple Dynamics



Simple Dynamics



System Identification: Nonlinear ODE Oscillator Model




$$\frac{dx_1}{dt} = \frac{k_1}{1 + x_n^\rho} - m_1 x_1$$

$$\frac{dx_2}{dt} = k_2 x_1 - m_2 x_2$$


⋮

$$\frac{dx_n}{dt} = k_n x_{n-1} - m_n x_n$$

Systems Identification - Posterior Inference



Mixing of Markov Chains



Geometric Concepts in MCMC

Geometric Concepts in MCMC

- ▶ Denote expected Fisher Information as $\mathbf{G}(\theta) = \text{cov}(\nabla_{\theta}\mathcal{L}(\theta))$

Geometric Concepts in MCMC

- ▶ Denote expected Fisher Information as $\mathbf{G}(\theta) = \text{cov}(\nabla_{\theta}\mathcal{L}(\theta))$
- ▶ Rao, 1945 to first order

$$\chi^2(\delta\theta) = \int \frac{|p(\mathbf{y}; \theta + \delta\theta) - p(\mathbf{y}; \theta)|^2}{p(\mathbf{y}; \theta)} d\mathbf{y} \approx \delta\theta^\top \mathbf{G}(\theta) \delta\theta$$

Geometric Concepts in MCMC

- ▶ Denote expected Fisher Information as $\mathbf{G}(\theta) = \text{cov}(\nabla_{\theta} \mathcal{L}(\theta))$
- ▶ Rao, 1945 to first order

$$\chi^2(\delta\theta) = \int \frac{|p(\mathbf{y}; \theta + \delta\theta) - p(\mathbf{y}; \theta)|^2}{p(\mathbf{y}; \theta)} d\mathbf{y} \approx \delta\theta^T \mathbf{G}(\theta) \delta\theta$$

- ▶ Jeffreys, 1948 to first order

$$D(\theta || \delta\theta) = \int p(\mathbf{y}; \theta + \delta\theta) \log \frac{p(\mathbf{y}; \theta + \delta\theta)}{p(\mathbf{y}; \theta)} d\mathbf{y} \approx \delta\theta^T \mathbf{G}(\theta) \delta\theta$$

Geometric Concepts in MCMC

- ▶ Denote expected Fisher Information as $\mathbf{G}(\theta) = \text{cov}(\nabla_{\theta} \mathcal{L}(\theta))$
- ▶ Rao, 1945 to first order

$$\chi^2(\delta\theta) = \int \frac{|p(\mathbf{y}; \theta + \delta\theta) - p(\mathbf{y}; \theta)|^2}{p(\mathbf{y}; \theta)} d\mathbf{y} \approx \delta\theta^T \mathbf{G}(\theta) \delta\theta$$

- ▶ Jeffreys, 1948 to first order

$$D(\theta || \delta\theta) = \int p(\mathbf{y}; \theta + \delta\theta) \log \frac{p(\mathbf{y}; \theta + \delta\theta)}{p(\mathbf{y}; \theta)} d\mathbf{y} \approx \delta\theta^T \mathbf{G}(\theta) \delta\theta$$

- ▶ Expected Fisher Information $\mathbf{G}(\theta)$ is metric tensor of a Riemann manifold

Geometric Concepts in MCMC

- ▶ Denote expected Fisher Information as $\mathbf{G}(\theta) = \text{cov}(\nabla_{\theta} \mathcal{L}(\theta))$
- ▶ Rao, 1945 to first order

$$\chi^2(\delta\theta) = \int \frac{|p(\mathbf{y}; \theta + \delta\theta) - p(\mathbf{y}; \theta)|^2}{p(\mathbf{y}; \theta)} d\mathbf{y} \approx \delta\theta^T \mathbf{G}(\theta) \delta\theta$$

- ▶ Jeffreys, 1948 to first order

$$D(\theta || \delta\theta) = \int p(\mathbf{y}; \theta + \delta\theta) \log \frac{p(\mathbf{y}; \theta + \delta\theta)}{p(\mathbf{y}; \theta)} d\mathbf{y} \approx \delta\theta^T \mathbf{G}(\theta) \delta\theta$$

- ▶ Expected Fisher Information $\mathbf{G}(\theta)$ is metric tensor of a Riemann manifold
- ▶ Non-Euclidean geometry - invariants, connections, curvature, geodesics

Geometric Concepts in MCMC

- ▶ Denote expected Fisher Information as $\mathbf{G}(\theta) = \text{cov}(\nabla_{\theta} \mathcal{L}(\theta))$
- ▶ Rao, 1945 to first order

$$\chi^2(\delta\theta) = \int \frac{|p(\mathbf{y}; \theta + \delta\theta) - p(\mathbf{y}; \theta)|^2}{p(\mathbf{y}; \theta)} d\mathbf{y} \approx \delta\theta^T \mathbf{G}(\theta) \delta\theta$$

- ▶ Jeffreys, 1948 to first order

$$D(\theta || \delta\theta) = \int p(\mathbf{y}; \theta + \delta\theta) \log \frac{p(\mathbf{y}; \theta + \delta\theta)}{p(\mathbf{y}; \theta)} d\mathbf{y} \approx \delta\theta^T \mathbf{G}(\theta) \delta\theta$$

- ▶ Expected Fisher Information $\mathbf{G}(\theta)$ is metric tensor of a Riemann manifold
- ▶ Non-Euclidean geometry - invariants, connections, curvature, geodesics
- ▶ Asymptotic statistical analysis. e.g. Amari, 1981; Murray & Rice, 1993; Critchley *et al*, 1993; Kass, 1989; Dawid, 1975; Lauritsen, 1989

Geometric Concepts in MCMC

- ▶ Denote expected Fisher Information as $\mathbf{G}(\theta) = \text{cov}(\nabla_{\theta}\mathcal{L}(\theta))$
- ▶ Rao, 1945 to first order

$$\chi^2(\delta\theta) = \int \frac{|p(\mathbf{y}; \theta + \delta\theta) - p(\mathbf{y}; \theta)|^2}{p(\mathbf{y}; \theta)} d\mathbf{y} \approx \delta\theta^T \mathbf{G}(\theta) \delta\theta$$

- ▶ Jeffreys, 1948 to first order

$$D(\theta || \delta\theta) = \int p(\mathbf{y}; \theta + \delta\theta) \log \frac{p(\mathbf{y}; \theta + \delta\theta)}{p(\mathbf{y}; \theta)} d\mathbf{y} \approx \delta\theta^T \mathbf{G}(\theta) \delta\theta$$

- ▶ Expected Fisher Information $\mathbf{G}(\theta)$ is metric tensor of a Riemann manifold
- ▶ Non-Euclidean geometry - invariants, connections, curvature, geodesics
- ▶ Asymptotic statistical analysis. e.g. Amari, 1981; Murray & Rice, 1993; Critchley *et al*, 1993; Kass, 1989; Dawid, 1975; Lauritsen, 1989
- ▶ Statistical shape analysis Kent *et al*, 1996; Dryden & Mardia, 1998

Geometric Concepts in MCMC

- ▶ Denote expected Fisher Information as $\mathbf{G}(\theta) = \text{cov}(\nabla_{\theta} \mathcal{L}(\theta))$
- ▶ Rao, 1945 to first order

$$\chi^2(\delta\theta) = \int \frac{|p(\mathbf{y}; \theta + \delta\theta) - p(\mathbf{y}; \theta)|^2}{p(\mathbf{y}; \theta)} d\mathbf{y} \approx \delta\theta^T \mathbf{G}(\theta) \delta\theta$$

- ▶ Jeffreys, 1948 to first order

$$D(\theta || \delta\theta) = \int p(\mathbf{y}; \theta + \delta\theta) \log \frac{p(\mathbf{y}; \theta + \delta\theta)}{p(\mathbf{y}; \theta)} d\mathbf{y} \approx \delta\theta^T \mathbf{G}(\theta) \delta\theta$$


- ▶ Expected Fisher Information $\mathbf{G}(\theta)$ is metric tensor of a Riemann manifold
- ▶ Non-Euclidean geometry - invariants, connections, curvature, geodesics
- ▶ Asymptotic statistical analysis. e.g. Amari, 1981; Murray & Rice, 1993; Critchley *et al*, 1993; Kass, 1989; Dawid, 1975; Lauritsen, 1989
- ▶ Statistical shape analysis Kent *et al*, 1996; Dryden & Mardia, 1998
- ▶ Can geometric structure be employed in Monte Carlo methodology?

Manifolds

A manifold is a smooth, curved surface: A set *embedded* in \mathbb{R}^d , that locally looks like \mathbb{R}^n ($n < d$).

Example: the unit sphere (2-sphere): $d = 3$, $n = 2$

$$\mathcal{S}_2 = \{x \in \mathbb{R}^3 : \sum_i x_i^2 = 1\}$$




Manifolds

A manifold is a smooth, curved surface: A set *embedded* in \mathbb{R}^d , that locally looks like \mathbb{R}^n ($n < d$).

Example: the unit sphere (2-sphere): $d = 3$, $n = 2$



$$\mathcal{S}_2 = \{x \in \mathbb{R}^3 : \sum_i x_i^2 = 1\}$$



$x \in \mathbb{R}^d$ are the **embedded coordinates**

Coordinate systems and Riemannian metrics



Can parameterise the manifold with a **coordinate system** in $q \in \mathbb{R}^n$



$$(\sin q_1 \sin q_2, \cos q_1 \sin q_2, \cos q_2), \quad q_1 \in [0, 2\pi], q_2 \in [0, \pi]$$

Coordinate systems and Riemannian metrics

Can parameterise the manifold with a **coordinate system** in $q \in \mathbb{R}^n$




$$(\sin q_1 \sin q_2, \cos q_1 \sin q_2, \cos q_2), \quad q_1 \in [0, 2\pi], q_2 \in [0, \pi]$$

The Euclidean metric $\|\cdot\|$ induces a **Riemannian metric** G in the coordinate system:

$$\|dx\|^2 = \sum_{i,j} G(q) dq_i dq_j$$



M.C. Escher, Heaven and Hell, 1960



Geodesics

Geodesics are the paths of shortest distance.



- ▶ On a sphere, these are the *great circles*



Geodesics

Geodesics are the paths of shortest distance.

- ▶ On a sphere, these are the *great circles*




Given an initial velocity $v(0) \perp x(0)$, we have a nice explicit form


$$[x(t) \quad v(t)] = [x(0) \quad v(0)] \begin{bmatrix} 1 & \\ & \alpha^{-1} \end{bmatrix} \begin{bmatrix} \cos(\alpha t) & -\sin(\alpha t) \\ \sin(\alpha t) & \cos(\alpha t) \end{bmatrix} \begin{bmatrix} 1 & \\ & \alpha \end{bmatrix}$$

where $\alpha = \|v(0)\|$.

Geometric Concepts in MCMC




Geometric Concepts in MCMC




- ▶ Tangent space - local metric defined by $\delta\theta^T \mathbf{G}(\theta) \delta\theta = g_{kl} \delta\theta_k \delta\theta_l$

Geometric Concepts in MCMC



- ▶ Tangent space - local metric defined by $\delta\theta^T \mathbf{G}(\theta) \delta\theta = g_{kl} \delta\theta_k \delta\theta_l$
- ▶ Christoffel symbols - characterise Levi-Civita connection on manifold


Geometric Concepts in MCMC



- ▶ Tangent space - local metric defined by $\delta\theta^T \mathbf{G}(\theta) \delta\theta = g_{kl} \delta\theta_k \delta\theta_l$
- ▶ Christoffel symbols - characterise Levi-Civita connection on manifold

$$\Gamma_{kl}^i = \frac{1}{2} \sum_m g^{im} \left(\frac{\partial g_{mk}}{\partial \theta^l} + \frac{\partial g_{ml}}{\partial \theta^k} - \frac{\partial g_{kl}}{\partial \theta^m} \right)$$

Geometric Concepts in MCMC




- ▶ Tangent space - local metric defined by $\delta\theta^T \mathbf{G}(\theta) \delta\theta = g_{kl} \delta\theta_k \delta\theta_l$
- ▶ Christoffel symbols - characterise Levi-Civita connection on manifold

$$\Gamma_{kl}^i = \frac{1}{2} \sum_m g^{im} \left(\frac{\partial g_{mk}}{\partial \theta^l} + \frac{\partial g_{ml}}{\partial \theta^k} - \frac{\partial g_{kl}}{\partial \theta^m} \right)$$

- ▶ Geodesics - shortest path between two points on manifold

Geometric Concepts in MCMC



- ▶ Tangent space - local metric defined by $\delta\theta^T \mathbf{G}(\theta) \delta\theta = g_{kl} \delta\theta_k \delta\theta_l$
- ▶ Christoffel symbols - characterise Levi-Civita connection on manifold

$$\Gamma_{kl}^i = \frac{1}{2} \sum_m g^{im} \left(\frac{\partial g_{mk}}{\partial \theta^l} + \frac{\partial g_{ml}}{\partial \theta^k} - \frac{\partial g_{kl}}{\partial \theta^m} \right)$$

- ▶ Geodesics - shortest path between two points on manifold

$$\frac{d^2\theta^i}{dt^2} + \sum_{k,l} \Gamma_{kl}^i \frac{d\theta^k}{dt} \frac{d\theta^l}{dt} = 0$$

Fisher–Rao metric

A family of probability densities $\{p(\cdot | \theta) : \theta \in \Theta\}$, the Fisher information forms a natural Riemannian metric, known as **Fisher–Rao metric**:


$$g_{ij} = E_{Y|\theta} \left[-\frac{\partial^2}{\partial \theta_i \partial \theta_j} \log p(Y | \theta) \right]$$

Fisher–Rao metric

A family of probability densities $\{p(\cdot | \theta) : \theta \in \Theta\}$, the Fisher information forms a natural Riemannian metric, known as **Fisher–Rao metric**:

$$g_{ij} = E_{Y|\theta} \left[-\frac{\partial^2}{\partial \theta_i \partial \theta_j} \log p(Y | \theta) \right]$$

Examples:



Fisher–Rao metric

A family of probability densities $\{p(\cdot | \theta) : \theta \in \Theta\}$, the Fisher information forms a natural Riemannian metric, known as **Fisher–Rao metric**:

$$g_{ij} = E_{Y|\theta} \left[-\frac{\partial^2}{\partial \theta_i \partial \theta_j} \log p(Y | \theta) \right]$$

Examples:





Illustration of Geometric Concepts

- ▶ Consider Normal density $p(x|\mu, \sigma) = \mathcal{N}_x(\mu, \sigma)$

Illustration of Geometric Concepts

- ▶ Consider Normal density $p(x|\mu, \sigma) = \mathcal{N}_x(\mu, \sigma)$
- ▶ Local inner product on tangent space defined by metric tensor, i.e. $\delta\theta^T \mathbf{G}(\theta) \delta\theta$, where $\theta = (\mu, \sigma)^T$

Illustration of Geometric Concepts

- ▶ Consider Normal density $p(x|\mu, \sigma) = \mathcal{N}_x(\mu, \sigma)$
- ▶ Local inner product on tangent space defined by metric tensor, i.e. $\delta\theta^T \mathbf{G}(\theta) \delta\theta$, where $\theta = (\mu, \sigma)^T$
- ▶ Metric is Expected Fisher Information

$$\mathbf{G}(\mu, \sigma) = \begin{bmatrix} \sigma^{-2} & 0 \\ 0 & 2\sigma^{-2} \end{bmatrix}$$

Illustration of Geometric Concepts

- ▶ Consider Normal density $p(x|\mu, \sigma) = \mathcal{N}_x(\mu, \sigma)$
- ▶ Local inner product on tangent space defined by metric tensor, i.e. $\delta\theta^T \mathbf{G}(\theta) \delta\theta$, where $\theta = (\mu, \sigma)^T$
- ▶ Metric is Expected Fisher Information

$$\mathbf{G}(\mu, \sigma) = \begin{bmatrix} \sigma^{-2} & 0 \\ 0 & 2\sigma^{-2} \end{bmatrix}$$

- ▶ Components of connection $\partial_\mu \mathbf{G} = \mathbf{0}$ and $\partial_\sigma \mathbf{G} = -\text{diag}(2\sigma^{-3}, 4\sigma^{-3})$

Illustration of Geometric Concepts

- ▶ Consider Normal density $p(x|\mu, \sigma) = \mathcal{N}_x(\mu, \sigma)$
- ▶ Local inner product on tangent space defined by metric tensor, i.e. $\delta\theta^T \mathbf{G}(\theta) \delta\theta$, where $\theta = (\mu, \sigma)^T$
- ▶ Metric is Expected Fisher Information

$$\mathbf{G}(\mu, \sigma) = \begin{bmatrix} \sigma^{-2} & 0 \\ 0 & 2\sigma^{-2} \end{bmatrix}$$

- ▶ Components of connection $\partial_\mu \mathbf{G} = \mathbf{0}$ and $\partial_\sigma \mathbf{G} = -\text{diag}(2\sigma^{-3}, 4\sigma^{-3})$
- ▶ Metric on tangent space

$$\delta\theta^T \mathbf{G}(\theta) \delta\theta = \frac{(\delta\mu^2 + 2\delta\sigma^2)}{\sigma^2}$$

Illustration of Geometric Concepts

- ▶ Consider Normal density $p(x|\mu, \sigma) = \mathcal{N}_x(\mu, \sigma)$
- ▶ Local inner product on tangent space defined by metric tensor, i.e. $\delta\theta^T \mathbf{G}(\theta) \delta\theta$, where $\theta = (\mu, \sigma)^T$
- ▶ Metric is Expected Fisher Information

$$\mathbf{G}(\mu, \sigma) = \begin{bmatrix} \sigma^{-2} & 0 \\ 0 & 2\sigma^{-2} \end{bmatrix}$$

- ▶ Components of connection $\partial_\mu \mathbf{G} = \mathbf{0}$ and $\partial_\sigma \mathbf{G} = -\text{diag}(2\sigma^{-3}, 4\sigma^{-3})$
- ▶ Metric on tangent space

$$\delta\theta^T \mathbf{G}(\theta) \delta\theta = \frac{(\delta\mu^2 + 2\delta\sigma^2)}{\sigma^2}$$

- ▶ Metric tensor for univariate Normal defines a Hyperbolic Space

Illustration of Geometric Concepts

- ▶ Consider Normal density $p(x|\mu, \sigma) = \mathcal{N}_x(\mu, \sigma)$
- ▶ Local inner product on tangent space defined by metric tensor, i.e. $\delta\theta^T \mathbf{G}(\theta) \delta\theta$, where $\theta = (\mu, \sigma)^T$
- ▶ Metric is Expected Fisher Information


$$\mathbf{G}(\mu, \sigma) = \begin{bmatrix} \sigma^{-2} & 0 \\ 0 & 2\sigma^{-2} \end{bmatrix}$$

- ▶ Components of connection $\partial_\mu \mathbf{G} = \mathbf{0}$ and $\partial_\sigma \mathbf{G} = -\text{diag}(2\sigma^{-3}, 4\sigma^{-3})$
- ▶ Metric on tangent space


$$\delta\theta^T \mathbf{G}(\theta) \delta\theta = \frac{(\delta\mu^2 + 2\delta\sigma^2)}{\sigma^2}$$

- ▶ Metric tensor for univariate Normal defines a Hyperbolic Space
- ▶ Consider densities $\mathcal{N}(0, 1)$ & $\mathcal{N}(1, 1)$ and $\mathcal{N}(0, 2)$ & $\mathcal{N}(1, 2)$

Normal Density - Euclidean Parameter space



Normal Density - Riemannian Functional space



Langevin Diffusion on Riemannian manifold

- Discretised Langevin diffusion on manifold defines proposal mechanism

$$\boldsymbol{\theta}'_d = \boldsymbol{\theta}_d + \frac{\epsilon^2}{2} \left(\mathbf{G}^{-1}(\boldsymbol{\theta}) \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}) \right)_d - \epsilon^2 \sum_{i,j}^D \mathbf{G}(\boldsymbol{\theta})_{ij}^{-1} \Gamma_{ij}^d + \epsilon \left(\sqrt{\mathbf{G}^{-1}(\boldsymbol{\theta})} \mathbf{z} \right)_d$$

Langevin Diffusion on Riemannian manifold

- Discretised Langevin diffusion on manifold defines proposal mechanism

$$\theta'_d = \theta_d + \frac{\epsilon^2}{2} \left(\mathbf{G}^{-1}(\theta) \nabla_{\theta} \mathcal{L}(\theta) \right)_d - \epsilon^2 \sum_{i,j}^D \mathbf{G}(\theta)_{ij}^{-1} \Gamma_{ij}^d + \epsilon \left(\sqrt{\mathbf{G}^{-1}(\theta)} \mathbf{z} \right)_d$$

- Manifold with constant curvature then proposal mechanism reduces to

$$\theta' = \theta + \frac{\epsilon^2}{2} \mathbf{G}^{-1}(\theta) \nabla_{\theta} \mathcal{L}(\theta) + \epsilon \sqrt{\mathbf{G}^{-1}(\theta)} \mathbf{z}$$

Langevin Diffusion on Riemannian manifold

- Discretised Langevin diffusion on manifold defines proposal mechanism

$$\boldsymbol{\theta}'_d = \boldsymbol{\theta}_d + \frac{\epsilon^2}{2} \left(\mathbf{G}^{-1}(\boldsymbol{\theta}) \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}) \right)_d - \epsilon^2 \sum_{i,j}^D \mathbf{G}(\boldsymbol{\theta})_{ij}^{-1} \Gamma_{ij}^d + \epsilon \left(\sqrt{\mathbf{G}^{-1}(\boldsymbol{\theta})} \mathbf{z} \right)_d$$

- Manifold with constant curvature then proposal mechanism reduces to

$$\boldsymbol{\theta}' = \boldsymbol{\theta} + \frac{\epsilon^2}{2} \mathbf{G}^{-1}(\boldsymbol{\theta}) \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}) + \epsilon \sqrt{\mathbf{G}^{-1}(\boldsymbol{\theta})} \mathbf{z}$$

- MALA proposal with preconditioning

$$\boldsymbol{\theta}' = \boldsymbol{\theta} + \frac{\epsilon^2}{2} \mathbf{M} \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}) + \epsilon \sqrt{\mathbf{M}} \mathbf{z}$$

Langevin Diffusion on Riemannian manifold

- Discretised Langevin diffusion on manifold defines proposal mechanism

$$\boldsymbol{\theta}'_d = \boldsymbol{\theta}_d + \frac{\epsilon^2}{2} \left(\mathbf{G}^{-1}(\boldsymbol{\theta}) \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}) \right)_d - \epsilon^2 \sum_{i,j}^D \mathbf{G}(\boldsymbol{\theta})_{ij}^{-1} \Gamma_{ij}^d + \epsilon \left(\sqrt{\mathbf{G}^{-1}(\boldsymbol{\theta})} \mathbf{z} \right)_d$$

- Manifold with constant curvature then proposal mechanism reduces to

$$\boldsymbol{\theta}' = \boldsymbol{\theta} + \frac{\epsilon^2}{2} \mathbf{G}^{-1}(\boldsymbol{\theta}) \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}) + \epsilon \sqrt{\mathbf{G}^{-1}(\boldsymbol{\theta})} \mathbf{z}$$

- MALA proposal with preconditioning

$$\boldsymbol{\theta}' = \boldsymbol{\theta} + \frac{\epsilon^2}{2} \mathbf{M} \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}) + \epsilon \sqrt{\mathbf{M}} \mathbf{z}$$

- Proposal and acceptance probability

$$p_p(\boldsymbol{\theta}'|\boldsymbol{\theta}) = \mathcal{N}(\boldsymbol{\theta}'|\boldsymbol{\mu}(\boldsymbol{\theta}, \epsilon), \epsilon^2 \mathbf{G}^{-1}(\boldsymbol{\theta}))$$

$$p_a(\boldsymbol{\theta}'|\boldsymbol{\theta}) = \min \left[1, \frac{\pi(\boldsymbol{\theta}') p_p(\boldsymbol{\theta}|\boldsymbol{\theta}')}{\pi(\boldsymbol{\theta}) p_p(\boldsymbol{\theta}'|\boldsymbol{\theta})} \right]$$

Langevin Diffusion on Riemannian manifold

- Discretised Langevin diffusion on manifold defines proposal mechanism

$$\boldsymbol{\theta}'_d = \boldsymbol{\theta}_d + \frac{\epsilon^2}{2} \left(\mathbf{G}^{-1}(\boldsymbol{\theta}) \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}) \right)_d - \epsilon^2 \sum_{i,j}^D \mathbf{G}(\boldsymbol{\theta})_{ij}^{-1} \Gamma_{ij}^d + \epsilon \left(\sqrt{\mathbf{G}^{-1}(\boldsymbol{\theta})} \mathbf{z} \right)_d$$

- Manifold with constant curvature then proposal mechanism reduces to

$$\boldsymbol{\theta}' = \boldsymbol{\theta} + \frac{\epsilon^2}{2} \mathbf{G}^{-1}(\boldsymbol{\theta}) \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}) + \epsilon \sqrt{\mathbf{G}^{-1}(\boldsymbol{\theta})} \mathbf{z}$$

- MALA proposal with preconditioning

$$\boldsymbol{\theta}' = \boldsymbol{\theta} + \frac{\epsilon^2}{2} \mathbf{M} \nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}) + \epsilon \sqrt{\mathbf{M}} \mathbf{z}$$


- Proposal and acceptance probability

$$p_p(\boldsymbol{\theta}'|\boldsymbol{\theta}) = \mathcal{N}(\boldsymbol{\theta}'|\boldsymbol{\mu}(\boldsymbol{\theta}, \epsilon), \epsilon^2 \mathbf{G}^{-1}(\boldsymbol{\theta}))$$


$$p_a(\boldsymbol{\theta}'|\boldsymbol{\theta}) = \min \left[1, \frac{\pi(\boldsymbol{\theta}') p_p(\boldsymbol{\theta}|\boldsymbol{\theta}')}{\pi(\boldsymbol{\theta}) p_p(\boldsymbol{\theta}'|\boldsymbol{\theta})} \right]$$

- Proposal mechanism diffuses approximately along the manifold

Langevin Diffusion on Riemannian manifold



Langevin Diffusion on Riemannian manifold



Second-Order Diffusions

Second-Order Diffusions

- ▶ Define a second-order SDE on a Riemann manifold with the diffusion defined on vector field \mathbf{v} to obtain

Second-Order Diffusions

- ▶ Define a second-order SDE on a Riemann manifold with the diffusion defined on vector field \mathbf{v} to obtain

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

Second-Order Diffusions

- ▶ Define a second-order SDE on a Riemann manifold with the diffusion defined on vector field \mathbf{v} to obtain

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Vector function γ has components $\gamma_k = \sum_{i,j} \Gamma_{ij}^k v^i v^j$

Second-Order Diffusions

- ▶ Define a second-order SDE on a Riemann manifold with the diffusion defined on vector field \mathbf{v} to obtain

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Vector function γ has components $\gamma_k = \sum_{i,j} \Gamma_{ij}^k v^i v^j$
- ▶ Each Γ_{ij}^k being the Christoffel symbols of the Levi-Civita connection corresponding to the metric tensor $\mathbf{G}(\mathbf{x}_t)$

Second-Order Diffusions

- ▶ Define a second-order SDE on a Riemann manifold with the diffusion defined on vector field \mathbf{v} to obtain

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Vector function γ has components $\gamma_k = \sum_{i,j} \Gamma_{ij}^k v^i v^j$
- ▶ Each Γ_{ij}^k being the Christoffel symbols of the Levi-Civita connection corresponding to the metric tensor $\mathbf{G}(\mathbf{x}_t)$
- ▶ Note $d\mathbf{v}_t + \gamma(\mathbf{x}_t, \mathbf{v}_t)dt$ is the covariant time derivative of velocity and the contravariant form of the potential gradient is simply $\mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)$

Second-Order Diffusions

- ▶ Define a second-order SDE on a Riemann manifold with the diffusion defined on vector field \mathbf{v} to obtain

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Vector function γ has components $\gamma_k = \sum_{i,j} \Gamma_{ij}^k v^i v^j$
- ▶ Each Γ_{ij}^k being the Christoffel symbols of the Levi-Civita connection corresponding to the metric tensor $\mathbf{G}(\mathbf{x}_t)$
- ▶ Note $d\mathbf{v}_t + \gamma(\mathbf{x}_t, \mathbf{v}_t)dt$ is the covariant time derivative of velocity and the contravariant form of the potential gradient is simply $\mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)$
- ▶ Physically defines acceleration of a particle across manifold under the influence of potential field

Second-Order Diffusions

- ▶ Define a second-order SDE on a Riemann manifold with the diffusion defined on vector field \mathbf{v} to obtain

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Vector function γ has components $\gamma_k = \sum_{i,j} \Gamma_{ij}^k v^i v^j$
- ▶ Each Γ_{ij}^k being the Christoffel symbols of the Levi-Civita connection corresponding to the metric tensor $\mathbf{G}(\mathbf{x}_t)$
- ▶ Note $d\mathbf{v}_t + \gamma(\mathbf{x}_t, \mathbf{v}_t)dt$ is the covariant time derivative of velocity and the contravariant form of the potential gradient is simply $\mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)$
- ▶ Physically defines acceleration of a particle across manifold under the influence of potential field
- ▶ A unit friction or viscous force is included by the term \mathbf{v} and the fluctuating forcing is modelled by the Brownian motion

Second-Order Diffusions

- ▶ Define a second-order SDE on a Riemann manifold with the diffusion defined on vector field \mathbf{v} to obtain

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Vector function γ has components $\gamma_k = \sum_{i,j} \Gamma_{ij}^k v^i v^j$
- ▶ Each Γ_{ij}^k being the Christoffel symbols of the Levi-Civita connection corresponding to the metric tensor $\mathbf{G}(\mathbf{x}_t)$
- ▶ Note $d\mathbf{v}_t + \gamma(\mathbf{x}_t, \mathbf{v}_t)dt$ is the covariant time derivative of velocity and the contravariant form of the potential gradient is simply $\mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)$
- ▶ Physically defines acceleration of a particle across manifold under the influence of potential field
- ▶ A unit friction or viscous force is included by the term \mathbf{v} and the fluctuating forcing is modelled by the Brownian motion
- ▶ What has this to do with MCMC?

Second-Order Diffusions

Second-Order Diffusions

- ▶ What is invariant measure of this diffusion?

Second-Order Diffusions

- ▶ What is invariant measure of this diffusion?
- ▶ The Fokker-Planck equation for the diffusion follows as

Second-Order Diffusions

- ▶ What is invariant measure of this diffusion?
- ▶ The Fokker-Planck equation for the diffusion follows as

$$\frac{dp}{dt} = - v^a \frac{\partial p}{\partial x^a} + \frac{\partial}{\partial v^a} \left[\left(\Gamma_{bc}^a v^b v^c + g^{ab} \frac{\partial \phi}{\partial x^b} + v^a \right) p \right] + g^{ab} \frac{\partial^2 p}{\partial v^a \partial v^b}$$

Second-Order Diffusions

- ▶ What is invariant measure of this diffusion?
- ▶ The Fokker-Planck equation for the diffusion follows as

$$\frac{dp}{dt} = - v^a \frac{\partial p}{\partial x^a} + \frac{\partial}{\partial v^a} \left[\left(\Gamma_{bc}^a v^b v^c + g^{ab} \frac{\partial \phi}{\partial x^b} + v^a \right) p \right] + g^{ab} \frac{\partial^2 p}{\partial v^a \partial v^b}$$

- ▶ Some schoolboy algebra shows that for $dp/dt = 0$

Second-Order Diffusions

- ▶ What is invariant measure of this diffusion?
- ▶ The Fokker-Planck equation for the diffusion follows as

$$\frac{dp}{dt} = - v^a \frac{\partial p}{\partial x^a} + \frac{\partial}{\partial v^a} \left[\left(\Gamma_{bc}^a v^b v^c + g^{ab} \frac{\partial \phi}{\partial x^b} + v^a \right) p \right] + g^{ab} \frac{\partial^2 p}{\partial v^a \partial v^b}$$

- ▶ Some schoolboy algebra shows that for $dp/dt = 0$

$$\frac{dp}{p} = \left[-\frac{1}{2} \frac{\partial g_{ab}}{\partial x^c} v^a v^b + \frac{\partial}{\partial x^c} \log \sqrt{|g|} - \frac{\partial \phi}{\partial x^x} \right] dx^c$$

Second-Order Diffusions

- ▶ What is invariant measure of this diffusion?
- ▶ The Fokker-Planck equation for the diffusion follows as

$$\frac{dp}{dt} = -v^a \frac{\partial p}{\partial x^a} + \frac{\partial}{\partial v^a} \left[\left(\Gamma_{bc}^a v^b v^c + g^{ab} \frac{\partial \phi}{\partial x^b} + v^a \right) p \right] + g^{ab} \frac{\partial^2 p}{\partial v^a \partial v^b}$$

- ▶ Some schoolboy algebra shows that for $dp/dt = 0$

$$\frac{dp}{p} = \left[-\frac{1}{2} \frac{\partial g_{ab}}{\partial x^c} v^a v^b + \frac{\partial}{\partial x^c} \log \sqrt{|g|} - \frac{\partial \phi}{\partial x^a} \right] dx^c$$

- ▶ Therefore the above second-order SDE is satisfied by invariant densities $t \rightarrow \infty$ of the form

Second-Order Diffusions

- ▶ What is invariant measure of this diffusion?
- ▶ The Fokker-Planck equation for the diffusion follows as

$$\frac{dp}{dt} = -v^a \frac{\partial p}{\partial x^a} + \frac{\partial}{\partial v^a} \left[\left(\Gamma_{bc}^a v^b v^c + g^{ab} \frac{\partial \phi}{\partial x^b} + v^a \right) p \right] + g^{ab} \frac{\partial^2 p}{\partial v^a \partial v^b}$$

- ▶ Some schoolboy algebra shows that for $dp/dt = 0$

$$\frac{dp}{p} = \left[-\frac{1}{2} \frac{\partial g_{ab}}{\partial x^c} v^a v^b + \frac{\partial}{\partial x^c} \log \sqrt{|g|} - \frac{\partial \phi}{\partial x^a} \right] dx^c$$

- ▶ Therefore the above second-order SDE is satisfied by invariant densities $t \rightarrow \infty$ of the form

$$p(\mathbf{x}, \mathbf{v}) \propto \det(\mathbf{G}(\mathbf{x})) \exp \left(-\frac{1}{2} \mathbf{v}^\top \mathbf{G}(\mathbf{x}) \mathbf{v} - \phi(\mathbf{x}) \right)$$

Second-Order Diffusions

- ▶ What is invariant measure of this diffusion?
- ▶ The Fokker-Planck equation for the diffusion follows as

$$\frac{dp}{dt} = -v^a \frac{\partial p}{\partial x^a} + \frac{\partial}{\partial v^a} \left[\left(\Gamma_{bc}^a v^b v^c + g^{ab} \frac{\partial \phi}{\partial x^b} + v^a \right) p \right] + g^{ab} \frac{\partial^2 p}{\partial v^a \partial v^b}$$

- ▶ Some schoolboy algebra shows that for $dp/dt = 0$

$$\frac{dp}{p} = \left[-\frac{1}{2} \frac{\partial g_{ab}}{\partial x^c} v^a v^b + \frac{\partial}{\partial x^c} \log \sqrt{|g|} - \frac{\partial \phi}{\partial x^a} \right] dx^c$$

- ▶ Therefore the above second-order SDE is satisfied by invariant densities $t \rightarrow \infty$ of the form

$$p(\mathbf{x}, \mathbf{v}) \propto \det(\mathbf{G}(\mathbf{x})) \exp \left(-\frac{1}{2} \mathbf{v}^\top \mathbf{G}(\mathbf{x}) \mathbf{v} - \phi(\mathbf{x}) \right)$$

- ▶ So for $\phi(\mathbf{x}) = -\log \pi(\mathbf{x}) + \frac{1}{2} \log \det(\mathbf{G}(\mathbf{x}))$ then it follows that marginally

$$p(\mathbf{x}) = \exp(-\phi(\mathbf{x})) = \pi(\mathbf{x})$$

- ▶ Numerical integration forms basis of MCMC scheme..... however

Second-Order Diffusions

Second-Order Diffusions

- ▶ Transform the system of SDE's from the configuration to phase space such that transformed variable $\mathbf{p}_t = \mathbf{G}(\mathbf{x}_t)\mathbf{v}_t$

Second-Order Diffusions

- ▶ Transform the system of SDE's from the configuration to phase space such that transformed variable $\mathbf{p}_t = \mathbf{G}(\mathbf{x}_t)\mathbf{v}_t$

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

Second-Order Diffusions

- ▶ Transform the system of SDE's from the configuration to phase space such that transformed variable $\mathbf{p}_t = \mathbf{G}(\mathbf{x}_t)\mathbf{v}_t$

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ is transformed to

Second-Order Diffusions

- ▶ Transform the system of SDE's from the configuration to phase space such that transformed variable $\mathbf{p}_t = \mathbf{G}(\mathbf{x}_t)\mathbf{v}_t$

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ is transformed to

$$d\mathbf{x}_t = \mathbf{G}^{-1}(\mathbf{x}_t)\mathbf{p}_t dt$$

$$d\mathbf{p}_t = -\nu(\mathbf{x}_t, \mathbf{p}_t)dt - \nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{p}_t dt + \sqrt{2\mathbf{G}(\mathbf{x}_t)}d\mathbf{W}_t$$

Second-Order Diffusions

- ▶ Transform the system of SDE's from the configuration to phase space such that transformed variable $\mathbf{p}_t = \mathbf{G}(\mathbf{x}_t)\mathbf{v}_t$

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ is transformed to

$$d\mathbf{x}_t = \mathbf{G}^{-1}(\mathbf{x}_t)\mathbf{p}_t dt$$

$$d\mathbf{p}_t = -\nu(\mathbf{x}_t, \mathbf{p}_t)dt - \nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{p}_t dt + \sqrt{2\mathbf{G}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ where vector function ν has elements $-\partial_i g^{ab} p_a p_b$

Second-Order Diffusions

- ▶ Transform the system of SDE's from the configuration to phase space such that transformed variable $\mathbf{p}_t = \mathbf{G}(\mathbf{x}_t)\mathbf{v}_t$

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ is transformed to

$$d\mathbf{x}_t = \mathbf{G}^{-1}(\mathbf{x}_t)\mathbf{p}_t dt$$

$$d\mathbf{p}_t = -\nu(\mathbf{x}_t, \mathbf{p}_t)dt - \nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{p}_t dt + \sqrt{2\mathbf{G}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ where vector function ν has elements $-\partial_i g^{ab} p_a p_b$
- ▶ Invariant measure with density

$$p(\mathbf{x}, \mathbf{p}) \propto \frac{1}{\det(\mathbf{G}(\mathbf{x}))} \exp\left(-\frac{1}{2}\mathbf{p}^\top \mathbf{G}^{-1}(\mathbf{x})\mathbf{p} - \phi(\mathbf{x})\right)$$

Second-Order Diffusions

- ▶ Transform the system of SDE's from the configuration to phase space such that transformed variable $\mathbf{p}_t = \mathbf{G}(\mathbf{x}_t)\mathbf{v}_t$

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ is transformed to

$$d\mathbf{x}_t = \mathbf{G}^{-1}(\mathbf{x}_t)\mathbf{p}_t dt$$

$$d\mathbf{p}_t = -\nu(\mathbf{x}_t, \mathbf{p}_t)dt - \nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{p}_t dt + \sqrt{2\mathbf{G}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ where vector function ν has elements $-\partial_i g^{ab} p_a p_b$
- ▶ Invariant measure with density

$$p(\mathbf{x}, \mathbf{p}) \propto \frac{1}{\det(\mathbf{G}(\mathbf{x}))} \exp\left(-\frac{1}{2}\mathbf{p}^\top \mathbf{G}^{-1}(\mathbf{x})\mathbf{p} - \phi(\mathbf{x})\right)$$

- ▶ Stochastic Hamiltonian on Manifold - using symplectic integrator samples drawn from invariant density via RMHMC

Geodesic flow as proposal mechanism

Geodesic flow as proposal mechanism

- ▶ Intuitive that proposal mechanism follow direct paths - geodesics

Geodesic flow as proposal mechanism

- ▶ Intuitive that proposal mechanism follow direct paths - geodesics
- ▶ Consider Thermodynamic Integral - basis for Path & Bridge sampling

Geodesic flow as proposal mechanism

- ▶ Intuitive that proposal mechanism follow direct paths - geodesics
- ▶ Consider Thermodynamic Integral - basis for Path & Bridge sampling
- ▶ For $\theta \in \mathbb{R}^D$ with $p(y|\theta) = q(y|\theta)z(\theta)^{-1}$ denote $U(y, \theta) = \partial_\theta \log q(y|\theta)$ and let $\theta(t)$ be a function of index t

Geodesic flow as proposal mechanism

- ▶ Intuitive that proposal mechanism follow direct paths - geodesics
- ▶ Consider Thermodynamic Integral - basis for Path & Bridge sampling
- ▶ For $\theta \in \mathbb{R}^D$ with $p(y|\theta) = q(y|\theta)z(\theta)^{-1}$ denote $U(y, \theta) = \partial_\theta \log q(y|\theta)$ and let $\theta(t)$ be a function of index t

$$\lambda = \log \left(\frac{z(\theta(t_2))}{z(\theta(t_1))} \right) = \int_{t_1}^{t_2} E_{y|\theta(t)} \left\{ \sum_d \dot{\theta}_d(t) U_d(y, \theta) \right\} dt$$

Geodesic flow as proposal mechanism

- ▶ Intuitive that proposal mechanism follow direct paths - geodesics
- ▶ Consider Thermodynamic Integral - basis for Path & Bridge sampling
- ▶ For $\theta \in \mathbb{R}^D$ with $p(y|\theta) = q(y|\theta)z(\theta)^{-1}$ denote $U(y, \theta) = \partial_\theta \log q(y|\theta)$ and let $\theta(t)$ be a function of index t

$$\lambda = \log \left(\frac{z(\theta(t_2))}{z(\theta(t_1))} \right) = \int_{t_1}^{t_2} E_{y|\theta(t)} \left\{ \sum_d \dot{\theta}_d(t) U_d(y, \theta) \right\} dt$$

- ▶ Quality of N -sample path estimate $\hat{\lambda}$ is $\text{var}(\hat{\lambda})$ i.e. independent sampling

Geodesic flow as proposal mechanism

- ▶ Intuitive that proposal mechanism follow direct paths - geodesics
- ▶ Consider Thermodynamic Integral - basis for Path & Bridge sampling
- ▶ For $\theta \in \mathbb{R}^D$ with $p(y|\theta) = q(y|\theta)z(\theta)^{-1}$ denote $U(y, \theta) = \partial_\theta \log q(y|\theta)$ and let $\theta(t)$ be a function of index t

$$\lambda = \log \left(\frac{z(\theta(t_2))}{z(\theta(t_1))} \right) = \int_{t_1}^{t_2} E_{y|\theta(t)} \left\{ \sum_d \dot{\theta}_d(t) U_d(y, \theta) \right\} dt$$

- ▶ Quality of N -sample path estimate $\hat{\lambda}$ is $\text{var}(\hat{\lambda})$ i.e. independent sampling

$$\frac{1}{N} \left[\int_{t_1}^{t_2} \sum_{i,j} \left(g_{ij} \dot{\theta}_i(t) \dot{\theta}_j(t) \right) dt - \lambda^2 \right] = \frac{1}{N} \left[\int_{t_1}^{t_2} \sum_{i,j} \left(g^{ij} p_i(t) p_j(t) \right) dt - \lambda^2 \right]$$

where $g_{ij} = E_{y|\theta(t)} \{ U_i(y, \theta) U_j(y, \theta) \}$ and $p_i(t) = \sum_j g_{ij} \dot{\theta}_j(t)$

Geodesic flow as proposal mechanism

- ▶ Variance of estimate can be minimised by employing path that minimises

$$\int_{t1}^{t2} \dot{\theta}(t)^T \mathbf{G}(\theta(t)) \dot{\theta}(t) dt = \int_{t1}^{t2} \mathbf{p}(t)^T \mathbf{G}^{-1}(\theta(t)) \mathbf{p}(t) dt = \int_{t1}^{t2} H(\theta(t), \mathbf{p}(t)) dt$$

Geodesic flow as proposal mechanism

- ▶ Variance of estimate can be minimised by employing path that minimises

$$\int_{t1}^{t2} \dot{\theta}(t)^\top \mathbf{G}(\theta(t)) \dot{\theta}(t) dt = \int_{t1}^{t2} \mathbf{p}(t)^\top \mathbf{G}^{-1}(\theta(t)) \mathbf{p}(t) dt = \int_{t1}^{t2} H(\theta(t), \mathbf{p}(t)) dt$$

where $H(\theta(t), \mathbf{p}(t))$ is Hamiltonian defined on the manifold with position specific metric tensor $g_{ij}(\theta)$

Geodesic flow as proposal mechanism

- ▶ Variance of estimate can be minimised by employing path that minimises

$$\int_{t1}^{t2} \dot{\theta}(t)^T \mathbf{G}(\theta(t)) \dot{\theta}(t) dt = \int_{t1}^{t2} \mathbf{p}(t)^T \mathbf{G}^{-1}(\theta(t)) \mathbf{p}(t) dt = \int_{t1}^{t2} H(\theta(t), \mathbf{p}(t)) dt$$

where $H(\theta(t), \mathbf{p}(t))$ is Hamiltonian defined on the manifold with position specific metric tensor $g_{ij}(\theta)$

- ▶ This is satisfied by Geodesic equations

$$\frac{d^2\theta^i}{dt^2} + \sum_{k,l} \Gamma_{kl}^i \frac{d\theta^k}{dt} \frac{d\theta^l}{dt} = 0$$

Geodesic flow as proposal mechanism

- ▶ Variance of estimate can be minimised by employing path that minimises

$$\int_{t1}^{t2} \dot{\theta}(t)^T \mathbf{G}(\theta(t)) \dot{\theta}(t) dt = \int_{t1}^{t2} \mathbf{p}(t)^T \mathbf{G}^{-1}(\theta(t)) \mathbf{p}(t) dt = \int_{t1}^{t2} H(\theta(t), \mathbf{p}(t)) dt$$

where $H(\theta(t), \mathbf{p}(t))$ is Hamiltonian defined on the manifold with position specific metric tensor $g_{ij}(\theta)$

- ▶ This is satisfied by Geodesic equations

$$\frac{d^2\theta^i}{dt^2} + \sum_{k,l} \Gamma_{kl}^i \frac{d\theta^k}{dt} \frac{d\theta^l}{dt} = 0$$

- ▶ Replace D 2-order ODEs with $2D$ 1-order ODEs in Hamilton-Jacobi form

$$\frac{d\theta}{dt} = \frac{\partial}{\partial \mathbf{p}} H(\theta, \mathbf{p}) \quad \frac{d\mathbf{p}}{dt} = -\frac{\partial}{\partial \theta} H(\theta, \mathbf{p})$$

Geodesic flow as proposal mechanism

- ▶ Variance of estimate can be minimised by employing path that minimises

$$\int_{t1}^{t2} \dot{\theta}(t)^T \mathbf{G}(\theta(t)) \dot{\theta}(t) dt = \int_{t1}^{t2} \mathbf{p}(t)^T \mathbf{G}^{-1}(\theta(t)) \mathbf{p}(t) dt = \int_{t1}^{t2} H(\theta(t), \mathbf{p}(t)) dt$$

where $H(\theta(t), \mathbf{p}(t))$ is Hamiltonian defined on the manifold with position specific metric tensor $g_{ij}(\theta)$

- ▶ This is satisfied by Geodesic equations

$$\frac{d^2\theta^i}{dt^2} + \sum_{k,l} \Gamma_{kl}^i \frac{d\theta^k}{dt} \frac{d\theta^l}{dt} = 0$$

- ▶ Replace D 2-order ODEs with $2D$ 1-order ODEs in Hamilton-Jacobi form

$$\frac{d\theta}{dt} = \frac{\partial}{\partial \mathbf{p}} H(\theta, \mathbf{p}) \quad \frac{d\mathbf{p}}{dt} = -\frac{\partial}{\partial \theta} H(\theta, \mathbf{p})$$

- ▶ Solving Hamilton equations defines lowest variance / shortest path between two densities on manifold

Second-Order Diffusions

Second-Order Diffusions

- ▶ Stochastic Hamiltonian on Manifold driven by conditional Ornstein-Uhlenbeck process

Second-Order Diffusions

- ▶ Stochastic Hamiltonian on Manifold driven by conditional Ornstein-Uhlenbeck process

$$d\mathbf{x}_t = \mathbf{G}^{-1}(\mathbf{x}_t)\mathbf{p}_t dt$$

$$d\mathbf{p}_t = -\nu(\mathbf{x}_t, \mathbf{p}_t)dt - \nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{p}_t dt + \sqrt{2\mathbf{G}(\mathbf{x}_t)}d\mathbf{W}_t$$

Second-Order Diffusions

- ▶ Stochastic Hamiltonian on Manifold driven by conditional Ornstein-Uhlenbeck process

$$d\mathbf{x}_t = \mathbf{G}^{-1}(\mathbf{x}_t)\mathbf{p}_t dt$$

$$d\mathbf{p}_t = -\nu(\mathbf{x}_t, \mathbf{p}_t)dt - \nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{p}_t dt + \sqrt{2\mathbf{G}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Employ symplectic integrator e.g. RMHMC for drift term

Second-Order Diffusions

- ▶ Stochastic Hamiltonian on Manifold driven by conditional Ornstein-Uhlenbeck process

$$d\mathbf{x}_t = \mathbf{G}^{-1}(\mathbf{x}_t)\mathbf{p}_t dt$$

$$d\mathbf{p}_t = -\nu(\mathbf{x}_t, \mathbf{p}_t)dt - \nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{p}_t dt + \sqrt{2\mathbf{G}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Employ symplectic integrator e.g. RMHMC for drift term
- ▶ Note that OU diffusion process corresponds to ad-hoc partial momentum updating

Second-Order Diffusions

- ▶ Stochastic Hamiltonian on Manifold driven by conditional Ornstein-Uhlenbeck process

$$d\mathbf{x}_t = \mathbf{G}^{-1}(\mathbf{x}_t)\mathbf{p}_t dt$$

$$d\mathbf{p}_t = -\nu(\mathbf{x}_t, \mathbf{p}_t)dt - \nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{p}_t dt + \sqrt{2\mathbf{G}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Employ symplectic integrator e.g. RMHMC for drift term
- ▶ Note that OU diffusion process corresponds to ad-hoc partial momentum updating
- ▶ Solid theoretical basis amenable to analysis of algorithm performance

Second-Order Diffusions

- ▶ Stochastic Hamiltonian on Manifold driven by conditional Ornstein-Uhlenbeck process

$$d\mathbf{x}_t = \mathbf{G}^{-1}(\mathbf{x}_t)\mathbf{p}_t dt$$

$$d\mathbf{p}_t = -\nu(\mathbf{x}_t, \mathbf{p}_t)dt - \nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{p}_t dt + \sqrt{2\mathbf{G}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Employ symplectic integrator e.g. RMHMC for drift term
- ▶ Note that OU diffusion process corresponds to ad-hoc partial momentum updating
- ▶ Solid theoretical basis amenable to analysis of algorithm performance
- ▶ Phase space defined on co-tangent bundle with symplectic forms in place

Second-Order Diffusions

- ▶ Stochastic Hamiltonian on Manifold driven by conditional Ornstein-Uhlenbeck process

$$d\mathbf{x}_t = \mathbf{G}^{-1}(\mathbf{x}_t)\mathbf{p}_t dt$$

$$d\mathbf{p}_t = -\nu(\mathbf{x}_t, \mathbf{p}_t)dt - \nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{p}_t dt + \sqrt{2\mathbf{G}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Employ symplectic integrator e.g. RMHMC for drift term
- ▶ Note that OU diffusion process corresponds to ad-hoc partial momentum updating
- ▶ Solid theoretical basis amenable to analysis of algorithm performance
- ▶ Phase space defined on co-tangent bundle with symplectic forms in place
- ▶ What of the configuration space?

Second-Order Diffusions

Second-Order Diffusions

- ▶ This SDE defined on co-tangent bundle - non-symplectic forms

Second-Order Diffusions

- ▶ This SDE defined on co-tangent bundle - non-symplectic forms

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

Second-Order Diffusions

- ▶ This SDE defined on co-tangent bundle - non-symplectic forms

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Drift function can be seen to describe Lagrangian dynamics

Second-Order Diffusions

- ▶ This SDE defined on co-tangent bundle - non-symplectic forms

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Drift function can be seen to describe Lagrangian dynamics
- ▶ OU diffusion driving overall process

Second-Order Diffusions

- ▶ This SDE defined on co-tangent bundle - non-symplectic forms

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Drift function can be seen to describe Lagrangian dynamics
- ▶ OU diffusion driving overall process
- ▶ Construct explicit integrator for dynamics - with volume correction

Second-Order Diffusions

- ▶ This SDE defined on co-tangent bundle - non-symplectic forms

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Drift function can be seen to describe Lagrangian dynamics
- ▶ OU diffusion driving overall process
- ▶ Construct explicit integrator for dynamics - with volume correction
- ▶ Preserves original Hamiltonian dynamics - high proposal acceptance rates

Second-Order Diffusions

- ▶ This SDE defined on co-tangent bundle - non-symplectic forms

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Drift function can be seen to describe Lagrangian dynamics
- ▶ OU diffusion driving overall process
- ▶ Construct explicit integrator for dynamics - with volume correction
- ▶ Preserves original Hamiltonian dynamics - high proposal acceptance rates
- ▶ Speed improvement over implicit symplectic integrator between 2x to 10x

Second-Order Diffusions

- ▶ This SDE defined on co-tangent bundle - non-symplectic forms

$$d\mathbf{x}_t = \mathbf{v}_t dt$$

$$d\mathbf{v}_t = -\gamma(\mathbf{x}_t, \mathbf{v}_t)dt - \mathbf{G}^{-1}(\mathbf{x}_t)\nabla_{\mathbf{x}_t}\phi(\mathbf{x}_t)dt - \mathbf{v}_t dt + \sqrt{2\mathbf{G}^{-1}(\mathbf{x}_t)}d\mathbf{W}_t$$

- ▶ Drift function can be seen to describe Lagrangian dynamics
- ▶ OU diffusion driving overall process
- ▶ Construct explicit integrator for dynamics - with volume correction
- ▶ Preserves original Hamiltonian dynamics - high proposal acceptance rates
- ▶ Speed improvement over implicit symplectic integrator between 2x to 10x
- ▶ Lagrangian Dynamical Monte Carlo
<http://arxiv.org/abs/1211.3759>

Gaussian Mixture Model

- ▶ Univariate finite mixture model

$$p(x|\mu, \sigma^2) = 0.7 \times \mathcal{N}(x|0, \sigma^2) + 0.3 \times \mathcal{N}(x|\mu, \sigma^2)$$

Gaussian Mixture Model

- Univariate finite mixture model

$$p(x|\mu, \sigma^2) = 0.7 \times \mathcal{N}(x|0, \sigma^2) + 0.3 \times \mathcal{N}(x|\mu, \sigma^2)$$
















Figure: Arrows correspond to the gradients and ellipses to the inverse metric tensor. Dashed lines are isocontours of the joint log density

Gaussian Mixture Model



Gaussian Mixture Model



Log-Gaussian Cox Point Process with Latent Field

- ▶ The joint density for Poisson counts and latent Gaussian field

$$p(\mathbf{y}, \mathbf{x} | \mu, \sigma, \beta) \propto \prod_{i,j}^{64} \exp\{y_{i,j}x_{i,j} - m \exp(x_{i,j})\} \exp(-(\mathbf{x} - \mu \mathbf{1})^T \Sigma_\theta^{-1} (\mathbf{x} - \mu \mathbf{1}) / 2)$$

Log-Gaussian Cox Point Process with Latent Field

- ▶ The joint density for Poisson counts and latent Gaussian field

$$p(\mathbf{y}, \mathbf{x} | \mu, \sigma, \beta) \propto \prod_{i,j}^{64} \exp\{y_{i,j}x_{i,j} - m \exp(x_{i,j})\} \exp(-(\mathbf{x} - \mu \mathbf{1})^T \Sigma_\theta^{-1} (\mathbf{x} - \mu \mathbf{1})/2)$$

- ▶ Metric tensors

$$\begin{aligned}\mathbf{G}(\boldsymbol{\theta})_{i,j} &= \frac{1}{2} \text{trace} \left(\boldsymbol{\Sigma}_\theta^{-1} \frac{\partial \boldsymbol{\Sigma}_\theta}{\partial \theta_i} \boldsymbol{\Sigma}_\theta^{-1} \frac{\partial \boldsymbol{\Sigma}_\theta}{\partial \theta_j} \right) \\ \mathbf{G}(\mathbf{x}) &= \Lambda + \Sigma_\theta^{-1}\end{aligned}$$

where Λ is diagonal with elements $m \exp(\mu + (\Sigma_\theta)_{i,i})$

Log-Gaussian Cox Point Process with Latent Field

- ▶ The joint density for Poisson counts and latent Gaussian field

$$p(\mathbf{y}, \mathbf{x} | \mu, \sigma, \beta) \propto \prod_{i,j}^{64} \exp\{y_{i,j}x_{i,j} - m \exp(x_{i,j})\} \exp(-(\mathbf{x} - \mu \mathbf{1})^T \Sigma_\theta^{-1} (\mathbf{x} - \mu \mathbf{1})/2)$$

- ▶ Metric tensors

$$\begin{aligned}\mathbf{G}(\boldsymbol{\theta})_{i,j} &= \frac{1}{2} \text{trace} \left(\boldsymbol{\Sigma}_\theta^{-1} \frac{\partial \boldsymbol{\Sigma}_\theta}{\partial \theta_i} \boldsymbol{\Sigma}_\theta^{-1} \frac{\partial \boldsymbol{\Sigma}_\theta}{\partial \theta_j} \right) \\ \mathbf{G}(\mathbf{x}) &= \Lambda + \Sigma_\theta^{-1}\end{aligned}$$

where Λ is diagonal with elements $m \exp(\mu + (\Sigma_\theta)_{i,i})$

- ▶ Latent field metric tensor defining flat manifold is 4096×4096 , $\mathcal{O}(N^3)$ obtained from parameter conditional

Log-Gaussian Cox Point Process with Latent Field

- ▶ The joint density for Poisson counts and latent Gaussian field

$$p(\mathbf{y}, \mathbf{x} | \mu, \sigma, \beta) \propto \prod_{i,j}^{64} \exp\{y_{i,j}x_{i,j} - m \exp(x_{i,j})\} \exp(-(\mathbf{x} - \mu \mathbf{1})^T \Sigma_\theta^{-1} (\mathbf{x} - \mu \mathbf{1})/2)$$

- ▶ Metric tensors

$$\begin{aligned}\mathbf{G}(\boldsymbol{\theta})_{i,j} &= \frac{1}{2} \text{trace} \left(\boldsymbol{\Sigma}_\theta^{-1} \frac{\partial \boldsymbol{\Sigma}_\theta}{\partial \theta_i} \boldsymbol{\Sigma}_\theta^{-1} \frac{\partial \boldsymbol{\Sigma}_\theta}{\partial \theta_j} \right) \\ \mathbf{G}(\mathbf{x}) &= \Lambda + \Sigma_\theta^{-1}\end{aligned}$$

where Λ is diagonal with elements $m \exp(\mu + (\Sigma_\theta)_{i,i})$

- ▶ Latent field metric tensor defining flat manifold is 4096×4096 , $\mathcal{O}(N^3)$ obtained from parameter conditional
- ▶ MALA requires transformation of latent field to sample - with separate tuning in transient and stationary phases of Markov chain

RMHMC for Log-Gaussian Cox Point Processes




Figure: Data, Latent Field, Poisson Mean Field

RMHMC for Log-Gaussian Cox Point Processes

Table: Sampling the latent variables of a Log-Gaussian Cox Process - Comparison of sampling methods

Method	Time	ESS (Min, Med, Max)	s/Min ESS	Rel. Speed
MALA (Transient)	31,577	(3, 8, 50)	10,605	×1
MALA (Stationary)	31,118	(4, 16, 80)	7836	×1.35
mMALA	634	(26, 84, 174)	24.1	×440
RMHMC	2936	(1951, 4545, 5000)	1.5	×7070

Conclusion and Discussion

- ▶ Geometry of statistical models harnessed in Monte Carlo methods

Conclusion and Discussion

- ▶ Geometry of statistical models harnessed in Monte Carlo methods
 - ▶ Diffusions that respect structure and curvature of space - Manifold MALA

Conclusion and Discussion

- ▶ Geometry of statistical models harnessed in Monte Carlo methods
 - ▶ Diffusions that respect structure and curvature of space - Manifold MALA
 - ▶ Geodesic flows on model manifold - RMHMC - generalisation of HMC

Conclusion and Discussion

- ▶ Geometry of statistical models harnessed in Monte Carlo methods
 - ▶ Diffusions that respect structure and curvature of space - Manifold MALA
 - ▶ Geodesic flows on model manifold - RMHMC - generalisation of HMC
 - ▶ Assessed on correlated & high-dimensional latent variable models

Conclusion and Discussion

- ▶ Geometry of statistical models harnessed in Monte Carlo methods
 - ▶ Diffusions that respect structure and curvature of space - Manifold MALA
 - ▶ Geodesic flows on model manifold - RMHMC - generalisation of HMC
 - ▶ Assessed on correlated & high-dimensional latent variable models
 - ▶ Promising capability of methodology

Conclusion and Discussion

- ▶ Geometry of statistical models harnessed in Monte Carlo methods
 - ▶ Diffusions that respect structure and curvature of space - Manifold MALA
 - ▶ Geodesic flows on model manifold - RMHMC - generalisation of HMC
 - ▶ Assessed on correlated & high-dimensional latent variable models
 - ▶ Promising capability of methodology
- ▶ Ongoing development

Conclusion and Discussion

- ▶ Geometry of statistical models harnessed in Monte Carlo methods
 - ▶ Diffusions that respect structure and curvature of space - Manifold MALA
 - ▶ Geodesic flows on model manifold - RMHMC - generalisation of HMC
 - ▶ Assessed on correlated & high-dimensional latent variable models
 - ▶ Promising capability of methodology
- ▶ Ongoing development
 - ▶ Theoretical analysis of convergence

Conclusion and Discussion

- ▶ Geometry of statistical models harnessed in Monte Carlo methods
 - ▶ Diffusions that respect structure and curvature of space - Manifold MALA
 - ▶ Geodesic flows on model manifold - RMHMC - generalisation of HMC
 - ▶ Assessed on correlated & high-dimensional latent variable models
 - ▶ Promising capability of methodology
- ▶ Ongoing development
 - ▶ Theoretical analysis of convergence
 - ▶ Geodesic Monte Carlo on Embedded Manifolds
<http://arxiv.org/pdf/1301.6064.pdf>

Conclusion and Discussion

- ▶ Geometry of statistical models harnessed in Monte Carlo methods
 - ▶ Diffusions that respect structure and curvature of space - Manifold MALA
 - ▶ Geodesic flows on model manifold - RMHMC - generalisation of HMC
 - ▶ Assessed on correlated & high-dimensional latent variable models
 - ▶ Promising capability of methodology
- ▶ Ongoing development
 - ▶ Theoretical analysis of convergence
 - ▶ Geodesic Monte Carlo on Embedded Manifolds
<http://arxiv.org/pdf/1301.6064.pdf>
 - ▶ Investigate alternative manifold structures

Conclusion and Discussion

- ▶ Geometry of statistical models harnessed in Monte Carlo methods
 - ▶ Diffusions that respect structure and curvature of space - Manifold MALA
 - ▶ Geodesic flows on model manifold - RMHMC - generalisation of HMC
 - ▶ Assessed on correlated & high-dimensional latent variable models
 - ▶ Promising capability of methodology
- ▶ Ongoing development
 - ▶ Theoretical analysis of convergence
 - ▶ Geodesic Monte Carlo on Embedded Manifolds
<http://arxiv.org/pdf/1301.6064.pdf>
 - ▶ Investigate alternative manifold structures
 - ▶ Design and effect of metric and connection

Conclusion and Discussion

- ▶ Geometry of statistical models harnessed in Monte Carlo methods
 - ▶ Diffusions that respect structure and curvature of space - Manifold MALA
 - ▶ Geodesic flows on model manifold - RMHMC - generalisation of HMC
 - ▶ Assessed on correlated & high-dimensional latent variable models
 - ▶ Promising capability of methodology
- ▶ Ongoing development
 - ▶ Theoretical analysis of convergence
 - ▶ Geodesic Monte Carlo on Embedded Manifolds
<http://arxiv.org/pdf/1301.6064.pdf>
 - ▶ Investigate alternative manifold structures
 - ▶ Design and effect of metric and connection
 - ▶ Optimality of Hamiltonian flows as local geodesics

Conclusion and Discussion

- ▶ Geometry of statistical models harnessed in Monte Carlo methods
 - ▶ Diffusions that respect structure and curvature of space - Manifold MALA
 - ▶ Geodesic flows on model manifold - RMHMC - generalisation of HMC
 - ▶ Assessed on correlated & high-dimensional latent variable models
 - ▶ Promising capability of methodology
- ▶ Ongoing development
 - ▶ Theoretical analysis of convergence
 - ▶ Geodesic Monte Carlo on Embedded Manifolds
<http://arxiv.org/pdf/1301.6064.pdf>
 - ▶ Investigate alternative manifold structures
 - ▶ Design and effect of metric and connection
 - ▶ Optimality of Hamiltonian flows as local geodesics
 - ▶ Alternative transition kernels

Conclusion and Discussion

- ▶ Geometry of statistical models harnessed in Monte Carlo methods
 - ▶ Diffusions that respect structure and curvature of space - Manifold MALA
 - ▶ Geodesic flows on model manifold - RMHMC - generalisation of HMC
 - ▶ Assessed on correlated & high-dimensional latent variable models
 - ▶ Promising capability of methodology
- ▶ Ongoing development
 - ▶ Theoretical analysis of convergence
 - ▶ Geodesic Monte Carlo on Embedded Manifolds
<http://arxiv.org/pdf/1301.6064.pdf>
 - ▶ Investigate alternative manifold structures
 - ▶ Design and effect of metric and connection
 - ▶ Optimality of Hamiltonian flows as local geodesics
 - ▶ Alternative transition kernels
- ▶ No silver bullet or cure all - new powerful methodology for MC toolkit

Funding Acknowledgment

- ▶ EPSRC Established Career Fellowship
- ▶ Royal Society Wolfson Research Merit Award