

# Lecture 1. Causal Inference in High-Dimensional Approximately Sparse Structural Linear Models

VC

May 12, 2016, MIT

# Introduction

- ▶ Richer data and methodological developments lead us to consider more elaborate econometric models than before.

# Introduction

- ▶ Richer data and methodological developments lead us to consider more elaborate econometric models than before.
- ▶ Focus discussion on the linear endogenous model

$$\underbrace{y_i}_{\text{outcome}} = \underbrace{d_i}_{\text{treatment}} \underbrace{\alpha}_{\text{effect}} + \underbrace{\sum_{j=1}^p x_{ij}\beta_j}_{\text{controls}} + \underbrace{\epsilon_i}_{\text{noise}}, \quad (1)$$

$$\mathbb{E}[\epsilon_i | \underbrace{x_i, z_i}_{\text{exogenous vars}}] = 0.$$

# Introduction

- ▶ Richer data and methodological developments lead us to consider more elaborate econometric models than before.
- ▶ Focus discussion on the linear endogenous model

$$\underbrace{y_i}_{\text{outcome}} = \underbrace{d_i}_{\text{treatment}} \underbrace{\alpha}_{\text{effect}} + \underbrace{\sum_{j=1}^p x_{ij}\beta_j}_{\text{controls}} + \underbrace{\epsilon_i}_{\text{noise}}, \quad (1)$$

$$\mathbb{E}[\epsilon_i | \underbrace{x_i, z_i}_{\text{exogenous vars}}] = 0.$$

- ▶ Controls can be richer as more features become available (Census characteristics, housing characteristics, geography, text data)

⇐ “big” data

# Introduction

- ▶ Richer data and methodological developments lead us to consider more elaborate econometric models than before.
- ▶ Focus discussion on the linear endogenous model

$$\underbrace{y_i}_{\text{outcome}} = \underbrace{d_i}_{\text{treatment}} \underbrace{\alpha}_{\text{effect}} + \underbrace{\sum_{j=1}^p x_{ij}\beta_j}_{\text{controls}} + \underbrace{\epsilon_i}_{\text{noise}}, \quad (1)$$

$$\mathbb{E}[\epsilon_i | \underbrace{x_i, z_i}_{\text{exogenous vars}}] = 0.$$

- ▶ Controls can be richer as more features become available (Census characteristics, housing characteristics, geography, text data)
  - ⇐ “big” data
- ▶ Controls can contain transformation of “raw” controls in an effort to make models more flexible
  - ⇐ nonparametric series modeling, “machine learning”

# Introduction

- ▶ This **forces** us to explicitly consider **model selection** to select controls that are “most relevant”.
- ▶ Model selection techniques:
  - ▶ CLASSICAL: **t and F tests**
  - ▶ MODERN: **Lasso**, Regression Trees, Random Forests, Boosting

# Introduction

- ▶ This **forces** us to explicitly consider **model selection** to select controls that are “most relevant”.
- ▶ Model selection techniques:
  - ▶ CLASSICAL: **t and F tests**
  - ▶ MODERN: **Lasso**, Regression Trees, Random Forests, Boosting

If you are using *any* of these MS techniques directly in (1),  
you are doing it *wrong*.

Have to do *additional selection* to make it right.

# An Example: Effect of Institutions on the Wealth of Nations

- ▶ Acemoglu, Johnson, Robinson (2001)
- ▶ Impact of institutions on wealth

$$\underbrace{y_i}_{\text{log gdp per capita today}} = \underbrace{d_i}_{\text{quality of institutions}} \overset{\text{effect}}{\alpha} + \underbrace{\sum_{j=1}^p x_{ij} \beta_j}_{\text{geography controls}} + \epsilon_i, \quad (2)$$

- ▶ Instrument  $z_i$ : the early settler mortality (200 years ago)
- ▶ Sample size  $n = 67$
- ▶ Specification of controls:
  - ▶ Basic: constant, latitude ( $p=2$ )
  - ▶ Flexible: + cubic spline in latitude, continent dummies ( $p=16$ )



# Example: The Effect of Institutions

	Institutions	
	Effect	Std. Err.
Basic Controls	<b>.96**</b>	0.21
Flexible Controls	<b>.98</b>	0.80

- Is it ok to drop the additional controls?

# Example: The Effect of Institutions

	Institutions	
	Effect	Std. Err.
Basic Controls	<b>.96**</b>	0.21
Flexible Controls	<b>.98</b>	0.80

- Is it ok to drop the additional controls?

Potentially Dangerous.

# Example: The Effect of Institutions

	Institutions	
	Effect	Std. Err.
Basic Controls	<b>.96**</b>	0.21
Flexible Controls	<b>.98</b>	0.80

- Is it ok to drop the additional controls?

Potentially Dangerous. Very.

# Analysis: things can go wrong even with $p = 1$

- ▶ Consider a very simple exogenous model

$$y_i = d_i\alpha + x_i\beta + \epsilon_i, \quad \mathbb{E}[\epsilon_i \mid d_i, x_i] = 0.$$

- ▶ Common practice is to do the following.
- ▶ **Post-single selection** procedure:

**Step 1.** Include  $x_i$  only if it is a significant predictor of  $y_i$  as judged by a conservative test (t-test, Lasso, etc.). Drop it otherwise.

**Step 2.** Refit the model after selection, use standard confidence intervals.

- ▶ This can **fail miserably**, if  $|\beta|$  is close to zero but not equal to zero, formally if

$$|\beta| \propto 1/\sqrt{n}$$

# What can go wrong? Distribution of $\sqrt{n}(\hat{\alpha} - \alpha)$ is not what you think

$$y_i = d_i\alpha + x_i\beta + \epsilon_i, \quad d_i = x_i\gamma + v_i$$


$$\alpha = 0, \quad \beta = .2, \quad \gamma = .8,$$

$$n = 100$$

$$\epsilon_i \sim N(0, 1)$$

$$(d_i, x_i) \sim N\left(0, \begin{bmatrix} 1 & .8 \\ .8 & 1 \end{bmatrix}\right)$$

- ▶ selection done by a **t-test**



Reject  $H_0 : \alpha = 0$  (the truth) about 50% of the time (with nominal size of 5%)

# What can go wrong? Distribution of $\sqrt{n}(\hat{\alpha} - \alpha)$ is not what you think

$$y_i = d_i\alpha + x_i\beta + \epsilon_i, \quad d_i = x_i\gamma + v_i$$


$$\alpha = 0, \quad \beta = .2, \quad \gamma = .8,$$

$$n = 100$$

$$\epsilon_i \sim N(0, 1)$$

$$(d_i, x_i) \sim N\left(0, \begin{bmatrix} 1 & .8 \\ .8 & 1 \end{bmatrix}\right)$$

- ▶ selection done by **Lasso**



Reject  $H_0 : \alpha = 0$  (the truth) of no effect about 50% of the time

# Solutions?

Pseudo-solutions:

- ▶ **Practical:** bootstrap (does not work),
- ▶ **Classical:** assume the problem away by assuming that either  $\beta = 0$  or  $|\beta| \gg 0$ ,
- ▶ **Conservative:** don't do selection

# Solution: Post-double selection

► **Post-double selection** procedure:

- Step 1. Include  $x_i$  if it is a significant predictor of  $y_i$  as judged by a conservative test (t-test, Lasso etc).
- Step 2. Include  $x_i$  if it is a significant predictor of  $d_i$  as judged by a conservative test (t-test, Lasso etc). [In the IV models must include  $x_i$  if it a significant predictor of  $z_i$ ].
- Step 3. Refit the model after selection, use standard confidence intervals.

## Theorem

*DS is theoretically valid in low-dimensional setting and in high-dimensional approximately sparse settings.*

- Refs: Belloni et al: WC ES 2010, ReStud 2013; Chernozhukov, Hansen, Spindler, ARE 2015.



# Double Selection Works

$$y_i = d_i\alpha + x_i\beta + \epsilon_i, \quad d_i = x_i\gamma + v_i$$


$$\alpha = \mathbf{0}, \quad \beta = .2, \quad \gamma = .8,$$

$$n = 100$$

$$\epsilon_i \sim N(0, 1)$$

$$(d_i, x_i) \sim N\left(0, \begin{bmatrix} 1 & .8 \\ .8 & 1 \end{bmatrix}\right)$$

- **double selection**  
done by **t-tests**



Reject  $H_0 : \alpha = 0$  (the truth) about 5% of the time (for nominal size = 5%)

# Double Selection Works

$$y_i = d_i\alpha + x_i\beta + \epsilon_i, \quad d_i = x_i\gamma + v_i$$


$$\alpha = \mathbf{0}, \quad \beta = .2, \quad \gamma = .8,$$

$$n = 100$$

$$\epsilon_i \sim N(0, 1)$$

$$(d_i, x_i) \sim N\left(0, \begin{bmatrix} 1 & .8 \\ .8 & 1 \end{bmatrix}\right)$$

- **double selection**  
done by **Lasso**



Reject  $H_0 : \alpha = 0$  (the truth) about 5% of the time (nominal size = 5%)

# Intuition

- ▶ The **Double Selection** — the selection among the controls  $x_i$  that predict *either*  $d_i$  or  $y_i$  — creates this robustness. It finds controls whose omission would lead to a “large” omitted variable bias, and includes them in the regression.
- ▶ In essence the procedure is a model selection version of Frisch-Waugh-Lovell partialling-out procedure for estimating linear regression.
- ▶ The double selection method is robust to moderate selection mistakes in the two selection steps.

# More Intuition via OMVB Analysis

Think about omitted variables bias:

$$y_i = \alpha d_i + \beta x_i + \zeta_i \quad ; \quad d_i = \gamma x_i + v_i$$

If we drop  $x_i$ , the short regression of  $y_i$  on  $d_i$  gives

$$\sqrt{n}(\hat{\alpha} - \alpha) = \text{good term} + \underbrace{\sqrt{n}(D'D/n)^{-1}(X'X/n)(\gamma\beta)}_{\text{OMVB}}.$$

- ▶ the good term is asymptotically normal, and we want

$$\sqrt{n}\gamma\beta \rightarrow 0.$$

- ▶ **single selection** can drop  $x_i$  only if  $\beta = O(\sqrt{1/n})$ , but

$$\sqrt{n}\gamma\sqrt{1/n} \not\rightarrow 0$$

- ▶ **double selection** can drop  $x_i$  only if *both*  $\beta = O(\sqrt{1/n})$  and  $\gamma = O(\sqrt{1/n})$ , that is, if

$$\sqrt{n}\gamma\beta = O(1/\sqrt{n}) \rightarrow 0.$$

# Example: The Effect of Institutions, Continued

Going back to Acemoglu, Johnson, Robinson (2001):

- **Double Selection:** include  $x_{ij}$ 's that are significant predictors of either  $y_i$  or  $d_i$  or  $z_i$ , as judged by Lasso. Drop otherwise.

	Intitutions	
	Effect	Std. Err.
Basic Controls	<b>.96**</b>	0.21
Flexible Controls	<b>.98</b>	0.80
<b>Double Selection</b>	<b>.78**</b>	0.19

# Application: Effect of Abortion on Murder Rates in the U.S.

Estimate the consequences of abortion rates on crime in the U.S.,  
Donohue and Levitt (2001)

$$y_{it} = \alpha d_{it} + x'_{it}\beta + \zeta_{it}$$

- ▶  $y_{it}$  = change in crime-rate in state  $i$  between  $t$  and  $t - 1$ ,
- ▶  $d_{it}$  = change in the (lagged) abortion rate,
- 1.  $x_{it}$  = basic controls (time-varying confounding state-level factors, trends;  $p=20$ )
- 2.  $x_{it}$  = flexible controls (basic + state initial conditions + two-way interactions of all these variables)
- ▶  $p = 251$ ,  $n = 576$

# Effect of Abortion on Murder, continued

Estimator	Abortion on Murder	
	Effect	Std. Err.
Basic Controls	<b>-0.204**</b>	0.068
Flexible Controls	-0.321	1.109
Single Selection	<b>- 0.202**</b>	0.051
Double Selection	-0.166	0.216

- Double selection by Lasso: 8 controls selected, including state initial conditions and trends interacted with initial conditions

- ▶ This is sort of a negative result, unlike in AJR (2011)
- ▶ Double selection does not always overturn results. Plenty of positive results confirming:
  - ▶ Barro and Lee's convergence results in cross-country growth rates;
  - ▶ Poterba et al results on positive impact of 401(k) on savings;
  - ▶ Acemoglu et al (2014) results on democracy causing growth;



# High-Dimensional Prediction Problems

- ▶ Generic prediction problem

$$u_i = \sum_{j=1}^p x_{ij}\pi_j + \zeta_i, \quad \mathbb{E}[\zeta_i \mid \mathbf{x}_i] = 0, \quad i = 1, \dots, n,$$

can have  $p = p_n$  small,  $p \propto n$ , or even  $p \gg n$ .

- ▶ In the double selection procedure,  $u_i$  could be outcome  $y_i$ , treatment  $d_i$ , or instrument  $z_i$ . Need to find good predictors among  $x_{ij}$ 's.
- ▶ APPROXIMATE SPARSITY: after sorting, absolute values of coefficients decay fast enough:

$$|\pi|_{(j)} \leq A j^{-a}, \quad a > 1, j = 1, \dots, p = p_n, \forall n$$

- ▶ RESTRICTED ISOMETRY: small groups of  $x'_{ij}$ s are not close to being collinear.

# Selection of Predictors by Lasso

Assuming  $x'_{ij}$ s normalized to have the second empirical moment to 1.

- Ideal (Akaike, Schwarz): minimize

$$\sum_{i=1}^n \left( u_i - \sum_{j=1}^p x_{ij} b_j \right)^2 + \lambda \left( \sum_{j=1}^p 1\{b_j \neq 0\} \right).$$

- Lasso (Bickel, Ritov, Tsybakov, Annals, 2009): minimize

$$\sum_{i=1}^n \left( u_i - \sum_{j=1}^p x_{ij} b_j \right)^2 + \lambda \left( \sum_{j=1}^p |b_j| \right), \quad \lambda = \sqrt{\mathbb{E}\zeta^2} 2\sqrt{2n\log(pn)}$$

# Selection of Predictors by Lasso

Assuming  $x'_{ij}$ s normalized to have the second empirical moment to 1.

- Ideal (Akaike, Schwarz): minimize

$$\sum_{i=1}^n \left( u_i - \sum_{j=1}^p x_{ij} b_j \right)^2 + \lambda \left( \sum_{j=1}^p 1\{b_j \neq 0\} \right).$$

- Lasso (Bickel, Ritov, Tsybakov, Annals, 2009): minimize

$$\sum_{i=1}^n \left( u_i - \sum_{j=1}^p x_{ij} b_j \right)^2 + \lambda \left( \sum_{j=1}^p |b_j| \right), \quad \lambda = \sqrt{\mathbb{E}\zeta^2} 2\sqrt{2n\log(pn)}$$

- Root Lasso (Belloni, Chernozhukov, Wang, Biometrika, 2011): minimize

$$\sqrt{\sum_{i=1}^n \left( u_i - \sum_{j=1}^p x_{ij} b_j \right)^2} + \lambda \left( \sum_{j=1}^p |b_j| \right), \quad \lambda = \sqrt{2n\log(pn)}$$

# Lasso provides high-quality model selection

## Theorem

*Under approximate sparsity and restricted isometry conditions, Lasso and Root-Lasso find parsimonious models of approximately optimal size*

$$s = n^{\frac{1}{2a}}.$$

*Using these models, the OLS can approximate the regression functions at the nearly optimal rates in the root mean square error:*

$$\sqrt{\frac{s}{n} \log(pn)}$$

*This is also the rate at which Lasso approximates the regression functions.*

- ▶ Ref (Lasso): Bickel, Ritov, Tsybakov (Annals 2010)
- ▶ Ref (Post-Lasso, Root-Lasso): Belloni and Cherozhukov: Bernoulli, 2013, Belloni et al , Annals, 2014)

# Double Selection in Approximately Sparse Regression

- ▶ Exogenous model

$$y_i = d_i\alpha + \sum_{j=1}^p x_{ij}\beta_j + \zeta_i, \quad \mathbb{E}[\zeta_i \mid d_i, x_i] = 0, \quad i = 1, \dots, n,$$

$$d_i = \sum_{j=1}^p x_{ij}\gamma_j + \nu_i, \quad \mathbb{E}[\nu_i \mid x_i] = 0, \quad i = 1, \dots, n,$$

can have  $p$  small,  $p \propto n$ , or even  $p \gg n$ .

- ▶ APPROXIMATE SPARSITY: after sorting absolute values of coefficients decay fast enough:

$$|\beta|_{(j)} \leq Aj^{-a}, \quad a > 1, \quad |\gamma|_{(j)} \leq Aj^{-a}, \quad a > 1.$$

- ▶ RESTRICTED ISOMETRY: small groups of  $x'_{ij}$ s are not close to being collinear.

# Double Selection Procedure

► **Post-double selection** procedure

Step 1. Include  $x_{ij}$ 's that are significant predictors of  $y_i$  as judged by LASSO or OTHER high-quality selection procedure.

Step 2. Include  $x_{ij}$ 's that are significant predictors of  $d_i$  as judged by LASSO or OTHER high-quality selection procedures.

Step 3. Refit the model by least squares after selection, use standard confidence intervals.

► Ref: Belloni et al, 2010, ES World Congress, ReStud 2013

# Double Selection Procedure 2

A closely related procedure is the following:

► **Double partialling out by Lasso/Post-Lasso procedure:**

Step 1. Partial out from  $y_i$  the effect of all  $x_{ij}$ 's that are significant predictors of  $y_i$  as judged by LASSO or OTHER high-quality selection procedure. Obtain the residual  $\tilde{y}_i$ .

Step 2. Partial out from  $d_i$  the effect of all  $x_{ij}$ 's that are significant predictors of  $d_i$  as judged by LASSO or OTHER high-quality selection procedure. Obtain the residual  $\tilde{d}_i$ .

Step 3. Regress  $\tilde{y}_i$  on  $\tilde{d}_i$  using least squares, use standard confidence intervals.

- Ref: Chernozhukov, Hansen, Spindler, 2015, Annual Review of Economics; Belloni et al, Annals of Stats, 2014.

# Uniform Validity of the Double Selection/Partialling Out for Regression

## Theorem

***Uniformly within a class of approximately sparse models with restricted isometry conditions***

$$\sigma_n^{-1} \sqrt{n}(\hat{\alpha} - \alpha_0) \rightarrow_d N(0, 1),$$

where  $\sigma_n^2$  is conventional variance formula for least squares. Under homoscedasticity, semi-parametrically efficient.

- ▶ Model selection mistakes are asymptotically negligible due to double selection.
- ▶ Ref: Belloni et al, WC 2010, ReStud 2013; Belloni et al, Annals of Stats, 2014



# Double Selection for IV Regression

► **Post-double selection** procedure (Belloni et al 2014, JEP):

- Step 1. Include  $x_{ij}$ 's that are significant predictors of  $y_i$  as judged by LASSO or OTHER high-quality selection procedure.
- Step 2. Include  $x_{ij}$ 's that are significant predictors of either  $d_i$  or  $z_i$  as judged by LASSO or OTHER high-quality selection procedures.
- Step 3. Refit the model by two-stage least squares (or other IV estimator) after selection, use standard confidence intervals.

# Double Partialling Out for IV Model

A closely related procedure is the following:

► **Partialling out with double selection** procedure:

- Step 1. Partial out from  $y_i$  the effect of all  $x_{ij}$ 's that are significant predictors of  $y_i$  using LASSO, Post-LASSO or OTHER high-quality regularization procedure. Obtain the residual  $\tilde{y}_i$ .
- Step 2. Partial out from  $d_i$  the effect of all  $x_{ij}$ 's that are significant predictors of  $d_i$  as judged by LASSO or OTHER high-quality selection procedure. Obtain the residual  $\tilde{d}_i$ . Partial out from  $z_i$  the effect of all  $x_{ij}$ 's that are significant predictors of  $z_i$  as judged by LASSO or OTHER high-quality selection procedure. Obtain the residual  $\tilde{z}_i$ .
- Step 3. Run IV regression of  $\tilde{y}_i$  on  $\tilde{d}_i$  using  $\tilde{z}_i$  the instrument, use standard confidence intervals.
- Ref. Chernozhukov, Hansen, Spindler, 2015, Annual Review of Economics.

# Monte Carlo Confirmation

- ▶ In this simulation we used:  $p = 200$ ,  $n = 100$ ,  $\alpha_0 = .5$

$$y_i = d_i\alpha + x_i'\beta + \zeta_i, \quad \zeta_i \sim N(0, 1)$$

$$d_i = x_i'\gamma + v_i, \quad v_i \sim N(0, 1)$$

- ▶ **approximately sparse model:**


$$|\beta_j| \propto 1/j^2, |\gamma_j| \propto 1/j^2$$

- ▶  $R^2 = .5$  in each equation
- ▶ regressors are correlated Gaussians:

$$x \sim N(0, \Sigma), \quad \Sigma_{kj} = (0.5)^{|j-k|}.$$


# Distribution of Post Double Selection Estimator

$$p = 200, n = 100$$



# Distribution of Post-Single Selection Estimator

$$p = 200 \text{ and } n = 100$$



# Generalization: Orthogonalized or “Doubly Robust” Moment Equations

- ▶ Goal:
  - inference on structural parameter  $\alpha$  (e.g., elasticity)
  - having done Lasso & **other ML** fitting of reduced forms  $\eta(\cdot)$
- ▶ Use orthogonalization methods to remove biases. This often amounts to solving auxiliary prediction problems.
- ▶ In a nutshell, we want to set up moment conditions


$$\mathbb{E}[g(\underbrace{W}_{\text{data}}, \underbrace{\alpha_0}_{\text{structural parameter}}, \underbrace{\eta_0}_{\text{reduced form}})] = 0$$

such that the orthogonality conditions hold:


$$\partial_{\eta} \mathbb{E}[g(W, \alpha_0, \eta)] \Big|_{\eta=\eta_0} = 0$$

- ▶ See: Chernozhukov, Hansen, Spindler, AER, 2015

# Inference on Structural/Treatment Parameters



Without Orthogonalization



With Orthogonalization

# Conclusion

- ▶ It is time to address model selection
- ▶ Mostly dangerous: naive (post-single) selection does not work
- ▶ Double selection works
- ▶ More generally, the key is to use orthogonalized moment conditions for inference